

ESTIMATION OF TRANSMITTING ABILITIES FOR LACTATION TRAITS USING THE ANIMAL MODEL FOR HOLSTEIN CATTLE RAISED UNDER COMMERCIAL FARM IN EGYPT

Afifi, E.A.¹; M.H. Khalil²; Samira A. Arafa³ and M.A. Salem¹

¹ Department of Animal Production, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt,

² Department of Animal Production and Breeding, College of Agriculture and Veterinary Medicine, King Saud University, Qassim, Burieda, P.O. Box, 1842, Kingdom of Saudi Arabia.

³ Animal Production Research Institute, Ministry of Agriculture and Land Reclamation, Dokki, Cairo, Egypt,

ABSTRACT

Records of 5662 normal lactations produced by 1029 cows were analyzed by a single-trait Animal Model to estimate transmitting abilities of cows and their dams and sires for some lactation traits of Holstein cattle raised in El-Salhia commercial farm in Egypt. Lactation traits studied were 90-day milk yield (M90), 305-day milk yield (M305), total milk yield (TMY) and length of lactation period (LP). Repeatability estimates were moderate or slightly high (0.388, 0.402 and 0.417) for M90, M305 and TMY, respectively and low (0.146) for LP. Transmitting abilities (TA) were estimated for cows (CTA) using their own records, along with those for their dams (DTA) and sires (STA) without using their own records. Ranges in estimates of CTA for M90, M305, TMY and LP were 321 kg, 1087 kg, 1388 kg and 12.1 days, respectively. The corresponding ranges were 182 kg, 600 kg, 801 kg and 6.7 days for DTA and 208 kg, 702 kg, 851 kg and 8.5 days for STA. The percentages of cows and their dams and sires that had positive transmitting abilities for M90, M305, TMY and LP were 47.9, 47.3, 47.5 and 54.3% for CTA, 46.3, 46.4, 45.7 and 53.3% for DTA and 50.4, 52.5, 50.4 and 51.8% for STA. It could be concluded that selecting cows according to CTA would be more efficient than selecting them according to STA or DTA.

Keywords: Holstein cattle, lactation traits, transmitting ability, Animal Model, commercial farm, Egypt.

INTRODUCTION

At the beginning of the eighties, a common trend has been raised, in Egypt, for establishing large-scale commercial herds of dairy cattle through introducing new standard breeds (Holstein, Brown Swiss, Friesian, Pinzgauer...etc.) in order to increase the national milk production. Since that time and till now, several investigators have screened some of the genetic aspects of productive efficiency of these breeds under the semi-arid conditions in such herds (e.g. Salem, 1991; Gad 1995; others).

Genetic improvement for milk production traits could be achieved through selection and this necessitates identifying the elite cows and superior sires through the evaluation of breeding animals to be selected.

The present study was set up in order: (1) to characterize the lactational performance of Holstein cattle raised in a commercial herd under

Afifi, E. A. et al.

the Egyptian climatic conditions, and (2) to estimate transmitting abilities of milk production traits for cows with records and their dams and sires without records.

MATERIALS AND METHODS

Animals and breeding plan

Animals used in the present study comprised locally born females and males of Holstein cattle (H) raised in a commercial herd since 1982. Holstein females were imported from USA as pregnant heifers. The herd belongs to the General Cooperative for Developing Animal Wealth and Products, Egypt and was located at El-Salhia, Ismailia Governorate (East to the south of Nile Delta).

Feeding and management

Cows were kept under similar system of feeding and management. All year round, all cows were fed on good quality concentrates and corn silage. During winter and spring months (from December to May), animals were supplied with Egyptian clover (*Trifolium alexandrinum*) while during summer and autumn months (from June to the end of November) beet, maize and green sorghum (*Sorghum Vulgar Var Saccharatum*) were available. Also, rice straw was available all the year round. Feed was supplied to cows according to their live weight, production and pregnancy status. Clean water and mineral mixture were available all the time. Cows were inseminated artificially using frozen semen imported from USA. Heifers were bred when reached 16-18 months of age (about 350-375 kg) and cow was served during the first heat period following the 45th day post-partum. Pregnancy was detected by rectal palpation 60 days after the last service. Calves were given colostrum for four days after birth and housed in calf-boxes where they were bucket fed on whole milk and/ or milk replacer until weaning at 90 kg weight for male calves and 100 kg weight for females. After weaning and up to six months of age, calves of the same age were housed in-groups in pens provided with yards for exercise. At six-months of age, male calves were separated from females and housed in open sheds up to their sexual maturity. Lactating cows were machine-milked twice daily in a milk parlor. Milk yield was recorded daily to the nearest 0.1 kg at each milking. Cows were dried off two months before the expected date of next calving.

Data

Data used in this study covered nine consecutive years (1983-1991). All abnormal records (aborted cows and records without pedigree and breeding dates) were excluded. A total number of 5662 complete lactations for 1029 cows sired by 139 bulls were used. Lactation records of cows were grouped into age subclasses of 3-months interval, while they were grouped into days-open subclasses of 20-day intervals. Productive traits studied were 90-day milk yield, percentage of milk yield in the first 30 days of lactation,

records of the first, second, third and across all lactations were 1029, 1029, 1001 and 5662, respectively.

Estimation of transmitting ability

Transmitting abilities of cows (CTA) and their dams (DTA) and sires (STA) were estimated for milk production traits of all lactations using single-trait Animal Model program written by Misztal (1990). Transmitting abilities for cows (CTA) were predicted using their own records, while dam (DTA) and sire (STA) transmitting abilities were estimated for these animals without their own records. Transmitting abilities were calculated for 1029 cows, fathered by 139 sires and mothered by 1029 dams. The utilized Animal Model (in matrix notation) was:

$$Y = XB + Z_a a + Z_p p + e,$$

where Y = vector of observations for any lactation trait, X = incidence matrix for fixed effects, B = vector of an overall mean and fixed effects (year-season, age of cow and days open), Z_a = incidence matrix for random effects (direct genetic effect), a = vector of direct genetic effects of the cows, Z_p = incidence matrix for permanent environmental effects, P = vector of permanent environmental effects and e = vector of random error.

Since $\text{var}(a) = A\sigma_a^2$, $\text{var}(p) = I_p\sigma_p^2$, and $\text{var}(e) = I_n\sigma_e^2$. The variance-covariance matrix of the random effects was:

$$\text{var} \begin{bmatrix} a \\ p \\ e \end{bmatrix} = \begin{bmatrix} A\sigma_a^2 & 0 & 0 \\ 0 & I_p\sigma_p^2 & 0 \\ 0 & 0 & I_n\sigma_e^2 \end{bmatrix}$$

where A is the numerator relationship matrix, I_p is an identity matrix with order equal to number of animals, and I_n is an identity matrix with order equal to number of records.

RESULTS AND DISCUSSION

Means and variation

Means, their standard deviations (SD) and coefficients of variation (CV%) for lactation traits (M90, M305, TMY and LP) across all lactations of Holstein-Friesian cattle are given in Table 1. Means of lactation traits in this study generally fall within the range of those obtained for Holstein cattle raised in Egypt by Gad (1995). These estimates are higher than those reported by Moharram (1988) and Mokhtar *et al.* (1993). On the other hand, the actual mean of M305 in this study (5275 kg) is lower than that reported on imported Holstein cattle (6506 kg) by Ahmed (1996).

Table 1: Means, their standard deviations (SD) and coefficients of variation (CV%) for lactation traits in Holstein cattle.

Number of iterations and equations for transmitting ability
 The number of equations for lactation traits were 7939, while the number of iterations for M90, M305, TMY and LP were 108, 153, 214, and 127, respectively. Ducroco *et al.* (1990) and Wiggins and VanRaden (1990) reported that number of iterations required to reach the adequate convergence criteria might not be met before 100 or more iteration.

** = $p < 0.01$, *** = $p < 0.001$
 ** Degrees of freedom were 1028 and 4555 for cows and remainder, respectively.

Traits as defined in Table 1.

Trait*	Cow		Remainder**	
	σ^2_c	V%	σ^2_e	V%
M90	34191***	38.8	53875	61.2
M305	407020**	40.2	606690	59.8
TMY	656339**	41.7	917412	58.3
LP	5936***	14.6	34718	85.4
				0.146+0.013

Table 2: Estimates of cow component of variance (σ^2_c) and the remainder variance component (σ^2_e), their percentages of variation (V%) and repeatability (t) for lactation traits in Holstein cattle.

Variance components and repeatabilities for lactation traits are presented in Table 2. Repeatability estimates for lactation traits were moderate or slightly high and ranged from 0.146 to 0.417. These estimates fall mostly within the range (0.22 to 0.55) obtained for Holstein-Friesian and/or Friesian cattle by Abubakar *et al.* (1986a), Boldman and Freeman (1990), Wiggins and VanRaden (1990), Salem (1991), Khalil *et al.* (1994) and Gad (1995). Moderately high estimates of repeatability found in the literature and in this work for milk yield traits indicate that records beyond the first lactation actually add little new information about the producing ability (Soliman and Khalil, 1993). Accordingly, culling policies of cows for milk yield traits based on single record would be efficient from a genetic standpoint and consequently, assessment of several records are not required before selecting cows for these traits.

Variance components and repeatabilities

+++ Coefficient of variation computed as the residual deviation divided by the mean of a given trait.
 ++ Number of records used was 5662 lactations.
 LP = lactation length.
 + M90 = 90-day milk yield; M305 = 305-day milk yield; TMY = total milk yield and

Trait*	Mean**	SD	CV%***
M90 (kg)	1574	310	14.7
M305(kg)	5275	1056	14.8
TMY (kg)	6496	1516	14.7
LP (day)	371.1	54.0	5.0

and 704.5, 736.3 and 587.5 kg for TMY, respectively. With Friesian cattle raised in Egypt, Sadek *et al.* (1993) indicated that the range in STA was 340 kg, while Abdel-Gill (1991) found that the ranges in STA for LP were 46.1, 49.3 and 56.3 days in the first, second and the third lactation, respectively. Gebriel (1996) found that the ranges in transmitting ability for dams of buffalo cows (animals without records) were 361.3 kg for M305 and 43.52 days for LP. The same author added that the range recorded for buffalo cows CTA was higher than that obtained for dams of buffalo cows.

Table 4: Minimum, maximum and ranges of transmitting abilities for dams (DTA) and sires (STA) without records predicted by Animal Model for lactation traits in Holstein cattle.

Trait*	DTA for dams without records			STA for sires without records		
	Minimum	Maximum	Range	Minimum	Maximum	Range
M90	-85	97	182	-96	112	208
M305	-277	323	600	-328	374	702
TMY	-364	437	801	-390	461	851
LP	-3.6	3.1	6.7	-4.7	3.8	8.5

* Traits as defined in Table 1.

Animals with positive transmitting ability

The percentages of cows and their dams and sires that have positive estimates of transmitting ability for M90, M305, TMY and LP, respectively were 47.9, 47.3, 47.5 and 54.3% for CTA, 46.3, 46.4, 45.7 and 53.3% for DTA and 50.4, 52.5, 50.4 and 51.8% for STA (Table 5). The percentages for milk yield traits were mostly less than 50% for CTA and DTA, but they were higher than 50% for STA, *i.e.* positive estimates of transmitting ability obtained for cows are nearly similar to those obtained for their dams, but they are slightly smaller than those estimated for their sires. Ahmed (1996) found that the percentage of the positive CTA was 53.0% for M305 of Holstein cattle raised in Egypt. Gebriel (1996) with the Egyptian buffaloes found that the percentages of positive estimates of buffalo CTA (animals without records) for M305 and LP were equal. The same author added that positive estimates for buffalo CTA (44%) were lower than those for buffalo DTA (51%). El-Chafie (1981) found that the percentage of sires with positive BLUP estimates were 22% for Friesian sires raised in Egypt. In north and south America, Abubakar *et al.* (1987) noted that 47% of the Holstein sires had positive predicted sire values (PSV) for M305 in both Mexico and Colombia, while 94% of the sires had positive PSV in USA. Abdel-Gill (1991) with Friesian cattle in Egypt, found that the positive percentages of STA for M305 in the first, second and the third lactation constituted 50.0, 45.6 and 51.0%, respectively, while they were 43.4, 43.2 and 51.0 for TMY and 49.1, 47.3 and 46.9% for LP.

Table 5: Numbers and percentages of Holstein animals with and without records with positive estimates of predicted transmitting abilities recorded by the Animal Model.

Trait*	Cows with records		Dams without records		Sires without records	
	No. of cows	% of cows	No. of dams	% of dams	No. of sires	% of sires
M90	469	47.9	476	46.3	70	50.4
M305	487	47.3	477	46.4	73	52.5
TMY	489	47.5	470	45.7	70	50.4
LP	559	54.3	548	53.3	72	51.8

* Traits as defined in Table 1.

CONCLUSION

In spite of the importance of sire transmitting ability in the herd, the present ranges in estimates of transmitting abilities indicate that Holstein cows (animals with records) had the highest ranges of transmitting ability for all milk traits, followed by their sires and dams (animals without records) in descending order. So, selecting cows from the herd of the study (which belongs to the General Cooperative for Developing Animal Wealth and Products, Egypt, EL-Salhia) for milk traits according to the cow transmitting ability would be more reasonable than selecting them according to either their sire transmitting ability or their dam transmitting ability.

REFERENCES

- Abdel-Glil, M.F. (1991). Sire differences for milk production traits in Friesian cattle. Ph.D. Thesis, Faculty of Agric., Mostohor, Zagazig Univ., Egypt.
- Abubakar, B.Y.; R.E. McDowell and L.D. Van Vleck (1986b). Genetic evaluation of Holstein in Colombia. *J. Dairy Sci.*, 69 (4): 1081-1086.
- Abubakar, B.Y.; R.E. McDowell; K.E. Wellington and L.D. VanVleck (1986a). Estimating genetic values for milk production in the Tropics. *J. Dairy Sci.*, 69(4): 1087-1092.
- Abubakar, B.Y.; R.E.McDowell and L.D.Van Vleck (1987). Interaction of genotype and environment for breeding efficiency and milk production of Holstein in Mexico and Colombia. *Tropical Agriculture*, 64(1):17-22.
- Ahmed, R.El.M. (1996). Actual and predicted performance of Holstein-Friesian cattle in Egypt. M.Sc. Thesis, Faculty of Agric., Ain Shams Univ., Cairo, Egypt.
- Boldman, K.G. and A.E. Freeman (1990). Adjustment of heterogeneity of variances by herd production level in dairy cow and sire evaluation. *J. Dairy Sci.*, 73(2): 503-512.

Afifi, E. A. et al.

- Ducroco, V.; D. Boichard; B. Bonaiti; A. Barbat, and M. Brieno (1990). A Pseudo-absorption strategy for solving Animal Model equations for large data files. *J. Dairy Sci.*, 73(7): 1945-1955.
- El-Chafie, O.M.B. (1981). Studies on cattle (cows and buffaloes) sire evaluation and genetic parameters of partial lactation, dry period and service period. M.Sc. Thesis, Faculty of Agric., Alex. Univ., Egypt.
- Gad, M.S. (1995). A study for some productive traits in dairy cattle. M.Sc. Thesis, Faculty of Agric., Moshtohor, Zagazig University, Egypt.
- Gebriel, E. A. A. (1996). Study on the prediction of buffalo cow transmitting ability. M.Sc. Thesis, Faculty of Agric., Ain Shams Univ., Cairo, Egypt.
- Hintez, R.L.; R.W. Everett, and L.D. Van Vleck (1978). Estimation of genetic trends from cow and sire evaluations. *J. Dairy Sci.*, 61(5): 607-613
- Khalil, M.H.; M.F. Abdel-Gliil and M.K. Hamed (1994). Genetic aspects and adjustment factors for lactation traits of Friesian cattle raised in Egypt. *Egyptian J. Anim. Prod.* 31(1): 65-84.
- Misztal, I. (1990). Continuous genetic evaluation of dairy cattle. *J. Dairy Sci.*, 73 (Sup II 1): 134.
- Moharram, A.A (1988). Reproduction and dairy performance of Holstein-Friesian cattle in Egypt. *Revue d'Elevage et de Medecine Veterinaire des pays Tropicaux.*, 41 (2): 209-213. (*A. B. A.*, 57(12): 991.
- Mokhtar, S.; M.A.El-Alamy and A. Amin (1993). Evaluation of productive and reproductive performance of some dairy cattle breeds in Egypt. *Egyptian J. Anim. Prod.*, 30 (2): 161-171.
- Rozzi, P.; L.R.Schaffer; E.B. Burnside and W. Schlote (1990). International evaluation of Holstein-Friesian dairy sires from three countries. *Livestock Prod. Sci.*, 24:15-28.
- Sadek, M.H.; A.A. Halawa and A.A. Ashmawy (1993). Effect of using the additive genetic relationship matrix of sire's proofs. *Egyptian J. Anim. Prod.*, 30(2):189-200.
- Salem, M. A. (1991). Effects of age at first calving and other environmental factors on milk production of imported and locally born Friesian cattle. M.Sc. Thesis, Faculty of Agric., Moshtohor, Zagazig Univ., Egypt.
- Soliman, A. M. and M.H. Khalil (1993). Estimation of genetic parameters for single and composite milk traits in Fleckvieh cattle and their uses in programs of early selection. *Egyptian J. Anim. Prod.* 30(1): 21-36.
- Szkotnicki, W. J.; A. K.W. Tong; M.A. Sharaby; K.M. Krotch ; L.P. Johanson, and L.D. Schaeffer (1978). Sire and cow evaluation in Brown Swiss, Canadian, and milking Shorthorn. *J. Dairy Sci.*, 61(4): 497-508.
- Wiggans, G.R. and P.M. VanRaden (1990). Animal Model evaluation within-herd linked to national evaluation. *J. Dairy Sci.*, 73(7): 1956-1963.

تقدير المقدرة التمريرية لصفات إدرار اللبن باستخدام نموذج الحيوان لماشية الهولشتين المرباه في مزرعه تجاريه في مصر عزت عطا عفيفي^١ - ماهر حسب النبي خليل^١ - سميره عبده عرفه^٢ - محمود أحمد سالم^٢

^١ كلية الزراعة بمشتهر - جامعة الزقازيق - جمهورية مصر العربية
^٢ قسم انتاج وتربية الحيوان - كلية الزراعة والطب البيطري بالقصيم - جامعة الملك سعود ص ب ١٤٨٢ بريدة - المملكة العربية السعوديه
^٢ معهد بحوث الانتاج الحيواني - وزارة الزراعة واستصلاح الاراضي جمهورية مصر العربية

تم تحليل ٥٦٦٢ سجل لبن طبيعي أعطتها ١٠٢٩ بقرة باستخدام نموذج الحيوان Animal model لتقدير المقدرة التمريرية للأبقار ولأمهاتها ولأبائهن لبعض صفات إدرار اللبن لماشية الهولشتين المرباه في مزرعة الصالحية التجارية في مصر . كانت الصفات المدروسة هي محصول اللبن لأول ٩٠ يوم، محصول اللبن في ٣٠٥ يوم، المحصول الكلي من اللبن في موسم الإدرار، بالإضافة إلى طول فترة الإدرار .

كانت قيم المعامل التكراري لهذه الصفات متوسطة أو مرتفعه قليلا (٠,٤٠٢، ٠,٣٨٨)، (٠,٤١٧) وذلك لصفات محصول اللبن لأول ٩٠ يوم، محصول اللبن في ٣٠٥ يوم، المحصول الكلي من اللبن على التوالي - ومنخفضه (٠,١٤٦) لفترة الإدرار . تم تقدير المقدرة التمريرية للأبقار باستخدام سجلاتها بالإضافة إلى تقدير هذه القيم لأمهاتها وأبائهن بدون استخدام سجلاتها . وكانت الفروق بين أعلى القيم التمريرية وأدناها لمحصول اللبن لأول ٩٠ يوم، محصول اللبن في ٣٠٥ يوم، المحصول الكلي من اللبن، طول فترة الإدرار على التوالي ٣٢١ كجم، ١٠٨٧ كجم، ١٣٨٨ كجم، ١٢,١ يوما . وكانت الفروق المناظره ١٨٢ كجم، ٦٠٠ كجم، ٨٠١ كجم، ٦,٧ يوما للأمهات ، ٢٠٨ كجم ، ٧٠٢ كجم ، ٨٥١ كجم ، ٨,٥ يوما للأباء . كانت نسب القيم التمريرية الموجبه لصفات محصول أول ٩٠ يوم، محصول ٣٠٥ يوم، المحصول الكلي، طول فترة الإدرار على التوالي ٤٧,٩ ، ٤٧,٣ ، ٤٧,٥ ، ٥٤,٣ % للأبقار ، ٤٦,٣ ، ٤٦,٤ ، ٤٥,٧ ، ٥٣,٣ % للأمهات ، ٥٠,٤ ، ٥٢,٥ ، ٥٠,٤ ، ٥١,٨ % للأباء .