

# **Use of RAPD and ISSR markers for the identification of Egyptian Henbane (*Hyoscyamus muticus* L.) genotypes.**

El-Shawaf\*, I.I.S.; Bekhit\*, M. M. M.; Hassan\*, A. M.; El-Saied\*\*, F. M. and Masoud\*\*, I. M.

\* Department of Genetics, Faculty of Agriculture, Banha University, Egypt.

\*\* Department of plant Genetic Resource, Desert Research Center, Egypt.

## **ABSTRACT**

Three Egyptian henbane genotypes from three different locations (Wadi Sudr, El Maghara and Shalateen) were screened using two different molecular Techniques; RAPD and ISSR. Both of these techniques were used to detect some molecular markers associated with the genotype identification. RAPD results, from using seven arbitrary primers, revealed 57 amplified fragments, 30 of them were polymorphic (52.63 %). The results revealed the presence of 21 positive and 9 negative molecular markers for the genotypic identification. ISSR results with five specific primers revealed 41 amplified fragments, 15 of them were polymorphic (36.59 %). The results revealed the presence of 13 positive and 2 negative molecular markers for the genotypic identification. The Similarity indices and dendrogram for the genetic distances of the combination between the two techniques revealed that the highest similarity was 68.90 % between the henbane genotypes from El Maghara and Shalateen regions. Meanwhile, the lowest similarity index was between the henbane genotypes from El Maghara and Wadi Sudr regions (8.90 %). The dendrogram resulting from the combination of the two techniques separated the three henbane genotypes into one cluster (including El Maghara and Shalateen genotypes) with the Wadi Sudr genotype alone.

## INTRODUCTION

Medicinal plants have acquired increasing significance over the last few years, not only because their healing ability, but also due to their minor side effects comparing to the chemical medicine. Worldwide, 35000 plant species are used for medicinal purposes but only 90 species are considered in the most important industrial medicines. In developing countries traditional medicine is spread because natural remedies are cheaper than chemical medicine and are often the only medicine available in the remote rural area. Besides serving medicinal functions, medicinal plants in developing countries have an important economic role because they have 90% of the earth genetic diversity exists in the developing countries.

Family Solanaceae is one of the most intriguing plant families in the world not only because it is one of the largest families in plant kingdom with more than 3,000 species. Also, because it includes species which are essential for life such as *Hyoscyamus muticus* (an important medicinal plant of the Solanaceous family).

The Solanaceae is known for possessing a diverse range of alkaloidal glucosides, or simply alkaloids. One of the most important groups of these compounds is called the tropane alkaloids. *Hyoscyamus* plants have been known from ancient time as a remedy for various diseases, and serve today as a source of their pharmaceutically active constituents; the tropane alkaloids. The medicinal importance of scopolamine, hyoscyamine and atropine is illustrated by their presence in the list of the ten substances of plant origin most used as drugs in the USA in the 1973.

Random amplified polymorphic DNA (RAPD) markers and inter - simple sequence repeats (ISSR) markers are two molecular approaches that have been used to detect variation among plants. Each method has been used extensively to identify and determine relationships at the species and cultivar levels (**Rajaseger *et al.*, 1997, Raina *et al.*, 2001 and Martins *et al.*, 2003**). These methods are widely applicable because they are rapid, inexpensive, simple to perform, do not require prior knowledge of DNA sequence and

require very little starting DNA template (**Esselman *et al.*, 1999**). The ISSR method has been reported to produce more complex marker patterns than the RAPD approach (**Parsons *et al.*, 1997**). In addition, ISSR markers are more reproducible than RAPD markers, because ISSR primers, designed to anneal to a micro-satellite sequence, are longer than RAPD primers, allowing higher annealing temperatures to be used (**Goulao and Oliveira, 2001**).

The present work aimed to identify some genotypes of Egyptian henbane (*Hyoscyamus muticus L.*) using (RAPD and ISSR) techniques as molecular genetic fingerprints. Also the result of the combination between the RAPD and ISSR techniques analysis revealing the genetic distance and the dendrogram of the henbane genotypes under study.

## MATERIALS AND METHODS

### A. Plant materials:

Three Egyptian henbane collections (*Hyoscyamus muticus L.*), (Table 1), were obtained from three different locations; Wadi Sudr, El Maghara and Shalateen, Egypt.

### B. Methods:

#### 1. DNA isolation:

Total DNA was isolated from young leaves of the Egyptian henbane genotypes (*Hyoscyamus muticus L.*) according to **Dellaporta *et al.*, (1983)**.

#### 2. RAPD analysis:

Reaction conditions for amplification were optimized according to **Williams *et al.*, (1990)** using seven arbitrary primers synthesized by Operon Technology, Inc. (USA). The amplification was performed for 42 cycles, using a Coy thermocycler programmed, as follows; initial denaturation at 94°C for 3 min, one cycle, denaturing at 94°C for 45 sec., annealing at 37°C for 30 sec., extension at 72°C for 2 min., (40 cycles) and final extension at 72°C for 12 min., (one cycle).

The product was resolved on agarose gel (1.2 %) in TAE buffer and stained with ethidium bromide. Two different DNA markers were used; a 100 bp ladder mix and a 1kb DNA ladder mix (Pharmacia Biotech). The seven

arbitrary primers used for RAPD amplification, their names and sequences are found in Table (2). The run was performed for about 1 h at 80 V in a mini gel agarose electrophoresis apparatus (Biorad). The products were visualized by UV light. PCR products were photographed by gel electrophoresis system (Gel Doc. BIORAD 2000) under transilluminator.

### 3. Inter simple sequence repeats (ISSRs) analysis:

The five used primers in PCR amplification of Inter simple sequence repeat regions (Table 3) were synthesized on an ABI 392 DNA/RNA synthesizer (Applied Biosystems) at AGERI, ARC.

The amplification was performed as RAPD but with 55°C for annealing instead of 37°C.

### 4. Genetic distance relationships:

The banding patterns of the seven RAPD and the five ISSR primers were scored. Data were feed to the PC computer as 1 and 0 for the presence and absence of bands, respectively. Pair-wise comparisons of genotypes, based on the presence or absence of unique and shared polymorphic products, were used to regenerate similarity coefficients, according to **Jaccard (1908)**. The similarity coefficients were then used to construct dendrograms, using the unweighted pair group method with arithmetic averages (UPGAMA) employing the SHAN (Sequantial, Agglomerative, Hierarchical, and Nested clustering) from the NTSYS-PC (Numerical Taxonomy and Multivariate Analysis System), version 2.1 Program (**Rohlf, 2000**).

## RESULTS AND DISCUSSION

In the present study, the genetic variability and relationships of the natural medicinal plant (*Hyoscyamus muticus* L.) which collected from three different locations have been studied. These were done based on molecular markers (RAPD and ISSR) techniques. The results obtained could be summarized as follows:

### 1. Randomly amplified polymorphic DNA (RAPD) analysis:

The banding patterns of RAPD fragments using the seven arbitrary primers with the three henbane genotypes (Fig. 1) showed 57 amplified fragments; 30 of them were polymorphic (52.63 %). The total number of amplified and polymorphic fragments obtained with each primer is found in Table (4). Results

revealed the presence of 21 positive RAPD molecular markers in addition to 9 negative molecular markers. The developing of RAPD approach (**Williams et al., 1990**) has allowed simple, easy and less time consuming genome analysis at DNA level compared with RFLP (Restriction Fragment Length Polymorphism). Similar results were obtained in wheat (**Abdel-Tawab et al., 2003**), in maize (**Abdel-Tawab et al., 2002**), in sorghum (**Abdel-Tawab et al., 1998**).

## 2. Inter simple sequence repeats (ISSR) analysis:

The banding patterns of ISSR fragments using the five specific primers with the three henbane genotypes (Fig. 2) revealed 41 amplified fragments; 15 of them were polymorphic (36.59 %). The total number of amplified and polymorphic fragments obtained with each primer is found in Table (4). ISSR data revealed 13 positive and 2 negative molecular markers for the three henbane genotypes. **Ratnaparkhe et al., (1998)**, **Reddy et al., (2002)** and **Fahmy et al., (2008)** showed that ISSR markers could be used as highly informative markers for genomic mapping and gene tagging because the evolutionary rate of change within microsatellites is considerably higher more than many other types of DNA markers.

3. Genetic relationships of the three henbane genotypes based on RAPD and ISSR molecular markers:

The identification of the wild type germplasm is necessary for varietal improvement. Recently, RAPD and ISSR techniques are used to measure the degree of similarity / dissimilarity among genotypes and also to estimate the genetic distance between these germplasm. The obtained results of the banding patterns of RAPD and ISSR were used to study the genetic relationships of the three henbane genotypes at the levels of the markers and in combination between the two techniques.

RAPD amplification revealed different degrees of polymorphisms between the henbane genotypes. The obtained 57 bands were treated with Ntsyspc2 software to release the similarity matrices (Table 5) and the dendrogram of the genetic distances (Fig. 3). The genetic similarity matrices depended on all possible pairs of genotypes ranged from 10.00 % to 66.70 %. The highest genetic similarity indices were found between henbane genotypes from El Maghara and Shalateen (66.70 %). The lowest genetic similarity indices were found between henbane genotypes from Wadi Sudr and El Maghara and Shalateen (10.00 %). Meanwhile, the genetic similarity indices between henbane genotypes from Wadi Sudr and Shalateen were (23.30 %).

The obtained results revealed broad genetic diversity within the three henbane genotypes, especially between genotypes from Wadi Sudr and Shalateen. As found in Fig. (3), the dendrogram based on RAPD similarity indices separated into one cluster including the two genotypes from El Maghara and Shalateen and Wadi Sudr genotype is alone. Moreover **Votava et al., (2005)** used random amplified polymorphic DNA (RAPD) molecular markers to compare the relative genetic diversity of native chili (*Capsicum Annuum* Var. *Annuum* L.) landraces from Northwestern Mexico. The results indicated that the molecular analysis of these accessions provided a powerful means by which their genetic structures were characterized. The same results was obtained by **Singh et al., (2006)** who used Random amplified polymorphic DNA (RAPD) technique as a tool for assessing genetic diversity and species relationships among 28 accessions of *eggplant* from different parts of the country. The results indicates presence of high level of genetic diversity in *eggplant* and a dendrogram shows that *Solanum incanum* is closest to *Solanum melongena* followed by *Solanum nigrum*.

The ISSR data revealed that the genetic similarity indices ranged from 6.70 % to 73.30 % (Table 6 and Fig. 4). The nearest relationship was detected between the henbane genotypes from El Maghara and Shalateen (73.30 %). On the other hand, the farthest relationship of similarity matrices was detected between the henbane genotypes from Wadi Sudr and El Maghara (6.70 %), followed by Wadi Sudr and Shalateen (20.00 %). As indicated by the Fig. 4, the dendrogram based on the similarity matrices of ISSR-PCR banding patterns separated the three henbane genotypes into one cluster (including El Maghara and Shalateen genotypes) and Wadi Sudr alone. ISSR amplification does not require genome sequence information but produce highly polymorphic patterns (**Zietkiewicz et al., 1994; Nagaoka and Ogihara 1997; Prevost and Wilkinson 1999**). It was applied to the study of genetic relationships and phylogenetic analysis of various crop plants (**Joshi et al. 2000; Martin and Yelmo 2000 and Iruela et al. 2002**).

The results indicated that it was possible to discriminate between *H. muticus* L. accessions which collected from three different locations, since each genotype banding pattern was not similar to the others. Based on ISSR analysis, the same results were obtained by **Kochieva et al., (2002)** who used Inter-simple sequence repeats (ISSR) analysis to study the genetic diversity and

phylogenetic relationships in 54 wild accessions and cultivars of the genus *Lycopersicon*.

Depending on the combination between the banding patterns of RAPD-PCR and ISSR-PCR data through using NTsyspc2 software, the similarity indices were 68.90 % between the henbane genotype from El Maghara and Shalateen regions, followed by 22.20 % between the henbane genotype from Wadi Sudr and Shalateen. The lowest similarity indices were detected between the henbane genotype from Wadi Sudr and El Maghara (8.90 %). The dendrogram resulted from the combination of the two techniques (Fig. 5) was found also in one cluster (including the henbane genotype from El Maghara and Shalateen regions) while the henbane genotype from Wadi Sudr found alone. **Petros *et al.*, (2007)** studied the genetic diversity among the niger populations grown in different regions of Ethiopia using ISSR markers in an effort to provide some information for future research that might be aimed for improving some of the agronomic traits of niger for efficient utilization.

**Table (1):** Name and origin of the three henbane genotypes collected from the three locations.

Number	Genotype name	Origin	Family
1	Henbane	Wadi Sudr	Solanaceae
2	Henbane	El Maghara	Solanaceae
3	Henbane	Shalateen	Solanaceae

**Table (2):** RAPD primers code and their sequences used for detection of banding patterns in the three henbane genotypes.

Primer code	Sequence (5 → 3)	Primer Code	Sequence (5 → 3)
OP A10	CTGCTGGGAC	OP D07	ACCGCGAAGC
OP A19	CAATCGCCGT	OP I15	AAGAGAGGGG
OP B12	CCTTGACGCA	OP L20	TGGTGGACCA
OP A16	TGCGCGATAG		

**Table (3):** ISSR primers code and their sequences used for detection of banding patterns in the three henbane genotypes.

Primer code	Sequence (5 → 3)	Primer Code	Sequence (5 → 3)
A 98	(CA) <sub>6</sub> GT	HB 11	(GT) <sub>6</sub> CC
HB 09	(GT) <sub>6</sub> GG	HB 13	(GA) <sub>6</sub> GC
HB 10	(GA) <sub>6</sub> CC		

Table (4): The total number of amplified and polymorphic fragments, percentage of polymorphism and specific markers in the three henbane genotypes using both of RAPD and ISSR data.

Primer Number	Primer code	TAF	PF	Polymorphism %	SM
<b>RAPD</b>					
1	OP A10	8	5	62.50 %	+3 and -2
2	OP A19	8	3	37.50 %	+3
3	OP B12	6	3	50.00 %	+3
4	OP A16	6	3	50.00 %	+2 and -1
5	OP D07	9	5	55.56 %	+2 and -3
6	OP I15	9	4	44.44 %	+2 and -2
7	OP L20	11	7	63.63 %	+6 and -1
		<b>57</b>	<b>30</b>	<b>52.63 %</b>	<b>+21 and -9</b>
<b>ISSR</b>					
1	A 98	8	5	62.50 %	+3 and -2
2	HB 09	9	2	22.22 %	-2
3	HB 10	8	2	25.00 %	+2
4	HB 11	8	2	25.00 %	+2
5	HB 13	8	4	50.00 %	+4
		<b>41</b>	<b>15</b>	<b>36.59 %</b>	<b>+13 and -2</b>

Table (5): Similarity index (Pair wise comparison) among the three henbane genotypes based on RAPD data.


Genotype	Wadi Sudr	El Maghara	Shalateen
Wadi Sudr	1		
El Maghara	0.100	1	
Shalateen	0.233	0.667	1

Table (6): Similarity index (Pair wise comparison) among the three henbane genotypes based on ISSR data.


Genotype	Wadi Sudr	El Maghara	Shalateen
Wadi Sudr	1		
El Maghara	0.067	1	
Shalateen	0.200	0.733	1

Table (7): Similarity index (Pair wise comparison) among the three henbane genotypes based on RAPD and ISSR data.

<b>Genotype</b>	<b>Wadi Sudr</b>	<b>El Maghara</b>	<b>Shalateen</b>
<b>Wadi Sudr</b>	1		
<b>El Maghara</b>	0.089	1	
<b>Shalateen</b>	0.222	0.689	1



**Fig. (1):** RAPD banding patterns of the three henbane genotypes with seven arbitrary primers; M = 100-bp ladder.



**Figure (2) :** ISSR banding patterns of the three henbane genotypes amplified with the five ISSR primers; M= 100 bp ladder.

Fig. 3: Dendrogram for the genetic relationships among the three henbane genotypes based on RAPD-PCR data.




Fig. 4: A dendrogram revealing the genetic distance among the three henbane genotypes using ISSR data.





Fig. 5: Dendrogram for the genetic relationships among three henbane genotypes based on RAPD-PCR and ISSR-PCR data.



## REFERENCES

- Abdel-Tawab, F. M.; E. M. Fahmy; A. Bahieeldin; A. A. Mohamed; H. T. Mahfouz; H. F. Eissa and O. Moseilhy (2003):**Marker-assisted selection for drought tolerance in Egyptian bread wheat (*Triticum aestivum L.*). Egyptian J. Genet. And Cytol., 31: 43-63.
- Abdel-Tawab, F. M.; E. M. Fahmy; M. A. Rashed; G. A. Gad El-Karim; S. H. Abdel-Aziz and S. E. Ibrahim (2002):**Development of molecular markers for salt and drought tolerance in maize (*Zea mays L.*). Egyptian J. Genet. And Cytol., 32: 355-371.
- Abdel-Tawab, F. M.; M. A. Rashed; A. Dhindsa; A. Bahieeldin and A. Abo Doma (1998):**Molecular markers for salt tolerance in *Sorghum bicolor*. International Congress on Molecular Genetics, 21-26 Feb., Cairo, Egypt.
- Dellaporta, S. L.; J. Wood and J. B. Hicks (1983):**  
A plant DNA preparation version II, Plant Mol. Biol. Rep. 4: 19-21.
- Esselman, E. J.; L. Jianqiang; D. J. Crawford; J. L. Windus; A. D. Wolfe (1999):** Clonal diversity in the rare *Calamagrostis porteri* ssp. *Insuperata* (Poaceae): comparative results for allozymes and random amplified polymorphic DNA (RAPD) and inter simple sequence repeat (ISSR) markers. Molecular Ecology, 8: 443–451.
- Fahmy, E. M.; N. M. Abd-El-Gawad; I. H. El-Geddawy; O. M. Saleh and N. M. El-Azab (2008):**Development of RAPD and ISSR markers for drought tolerance in sugarcane (*Saccharum officinarum L.*). Egyptian J. Genet. And Cytol., 37 (1): 1-15.
- Goulao, L. and C. M. Oliveira (2001):**Molecular characterization of cultivars of apple (*Malus x domestica* Borkh) using microsatellite (SSR and ISSR) markers. Euphytica, 122: 81–89.
- Iruela, M.; J. Rubio and J. J. Cubera (2002):**Phylogenetic analysis in the genus *Cicer* and cultivated chickpea using RAPD and ISSR markers, Theor. Appl. Genet, 104:643\_651.
- Jaccard, P. (1908):**Nouvelles recherches sur la distribution locale. Bull. Soc. Vaud. Sci. Nat., 44: 223-270.
- Joshi, S. P.; V. S. Gupta and R. K. Agarwal (2000):**Genetic diversity and phylogenetic relationship as revealed by inter simple sequence repeat (ISSR) polymorphism in the genus *Oryza*. Theor. Appl. Genet. 100: 1311-1320.
- Kochieva, E. Z.; N. N. Ryzhova; I. A. Khrapalova; V. A. Pukhalskyi (2002):** Using RAPD for Estimating Genetic Polymorphism in and Phylogenetic Relationships among Species of the Genus *Lycopersicon* (Tourn.) Mill. Russian Journal of Genetics, (38) 9: 1104-1108.
- Martin, J. P. and S. M. D. Yelema (2000):**Genetic relationships among species of the genus *Diplotaxis* (Brassicaceae) using inter simple sequence repeat markers. Theor. Appl. Genet, 101: 1234-1241.

- Martins, M; R. Tenreiro; and M. M. Oliveira (2003):**Genetic relatedness of Portuguese almond cultivars assessed by RAPD and ISSR markers. *Plant Cell Reports*, 22: 71–78.
- Nagaoka, T. and Y. Ogihara (1997):**Applicability of inter-simple sequence repeat polymorphisms in wheat for use as DNA markers in comparison to RFLP and RAPD markers. *Theor. Appl. Genet*, 94: 597-602.
- Parsons, B. J; H. J. Newbury; M. T. Jackson and B. V. Ford-Lloyd (1997):** Contrasting genetic diversity relationships are revealed in rice (*Oryza sativa L.*) using different marker types, *Mol. Breed* 3: 115-125.
- Petros, Y.; A. Merker; and H. Zeleke (2007):**Analysis of genetic diversity of *Guizotia abyssinica* from Ethiopia using inter simple sequence repeat markers. *Hereditas* 144: 18-24.
- Prevost, A. and M. J. Wilkinson (1999):**A new system of comparing PCR primers applied to ISSR fingerprinting of potato cultivars. *Theor. Appl. Genet.* 98: 107–112.
- Raina, S. N.; V. Rani; T. Kojima; Y. Ogihara; K. P. Singh; R. M. Devarumath (2001):**RAPD and ISSR fingerprints as useful genetic markers for analysis of genetic diversity, varieties identification, and phylogenetic relationships in peanut (*Arachis hypogaea*) cultivars and wild species. *Genome*, 44: 763–772.
- Rajaseger, G.; H. T. W. Tan; I. M. Turner; P. P. Kumar (1997):** Analysis of genetic diversity among *Ixora* cultivars (Rubiaceae) using random amplified polymorphic DNA. *Annals of Botany*, 80: 355–361.
- Ratnaparkhe, M. B.; M. Tekeoglu and J. Muehlbaure (1998):**Inter-simple sequence repeats (ISSR) polymorphisms are useful for finding markers associated with disease resistance gene clusters. *Theor. Appl. Genet.*, 97: 515-519.
- Reddy, M. P.; N. Sarla and E. A. Siddiq (2002):**Fourth International Rice Genetics Symposium, IRRI, Philippines, Abstract, pp. 212.
- Rohlf, F.J. (2000):**NTSYS-pc: Numerical taxonomy and multivariate analysis system. Version 2.02 Exeter Software, Setauket, N.Y.
- Singh A. K., Singh, M.; Singh, A. K.; Singh, R.; Kumar, K. and Kalloo, G. (2006):**Genetic diversity within in the genus *Solanum* (Solanaceae) as revealed by RAPD markers. *Current Science*, 90 (5): 711-716.
- Votava, E. J.; Baral, J. B. and Bosland, P. W. (2005):**Genetic diversity of Chile (*Capsicum Annuum* Var. *Annuum* L.) Landraces from Northern New Mexico, Colorado, and Mexico, *59(1)*: 8-17.
- Williams, J. G. K.; A. R. Kubelik; K. J. A. Rafalski and S. V. Tingey (1990):**DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Research*. 18: 6531-6535.
- Zietkiewicz, E., Rafalski, A. and Labuda, D. (1994):**Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20:176\_183.

## استخدام تقنيات الـ RAPD-PCR و الـ ISSR كدلائل جزيئية بهدف تعريف التراكيب الوراثية لنبات السكران المصرى (*Hyoscyamus muticus* L.).

ابراهيم سليمان الشواف<sup>١</sup>، مخلوف محمد محمود بخيت<sup>١</sup>، عبدالوهاب محمد حسن<sup>١</sup>، فريدة محمد السعيد<sup>٢</sup> و أنجى محمد مسعود<sup>٢</sup>

١ قسم الوراثة، كلية الزراعة، جامعة بنها، مصر.

٢ قسم المصادر الوراثية النباتية، مركز بحوث الصحراء، مصر.

تم تقييم ثلاثة تراكيب وراثية لنبات السكران المصرى من ثلاثة مواقع مختلفة (وادي سدر، المغارة و شلاتين) و ذلك باستخدام تقنية كل من الـ RAPD و الـ ISSR. تم استخدام كل من الطريقتين للحصول على دلائل جزيئية مرتبطة بتعريف هذه التراكيب الوراثية. كانت نتائج طريقة الـ RAPD المستخلصة من استخدام ٧ بادئات جزيئية عشوائية ٥٧ شظية مكبرة، ٣٠ شظية منها كانت متباينة بنسبة ٥٢،٦٣ % . أظهرت النتائج وجود ٢١ دليل جزيئى موجب و ٩ دلائل جزيئية سالبة بالنسبة لتعريف التراكيب الوراثية. أوضحت نتائج الـ ISSR باستخدام ٥ بادئات متخصصة وجود ٤١ شظية مكبرة، ١٥ شظية منها كانت متباينة بنسبة ٣٦،٥٩ % حيث أوضحت النتائج وجود ١٣ دليل جزيئى موجب و دليلين جزيئيين سالبين بالنسبة لتعريف التراكيب الوراثية لنبات السكران المصرى. بينت نتائج القراية الوراثية والشجرة الوراثية بالنسبة للمسافات الوراثية من التحليل المشترك لكل من الـ RAPD و الـ ISSR أن أعلى نسبة تشابه كانت ٦٨،٩٠ % بين التراكيب الوراثية لنبات السكران من مناطق المغارة و شلاتين. بينما كانت أقل نسبة تشابه ٨،٩٠ % بين التراكيب الوراثية من مناطق وادي سدر و المغارة. أوضحت نتائج الشجرة الوراثية الناتجة من التحليل المشترك بين الـ RAPD و الـ ISSR حدوث انفصال الى عنقود واحد يضم كل من التراكيب الوراثية من مناطق المغارة و شلاتين و وجود التركيب الوراثى من منطقة وادي سدر منفردا.