

Fatty Acids in Heterocyclic Synthesis. Part XIV: Synthesis of Surface Active Agents from Some Novel Class of Oxadiazole, Thiadiazole and Triazole Derivatives Having Microbiological Activities

Iman A. Gad El-Karim · Mahasen S. Amine ·
Amal A. Mahmoud · Alaa S. Gouda

Received: 17 March 2013 / Accepted: 29 August 2013
© AOCs 2013

Abstract Reaction of stearic acid with semicarbazide in refluxing POCl_3 afforded 2-amino-5-heptadecyl 1,3,4-oxadiazole. Acylation of the amino group with acetic anhydride, ethyl chloroacetate and chloroacetic acid gave amide and β -amino acid derivatives. These compounds were cyclized to imidazo[2,1-*b*]oxadiazole derivatives by two different techniques. Treating the starting oxadiazole compound with P_2S_5 , hydroxyl amine and hydrazine hydrate in benzene afforded thiadiazole and triazole derivatives. Unexpectedly, triazolo[3,4-*b*][1,3,4]oxadiazole derivative was obtained when 1,3,4-oxadiazole derivative was refluxed with hydrazine hydrate in ethanol. The biological activities of the synthesized compounds were screened in vitro against some gram positive and gram negative bacteria and fungi. Addition of quantitative amount of propylene oxide units (3, 5, 7 mol) to the synthesized compounds afforded new nonionic surfactants. The physico-chemical and surface properties of the novel synthesized surfactants such as surface and interfacial tension, cloud point, wetting time, emulsion stability, foam height, CMC, resistance to hydrolysis and their biodegradability were investigated. In addition, surface parameters including effectiveness (π_{CMC}), efficiency (PC_{20}), maximum surface excess (Γ_{max}) and (A_{min}) were examined.

Keywords Stearic acid · 2-Aminooxadiazole · Nonionic surfactant · Antimicrobial · Triazolo[3,4-*b*]oxadiazole · Imidazo[3,1-*b*]oxadiazole · Thiadiazole · Triazole · Surface properties

Introduction

2-Amino oxadiazoles have a great importance in biological and industrial applications. They have bactericidal, fungicidal [1], herbicidal, and insecticidal activities [2]. In addition, they are employed as anti-inflammatory [3], analgesic [4], anticonvulsant, nervous system depressant [5], antiallergic [6] and antiviral drugs [7]. Conjugated systems of 2,5-disubstituted-[1,3,4]oxadiazoles often fluoresce, such a property makes them potentially useful as laser dyes, optical brighteners, scintillators, luminescent dyes and photosensitizing materials [8]. Having a long alkyl chain (hydrophobic part) attached to the oxadiazole nucleus increases their usefulness even further. Moreover, the introduction of a quantitative amount of hydrophilic part such as polyoxyethylene, or polyoxypropylene increases the solubility of 1,3,4-oxadiazoles in water and enhances the potential of their usage as nonionic surfactants. Generally, surfactants are one of the most important and widely used products in nearly every industrial sector. The growing and diverse applications of nonionic surfactants in therapy [9], industry [10], agriculture [11] and housing [12] prompted us to continue our research program on the utilization of fatty acids in heterocyclic synthesis, to synthesize novel nonionic surfactants [13–17]. Herein, we report the synthesis and some reactions of 2-amino-5-heptadecyl[1,3,4]oxadiazole (**1**) and the physico-chemical and surface properties of the novel synthesized surfactants.

Electronic supplementary material The online version of this article (doi:10.1007/s11743-013-1530-9) contains supplementary material, which is available to authorized users.

I. A. Gad El-Karim (✉) · M. S. Amine ·
A. A. Mahmoud · A. S. Gouda
Department of Chemistry, Faculty of Science, Benha University,
P.O.:13518, Benha, Egypt
e-mail: i.elkarim@gmail.com

Experimental Protocols

Materials

Stearic acid (Chemajet Chemical Company, 99 % GPR), semicarbazide hydrochloride (Winlab, 99 %), POCl₃ (Alpha Chemie, ≥98 %), 4-chlorobenzaldehyde [Fluka Chemika, ≥98 % (NT), piperidine (Merck Schuchardt, >98 % (GC)], acetic anhydride (Adwic, 98 %), ethyl chloroacetate (Loba Chemie, >98 %), chloroacetic acid [Aldrich, 99 % (T)], pyridine (LR) [SD Fine-Chem Limited, 99 % (GC)], glacial acetic acid (S D Fine-Chem Limited, 99 %), phosphorous pentasulfide (Sigma-Aldrich, 95 %), hydroxylamine hydrochloride (Winlab, 98 %), hydrazine hydrate (Loba Chemie, 80 %), benzene, toluene (Adwic), ethanol abs. (Adwic, 99 %), methanol (S D Fine-Chem Limited, 99 %), paraffin oil (Aldrich, dynamic viscosity; 110–230 mPa s, density; 0.827–0.890 g/mL at 20 °C).

Structure Confirmation

The structural assignments of new compounds are based on their elemental analysis and spectral data (IR, ¹H NMR, ¹³C NMR, and Mass spectra) (Full description of the obtained data are presented in the supplementary material section). All melting points are uncorrected and determined by the open capillary method using a Gallenkamp melting point apparatus.

IR spectra (KBr disk) of the synthesized compounds were recorded on FT/IR-BRUKER, Vector 22 (Germany), JASCO FT/IR-4100 (Japan), and JASCO FT/IR-460+ (Japan). ¹H- and ¹³C-NMR spectra were recorded in deuterated chloroform (CDCl₃) or dimethyl sulfoxide (DMSO-d₆) as a solvent on a Varian Mercury VXR-300 spectrometer (200 MHz for ¹H NMR and 75 MHz for ¹³C NMR) using TMS as internal reference. Chemical shifts are expressed in ppm. The mass spectra of compounds **1**, **7**, **11** and **12** were recorded on a Shimadzu GCMS-QP-1000EX mass spectrophotometer and those of **2–6** and **8–10** were recorded on a Shimadzu QP-2010 Plus mass spectrophotometer at 70 eV.

Homogeneity of all compounds synthesized was checked by TLC. All the synthesized compounds gave satisfactory elemental analyses. Surface active properties were carried out at the Chemistry Department, Faculty of Science, Benha University, Egypt. Antibacterial and antifungal activities were carried out in the Micro Analytical Center, Faculty of Science, Cairo University.

Synthesis

Synthesis of 2-amino-5-heptadecyl-1,3,4-oxadiazole (1). A mixture of stearic acid (0.01 mol) and semicarbazide

hydrochloride (0.01 mol) in (20 mL) POCl₃ was refluxed for 8 h. The reaction mixture was concentrated, cooled, and then poured onto ice while stirring. The produced solid was filtered off and recrystallized from ethanol. Compound **1** was obtained as a pale yellow powder in 77 % yield, mp 102–104 °C. Anal. Calc. (%) for C₁₉H₃₇N₃O: C; 70.54, H; 11.53, N; 12.99. Found: C; 70.23, H; 11.29, N; 12.81.

Synthesis of (4-chloro-benzylidene)-(5-heptadecyl-[1,3,4]oxadiazol-2-yl)-amine (2). A mixture of **1** (0.01 mol) and 4-chlorobenzaldehyde (0.01 mol) in ethanol (30 mL) in the presence of piperidine (0.5 mL) was refluxed for 6 h. The reaction mixture was concentrated, cooled and poured into ice-HCl. The obtained solid product was filtered off, dried and recrystallized from ethanol. Compound **2** was obtained as a white powder in 81 % yield, mp 136–138 °C. Anal. Calc. (%) for C₂₆H₄₀ClN₃O: C; 70.01, H; 9.04, N; 9.42, Cl; 7.95. Found: C; 70.13, H; 9.29, N; 9.61, Cl; 8.05.

Synthesis of N-(5-heptadecyl-[1,3,4]oxadiazol-2-yl)-acetamide (3). First, 0.01 mol of **1** was refluxed in 30 mL of acetic anhydride for 3 h. The reaction mixture was concentrated, poured into cold water (30 mL). The solid product obtained was filtered off, dried and recrystallized from acetic acid. Compound **3** was obtained as a deep brown powder in 75 % yield, mp 85–87 °C. Anal. Calc. (%) for C₂₁H₃₉N₃O₂: C; 69.00, H; 10.75, N; 11.49. Found: C; 69.13, H; 10.29, N; 11.61.

Synthesis of 2-chloro-N-(5-heptadecyl-[1,3,4]oxadiazol-2-yl)-acetamide (4). A mixture of **1** (0.01 mol) and ethyl chloroacetate (0.01 mol) in ethanol (30 mL) was refluxed for 8 h. The reaction mixture was concentrated and cooled. The solid product was filtered off, dried and recrystallized from ethanol. Compound **4** was obtained as a pale brown powder in 85 % yield, mp 132–134 °C. Anal. Calc. (%) for C₂₁H₃₈ClN₃O₂: C; 63.06, H; 9.58, N; 10.51, Cl; 8.86. Found: C; 63.23, H; 10.19, N; 10.62, Cl; 8.65.

Synthesis of 2-[(5-heptadecyl-[1,3,4]oxadiazol-2-yl)amino]acetic acid (5). A mixture of **1** (0.01 mol) and chloroacetic acid (0.01 mol) in benzene (30 mL) in the presence of piperidine (0.5 mL) was refluxed for 7 h. The reaction mixture was concentrated, cooled and poured into ice-HCl. The solid product obtained was filtered off, dried and recrystallized from benzene. Compound **5** was obtained as a pale gray powder in 71 % yield, mp 126–128 °C. Anal. Calc. (%) for C₂₁H₃₉N₃O₃: C; 66.14, H; 10.24, N; 11.02. Found: C; 66.45, H; 10.51, N; 10.69.

Synthesis of 2-heptadecylimidazo[2,1-b][1,3,4]oxadiazol-6(5H)-one (6). Initially, 0.01 mol of **4** was refluxed with benzene (30 mL) in the presence of pyridine (0.5 mL) for 5 h or subjected to fusion in an oil bath at 170 °C for 2 h. The reaction mixture was concentrated, cooled and poured into ice-HCl. The obtained solid product was filtered off, dried and recrystallized from ethanol. Compound

6 was obtained as yellow powder in 70 % yield, mp 149–151 °C. Anal. Calc. (%) for $C_{21}H_{37}N_3O_2$: C; 69.38, H; 10.26, N; 11.56. Found: C; 69.55, H; 10.41, N; 11.68.

Synthesis of 2-heptadecylimidazo[2,1-b][1,3,4]oxadiazol-5(6H)-one (7). (0.01 mol) of **5** was refluxed in (30 mL) glacial acetic acid containing sodium acetate (0.01 mol) for 5 h or fused in an oil bath at 170 °C for 2 h. The reaction mixture was concentrated, cooled and poured into ice with stirring. The solid product obtained was filtered off, washed several times with water and recrystallized from ethanol. Compound **7** was obtained as a gray powder in 68 % yield, mp 161–163 °C. Anal. Calc. (%) for $C_{21}H_{37}N_3O_2$: C; 69.38, H; 10.26, N; 11.56. Found: C; 69.55, H; 10.41, N; 11.68.

Synthesis of 2-amino-5-heptadecyl-1,3,4-thiadiazole (8). A mixture of **1** (0.01 mol) and phosphorous pentasulfide (0.02 mol) was refluxed in (30 mL) dry xylene until the color changes into orange. The reaction mixture was filtered while hot and the filtrate was left to cool at room temperature. The obtained solid was filtered off, dried and recrystallized from benzene. Compound **8** was obtained as a yellow powder in 80 % yield, mp 110–112 °C. Anal. Calc. (%) for $C_{19}H_{37}N_3S$: C; 67.20, H; 10.98, N; 12.37, S; 9.44. Found: C; 66.91, H; 10.64, N; 12.08, S; 9.13.

Synthesis of 3-amino-5-heptadecyl[1,2,4]triazol-4-ol (9). A mixture of **1** (0.01 mol) and hydroxylamine hydrochloride (0.01 mol) was refluxed in (30 mL) pyridine for 4–5 h. The reaction mixture was cooled and poured into ice-HCl. The solid product obtained was filtered off, dried and recrystallized from benzene. Compound **9** was obtained as a greenish black powder in 69 % yield, mp 104–106 °C. Anal. Calc. (%) for $C_{19}H_{38}N_4O$: C; 67.41, H; 11.31, N; 16.55. Found: C; 67.53, H; 11.22, N; 16.78.

Synthesis of 5-heptadecyl[1,2,4]triazole-3,4-diamine (10). A mixture of **1** (0.01 mol) and hydrazine hydrate (0.01 mol) was refluxed in 30 mL of benzene for 6 h. The reaction mixture was concentrated and cooled. The solid was filtered off, dried and recrystallized from benzene. Compound **10** was obtained as a yellowish brown powder in 78 % yield, mp 94–96 °C. Anal. Calc. (%) for $C_{19}H_{39}N_5$: C; 67.61, H; 11.65, N; 20.75. Found: C; 67.54, H; 11.41, N; 20.58.

Synthesis of 6-heptadecyl[1,2,4]triazolo[3,4-b][1,3,4]oxadiazol-3-amine (11) and ethyl-N-[amino(hydrazinylidene)methyl]heptadecylhydrazonoate (12). A mixture of **1** (0.01 mol) and hydrazine hydrate (0.01 mol) was refluxed in 30 mL of ethanol for 5–6 h. Compound **11** was obtained directly on heating and separated, filtered off, dried and recrystallized from acetic acid. While the filtrate was concentrated, cooled then the solid **12** was filtered off, dried and recrystallized from ethanol. Compound **11** was obtained as a white powder in 69 % yield, mp 130–132 °C;

Anal. Calc. (%) for $C_{20}H_{37}N_5O$: C; 66.08, H; 10.26, N; 19.26. Found: C; 66.14, H; 10.31, N; 19.31.

Compound **12** was obtained as a pale yellow powder in 73 % yield, mp 98–100 °C. Anal. Calc. (%) for $C_{21}H_{45}N_5O$: C; 65.75, H; 11.82, N; 18.26. Found: C; 65.58, H; 12.01, N; 18.39.

Antimicrobial Activities

The antimicrobial susceptibility or resistance profile of the synthesized compounds was determined with the modified Kirby-Bauer disc diffusion technique [18–21], using Mueller–Hinton agar.

One hundred microliters of the test bacteria/fungi were grown in 10 mL of fresh media until they reached a count of approximately 108 cells/mL for bacteria and 105 cells/mL for fungi. The microbial suspension (100 μ L) was spread onto agar plates that are relevant to the broth in which they are maintained. Standard discs of Tetracycline (Antibacterial agent), Amphotericin B (Antifungal agent) were used as positive control for antimicrobial activity, while filter discs impregnated with 10 μ L of solvent (distilled water, chloroform, DMSO) were used as a negative control. Blank paper discs (Schleicher and Schuell, Spain) with a diameter of 8.0 mm were impregnated with 10 μ L of the tested concentration.

When the filter paper disc impregnated with the tested compound (of known concentration) was placed on agar, the compound diffused from the disc into the agar in the area around the disc. The solubility of the tested compound and its molecular size determine the size of the area of its infiltration around the disc. An organism that is susceptible to the tested compound will not grow in the area around the disc. The size of this inhibition zone is used as a measure of the antimicrobial activity “A” of a tested compound at its minimum inhibitory concentration (MIC). The observed data on the antimicrobial activity of the compounds are given in Table 1.

Preparation of Nonionic Surfactants from the Synthesized Heterocyclic Compounds

Propoxylation (hydroxylation)

Following Morgos procedure [22], 0.5 wt% KOH solution containing 0.01 mol of the synthesized compound was stirred and heated to 70 °C while passing a slow stream of nitrogen through the system to flush out oxygen. The nitrogen stream was stopped and propylene oxide was added drop-wise with continuous stirring and heating under an efficient reflux system to retain the propylene oxide. The reaction was conducted for different intervals of time ranging from 1–10 h. The apparatus was then filled with

Table 1 Antimicrobial activity of compounds **1–12**

Sample	Inhibition zone diameter (mm/mg sample)							
	<i>Escherichia coli</i> G ⁻		<i>Staphylococcus aureus</i> G ⁺		<i>Aspergillus flavus</i> Fungus		<i>Candida albicans</i> Fungus	
	MIC	A	MIC	A	MIC	A	MIC	A
Control: DMSO	–	0.0	–	0.0	–	0.0	–	0.0
Positive reference								
<i>Tetracycline</i>	200	31	200	30	–	–	–	–
<i>Amphotericin B</i>	–	–	–	–	200	16	200	19
1	100	14	200	13	400	0.0	400	0.0
2	200	11	200	11	400	0.0	400	0.0
3	200	10	100	11	400	0.0	400	0.0
4	400	12	200	11	400	0.0	400	0.0
5	400	11	100	10	400	0.0	400	0.0
6	400	15	100	14	400	11	400	0.0
7	400	14	200	13	400	12	400	0.0
8	100	14	200	15	400	0.0	400	0.0
9	100	13	400	13	400	0.0	400	0.0
10	100	18	200	15	400	0.0	400	12
11	100	16	200	14	400	13	400	11
12	200	12	100	11	400	0.0	400	0.0

A = 0.0, Not active; A < 7 mm, slightly active; A > 15 mm, moderately active; A > 20 mm, highly active

MIC minimum inhibitory, A antimicrobial activity of tested compounds; concentration

nitrogen and cooled. The reaction vessel was weighed. The amount of reacted propylene oxide and the average degree of propoxylation were determined from the increment in the mass of the reaction mixture [23]. The selected average numbers of moles, n , are 3, 5 and 7.

Surface Active Properties of Surfactants

Surface and Interfacial Tensions

Surface and interfacial tension measurements on **I(a–c)**–**IX(a–c)** were carried out according to Findlay [24] with a Krüss tensiometer [25] (Krüss GmbH, Hamburg, Instrument Nr. K6) for different concentrations of the synthesized surfactants ($0.05\text{--}10^{-6}$ mol/L), using a platinum-iridium ring at constant temperature (25 ± 1 °C). Paraffin oil was used for the interfacial tension measurements. The tensiometer was calibrated using the method described in ASTM Designation: D1331-01 [26].

Cloud Point

The cloud point is a measure of the inverse solubility of a nonionic surface active agent. In a temperature-controlled bath, a 1-wt% solution of the tested compound was gradually heated until the clear or nearly clear solution became definitely turbid [27]. The temperature was then recorded

and the solution was allowed to cool down until it became clear again. The process was repeated to check the reproducibility of the recorded temperature.

Wetting Time

Wetting time was measured by immersing a cotton skein (1 g) in a 0.1 wt% solution of the prepared surfactants in distilled water at 25 °C according to the Draves technique [28]. The sinking time was measured in seconds.

Foaming Properties

Foam height was measured by the Ross Miles method [29]. In this procedure a given surfactant solution was allowed to fall from a set height into the same surfactant solution in a volumetric cylinder, hence creating foam. The height of the foam was visually assessed.

Emulsion Stability

The emulsifying property of the prepared surfactants was determined as follows: In a 100-mL graduated stoppered tube; an aqueous solution of the surfactant (10 mL, 20 mol) was mixed with light paraffin oil (6 mL). The mixture was shaken vigorously by magnetic stirring [Thermo scientific CimarecTM stirring hot plate, model

no: sp131320-30, estimated stirring speed (1,100 rpm)] for 2 min at 25 °C. The tube was placed upright and the separation of the formed emulsion was observed. The time taken for the separation of (9 mL) of the aqueous layer indicates the emulsion stability of the surfactant [30].

Critical Micelle Concentration (CMC) Measurements

The critical micelle concentration (CMC) is the minimum concentration at which surfactants molecules begin to form micelles [31]. CMC values were obtained through a conventional plot of the surface tension versus the logarithm of concentration of surfactant. The CMC concentration corresponds to the point where the surfactant first shows the lowest surface tension, and after which the surface tension remains nearly constant.

Effectiveness (π_{CMC})

The effectiveness of a certain surfactant π_{CMC} is expressed in terms of the decrease in the surface tension that is induced by this surfactant at the critical micelle concentration [32]. It is calculated from the difference between the surface tension of pure water (γ_0) and the surface tension of the surfactant solution at the critical micelle concentration (γ_{CMC}), Eq. (1).

$$\pi_{CMC} = \gamma_0 - \gamma_{CMC} \quad (1)$$

Efficiency (PC_{20})

Efficiency of a surfactant (PC_{20}) is defined by the values of the negative logarithm of the bulk concentration necessary to reduce surface tension by 20 mN/m [33]. It can be calculated from the following equation, Eq. (2).

$$PC_{20} = \frac{\gamma - 20 - \gamma_{CMC}}{2.303nRT} - \log C_{CMC} \quad (2)$$

where R universal gas constant 8.31×10^7 ergs $\text{mol}^{-1} \text{K}^{-1}$. T absolute temperature K.

Maximum Surface Excess Γ_{max}

The values of the maximum surface excess Γ_{max} —expressed in mol/cm^2 were calculated from surface or interfacial data by the use of Gibbs equation [34], Eq. (3):

$$\Gamma_{max} = \frac{-1}{2.303} RT \left(\frac{\delta\gamma}{\delta \log C} \right)_T \quad (3)$$

where $\delta\gamma$ surface pressure in mN/m. C surfactant concentration. $(\delta\gamma/\delta \log C)_T$ is the slope of a plot of surface tension versus concentration curves below CMC at constant temperature.

Minimum Surface Area (A_{min})

Knowing Γ_{max} , it is easy to calculate the effective area occupied by each surfactant molecule adsorbed at the air/water interface at surface saturation [35, 36]. The average area A_{min} (in $\text{\AA}^2/\text{mol}$) is given by Eq. (4):

$$A_{min} = 10^{16} / \Gamma_{max} N \quad (4)$$

where N Avogadro's number 6.023×10^{23} .

Resistance to Hydrolysis

The resistance of a certain surfactant towards acid and base hydrolysis was established by measuring the surface tension of that surfactant in acidic and alkaline media. Thus, the surface tension of a 0.1 % solution of the surfactant in 5 % sulfuric acid or in 1 % sodium hydroxide was measured at room temperature after boiling for 30 and 60 min.

Biodegradability of the Synthesized Surfactants

The biodegradation tests of the synthesized nonionic surfactants were performed according to the River Water Die-Away method [37]. The river water for testing was sampled from the River Nile. In this test, a stirred solution containing the tested surfactant (1,000 ppm) was incubated at 25 °C. Samples were withdrawn daily, filtered using Whatman filter paper and the surface tension was measured using a Du-Nouy tensiometer (Kruss type K6). The process was repeated for 7 days. The biodegradation percentage $D\%$ was calculated in terms of the measured surface tension according to Eq. (5):


$$D = [(\gamma_t - \gamma_0) / (\gamma_{bt} - \gamma_0)] \times 100 \quad (5)$$

where γ_t surface tension at time t . γ_0 surface tension at time zero (initial surface tension). γ_{bt} surface tension of the blank experiment at time t .

Results and discussion

In continuation of our research program on the utilization of fatty acids in the synthesis of heterocyclic compounds as precursors for novel nonionic surfactant [13–17], herein we report the synthesis and some reactions of 2-amino-5-heptadecyl[1,3,4]oxadiazole (**1**) (Scheme 1). Thus, the reaction of stearic acid with semicarbazide hydrochloride in refluxing POCl_3 afforded **1** in good yield. Formation of **1** was confirmed by spectroscopic tools, for instance, the mass spectra showed $[\text{M}^+]$ at 323 (1.0 %).

Condensation of compound **1** with *p*-chlorobenzaldehyde produced a Schiff's base, namely, (4-chloro-benzylidene)-(5-heptadecyl-[1,3,4]oxadiazol-2-yl)-amine (**2**), while



Scheme 1 Synthetic routes of compounds 1–12 and surfactants I–IX (i) *p*-Chlorobenzaldehyde, ethanol, piperidine, reflux 6 h; (ii) acetic anhydride, reflux 3 h; (iii) Cl–CH₂COOEt, EtOH, reflux 8 h; (iv) Cl–CH₂COOH, benzene, piperidine, reflux 7 h; (v) benzene, pyridine, reflux 5 h/or fusion in oil-bath for 2 h; (vi) AcOH, AcONa, reflux 5 h

or fusion for 2 h in oil-bath; (vii) P₂S₅, dry xylene, reflux till orange color; (viii) NH₂OH.HCl, pyridine, reflux 4–5 h; (ix) N₂H₄·H₂O, benzene, reflux 6 h; (x) N₂H₄·H₂O, EtOH, reflux 5–6 h; (xi) *n* = 3, 5 and 7 mol of propylene oxide (P.O.)

acetylation of compound 1 with Ac₂O produced *N*-(5-heptadecyl[1,3,4]oxadiazol-2-yl)acetamide (3).

Reaction of 2-amino oxadiazole 1 with ethyl chloroacetate in refluxing ethanol and with chloroacetic acid in benzene in the presence of catalytic amount of piperidine, produced 2-chloro-*N*-(5-heptadecyl[1,3,4]oxadiazol-2-yl)acetamide (4) and 2-[(5-heptadecyl[1,3,4]oxadiazol-2-yl)amino]acetic acid (5), respectively.


Furthermore, in order to increase the utility of oxadiazoles, compounds 4 and 5 were subjected to cyclization. Thus, cyclization of compound 4 either by fusion or refluxing in benzene in the presence of pyridine produced 2-heptadecylimidazo[2,1-*b*][1,3,4]oxadiazol-6(5*H*)-one (6); the formation of which was confirmed by spectroscopic analysis. IR of compound 6 lacked any ν_{NH} but showed $\nu_{\text{C=O}}$ of α - β unsaturated five-membered ketone at 1,674 cm⁻¹. The mass

spectrum showed [M⁺] and [M⁺]+1 at 363 (0.68 %) and 364 (0.81 %), respectively.

Cyclization of compound 5 using sodium acetate in acetic acid afforded 2-heptadecylimidazo[2,1-*b*][1,3,4]oxadiazol-5(6*H*)-one (7). The formation of 7 was evident from its IR which lacked any bands for the carboxy hydroxyl group, and showed the expected $\nu_{\text{C=O}}$ of unconjugated cyclic five membered ketone at 1,681 cm⁻¹. ¹H NMR showed singlet band at 2.88 ppm for cyclic CH₂ moiety which supported the proposed structure.

Substitution of the oxygen atom of oxadiazoles by a sulfur or nitrogen atom represents a vital route to obtain thiadiazole or triazole derivatives, respectively. Thus, when oxadiazole 1 was allowed to react with P₂S₅ in refluxing xylene, 2-amino-5-heptadecyl[1,3,4]-thiadiazole (8) was obtained.

Scheme 2 Proposed mechanism for the formation of compounds **11** and **12**



The synthesis of compound **8** was previously achieved by the reaction of thiosemicarbazide and stearic acid [38].

On the other hand, reaction of **1** with hydroxylamine hydrochloride in pyridine afforded 3-amino-5-heptadecyl[1,2,4]triazol-4-ol (**9**). Similarly, oxadiazole **1** reacted with hydrazine hydrate in refluxing benzene to give 5-heptadecyl [1,2,4]triazole 3,4-diamine (**10**).

Contrary to what was expected, carrying out the reaction in boiling ethanol afforded two different compounds, none of them being the expected triazole **10**. These compounds were identified as 6-heptadecyl[1,2,4]triazolo[3,4-*b*][1,3,4]oxadiazol-3-amine (**11**) and ethyl-*N*-[amino(hydrazinylidene)methyl]heptadecylhydrazonoate (**12**). The mechanism proposed to account for the formation of **11** and **12** involved an initial nucleophilic attack of ethanol on the oxadiazole nucleus followed either by cycloelimination [path a] with the subsequent 1,3-dipolar cycloaddition of intermediate (**I**) to **1** leading to the formation of the thermodynamically more stable triazole **11** rather than 5-heptadecyl[1,2,3]triazolo[5,1-*b*][1,3,4]oxadiazol-3-amine (**13**); or by ring opening [path b] with the subsequent reaction with hydrazine to produce (**12**) (Scheme 2). The formation of the isomeric triazole (**13**) is ruled out since its formation necessitates 1,3-dipolar cycloaddition of intermediate (**I**) with inverse electron demand (Scheme 2), in addition, ab initio calculations at the RHF/6-31G level indicated that the triazole **11** is more stable than **13** by 22.24 kcal/mol. The structure of both compounds was confirmed by spectroscopic analysis. Similar heterolytic addition of low molecular weight alcohols to the C=N of oxadiazole has been reported for the photochemical reaction of 2-phenyl- and 2,5-diphenyl-1,3,4-oxadiazoles [39, 40].

Antimicrobial Activities of the Synthesized Oxadiazole Derivatives

All the synthesized compounds (**1–12**) were screened in vitro (using a modified Kirby-Bauer disc diffusion

method [18–21] against Gram-negative bacteria, namely, *Escherichia coli*, Gram-positive bacteria *Staphylococcus aureus* and fungi namely, *Aspergillus flavus* and *Candida albicans*. Tetracycline and Amphotericin B were taken as positive references for antibacterial and antifungal agents, respectively. The antimicrobial activity A of a tested compound is determined at its minimum inhibitory concentration (MIC), i.e. at the lowest concentration of the compound that completely inhibited visible growth of a specific microorganism.

As outlined in Table 1, the synthesized compounds show varying degrees of inhibition against the tested microorganisms. Thus, while 5-heptadecyl-[1, 3, 4]oxadiazole derivatives (**1–5**), 2-amino-5-heptadecyl[1,3,4]-thiadiazole (**8**), 5-heptadecyl-[1,3,4]triazole derivative (**9**), and hydrazinylidene hydrazonoate derivative (**12**) are inactive toward the tested fungi, they have exhibited moderate activity against both bacteria strains. On the other hand, imidazo[2, 1-*b*][1,3,4]oxadiazolone derivatives (**6,7**) exhibit moderate to high activity towards the tested bacteria strains and exhibit good antifungal activity against *Aspergillus flavus*, while, 5-heptadecyl [1,2,4]triazole 3, 4-diamine (**10**) reveals good activity toward both bacteria strains and exhibits moderate antifungal activity against *Candida albicans*. Triazolo[3,4-*b*][1,3,4]oxadiazole (**11**) is considered to be the most active compound in this series since it shows good to very good activity against both bacterial and fungal strains. These results are in agreement with previously reported results for oxadiazole, oxathiazole and triazoles derivatives [41–44].

Nonionic Surfactants from Some Synthesized Compounds

Propoxylation of some of the new compounds (**1, 3, 4, 5** and **8–12**) with various quantities of propylene oxide (3, 5, and 7 mol) produced nonionic surfactants **I(a–c)**- **IX(a–c)**, the

structure of which was confirmed via IR and $^1\text{H-NMR}$ spectra. Thus, in addition to the original bands reported for these compounds, IR spectrum showed a broad band in the region (3,500–2,500) cm^{-1} (νOH) and two other bands in the region (1,100–1,000) and (950–900) cm^{-1} for ($\nu\text{C-O-C}$ ether linkage of polypropoxy chain). The $^1\text{H-NMR}$ spectrum showed the protons of the propyleneoxy groups which appear as broad multiple signals in the region (3.2–3.7) ppm. The physical properties of these compounds are shown in Table 2.

Surface Active Properties

The surface active properties of the prepared propoxylated compounds **I(a–c)** to **IX(a–c)** were measured in a neutral medium. The data obtained are outlined in Table 3a, b.

Surface and Interfacial Tensions

Surfactants are used in a large number of applications due to their ability to alter the energy relationships at interfaces and to lower the surface and interfacial tension [45]. As

Table 2 Physicochemical properties of the synthesized surfactants

Compd.	M. F.	M.wt	Color	Shape
Ia	$\text{C}_{28}\text{H}_{55}\text{N}_3\text{O}_4$	497	Yellow	Semi-solid
Ib	$\text{C}_{34}\text{H}_{67}\text{N}_3\text{O}_6$	613	Yellow	Semi-solid
Ic	$\text{C}_{40}\text{H}_{79}\text{N}_3\text{O}_8$	729	Yellow	Semi-solid
IIa	$\text{C}_{30}\text{H}_{57}\text{N}_3\text{O}_5$	539	Black	Powder
IIb	$\text{C}_{36}\text{H}_{69}\text{N}_3\text{O}_7$	655	Black	Powder
IIc	$\text{C}_{42}\text{H}_{81}\text{N}_3\text{O}_9$	771	Black	Powder
IIIa	$\text{C}_{30}\text{H}_{56}\text{N}_3\text{O}_5\text{Cl}$	573	Brown	Semi-solid
IIIb	$\text{C}_{36}\text{H}_{68}\text{N}_3\text{O}_7\text{Cl}$	689	Brown	Semi-solid
IIIc	$\text{C}_{42}\text{H}_{80}\text{N}_3\text{O}_9\text{Cl}$	805	Brown	Semi-solid
IVa	$\text{C}_{30}\text{H}_{57}\text{N}_3\text{O}_6$	555	Dark-gray	Semi-solid
IVb	$\text{C}_{36}\text{H}_{69}\text{N}_3\text{O}_8$	671	Dark-gray	Semi-solid
IVc	$\text{C}_{42}\text{H}_{81}\text{N}_3\text{O}_{10}$	787	Dark-gray	Semi-solid
Va	$\text{C}_{28}\text{H}_{55}\text{N}_3\text{O}_3\text{S}$	513	Pale-yellow	Powder
Vb	$\text{C}_{34}\text{H}_{67}\text{N}_3\text{O}_5\text{S}$	629	Pale-yellow	Powder
Vc	$\text{C}_{40}\text{H}_{79}\text{N}_3\text{O}_7\text{S}$	745	Pale-yellow	Powder
VIa	$\text{C}_{28}\text{H}_{56}\text{N}_4\text{O}_4$	512	Dark-green	Semi-solid
VIb	$\text{C}_{34}\text{H}_{68}\text{N}_4\text{O}_6$	628	Dark-green	Semi-solid
VIc	$\text{C}_{40}\text{H}_{80}\text{N}_4\text{O}_8$	744	Dark-green	Semi-solid
VIIa	$\text{C}_{28}\text{H}_{57}\text{N}_5\text{O}_3$	511	Brown	Semi-solid
VIIb	$\text{C}_{34}\text{H}_{69}\text{N}_5\text{O}_5$	627	Brown	Semi-solid
VIIc	$\text{C}_{40}\text{H}_{81}\text{N}_5\text{O}_7$	743	Brown	Semi-solid
VIIIa	$\text{C}_{29}\text{H}_{55}\text{N}_5\text{O}_4$	537	White	Powder
VIIIb	$\text{C}_{35}\text{H}_{67}\text{N}_5\text{O}_6$	653	White	Powder
VIIIc	$\text{C}_{41}\text{H}_{79}\text{N}_5\text{O}_8$	769	White	Powder
IXa	$\text{C}_{30}\text{H}_{63}\text{N}_5\text{O}_4$	557	Yellow	Semi-solid
IXb	$\text{C}_{36}\text{H}_{75}\text{N}_5\text{O}_6$	673	Yellow	Semi-solid
IXc	$\text{C}_{42}\text{H}_{87}\text{N}_5\text{O}_8$	789	Yellow	Semi-solid

surfactant molecules dissolve in water, they orientate themselves in-between the water molecules and weaken the hydrogen bond between them. This in turn decreases the holding forces and lowers the surface and interfacial tension. The extent by which the surfactant decreases the surface tension depends on two competing forces. These are the repulsion forces between water molecules and the hydrophobic part of the surfactant; and the attraction forces between water molecules and the hydrophilic part of the surfactant molecules. In the case at hand, the hydrophobic part (heptadecyl side chain) is constant, while the hydrophilic part changes as the number of the propylene oxide units changes. Consequently, for a certain molecule, at a certain concentration, while the abovementioned repulsive force is unchanged, increasing the number of propylene oxide units increases the attractive forces. As a consequence and in accordance with the previously reported results [46], the surface and interfacial tensions of the solution increases by increasing the number of propylene oxide units (n) included in the molecule, Table 3a. In addition, comparing structurally related surfactants **I(a–c)** and **V(a–c)**–**VI I(a–c)** indicated that, at a certain (n) thiadiazole derivative **V(a–c)** is slightly more effective in decreasing the surface tension than oxadiazole **I(a–c)** and triazole derivatives **VII(a–c)** and **VI I(a–c)**. Within the oxadiazole derivatives **I(a–c)**–**IV(a–c)**, surfactants **III(a–c)** induce the highest reduction in surface and interfacial tensions.

Cloud Point

For nonionic surfactants, a common and characteristic observation is that they exhibit a reverse solubility versus temperature behavior in water, consequently their solutions tend to become visibly turbid at a well-defined temperature. This is known as the cloud point, above which the surfactant solution phase separates into two phases. The cloud point is a critical factor in the performance of nonionic surfactants; generally, nonionic surfactants show optimal effectiveness when used near or below their cloud point. In addition, knowing the cloud point helps us to determine the storage stability since storing at temperatures significantly higher than the cloud point may result in phase separation and instability. The cloud point depends on the chemical structure [47, 48]. For a given hydrophobic group the cloud point is found to increase with increasing the length of the hydrophilic moiety [49, 50]. As outlined in Table 3a, the cloud point for the synthesized surfactants increases by increasing the number of propylene oxide groups.

Wetting Time

The degree of wetting is the resultant of two competing forces. These are, adhesive forces between a liquid and

Table 3 Surface properties of some synthesized surfactants

Compd.	No. of moles ^a	Surface tension (mN/m) 0.1 wt%	Interfacial tension (mN/m) 0.1 wt%	Cloud point (°C) 1.0 wt%	Wetting time (s) 0.1 wt%	Foam height (mm) 1.0 wt%	Emulsion stability (min) 20 mol
(a)							
I(a-c)	3	33	10	86	52	70	45
	5	35	13	93	47	80	43
	7	38	15	>100	42	100	40
II(a-c)	3	34	9	82	50	50	44
	5	36	12	98	47	70	42
	7	39	13	>100	48	90	38
III(a-c)	3	29	8	81	44	90	49
	5	30	10	95	39	100	46
	7	33	13	>100	38	120	42
IV(a-c)	3	31	11	86	53	100	52
	5	33	14	94	48	120	49
	7	36	16	>100	45	150	44
V(a-c)	3	31	10	85	50	60	44
	5	32	12	90	46	80	42
	7	34	14	>100	40	100	38
VI(a-c)	3	34	12	86	54	90	42
	5	36	14	95	50	110	38
	7	37	16	>100	46	120	36
VII(a-c)	3	37	13	89	52	90	55
	5	41	15	97	48	100	52
	7	43	16	>100	45	120	46
VIII(a-c)	3	30	11	88	50	100	52
	5	31	13	96	48	120	49
	7	33	15	>100	46	150	44
IX(a-c)	3	33	10	82	43	90	46
	5	34	12	91	40	110	43
	7	36	13	>100	38	120	40
Compd.	No. of moles ^a	CMC (mmol/L)	γ_{CMC} (mN/m)	π_{CMC} mN/m	PC ₂₀ (mmol/L)	Γ_{max} (mol/cm ²)	A_{min} (Å ² /mol)
(b)							
I(a-c)	3	5.0	34	38	2.30	0.99	1.66
	5	7.8	37	35	2.11	1.89	0.88
	7	10.0	39	33	2.00	2.27	0.73
II(a-c)	3	10.0	34.5	37.5	2.00	0.95	1.73
	5	12.0	37.5	34.5	1.92	1.82	0.91
	7	19.9	39	33	1.70	2.19	0.75
III(a-c)	3	3.1	29	43	2.51	0.84	1.97
	5	3.9	31	41	2.41	1.62	1.02
	7	5.4	34	38	2.27	1.97	0.84
IV(a-c)	3	0.64	32	40	3.19	0.98	1.65
	5	0.95	35.5	36.5	3.02	1.88	0.87
	7	1.9	38	34	2.72	2.26	0.72
V(a-c)	3	4.0	31	41	2.40	0.80	2.06
	5	5.6	33	39	2.25	1.55	1.07
	7	9.0	35	37	2.05	1.89	0.87
VI(a-c)	3	2.0	35	37	2.70	0.97	1.70
	5	4.5	36.5	35.5	2.35	1.85	0.89
	7	11.0	39	33	1.96	2.23	0.74
VII(a-c)	3	14.0	38	34	1.85	1.14	1.44
	5	59.0	42	30	1.23	2.15	0.77
	7	81.0	45	27	1.09	2.57	0.64

Table 3 continued

Compd.	No. of moles ^a	CMC (mmol/L)	γ_{CMC} (mN/m)	π_{CMC} mN/m	PC ₂₀ (mmol/L)	Γ_{max} (mol/cm ²)	A_{min} (Å ² /mol)
VIII(a-c)	3	3.8	30	42	2.42	0.79	2.05
	5	5.1	33	39	2.29	1.59	1.09
	7	6.0	34	38	2.22	1.89	0.87
IX(a-c)	3	5.3	34	38	2.28	0.98	1.68
	5	7.7	35	37	2.11	1.87	0.88
	7	10.7	38	34	1.97	2.25	0.73

^a Number of propylene oxide units

solid that causes a liquid drop to spread across the surface; and cohesive forces within the liquid which causes the drop to ball up and avoid contact with the surface. Addition of surfactants to water lowers the surface and interfacial tension; this in turn lowers the contact angle between the

liquid–vapor interface and the solid–liquid interface; resulting in more surface wetting. Consequently, surfactants are commonly used as wetting agents. In the present work, the efficiency of the synthesized surfactants as wetting agents was measured in terms of wetting time

Table 4 Resistance of the synthesized surfactants towards acidic and alkaline hydrolysis

Compd	Surface tension (mN/m) 0.1 wt	Surface tension of 0.1 % surfactant (stability to hydrolysis)					
		H ₂ SO ₄ (5 %) 25 °C	After boiling (5 %)H ₂ SO ₄		NaOH (1 %) 25 °C	After boiling (1 %NaOH)	
			30 min	60 min		30 min	60 min
Ia	33	34	34	36	33	34	34
Ib	35	36	37	40	36	36	37
Ic	38	38	39	40	38	38	38
IIa	34	35	36	38	34	35	35
IIb	36	37	37	38	37	37	37
IIc	39	39	39	42	39	40	40
IIIa	29	30	31	34	29	30	30
IIIb	30	31	33	34	30	30	31
IIIc	33	34	34	36	33	33	34
IVa	31	32	33	34	31	31	31
IVb	33	35	36	36	33	34	34
IVc	36	37	38	40	37	37	38
Va	31	33	33	36	32	32	33
Vb	32	34	34	35	32	33	33
Vc	34	36	36	38	34	35	35
VIa	34	35	35	37	34	34	35
VIb	36	36	37	40	37	37	37
VIc	37	39	40	41	38	38	39
VIIa	37	39	40	43	37	38	38
VIIb	41	41	43	44	41	41	42
VIIc	43	44	46	46	43	44	44
VIIIa	30	31	33	34	30	30	31
VIIIb	31	32	32	34	32	33	33
VIIIc	33	33	34	37	34	35	35
IXa	33	34	36	37	33	33	34
IXb	34	35	36	39	34	35	35
IXc	36	38	41	42	37	37	37

according to Draves technique [28], which measures the ability of a surfactant solution to displace air from a weighted skein of cotton by spreading wetting. Shorter wetting times indicate more efficient wetting. Surfactants **I(a–c)** to **IX(a–c)** demonstrate varying wetting abilities depending on the number of propylene oxide units (n) (Table 3a). For a given surfactant, increasing (n) increases the adhesive forces relative to the cohesive forces [51] with the concomitant decrease of wetting time. Surfactants **IIIc** and **IXc** exhibited the shortest sinking time, consequently, are the most efficient wetting agents among the studied group. This may be attributed to the increase in the adhesive forces due to the presence of chlorine atom and ethoxy group in **III** and **IX**, respectively.

Foaming Properties

Foam consists of gas covered with thin liquid film. It is thermodynamically unstable and tends to collapse to liquid which is the lowest energy state. In solution, the presence of a surfactant lowers the surface tension and eases foam generation with simultaneous adsorption of the surfactant molecules onto the interface between gas and liquid. The adsorption of surfactant molecules at the foam walls opposes their collapse. The foamability of the synthesized surfactant was examined using the Ross Miles method [29]. As indicated in Table 3a, and in agreement with previous reports [51–53], the foam height of the synthesized surfactants increases by increasing the propylene oxide units per molecule of surfactant. These results suggest that increasing the adhesive forces—brought upon by increasing (n)- draws more surfactant to the interface which in turn enhances the formation of foam.

Emulsion Stability

Emulsification is an important property of surfactants that broadens their applications. For instance, tiny fragments of oil suspended in pure water will spontaneously assemble themselves into much larger masses. In the presence of an appropriate surfactant, the interfacial tension between the two immiscible liquids is reduced to a sufficiently low value, thus, emulsification takes place and permits stability of minute droplets of oil in the bulk of water (or vice versa). The emulsifying power of the prepared surfactants in terms of time needed for the separation of 9 mL of the solution is presented in Table 3a. In accordance with the interfacial tension measurements (*vide supra*), the data show that, the stability of the formed emulsion decreases as the number of propylene oxide units increases. Obviously, the solubility of the surfactant in the oil phase decreases as the hydrophilic part (n) increases, resulting in weaker

emulsion [53, 54]. Generally, the measured time ranged between 36 and 55 min, thus, indicating moderate emulsifying properties for the synthesized surfactants.

CMC Measurements

The CMC measures the efficiency of surfactants, it indicates the amount of surfactant required to reach maximum surface tension reduction. A surfactant having a low CMC value enjoys excellent wetting, emulsifying, solubilizing and detergency properties. Within a given series of structurally related surfactants, factors that lead to a decrease in the surface tension are expected to decrease the CMC value. As outlined in Table 3a, b, the surface tension, CMC values and the surface tension at the critical micelle concentration (γ_{CMC}) measured for **I(a–c)**–**IX(a–c)** increase as the propylene oxide units increases. Within each series of surfactants, noting that the hydrophobic part is unchanged, the smallest CMC value is reached where $n = 3$, i.e., when the adhesive force is at its least (Graphs of surface tension vs log of concentration of added surfactant, used in the determination of the CMC are presented in the supplementary material section). Previous reports agree with our results [55].

Effectiveness (π_{CMC})

The effectiveness of a surfactant (π) is measured by its ability to induce the maximum reduction in the surface tension. Since the CMC point presents the minimum surfactant concentration needed to produce the maximum reduction in surface tension; consequently, the effectiveness of a surfactant (π_{CMC}) can be measured from the decrease in the surface tension of pure water (γ_0) that is induced by this surfactant at the CMC [32]. Logically, effectiveness of the prepared surfactants decreases as (γ_{CMC}) increases, Table 3b.

Efficiency (PC_{20})

In solution, surfactants lower the surface tension due to their adsorption in between water molecules. Thus, the performance of a surfactant can also be discussed in terms of its efficiency of adsorption. This is usually defined as the surfactant adsorption efficiency, which is the concentration of a surfactant required to produce a 20 mN/m reduction in surface tension. It is designated by PC_{20} and can be calculated by Eq. 2 (*vide supra*) [33]. The larger PC_{20} the more efficiently the surfactant is adsorbed at the interface and the more efficiently it reduces surface tension. In accordance of the above mentioned results, values of PC_{20} for the prepared

surfactants are found to decrease by increasing the number of propylene oxide units (cf Table 3a, b).

Maximum Surface Excess Γ_{max}

The extent of surfactant adsorption at a liquid surface is expressed in terms of its surface excess concentration, Γ . This is defined as the excess of surfactant present per unit area of surface over what would be present if uniform surfactant concentration existed right up to the surface. Surface excess concentration Γ_{max} , is related to surface tension by the Gibbs equation, Eq. 3 [33] and to its maximum surface concentration CMC, and efficiency by the following equation (Eq. 6) by Rosen [56].

$$\pi_{CMC} = 20 + 2.303nRT\Gamma_{max} \log \left[\frac{C_{CMC}}{C_{\pi} - 20} \right] \quad (6)$$

The data obtained from applying Eq. 3 are presented in Table 3b. It indicated that the maximum surface excess

increased by increasing the number of propylene oxide units and ranged between 0.79 and 2.57 mol/cm².

Minimum Surface Area (A_{min})

The area per surfactant molecule A_{min} at the air/water interface at surface saturation provides information about the degree of packing and the orientation of the adsorbed surfactant molecule [35, 36]. The calculated average areas A_{min} are given in Table 3b. Within a series of surfactants, increasing the number of propylene oxide units results in a significant decrease in A_{min} . This indicates that higher order of packing is induced upon increasing (n).

Resistance to Hydrolysis

Surfactants are intensively used in the manufacture of detergents, consequently, the stability of a certain surfactant towards acids and bases is a crucial factor in its utilization. Table 4 presents the results obtained from testing the

Table 5 Biodegradability of the synthesized surfactants

Compd.	No. of moles ^a	1st Day	2nd day	3rd Day	4th Day	5th day	6th day	7th day
I(a-c)	3	63	67	75	84	92	–	–
	5	55	64	69	80	88	91	–
	7	47	58	66	77	45	94	–
II(a-c)	3	51	62	77	84	94	–	–
	5	46	54	68	71	84	91	–
	7	41	52	65	67	81	88	–
III(a-c)	3	49	58	67	75	82	87	–
	5	46	51	58	69	76	84	–
	7	41	52	56	67	72	83	91
IV(a-c)	3	53	69	74	83	97	–	–
	5	49	62	67	78	85	92	–
	7	46	55	57	66	76	89	–
V (a-c)	3	53	66	75	85	95	–	–
	5	48	60	68	78	91	–	–
	7	45	50	62	74	83	93	–
VI(a-c)	3	56	68	79	86	96	–	–
	5	53	62	69	85	94	–	–
	7	51	60	66	84	93	–	–
VII(a-c)	3	50	60	77	84	94	–	–
	5	46	54	68	71	84	95	–
	7	41	52	65	67	80	93	–
VIII(a-c)	3	51	62	78	86	93	–	–
	5	46	55	69	73	85	91	–
	7	42	54	66	69	80	90	–
IX(a-c)	3	54	61	73	81	90	–	–
	5	51	60	77	84	91	–	–
	7	47	58	66	78	85	92	–

^a Number of propylene oxide units

resistance of the synthesized surfactants towards acidic and alkaline hydrolysis. In an acidic medium, compounds **I(a–c)** to **IX(a–c)** show high stability when boiled for 30 min, but they are less stable upon boiling for 60 min. On the other hand, in an alkaline medium, the synthesized surfactants are only slightly affected even after boiling for 60 min. Therefore, we concluded that, it would be safe to use the synthesized surfactants in the manufacture of detergents.

Biodegradability of the Synthesized Surfactants

Biodegradation is the destruction of a chemical by the metabolic activity of microorganisms. Since surfactants are susceptible to biodegradation and in order to examine their effect on water pollution, the biodegradation of the synthesized nonionic surfactants was evaluated by the conventional River Die-Away test employing the surface-tension technique as an analytical tool [57, 58]. The biodegradability data are presented in Table 5. Within experimental accuracy, all the prepared nonionic surfactants seem to degrade easily. The results showed that, 40–50 % of the surfactants was biodegradable within the first day, further degradation occurred on the next days until it died away on the 6th day. Consequently, these compounds are safe for human beings and the environment. It should be noted that, since the studied compounds share the same hydrophobic part, then, the extent of biodegradation encountered for these compounds depends mainly on the number of propylene oxide units. Consequently, it can be concluded that the biodegradability of a certain surfactant decreases by increasing the number of propylene oxide units.

Conclusions

New oxadiazole, thiadiazole and triazole derivatives (**1–12**) have been successfully synthesized. All the synthesized compounds (**1–12**) exhibited moderate to high activity toward tested G^- , G^+ bacteria, while some of them (**6**, **7**, **10**, **11**) exhibited good antifungal activity. New classes of nonionic surface active agents containing heterocyclic moieties were synthesized via subjecting the synthesized compounds to different quantities of propylene oxide. All the new nonionic surfactants displayed good surface active properties that are affected by the size of the hydrophilic part. The lower the number of propylene oxide units, the more effective is the surfactant in decreasing the surface and interfacial tension. Consequently, the lower the values of CMC and Γ_{\max} are, the higher is the emulsion stability, effectiveness (π_{CMC}), efficiency (PC_{20}) and minimum surface area (A_{\min}). In addition, the synthesized surfactants showed good tolerance towards acidic and alkaline media and good degradation susceptibility within 7 days. As a

consequence, the synthesized compounds and surfactants are safe for human beings as well as the environment. They can be used in the manufacture of drugs, cosmetics, moderate emulsifiers, wetting agents, textiles and dyes.

References

- Soares de Oliveira C, Lira BF, Barbosa-Filho JM, Lorenzo JGF, Filgueiras de Athayde-Filho P (2012) Synthetic approaches and pharmacological activity of 1,3,4-Oxadiazoles: a review of the literature from 2000–2012. *Molecules* 17:10192–10231
- Chauviere G, Bouteille B, Enanga B, Croft SL, Dumas M, Perie J, de Albuquerque C (2003) Synthesis and biological activity of nitro heterocycles analogous to megazol, a trypanocidal lead. *J Med Chem* 46:427–440
- Mazzone G, Bonina F, Pulglisi G, Panico AM, Arrigo RR (1984) 2,5-Diaryl-substituted 1,3,4-oxadiazoles: synthesis and preliminary pharmacological investigation. *Farmaco Ed Sci* 39:414–420
- Raman K, Parmar SS, Salzman SK (1989) Anti-inflammatory activity of substituted 1,3,4-oxadiazoles. *J Pharm Sci* 78:999–1002
- Ergenc N, Buyuktimkin S, Capan G, Baktir G, Rollas S (1991) Quinazolinones. 19. Communication [1]: synthesis and evaluation of some CNS depressant properties of 3-(2-[(5-aryl-1,3,4-oxadiazole-2-yl)amino]acetamido)-2-methyl-4(3H)-quinazolinones. *Pharmazie* 46:290–291
- Musser JH, Richard EB, Bernard L, Kevin B, Howard J, Robert K, Fuchih H, Atul K, Mitchell L (1984) Synthesis of 2-(2,3-dihydro-2-oxo-1,3,4-oxadiazol-5-yl)benzo heterocycles. A novel series of orally active antiallergic agents. *J Med Chem* 27:121–125
- Rostoma SAF, Shalaby MA, El-Demellawy MA (2003) Poly-substituted pyrazoles, part 5. Synthesis of new 1-(4-chlorophenyl)-4-hydroxy-1H-pyrazole-3-carboxylic acid hydrazide analogs and some derived ring systems. A novel class of potential antitumor and anti-HCV agents. *Eur J Med Chem* 38:959–974
- Suwiński J, Szczepankiewicz W (2008) Five-membered rings: triazoles, oxadiazoles, thiadiazoles and their fused carbocyclic derivatives. In: Katritzky AR (ed) *Comprehensive heterocyclic chemistry III*, vol 5. Elsevier Ltd, Oxford, pp 397–466
- Banerjee R (2005) Novel surfactant therapy for developing countries: current status and future directions. *Lung Biology Health Disease* 201:449–470
- Zhang X, Jin Z, Wang R, Tao Z, Zhong-fan L (2004) Surfactant-directed polypyrrole/CNT nanocables: synthesis, characterization, and enhanced electrical properties. *Chem Phys Chem* 5(7):998–1002
- Lakhrissi L, Ejjiyar S, Massoui M, Comelles F, Garcia MT, Ribosa I, Azemar N, Solans C (2002) Surface and self-aggregation properties of bis-benzimidazolones derivatives of D-glucose. *Comunicaciones Jornadas Comité Espanol Detergencia* 32:365–373
- Pegiadou S, Varka EM, Coutouli-Argyropoulou E (2004) Histidine as hydrophile in a new homologous series of glycerol ether surfactants. Synthesis, characterization, surface properties. *Comunicaciones Jornadas Comité Espanol Detergencia* 34:311–320
- Amine MS, Eissa AMF, Elsayy AA, Shaaban AF, El sayed R (2004) New heterocycles having a double characters; as antimicrobial and surface active agents. Part 3: nonionic compounds from fatty acid hydrazide. *OLAJ, Szappan Kozmetika* 53:124–128
- Amine MS, Aly AA, El-Sayed R (2006) Synthesis and surface active properties of condensed and non-condensed quinazoline

- derivatives of industrial application. *Indian J Chem* 45B:1020–1027
15. Amine MS, Eissa AMF, Elsayy AA, Shaaban AF, El sayed R (2004) New heterocycles having a double characters as antimicrobial and surface active agents. Part1: nonionic compounds from fatty acid isothiocyanate. *Grasas Aceites* 55:370–376
 16. Amine MS, Mahmoud AA, Badr SK, Gouda AS (2011) Fatty acids in heterocyclic synthesis. Part XIII: 2-chloro-5-heptadecyl[1,3,4]thiadiazole as a building block for synthesis of S-triazolo[2,4-b]thiadiazole, benzimidazo[2, 1-b]thiadiazole, thiadiazolo[2,3-b]quinazolin-5-one as well as thiadiazol-2-yl (piperidine, pyridazine and/or phthalazine). *OLAJ Szappan, Kozmetika* 60:51–57
 17. Amine MS, Mahmoud AA, Badr SK, Gouda AS (2012) Fatty acids in heterocyclic synthesis part xiii: synthesis of surfactants from pyrazole, isoxazole, pyrimidine and triazine, incorporating the 1,3,4-thiadiazole moiety having dyeing and antimicrobial activities. *J Surf Deterg* 15:179–190
 18. Bauer AW, Kirby WM, Sherris C, Turck M (1966) Antibiotic susceptibility testing by a standardized single disk method. *Am J Clin Pathol* 45:493–496
 19. Pfaller MA, Burmeister L, Bartlett MA, Rinaldi MG (1988) Multicenter evaluation of four methods of yeast inoculum preparation. *J Clin Microbiol* 26:1437–1441
 20. Jorgensen JH, Turnidge JD (2007) Susceptibility test methods: dilution and disk diffusion methods. In: Murray PR, Baron EJ, Jorgensen JH, Landry ML, Pfaller MA (eds) *Manual of clinical microbiology*, 9th edn. ASM Press, Washington DC, pp 1152–1172
 21. Thornsberry C, Sweson JM, Baker CN, Mc Dougal LK, Stocker SA, Hill BC (1988) Methods for determining susceptibility of fastidious and unusual pathogens to select antimicrobial agents. *Diag Microbiol Infect Dis* 9:139–153
 22. Morgos J, Sallay P, Farkas L, Rusznak I (1983) A novel way for ethoxylation with basic catalyst. *J Am Oil Chem Soc* 60:1905–1907
 23. Ahmed MHM, Sallay P, Rusznak I, Farkas L (1996) Determination of the average ethoxylation degree. *Tenside Surf Deterg* 33:410–411
 24. Findlay A (1963) *Practical physical chemistry*, 6th edn. Longmans, London, pp 1040–1069
 25. Weil JK, Stirton AJ, Nunez-Ponzoa MV (1966) Ether alcohol sulfates. The effect of oxypropylation and oxybutylation on surface active properties. *J Am Oil Soc* 43:603–609
 26. ASTM D1331-01 (2001) Standard Test method for surface and interfacial tension of solutions of surface active agents
 27. Durham K (1961) Properties of detergent solutions-amphipathy and adsorption, vol 1. *Surf Activity Deterg*. MacMillan &Co. Ltd., London, pp 1–28
 28. Draves CZ, Clarkson R (1931) A new method for the evaluation of wetting agents. *J Am Dye Stuff Report* 20:201
 29. Ross J, Milles GD (1941) Apparatus for comparison of foaming properties of soaps and detergents. *Oil Soap* 18:99–102
 30. Scholnick F, Linfield WM (1977) Lactose-derived surfactants (III): fatty esters of oxyalkylated lactitol. *J Am Oil Chem Soc* 54:430–435
 31. Colwell CE, Rixon WE (1961) Consideration in the use of nonionic surface-active agents. *Am Dyest Re* 50:679–682
 32. Lu JR, Lee EM, Thomas RK, Penfold J, Flitsch SL (1993) Direct determination by neutron reflection of the structure of triethylene oxide monododecyl ether layers at the air/water interface. *Langmuir* 9:1352–1358
 33. Matsuoka K, Moroi Y (2003) Micellization of fluorinated amphiphiles. *Curr Opin Colloids Interface Sci* 8:227–234
 34. Zhigang Xu, Li Pengfei, Qi Weihong, Li Zongshi, Lubo Ch (2006) Effect of aromatic ring in the alkyl chain on surface properties of aryl alkyl surfactant solution. *J Surf Deterg* 9:245–248
 35. Karakashev SI, Nguyen AV, Miller JD (2008) Equilibrium adsorption of surfactants at the gas-liquid interface. *Adv Polymer Sci* 218:25–55
 36. Karakashev SI, Manev ED (2002) Effect of interactions between the adsorbed species on the properties of single and mixed surfactant monolayers at air/water interface. *J Colloid Interface Sci* 248:477–486
 37. Falbe J (1986) *Surfactants for consumer*, Chap 4. Springer Verlag, Heidelberg
 38. Amine MS, Ahmed MHM, Hebash KA, El-Sheikh AA (2007) New heterocycles having a double character as antimicrobial and surface active agents. Part 6: novel thiadiazole from stearic acid. *OLAJ, Szappan, Kozmetika* 56:57–61
 39. Tsuge O, Oe K, Tashiro M (1977) Photolysis of 1,3,4-oxadiazoles in alcohols. *Chem Lett* 6:1207–1210
 40. Hill J (1984) In: Katritzky AR, Rees CW (eds) *Comprehensive heterocyclic chemistry I*, vol 6. Pergamon, Oxford, p 427
 41. Padmavathi V, Thriveni P, Sudhakar RG, Deepti D (2008) Synthesis and antimicrobial activity of novel sulfone-linked bis heterocycles. *Eur J Med Chem* 43:917–924
 42. Kumar H, Javad SA, Khan A, Amir M (2008) 1,3,4-Oxadiazole/thiadiazole and 1,2,4-triazole derivatives of biphenyl-4-yloxy acetic acid: synthesis and preliminary evaluation of biological properties. *Eur J Med Chem* 43:2688–2698
 43. El-Emam AA, Al-Deeb OA, Al-Omara M, Lehmann J (2004) Synthesis, antimicrobial, and anti-HIV-1 activity of certain 5-(1-adamantyl)-2- substituted thio-1,3,4-oxadiazoles and 5-(1-adamantyl)-3-substituted aminomethyl-1,3,4-oxadiazoline-2-thiones. *Bioorg Med Chem* 12:5107–5113
 44. Shaoyong K, Fengyi L, Wang N, Yang Q, Qian X (2009) 1,3,4-Oxadiazoline derivatives as novel potential inhibitors targeting chitin biosynthesis: design, synthesis and biological evaluation. *Bioorg Med Chem* 19:332–335
 45. Miller P, Westra P (1998) How surfactants work. *Colorado State Univ. Coop. Ex., Prod. Crop Ser. No. 0.564*
 46. Chlebicki J, Sokołowski A (1991) Adsorption of n-alkylthiooligooxypropylene glycols at aqueous solution-air interface. *Colloids Surf* 56:251–261
 47. Chen ML, Wang ZW, Zhang GX, Gu J, Cun Z, Tao FM (2007) Studies on the cloud points of nonionic surfactants with QSPR. *Chem Res Chinese U* 23:715–719
 48. Mitchell DJ, Tiddy GJT, Waring L, Bostock T, McDonald MP (1983) Phase behaviour of polyoxyethylene surfactants with water. *J Chem Soc, Faraday Trans* 79:975–1000
 49. Schott HJ (1969) Hydrophile-lipophile balance and cloud points of nonionic surfactants. *J Pharm Sci* 58:1443–1449
 50. Ahmed MHM (2004) Preparation and surface active properties of novel succinic acid based surfactants. *OLAJ Szappan Kozmetika* 53:23–28
 51. Eissa AMF, El-Sayed R (2007) Synthesis and evaluation of α -hydroxy fatty acid-derived heterocyclic compounds with potential industrial interest. *Grasas Aceites* 58:20–28
 52. Kim TS, Kida T, Nakatsuji Y, Hirao T, Ikeda JI (1996) Surface active properties of novel cationic surfactants with two alkyl chains and two ammonia groups. *J Am Oil Chem Soc* 73:907–912
 53. El-Sukkary MMA, Syed NA, Aiad I, El-Azab WIM (2008) Synthesis and characterization of some alkyl polyglycosides surfactants. *J Surf Deterg* 11:129–137
 54. Takeshi H (1970) Studies of ester containing surfactant: preparation and properties of sodium sulfalkanoates. *Bull Chem Soc* 43:2236–2239
 55. Zhigang Xu, Li Pengfei, Qi Weihong, Li Zongshi, Lubo Ch (2006) Effect of aromatic ring in the alkyl chain on surface

properties of aryl alkyl surfactant solution. *J Surf Deterg* 9:245–248

56. Rosen MJ (2004) *Surfactants and interfacial phenomena*, 3rd edn. Wiley, New York
57. Eter ET, Richard RE, David A (1974) Biodegradable surfactants derived from corn starch. *J Am Oil Chem Soc* 51:486–494
58. Balson T, Felix MSB (1995) The biodegradability of non-ionic surfactants. *Biodegradability of surfactants*. Springer, The Netherlands, pp 204–230

Author Biographies

Iman A. Gad El-Karim received her Ph.D. in organic chemistry from the Imperial College of Science and Technology—London University. Currently, she is an assistant professor of organic chemistry at the Faculty of Science, Benha University, Egypt. Her research interests relate to the synthesis of novel organic compounds having biological activities, novel surfactants and theoretical organic chemistry.

Mahasen S. Amine graduated from Ain Shams University, Egypt in 1973. She received her M.Sc. and Ph.D. degrees in 1983 and 1988 respectively from Ain Shams University. Since 1998 she has held the

position of full professor in organic chemistry at the Faculty of Science, Benha University. Her research interests relate to the synthesis of heterocyclic compounds of biological interest and using fatty acids to synthesize heterocycles having surface and biological properties.

Amal A. Mahmoud is currently a professor of organic chemistry at the Faculty of Science at Benha University, Egypt. She received her M.Sc. and Ph.D. degrees from Zagazig University. Her research interests relate to organic synthesis and polymeric surfactants having heterocyclic moiety.

Alaa S. Gouda is an associate lecturer at the department of Chemistry in the Faculty of Science at Benha University, Egypt. He received his B.Sc. in 2006 and M.Sc. in 2010 from the same university. His research interests relate to the synthesis of heterocyclic compounds of biological interest and using fatty acids for synthesizing heterocycles having surface and biological activities.