



# Insulators and domains of gene expression

Tamer Ali<sup>1</sup>, Rainer Renkawitz and Marek Bartkuhn

The genomic organization into active and inactive chromatin domains imposes specific requirements for having domain boundaries to prohibit interference between the opposing activities of neighbouring domains. These boundaries provide an insulator function by binding architectural proteins that mediate long-range interactions. Among these, CTCF plays a prominent role in establishing chromatin loops (between pairs of CTCF binding sites) through recruiting cohesin. CTCF-mediated long-range interactions are integral for a multitude of topological features of interphase chromatin, such as the formation of topologically associated domains, domain insulation, enhancer blocking and even enhancer function.

## Address

Institute for Genetics, Justus-Liebig-University, Heinrich-Buff-Ring 58, D35392 Giessen, Germany

Corresponding author: Renkawitz, Rainer ([rainer.renkawitz@gen.bio.uni-giessen.de](mailto:rainer.renkawitz@gen.bio.uni-giessen.de))

<sup>1</sup> Permanent address: Faculty of Science, Benha University, Benha, Egypt.

**Current Opinion in Genetics & Development** 2016, **37**:17–26

This review comes from a themed issue on **Genome architecture and expression**

Edited by **Frederic Berger** and **Pamela Geyer**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 20th January 2016

<http://dx.doi.org/10.1016/j.gde.2015.11.009>

0959-437X/© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

The concept of inactive and active chromatin domains was suggested quite early on as a way to interpret compact and less dense chromatin packaging in diploid interphase nuclei or in polytene chromosomes. The existence of such domains specifically requires the presence of domain boundaries to insulate the opposite activities of neighbouring domains. Such shielding elements, known as insulators, have been functionally identified by a position-independent high-level expression of a transgene in mice and flies [1,2]. In contrast to this barrier effect of an insulator, another shielding activity was called enhancer blocking [3], since it interferes with the action of an enhancer on a specific promoter when the insulator is positioned between the two.

Following the discovery of several *Drosophila*-specific insulator binding proteins (IBPs), such as BEAF32 [4], Su(Hw) [5,6] and Zw5 [7], the vertebrate factor CTCF [8,9] was shown to mediate insulation [10]. Later, the high conservation of chromatin insulation was demonstrated

by the identification of CTCF in *Drosophila* (dCTCF) [11–13] and by comparing shared features (Table 1). Here, we summarize recent results on the genome-wide binding of these and more recently discovered insulator factors, and the projection of these binding sites onto the three-dimensional chromatin structure. These observations and results from high throughput analyses and functional tests are discussed with respect to a unifying mechanism for insulator-mediated barrier function and enhancer blocking activity.

## CTCF: inhibitor and facilitator of enhancer function

Enhancer blocking activity of an insulator depends on its arrangement, that is, it has to be situated between the enhancer and promoter. This fact alone implies that the enhancer blocking activity is achieved by interfering with the chromatin looping required for enhancer/promoter contact. Detailed analysis of three-dimensional looping and the role played by the insulator protein CTCF revealed that CTCF not only possesses interference (enhancer blocking) activity, but also additionally mediates chromatin contacts or loops required for enhancer function. Examples for such bivalent consequences of loop formation are discussed below.

Bioinformatics evaluation of genome-wide chromatin interaction data led to the construction of a genome-wide interaction map of regulatory elements, which indicated that enhancer–promoter interactions are highly cell-type specific. Key interacting components are CTCF and cohesin [14]. This is exemplified by the MHC-II locus, which is active in B cells and bound by CTCF at 15 sites. In plasmablasts, this locus is inactive and only one third of the CTCF sites are bound. This correlates with the finding that CTCF is required for the cell type specific three-dimensional architecture of the locus and for maximal MHC-II gene expression in B cells [15\*].

Another example is where CTCF/cohesin organizes a loop pattern that includes the promoter of the PTGS2 gene such that the PTGS2 gene is activated. In cancer cells the CpG island at the PTGS2 promoter is methylated and the gene is turned off. This silencing mechanism is in part caused by the methylation-induced loss of CTCF binding, which results in a change in chromatin looping and abrogation of gene activity [16].

Regulation of dCTCF binding in *Drosophila* development is seen at the homeotic gene *Ultrabithorax* (*Ubx*), which is activated by *Ubx* enhancer elements in the third thoracic leg imaginal disc. Here, a dCTCF site at the enhancer generates a loop with the gene promoter. In inactive tissues

Table 1

Insulator components with conserved features in vertebrates and *Drosophila*.

Factor	Organism	Description	References
CTCF	<i>H. sap.</i>	Enhancer blocking activity of the chicken beta-globin insulator	[10]
dCTCF	<i>D. mel.</i>	Enhancer blocking of Fab-8 insulator	[13]
GAGA	<i>D. mel.</i>	Enhancer blocking of the eve promoter	[77]
Th-POK	<i>M. mus.</i>	Binding to enhancer-blocking elements in murine Hox clusters	[78]
Cohesin	<i>H. sap.</i>	Cohesin is required for enhancer blocking of the H19 ICR	[79]
	<i>D. mel.</i>	Enriched at TAD borders	[54**]
TFIIIC	<i>H. sap.</i>	Loss of binding to tDNA promoters reduces their enhancer blocking activity	[80]
	<i>D. mel.</i>	Binding to borders of topological domains (ChIP-seq)	[54**]
Condensin	<i>M. mus.</i>	Binding correlates with enhancer blocking capacity of TAD borders	[54**]
	<i>D. mel.</i>	Enriched at TAD borders	[54**]
Rm62	<i>D. mel.</i>	Interacts with CP190 and mutations affect gypsy-mediated insulation in ct and y2-loci	[81]
p68	<i>H. sap.</i>	Along with SRA required for CTCF to perform proper insulation	[82]
PARP1	<i>D. mel.</i>	Modifies insulator functions	[83]
	<i>H. sap.</i>	Prevents DNA methylation of CTCF target sites. Controls circadian transcription	[84,85**]
dMes-4	<i>D. mel.</i>	BEAF-32 co-factor, involved in gene regulation	[86]
PRDM5	<i>M. mus.</i>	Interacts and overlaps with CTCF and Cohesin; recruits G9a (HMT)	[87]
Nurf-301	<i>D. mel.</i>	Regulates Fab-8 enhancer blocking activity	[72*]
Bptf	<i>H. sap.</i>	Interacts with CTCF; regulates nucleosomal arrays around CTS	[74]
TGF- $\beta$ signal-ling	<i>D. mel.</i>	Genome-wide overlap with and dependency, to some extent, on dCTCF	[88]
	<i>H. sap.</i>	CTCF physically interacts with Smad3 and recruits Smad to H19 ICR	[89]
CP190	<i>D. mel.</i>	enhancer-blocking activity, mediates long-range interactions	[25,37]
Kaiso	<i>H. sap.</i>	Similar to BTB domain of CP190; mediates enhancer-blocking activity; physically interacts with CTCF	[90]

this dCTCF site is not occupied and enhancer/promoter interaction is lost [17], clearly demonstrating that this dCTCF site is a facilitator of enhancer action.

Not only can enhancer/promoter interactions be facilitated by CTCF/cohesin, but also other 3D-interactions may depend on CTCF sites. For example, CTCF-dependent enhancer/enhancer clustering in the nucleus was observed in thymocytes. Targeted 3C analysis demonstrated that interactions between the *Cd3* super-enhancers as well as with other enhancers were significantly weakened in cohesin-deficient thymocytes [18\*]. Furthermore, CTCF/cohesin dependent inter-chromosomal contacts control enhancer inhibition in case of the Sox-2 and Sox-17 genes [19\*\*]. Similarly, enhancer inhibition and facilitation was observed in erythroid cells, where together with other factors, CTCF bound to several sites mediates an intra-chromosomal interaction on chromosome 1 between the TAL1 promoter and its downstream enhancer, allowing for regulated TAL1-expression. However, in T-cell acute lymphoblastic leukemia, these interactions are altered, resulting in aberrant expression of the TAL1 oncogene [20\*\*].


### Barrier function and topologically associated domains

The identification of chromatin domains with either active histone marks or silencing modifications led to

the concept of barriers that prevent one domain interfering with the neighbouring one. Loss of barrier function has often been related to inactive marks spreading into the active domain [21–29], although activation of inactive domains is possible as well (see below). Initial analyses revealed CTCF binding and loop formation at barrier sites flanked by opposite chromatin states [30–32]. In *Drosophila*, depletion of dCTCF results in a small change in H3K27me3 spreading [33], when testing genome-wide effects. Additional architectural proteins are also present at chromatin barriers and may compensate for the loss of CTCF-dependent barrier functions (see below). Therefore, only a few barrier sites, which are primarily dependent on CTCF, showed an expansion of the H3K27me3 mark into the flanking region [34–38]. As discussed below, the role, played by insulators and CTCF in barrier function, is further supported by the analysis of homeotic genes, and illustrated by the concept of topologically associated domains [39] (TADs) (Figure 1).

Homeotic genes are expressed during development in a cell type, and stage specific manner. The collinear genomic arrangement and expression of the gene clusters specify the segmental identities along the body axis of *Drosophila* and mammals. Thus, in a given cell type or specific developmental stage one group of Hox genes may be turned off by Polycomb function, resulting in H3K27me3 modification of the respective gene locus.

Figure 1



Insulators, chromatin domains and topologically associated domains (TADs). Interaction matrix representing a virtual Hi-C experiment (top). The grey scale above indicates interaction frequencies. Interactions occur predominantly within TADs (e.g. enhancer–promoter interactions), which are often grouped in subdomains. Interactions between TAD boundaries are thought to depend on the binding of CTCF (shown as a schematic ChIP-seq track in red) to its cognate DNA-binding motif (black arrows). CTCF sites not involved in binding to TAD boundaries are shown in pale red and grey motifs, respectively. Motifs involved in long-range chromatin interactions show an inverted repeat orientation (see Figure 2). As not all TAD boundaries are bound by CTCF it is likely that additional factors may be involved in their function (indicated by question mark). TADs are often coincident with chromatin domains represented by a schematic ChIP-seq track for an active (H3K36me3; green) and a repressive (H3K27me3; blue) histone modification. Active TADs are gene-rich (black bars for active genes) in contrast to gene-poor repressed domains (grey bars).

In both mammals and *Drosophila*, *Hox* gene clusters are marked by CTCF/dCTCF binding at the borders between individual regulatory elements of the *Hox* genes [12,40–42]. In *Drosophila*, developmental expression is specific for each parasegment. Chromatin purified from single parasegments revealed a ‘step-wise’ pattern of acetylated H3K27 (active gene domain) or of H3K27me3 (inactive gene domain) with sharp, dCTCF-bound boundaries at the bithorax complex (BX-C) regulatory domains [43\*\*]. This suggests that functional boundaries associated with dCTCF binding restrict H3K27me3 or H3K27 acetylation to one domain, preventing spreading into the neighbouring domain.

A similar situation is found with the mouse and human *HoxA* genes. Kinetic analyses of myelomonocytes differentiating into monocytes/macrophages revealed a dynamic change in *HoxA* cluster topology [44]. *HoxA* expression in ES cells is silenced by H3K27me3, whereas differentiation into neuronal cells is marked by activation of the

rostral group of the *HoxA* cluster, while the caudal group of genes remain silenced [45,46]. Again, the pattern of gene activity is associated with a ‘step-wise’ pattern of H3K27 modification [47\*\*] with H3K27me3 enriched at silent genes. To test the requirement for CTCF at functional boundaries, the CRISPR/Cas technique was used to delete a CTCF binding site separating the active gene groups from the repressed genes within the *HoxA* cluster. CTCF loss at these sites resulted in spreading of H3K4 methylation, an active chromatin modification, into the repressed region, thereby activating a caudal *Hox* gene [46,47\*\*]. This clearly shows that CTCF acts as a barrier, in this case for active marks spreading into a silenced region.

Analysis of *Hox* genes suggests that barrier function is linked with TAD organization (Figure 1). In wildtype, *Hox* gene expression and H3K27 modification correlate with two TADs in motor neurons. Deletion of the CTCF site at the boundary not only removes the barrier, but also

shifts the TAD boundary further into the caudal TAD, up to the next CTCF site [47\*\*]. Upon removing this site as well, the barrier and TAD boundary shifted even further into the caudal gene region. Thus, both barrier function and TAD boundary function are controlled by CTCF and are probably two features of the same phenomenon.

How do these structural units relate to genomic functions such as control of transcription? In one study, ChIA-PET was used to generate a map of enhancer–promoter interactions in ES cells. Genes controlled by super-enhancers were found to reside within a super-enhancer domain structure with the flanking, protein-bound CTCF/cohesin sites forming a loop. Consequently, these loops generate insulated neighbourhoods that are preserved in multiple cell types. Similarly, Polycomb repressed genes are organized in insulated neighbourhoods flanked by CTCF/cohesin, thereby forming a Polycomb domain [48\*\*].

Another strategy to identify functional domains was to insert regulatory sensor transposons into hundreds of sites within the mouse genome. The enhancers identified in this screen acted along broad regions that correlated strongly with TADs [49\*\*]. This suggests that three-dimensional enhancer action is restricted to the genomic region defined by TADs, and therefore the functional domain structure concurs with the topological features identified by 3D mapping.

Which features of TAD domain boundaries are required for TAD formation? There is good evidence that interactions within TADs contribute to boundary function [50]. In addition, TAD domain boundaries are strongly enriched for CTCF/Cohesin binding. The importance of CTCF/Cohesin for the structural and functional integrity of TADs has been documented in several cases. The consequence of CTCF loss for the 3D structure of chromatin was tested by depleting CTCF [51,52\*]. Some changes were observed, such as a mild reduction in intra-domain interactions as well as a gain in inter-domain interactions. Nevertheless, the overall organization and long distance interaction remained. This, and the fact that many more CTCF sites exist that are not located at TAD boundaries (Figure 1), argues for additional factors involved in the 3D landscape of chromatin. In fact, besides CTCF/dCTCF, many architectural factors have been found at TAD borders. These factors are cohesin components SMC3 and RAD21, TFIIC subunits, condensin subunits and PRDM5, a SET domain protein [53\*\*,54\*\*]. Furthermore, there is evidence that CTCF is an RNA binding protein and that RNA is involved in CTCF recruitment and long-range interaction [55]. Another *Drosophila* IBP, Su(Hw) has been shown to interact with RNA as well [56].


The functional importance of the TAD organization becomes evident when TAD borders are deleted. Human families with rare limb malformations show rearrangements

in the extended *WNT6/IHH/EPHA4/PAX3* region. Comparable rearrangements were generated in mice using the CRISPR/Cas technique [57\*\*]. These mutations resulted in disease-relevant changes in interactions between promoters and non-coding DNA as well in aberrant gene expression. Furthermore, these mice developed digital malformations similar to phenotypes observed in patients. These changes in chromatin interaction and function only occurred if the rearrangement disrupted a CTCF-associated TAD boundary [57\*\*].

The relevance of TAD organization is further underscored by its evolutionary conservation. A group of homeobox genes, called the *Six* cluster, is highly conserved from sea urchins and zebrafish to mice and humans. Similarly, the TAD organization is conserved with two largely independent regulatory landscapes contained within two adjacent TADs [58\*]. Interestingly, CTCF binding sites at the TAD borders are found in opposite orientations, also a highly conserved feature. CTCF sites divergent between species correlate with divergence of an internal domain structure. Comparing genomes and domain structures of mouse and dog revealed insertions, inversions and duplications. Interestingly, in each case the rearrangement occurred at the border between two TADs [59\*\*].

In *Drosophila*, functional tests revealed that inverted insulators form loops more efficiently than insulators in identical orientation [60]. Similarly, in vertebrates it became obvious that the direction of CTCF binding sites plays a major role in determining which combinations of CTCF binding sites are compatible for interacting and subsequently generating loops. First, the orientation of CTCF binding motifs is strongly conserved across evolution [59\*\*]. Second, genome-wide analysis revealed that 72% or 48% of the mouse or human TAD borders, respectively, contain a pair of convergent CTCF sites [53\*\*,58\*]. This suggests a functional role of pairing between CTCF bound TAD borders, and that the selection of sites involved in pairing may be driven by the orientation of the CTCF binding sequences. Indeed, CRISPR/Cas mediated inversion of one of the CTCF binding sites in the *Pcdh* and *beta-globin* gene clusters induced directional switching of genome topology or partial merging of neighbouring chromatin domains [61\*\*]. But how can the direction of CTCF binding motifs influence pairing between insulators often separated by several hundred kilobases of DNA? A hypothetical model includes the biophysical ability of CTCF to bend DNA by 90° [62]. This causes a structure with an orientation that may be more accessible to pairing with another CTCF molecule bound to an inverted binding site (Figure 2). Furthermore, such a three-dimensional arrangement may have sterical consequences for nucleosome formation and for binding of cohesin and additional factors. Physical modelling suggested a loop extrusion model explaining why loops tend not to overlap and why

Figure 2



The structure and orientation of the CTCF/DNA complex may guide pairing of TAD boundaries. A hypothetical model includes the biophysical ability of CTCF to bend DNA by 90° [62]. When DNA binding motifs are convergent, this may facilitate homodimeric CTCF interaction and formation of the bent conformation at the bottom of the loop. The DNA bend is found at the DNA spacer between zinc finger groups 1 to 7 and 8 to 11, which has been identified by DNase I hypersensitivity [91]. Physical modelling supports a loop extrusion model in the context of paired CTCF binding sites in convergent orientation [63\*\*].

the CTCF-binding motifs at pairs of loop anchors lie in the convergent orientation [63\*\*]. This model was nicely supported by genome editing altering CTCF-binding sites. In every case the convergent rule correctly predicted loop formation [63\*\*].

### The specific role of *Drosophila* CP190

As described above, the general functions of insulator factors are highly conserved between vertebrates and *Drosophila* (Table 1). Nevertheless, an insect specific factor crucial for insulator function is the centrosomal protein 190 (CP190). Although identified in the context of centrosomes [64], a functional role was found in insulation mediated by Su(Hw) [65]. Subsequently, other IBPs have been identified that also bind CP190, for example BEAF-32, GAF, Zw5 and dCTCF, which frequently co-localize with CP190 throughout the *Drosophila* genome [33]. From these data it became evident that a class of potential


insulator sites was bound by CP190 in the absence of any known DNA binding factors.

Recent searches for additional, DNA binding and CP190 interacting factors identified insulator binding factors 1 and 2 (Ibf1, Ibf2) [66], a zinc finger protein interacting with CP190 (ZIPIC) and Pita [67\*]. All four factors mediate enhancer blocking of transgenes in *Drosophila*. Genome-wide binding was frequently found to be clustered with other IBPs and with TAD borders [54\*\*]. In addition to this correlation, many IBP binding sites are found within TADs and many TAD boundaries are not associated with IBPs (Figure 3). Mapping of the CP190 protein revealed separate interaction domains with Pita and ZIPIC [67\*]. This suggests that CP190 has a bridging function, simultaneously contacting several proteins. Such a feature was implicated when deletion of the CP190 interaction domain of BEAF-32 [68\*] resulted in BEAF-32 located at distant sites failing to interact with GAF or dCTCF bound promoters. The biophysical capacity of CP190/BEAF-32 to mediate long-range interactions *in vitro* further supports the bridging function of CP190 [69].

Assuming that CP190 is a bridging factor and that IBPs are frequently clustered [33,54\*\*,67\*,70], one can envision a concept of several IBPs targeting CP190 more efficiently (Figure 3). On the other hand, this multitude of clustered and CP190 interacting factors causes some kind of redundancy. This is evident from a dCTCF mutant lacking the CP190 interaction domain, which is still able to function similarly to wildtype dCTCF [71]. Based on synergistic recruitment one would expect CP190 binding to scale with IBP binding, which indeed could be shown [67\*]. Furthermore, insulator function and topological domain border strength both correlate with IBP protein occupancy [54\*\*].

The search for additional architectural proteins involved in insulator function revealed many more factors contributing to insulator strength, such as TFIIC, Rad21 (cohesin), Chromator, DREF, L(3)mbt and condensin factors CAP-H2 and Barren [54\*\*]. Occupancy and clustering of these factors to individual sites correlates with enhancer-blocking activity and TAD border strength. Thus, more architectural factors binding to insulators increase the insulator function. When re-analysing binding and Hi-C data from mouse and human ESCs and IMR90 fibroblasts, a similar conclusion could be drawn: mammalian TAD borders are enriched for the architectural factors CTCF, TFIIC, cohesin and condensin components and binding correlates with topological structure and regulatory potential [54\*\*]. Thus, a highly conserved molecular mechanism for TAD boundary function and insulation (Table 1) arises from the binding strength of factors connected by protein/protein interactions mediated by CP190, cohesin and condensin and possibly many others.

Figure 3



*Drosophila* CP190 recruitment and strength of TAD boundaries/insulators correlate with combinatorial binding of architectural proteins. **(a)** The interaction matrix represents TADs. Boundaries between TADs are often marked by CP190 binding (schematic ChIP-seq track, blue). CP190 is recruited to chromatin by a wide variety of insulator binding factors (IBPs, as exemplified by CTCF, BEAF32 and Pita in schematic ChIP-seq tracks). Frequently, different insulator binding factors cluster together, suggesting a cooperative recruitment mode for targeting CP190 to chromatin. Combinatorial recruitment of CP190 to TAD boundaries may be functionally important since high occupancy of IBPs and other architectural proteins such as cohesin, condensin and TFIIIC predict the strength of insulator function as well as TAD borders [54\*\*]. It should be noted that not all TAD boundaries are bound by known IBPs (?) and that many IBP binding sites are found within TADs. **(b)** The physical DNA string model summarizes the contact and binding data illustrated in (a).

In addition to the architectural and looping functions, an enzymatic activity in nucleosomal depletion was postulated due to the finding that dCTCF/CP190 binding sites show reduced nucleosomal occupancy, whereas dCTCF sites devoid or depleted of CP190 are loaded with nucleosomes [37]. A functional siRNA screen identified NURF and dREAM complexes binding to CP190 and being required for enhancer blocking [72\*,73]. Probably, the nucleosomal remodelling activity of ISWI, a component of NURF, causes nucleosomal depletion at CP190/dCTCF sites. Interestingly, a NURF and CTCF connection has also been found in vertebrates (Table 1) [74].

Testing chromatin conformation at a synthetic cluster of hundreds of binding sites for a LacI-CP190 fusion revealed a general opening and expansion of chromatin

in *Drosophila* cells [75]. A similar function was mediated by vertebrate CTCF in vertebrate cells [36,76]. Analysis of chromatin before and after CTCF recruitment revealed active removal of the H3K27me3 mark, likely by incorporating the H3.3 variant [36]. This variant is often associated with unstable nucleosomes and may explain that insulators are depleted of nucleosomes, and that flanking nucleosomes are free of the repressive histone mark H3K27me3.

When comparing vertebrate and *Drosophila* in respect to chromatin domains and insulation, many observations are comparable, as discussed above. Nevertheless, there are many more IBPs in *Drosophila*, not found in vertebrates as is CP190. Potentially, the demand for efficient, and maybe locus-specific insulation may be much higher in

case of the very compact *Drosophila* genome. This diversity of IBPs seems to be functionally merged by CP190.

## Conclusions and perspectives

Recent advances in determining the three-dimensional folding and interaction of chromatin at high resolution have highlighted the impact of higher-order chromatin structure on genome function. This is supported by the emerging concept of topologically associated domains separating the genome into conserved chromosomal neighbourhoods encompassing blocks of similarly regulated genomic regions. Architectural proteins, including CTCF, are the determinants for the strength of TAD formation and insulator function. The selection of interacting regions, in the case of CTCF, is dictated by the binding site orientation. It is obvious that CTCF and its orientation only partly account for the determinants selecting and mediating proper interactions. About 30 000 sites in the vertebrate genome are bound by CTCF, but only a fraction is found at TAD borders. What are the factors or combinations of factors determining the specificity of interacting elements? Furthermore, not all TAD borders have CTCF sites. Which factors or features are mediating the boundary function in these cases?

Despite the fact that many more insulator proteins are known in *Drosophila* than in vertebrates, the general features and many of the components are highly conserved (Table 1). Does this mean there are many more vertebrate factors involved in insulation that are yet to be found in vertebrates? And if so, will they help in solving the specificity problem?

In addition to long-range interaction and looping functions, characteristic chromatin modifications are found at insulators and are required for insulator activity. Furthermore, RNA molecules are involved in CTCF function. It remains to be seen whether these activities are fundamental to insulator function, or whether they support efficient binding of the architectural proteins, thereby maintaining long-range interactions.

## Acknowledgements

We apologize to those whose work could not be discussed due to space limitations. This work was funded by the DFG: TRR81 (<http://www.dfg.de>). TA is supported by a DAAD grant.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Grosveld F, van Assendelft GB, Greaves DR, Kollias G: **Position-independent, high-level expression of the human beta-globin gene in transgenic mice.** *Cell* 1987, **51**:975-985.
  2. Kellum R, Schedl P: **A position-effect assay for boundaries of higher order chromosomal domains.** *Cell* 1991, **64**:941-950.
  3. Kellum R, Schedl P: **A group of scs elements function as domain boundaries in an enhancer-blocking assay.** *Mol Cell Biol* 1992, **12**:2424-2431.
  4. Zhao K, Hart CM, Laemmli UK: **Visualization of chromosomal domains with boundary element-associated factor BEAF-32.** *Cell* 1995, **81**:879-889.
  5. Geyer PK, Corces VG: **DNA position-specific repression of transcription by a Drosophila zinc finger protein.** *Genes Dev* 1992, **6**:1865-1873.
  6. Holdridge C, Dorsett D: **Repression of hsp70 heat shock gene transcription by the suppressor of hairy-wing protein of Drosophila melanogaster.** *Mol Cell Biol* 1991, **11**:1894-1900.
  7. Gaszner M, Vazquez J, Schedl P: **The Zw5 protein, a component of the scs chromatin domain boundary, is able to block enhancer-promoter interaction.** *Genes Dev* 1999, **13**:2098-2107.
  8. Baniahmad A, Steiner C, Kohne AC, Renkawitz R: **Modular structure of a chicken lysozyme silencer: involvement of an unusual thyroid hormone receptor binding site.** *Cell* 1990, **61**:505-514.
  9. Lobanenkov VV, Nicolas RH, Adler VV, Paterson H, Klenova EM, Polotskaja AV, Goodwin GH: **A novel sequence-specific DNA binding protein which interacts with three regularly spaced direct repeats of the CCCTC-motif in the 5'-flanking sequence of the chicken c-myc gene.** *Oncogene* 1990, **5**:1743-1753.
  10. Bell AC, West AG, Felsenfeld G: **The protein CTCF is required for the enhancer blocking activity of vertebrate insulators.** *Cell* 1999, **98**:387-396.
  11. Gerasimova TI, Lei EP, Bushey AM, Corces VG: **Coordinated control of dCTCF and gypsy chromatin insulators in Drosophila.** *Mol Cell* 2007, **28**:761-772.
  12. Mohan M, Bartkuhn M, Herold M, Philippen A, Heini N, Bardenhagen I, Leers J, White RAH, Renkawitz-Pohl R, Saumweber H *et al.*: **The Drosophila insulator proteins CTCF and CP190 link enhancer blocking to body patterning.** *Embo J* 2007, **26**:4203-4214.
  13. Moon H, Filippova G, Loukinov D, Pugacheva E, Chen Q, Smith ST, Munhall A, Grewe B, Bartkuhn M, Arnold R *et al.*: **CTCF is conserved from Drosophila to humans and confers enhancer blocking of the Fab-8 insulator.** *Embo Rep* 2005, **6**:165-170.
  14. Heidari N, Phanstiel DH, He C, Grubert F, Jahanbani F, Kasowski M, Zhang MQ, Snyder MP: **Genome-wide map of regulatory interactions in the human genome.** *Genome Res* 2014, **24**:1905-1917.
  15. Majumder P, Schärer CD, Choi NM, Boss JM: **B cell differentiation is associated with reprogramming the CCCTC binding factor-dependent chromatin architecture of the murine MHC class II locus.** *J Immunol* 2014, **192**:3925-3935.
  - The MHC-II locus is marked by 15 CTCF binding sites. In B-cells, all sites are bound by CTCF, the locus is folded into a three-dimensional architecture and the genes show CTCF-dependent maximal expression. In non-expressing plasmablasts, only one third of the CTCF sites are significantly bound, correlating with a novel chromatin architecture.
  16. Kang JY, Song SH, Yun J, Jeon MS, Kim HP, Han SW, Kim TY: **Disruption of CTCF/cohesin-mediated high-order chromatin structures by DNA methylation downregulates PTGS2 expression.** *Oncogene* 2015, **34**:5677-5684.
  17. Magbanua JP, Runneburger E, Russell S, White R: **A variably occupied CTCF binding site in the ultrabithorax gene in the Drosophila bithorax complex.** *Mol Cell Biol* 2015, **35**:318-330.
  18. Ing-Simmons E, Seitan VC, Faure AJ, Flicek P, Carroll T, Dekker J, Fisher AG, Lenhard B, Merckenschlager M: **Spatial enhancer clustering and regulation of enhancer-proximal genes by cohesin.** *Genome Res* 2015, **25**:504-513.
  - Binding of CTCF and cohesin is highly enriched at enhancers, and in particular at enhancer arrays or 'super-enhancers'. Spatial enhancer clustering is facilitated by cohesin and primarily regulates genes in the vicinity of enhancer elements.
  19. Abboud N, Moore-Morris T, Hiriart E, Yang H, Bezerra H, Gualazzi MG, Stefanovic S, Guenantin AC, Evans SM, Puceat M: **A cohesin-OCT4 complex mediates Sox enhancers to prime an early embryonic lineage.** *Nat Commun* 2015, **6**:6749.

Cell fate switching from pluripotent ES cells to mesendoderm determination involves OCT4 binding to cohesin. This changes the higher-order chromatin structure at the Sox-2 and Sox-17 genes. These alterations involve intra-chromosomal and inter-chromosomal contacts, resulting in active enhancer/promoter contacts.

20. Patel B, Kang Y, Cui K, Litt M, Riberio MSJ, Deng C, Salz T, ●● Casada S, Fu X, Qiu Y *et al.*: **Aberrant TAL1 activation is mediated by an interchromosomal interaction in human T-cell acute lymphoblastic leukemia.** *Leukemia* 2014, **28**:349-361.
- In erythroid cells, CTCF mediates an intra-chromosomal interaction between the TAL1 promoter and its downstream enhancer. Three-dimensional chromatin interactions are changed in T-cell acute lymphoblastic leukemia (T-ALL), such that the TAL1 oncogene is activated.
21. Barkess G, West AG: **Chromatin insulator elements: establishing barriers to set heterochromatin boundaries.** *Epigenomics* 2012, **4**:67-80.
22. Gaszner M, Felsenfeld G: **Insulators: exploiting transcriptional and epigenetic mechanisms.** *Nat Rev Genet* 2006, **7**:703-713.
23. Herold M, Bartkuhn M, Renkawitz R: **CTCF: insights into insulator function during development.** *Development* 2012, **139**:1045-1057.
24. Matzat LH, Lei EP: **Surviving an identity crisis: a revised view of chromatin insulators in the genomics era.** *Biochim Biophys Acta-Gene Regul Mech* 2014, **1839**:203-214.
25. Negre N, Brown CD, Ma LJ, Bristow CA, Miller SW, Wagner U, Kheradpour P, Eaton ML, Loriaux P, Sealfon R *et al.*: **A cis-regulatory map of the Drosophila genome.** *Nature* 2011, **471**:527-531.
26. Ong CT, Corces VG: **CTCF: an architectural protein bridging genome topology and function.** *Nat Rev Genet* 2014, **15**:234-246.
27. Vogelmann J, Valeri A, Guillou E, Cuvier O, Nollmann M: **Roles of chromatin insulator proteins in higher-order chromatin organization and transcription regulation.** *Nucleus-Austin* 2011, **2**:358-369.
28. Wallace JA, Felsenfeld G: **We gather together: insulators and genome organization.** *Curr Opin Genet Dev* 2007, **17**:400-407.
29. Yang J, Corces VG: **Insulators, long-range interactions, and genome function.** *Curr Opin Genet Dev* 2012, **22**:86-92.
30. Cuddapah S, Jothi R, Schones DE, Roh TY, Cui KR, Zhao KJ: **Global analysis of the insulator binding protein CTCF in chromatin barrier regions reveals demarcation of active and repressive domains.** *Genome Res* 2009, **19**:24-32.
31. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B: **Topological domains in mammalian genomes identified by analysis of chromatin interactions.** *Nature* 2012, **485**:376-380.
32. Handoko L, Xu H, Li GL, Ngan CY, Chew E, Schnapp M, Lee CWH, Ye CP, Ping JLH, Mulawadi F *et al.*: **CTCF-mediated functional chromatin interactome in pluripotent cells.** *Nat Genet* 2011, **43**:630-U198.
33. Schwartz YB, Linder-Basso D, Kharchenko PV, Tolstorukov MY, Kim M, Li HB, Gorchakov AA, Minoda A, Shanower G, Alekseyenko AA *et al.*: **Nature and function of insulator protein binding sites in the Drosophila genome.** *Genome Res* 2012, **22**:2188-2198.
34. Essafi A, Webb A, Berry RL, Slight J, Burn SF, Spraggon L, Velecela V, Martinez-Estrada OM, Wiltshire JH, Roberts SG *et al.*: **A wt1-controlled chromatin switching mechanism underpins tissue-specific wnt4 activation and repression.** *Dev Cell* 2011, **21**:559-574.
35. Soto-Reyes E, Recillas-Targa F: **Epigenetic regulation of the human p53 gene promoter by the CTCF transcription factor in transformed cell lines.** *Oncogene* 2010, **29**:2217-2227.
36. Weth O, Paprotka C, Gunther K, Schulte A, Baierl M, Leers J, Galjart N, Renkawitz R: **CTCF induces histone variant incorporation, erases the H3K27me3 histone mark and opens chromatin.** *Nucleic Acids Res* 2014, **42**:11941-11951.
37. Maksimenko O, Kyrchanova O, Bonchuk A, Stakhov V, Parshikov A, Georgiev P: **Highly conserved ENY2/Sus1 protein binds to Drosophila CTCF and is required for barrier activity.** *Epigenetics* 2014, **9**:1261-1270.
38. Bartkuhn M, Straub T, Herold M, Herrmann M, Rathke C, Saumweber H, Gilfillan GD, Becker PB, Renkawitz R: **Active promoters and insulators are marked by the centrosomal protein 190.** *Embo J* 2009, **28**:877-888.
39. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO *et al.*: **Comprehensive mapping of long-range interactions reveals folding principles of the human genome.** *Science* 2009, **326**:289-293.
40. Heger P, Marin B, Bartkuhn M, Schierenberg E, Wiehe T: **The chromatin insulator CTCF and the emergence of metazoan diversity.** *Proc Natl Acad Sci U S A* 2012, **109**:17507-17512.
41. Holohan EE, Kwong C, Adryan B, Bartkuhn M, Herold M, Renkawitz R, Russell S, White R: **CTCF genomic binding sites in Drosophila and the organisation of the bithorax complex.** *PLoS Genet* 2007, **3**:1211-1222.
42. Soshnikova N, Montavon T, Leleu M, Galjart N, Duboule D: **Functional analysis of CTCF during mammalian limb development.** *Dev Cell* 2010, **19**:819-830.
43. Bowman SK, Deaton AM, Domingues H, Wang PI, Sadreyev RI, ●● Kingston RE, Bender W: **H3K27 modifications define segmental regulatory domains in the Drosophila bithorax complex.** *Elife* 2014, **3**:e02833.
- The *Drosophila* bithorax-complex contains homeotic genes with their segment-specific regulatory elements collinearly arranged with the order of parasegments along the body axis of the fly. CTCF bound insulators separate the segment-specific control regions. The authors isolated chromatin from individual parasegments and found that the repressive chromatin mark H3K27me3 precisely follows a sharp 'step-wise' pattern with CTCF marking the steps.
44. Rousseau M, Crutchley JL, Miura H, Suderman M, Blanchette M, Dostie J: **Hox in motion: tracking HoxA cluster conformation during differentiation.** *Nucleic Acids Res* 2014, **42**:1524-1540.
45. Kim YJ, Cecchini KR, Kim TH: **Conserved, developmentally regulated mechanism couples chromosomal looping and heterochromatin barrier activity at the homeobox gene A locus.** *Proc Natl Acad Sci U S A* 2011, **108**:7391-7396.
46. Xu M, Zhao GN, Lv X, Liu G, Wang LY, Hao DL, Wang J, Liu DP, Liang CC: **CTCF controls HOXA cluster silencing and mediates PRC2-repressive higher-order chromatin structure in NT2/D1 cells.** *Mol Cell Biol* 2014, **34**:3867-3879.
47. Narendra V, Rocha PP, An D, Raviram R, Skok JA, Mazzoni EO, ●● Reinberg D: **Transcription. CTCF establishes discrete functional chromatin domains at the Hox clusters during differentiation.** *Science* 2015, **347**:1017-1021.
- The authors studied the *HoxA* gene cluster during ES cell differentiation into motor neurons. CRISPR/Cas mediated deletion of CTCF binding sites resulted in the expansion of active chromatin into repressive domains. Spreading of active chromatin was confined to the region up to the next CTCF site.
48. Downen JM, Fan ZP, Hnisz D, Ren G, Abraham BJ, Zhang LN, ●● Weintraub AS, Schuijers J, Lee TI, Zhao K *et al.*: **Control of cell identity genes occurs in insulated neighborhoods in mammalian chromosomes.** *Cell* 2014, **159**:374-387.
- The authors used the cohesin ChIA-PET technique, which combines immunoprecipitation of chromatin bound cohesin with chromosome conformation analysis. Super-enhancer-driven cell identity genes and Polycomb-bound lineage-specifying genes were found to occur in insulated chromosome loops. These were generally flanked by interacting CTCF/cohesin sites.
49. Symmons O, Uslu VV, Tsujimura T, Ruf S, Nassari S, Schwarzer W, ●● Ettwiller L, Spitz F: **Functional and topological characteristics of mammalian regulatory domains.** *Genome Res* 2014, **24**:390-400.
- The authors inserted and mapped more than 1000 integration sites of a regulatory sensor transposon within the mouse genome. Transposon activation identified enhancers acting along broad regions, which correlated strongly with TADs. This suggests that TADs confine regulatory activities to regulatory domains.

50. Giorgetti L, Galupa R, Nora EP, Piolot T, Lam F, Dekker J, Tiana G, Heard E: **Predictive polymer modeling reveals coupled fluctuations in chromosome conformation and transcription.** *Cell* 2014, **157**:950-963.
51. Seitan VC, Faure AJ, Zhan Y, McCord RP, Lajoie BR, Ing-Simmons E, Lenhard B, Giorgetti L, Heard E, Fisher AG *et al.*: **Cohesin-based chromatin interactions enable regulated gene expression within preexisting architectural compartments.** *Genome Res* 2013, **23**:2066-2077.
52. Zuin J, Dixon JR, van der Reijden MI, Ye Z, Kolovos P, Brouwer RW, van de Corput MP, van de Werken HJ, Knoch TA, van IJcken WFJ *et al.*: **Cohesin and CTCF differentially affect chromatin architecture and gene expression in human cells.** *Proc Natl Acad Sci U S A* 2014, **111**:996-1001.
- Depletion of cohesin or CTCF resulted in a general loss of local chromatin interactions. Depletion of CTCF not only reduced intra-domain interactions but also increased inter-domain interactions, supporting the role of CTCF at TAD borders.
53. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping.** *Cell* 2014, **159**:1665-1680.
- The authors established an *in situ* Hi-C method where DNA-DNA proximity ligation is performed in intact nuclei. This allowed them to map the genomic architecture at 1 kb resolution. Loop anchors typically occur at domain boundaries and bind CTCF. These CTCF sites occur predominantly (>90%) in a convergent orientation.
54. Van Bortle K, Nichols MH, Li L, Ong CT, Takenaka N, Qin ZS, Corces VG: **Insulator function and topological domain border strength scale with architectural protein occupancy.** *Genome Biol* 2014, **15**:R82.
- Mapping genome-wide binding for several *Drosophila* architectural proteins identified an extensive pattern of colocalization. Architectural proteins in *Drosophila* as well as in mouse and human stem cells established dense clusters at the borders of topological domains. Insulator function and topological domain border strength correlated with IBP protein occupancy.
55. Kung JT, Kesner B, An JY, Ahn JY, Cifuentes-Rojas C, Colognori D, Jeon Y, Szanto A, del Rosario BC, Pinter SF *et al.*: **Locus-specific targeting to the X chromosome revealed by the RNA interactome of CTCF.** *Mol Cell* 2015, **57**:361-375.
56. Matzat LH, Dale RK, Lei EP: **Messenger RNA is a functional component of a chromatin insulator complex.** *EMBO Rep* 2013, **14**:916-922.
57. Lupianez DG, Kraft K, Heinrich V, Krawitz P, Brancati F, Klopocki E, Hom D, Kayserili H, Opitz JM, Laxova R *et al.*: **Disruptions of topological chromatin domains cause pathogenic rewiring of gene-enhancer interactions.** *Cell* 2015, **161**:1012-1025.
- Several unrelated families with syndactyly and brachydactyly were found to possess genomic disruptions of the TAD structure at the EPHA4 locus. Using CRISPR/Cas genome editing, mice were generated with corresponding rearrangements. In both mouse limb tissue and patient-derived fibroblasts, disease-relevant structural changes cause ectopic interactions between promoters and non-coding DNA. This rewiring only occurred if the variant disrupted a CTCF-associated boundary domain.
58. Gomez-Marin C, Tena JJ, Acemel RD, Lopez-Mayorga M, Naranjo S, de la Calle-Mustienes E, Maeso I, Beccari L, Aneas I, Vielas E *et al.*: **Evolutionary comparison reveals that diverging CTCF sites are signatures of ancestral topological associating domains borders.** *Proc Natl Acad Sci U S A* 2015, **112**:7542-7547.
- Chromosome conformation analysis identified two largely independent regulatory landscapes at the conserved homeotic *Six* genes, which are contained within two adjacent TADs. Evolutionary comparison of these TAD borders revealed the presence of CTCF sites with convergent orientations in all studied deuterostomes.
59. Vietri Rudan M, Barrington C, Henderson S, Ernst C, Odom DT, Tanay A, Hadjur S: **Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain architecture.** *Cell Rep* 2015, **10**:1297-1309.
- Comparative Hi-C analysis of liver cells from mouse, macaque, rabbit and dog showed a robust conservation within syntenic regions. Similarly, CTCF/cohesin binding sites are conserved and enriched at TAD borders in a convergent orientation. Genomic reorganization involved intact modules.
60. Kyrchanova O, Chetverina D, Maksimenko O, Kullyev A, Georgiev P: **Orientation-dependent interaction between *Drosophila* insulators is a property of this class of regulatory elements.** *Nucleic Acids Res* 2008, **36**:7019-7028.
61. Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y *et al.*: **CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter function.** *Cell* 2015, **162**:900-910.
- The authors used the CRISPR technique to invert regulatory elements with CTCF binding sites. This resulted in directional changes of 3D-topology at *Pcdh* and beta-globin genes. Together with the finding that the vast majority of genome-wide chromatin loops occur between convergent CTCF binding sites, these results suggest that TAD formation can be predicted and manipulated.
62. Arnold R, Burcin M, Kaiser B, Muller M, Renkawitz R: **DNA bending by the silencer protein NeP1 is modulated by TR and RXR.** *Nucleic Acids Res* 1996, **24**:2640-2647.
63. Sanborn AL, Rao SS, Huang SC, Durand NC, Huntley MH, Jewett AI, Bochkov ID, Chinnappan D, Cutkosky A, Li J *et al.*: **Chromatin extrusion explains key features of loop and domain formation in wild-type and engineered genomes.** *Proc Natl Acad Sci U S A* 2015.
- In this publication a physical simulation is presented, using the data from high-resolution spatial proximity maps. This model is consistent with the formation of loops by a process of extrusion and explains why the CTCF-binding motifs at pairs of loop anchors lie in the convergent orientation. The authors tested their model by using the CRISPR/Cas technology to delete or invert specific CTCF motifs. In every of the 13 mutations generated, the convergent rule was in agreement with loop formation.
64. Whitfield WG, Millar SE, Saumweber H, Frasch M, Glover DM: **Cloning of a gene encoding an antigen associated with the centrosome in *Drosophila*.** *J Cell Sci* 1988, **89**(Pt 4):467-480.
65. Pai CY, Lei EP, Ghosh D, Corces VG: **The centrosomal protein CP190 is a component of the gypsy chromatin insulator.** *Mol Cell* 2004, **16**:737-748.
66. Cuartero S, Fresan U, Reina O, Planet E, Espinas ML: **Ibf1 and Ibf2 are novel CP190-interacting proteins required for insulator function.** *EMBO J* 2014, **33**:637-647.
67. Maksimenko O, Bartkuhn M, Stakhov V, Herold M, Zolotarev N, Jox T, Buxa MK, Kirsch R, Bonchuk A, Fedotova A *et al.*: **Two new insulator proteins, Pita and ZIPIC, target CP190 to chromatin.** *Genome Res* 2015, **25**:89-99.
- The search for multi-zinc finger proteins involved in insulation identified Pita and ZIPIC in *Drosophila*. Both bind to CP190 and mediate enhancer blocking as well as protection from PRE-mediated silencing.
68. Liang J, Lacroix L, Gamot A, Cuddapah S, Queille S, Lhoumaud P, Lepetit P, Martin PG, Vogelmann J, Court F *et al.*: **Chromatin immunoprecipitation indirect peaks highlight long-range interactions of insulator proteins and Pol II pausing.** *Mol Cell* 2014, **53**:672-681.
- Chromatin immunoprecipitation of insulator-binding proteins (IBPs) not only detected the interaction with their cognate binding site, but also identified DNA binding sites of other factors contacting IBPs. These indirect peaks are therefore long-range contact sites. The authors identified CP190 as mediating this interaction, which in many cases links IBPs with RNAPII pausing sites.
69. Vogelmann J, Le Gall A, Dejardin S, Allemand F, Gamot A, Labesse G, Cuvier O, Negre N, Cohen-Gonsaud M, Margeat E *et al.*: **Chromatin insulator factors involved in long-range DNA interactions and their role in the folding of the *Drosophila* genome.** *PLoS Genet* 2014, **10**:e1004544.
70. Van Bortle K, Ramos E, Takenaka N, Yang J, Wahi JE, Corces VG: ***Drosophila* CTCF tandemly aligns with other insulator proteins at the borders of H3K27me3 domains.** *Genome Res* 2012, **22**:2176-2187.
71. Bonchuk A, Maksimenko O, Kyrchanova O, Ivlieva T, Mogila V, Deshpande G, Wolle D, Schedl P, Georgiev P: **Functional role of dimerization and CP190 interacting domains of CTCF protein in *Drosophila melanogaster*.** *BMC Biol* 2015, **13**:63.
72. Bohla D, Herold M, Panzer I, Buxa MK, Ali T, Demmers J, Kruger M, Scharfe M, Jarek M, Bartkuhn M *et al.*: **A functional insulator screen identifies NURF and dREAM components to be required for enhancer-blocking.** *PLoS One* 2014:9.
- Reporter gene activity was determined to identify components required for maximal enhancer blocking. RNAi depletion identified 78 genes

required for optimal Fab-8-mediated enhancer blocking. These included all four components of the NURF complex as well as several subunits of the dREAM complex.

73. Korenjak M, Kwon E, Morris RT, Anderssen E, Amzallag A, Ramaswamy S, Dyson NJ: **dREAM co-operates with insulator-binding proteins and regulates expression at divergently paired genes.** *Nucleic Acids Res* 2014, **42**:8939-8953.
  74. Qiu Z, Song C, Malakouti N, Murray D, Hariz A, Zimmerman M, Gygas D, Alhazmi A, Landry JW: **Functional interactions between NURF and Ctcf regulate gene expression.** *Mol Cell Biol* 2015, **35**:224-237.
  75. Ahanger SH, Gunther K, Weth O, Bartkuhn M, Bhone RR, Shouche YS, Renkawitz R: **Ectopically tethered CP190 induces large-scale chromatin decondensation.** *Scient Rep* 2014, **4**.
  76. Kitchen NS, Schoenherr CJ: **Sumoylation modulates a domain in CTCF that activates transcription and decondenses chromatin.** *J Cell Biochem* 2010, **111**:665-675.
  77. Ohtsuki S, Levine M: **GAGA mediates the enhancer blocking activity of the eve promoter in the Drosophila embryo.** *Genes Dev* 1998, **12**:3325-3330.
  78. Srivastava S, Puri D, Garapati HS, Dhawan J, Mishra RK: **Vertebrate GAGA factor associated insulator elements demarcate homeotic genes in the HOX clusters.** *Epigenet Chromatin* 2013, **6**:8.
  79. Wendt KS, Yoshida K, Itoh T, Bando M, Koch B, Schirghuber E, Tsutsumi S, Nagae G, Ishihara K, Mishiro T *et al.*: **Cohesin mediates transcriptional insulation by CCCTC-binding factor.** *Nature* 2008, **451**:796-801.
  80. Raab JR, Chiu J, Zhu J, Katzman S, Kurukuti S, Wade PA, Haussler D, Kamakaka RT: **Human tRNA genes function as chromatin insulators.** *EMBO J* 2012, **31**:330-350.
  81. Lei EP, Corces VG: **RNA interference machinery influences the nuclear organization of a chromatin insulator.** *Nat Genet* 2006, **38**:936-941.
  82. Yao H, Brick K, Evrard Y, Xiao T, Camerini-Otero RD, Felsenfeld G: **Mediation of CTCF transcriptional insulation by DEAD-box RNA-binding protein p68 and steroid receptor RNA activator SRA.** *Genes Dev* 2010, **24**:2543-2555.
  83. Ong CT, Van Bortle K, Ramos E, Corces VG: **Poly(ADP-ribosyl)ation regulates insulator function and intrachromosomal interactions in Drosophila.** *Cell* 2013, **155**:148-159.
  84. Zampieri M, Guastafierro T, Calabrese R, Ciccarone F, Bacalini MG, Reale A, Perilli M, Passananti C, Caiafa P: **ADP-ribose polymers localized on Ctcf-Parp1-Dnmt1 complex prevent methylation of Ctcf target sites.** *Biochem J* 2012, **441**:645-652.
  85. Zhao H, Sifakis EG, Sumida N, Millan-Arino L, Scholz BA, Svensson JP, Chen X, Ronnegren AL, Mallet de Lima CD, Varnoosfaderani FS *et al.*: **PARP1- and CTCF-mediated interactions between active and repressed chromatin at the lamina promote oscillating transcription.** *Mol Cell* 2015, **59**:984-997.
- PARP1 and CTCF regulate the contacts between circadian genes and nuclear lamina. Oscillating binding to the lamina promotes oscillating transcriptional attenuation of clock-controlled genes.
86. Lhoumaud P, Hennion M, Gamot A, Cuddapah S, Queille S, Liang J, Micas G, Morillon P, Urbach S, Bouchez O *et al.*: **Insulators recruit histone methyltransferase dMes4 to regulate chromatin of flanking genes.** *EMBO J* 2014, **33**:1599-1613.
  87. Galli GG, Carrara M, Francavilla C, de Lichtenberg KH, Olsen JV, Calogero RA, Lund AH: **Genomic and proteomic analyses of Prdm5 reveal interactions with insulator binding proteins in embryonic stem cells.** *Mol Cell Biol* 2013, **33**:4504-4516.
  88. Van Bortle K, Peterson AJ, Takenaka N, O'Connor MB, Corces VG: **CTCF-dependent co-localization of canonical Smad signaling factors at architectural protein binding sites in D. melanogaster.** *Cell Cycle* 2015, **14**:2677-2687.
  89. Bergstrom R, Savary K, Moren A, Guibert S, Heldin CH, Ohlsson R, Moustakas A: **Transforming growth factor beta promotes complexes between Smad proteins and the CCCTC-binding factor on the H19 imprinting control region chromatin.** *J Biol Chem* 2010, **285**:19727-19737.
  90. Defossez PA, Kelly KF, Filion GJ, Perez-Torrado R, Magdinier F, Menoni H, Nordgaard CL, Daniel JM, Gilson E: **The human enhancer blocker CTC-binding factor interacts with the transcription factor Kaiso.** *J Biol Chem* 2005, **280**:43017-43023.
  91. Burcin M, Arnold R, Lutz M, Kaiser B, Runge D, Lottspeich F, Filippova GN, Lobanenko VV, Renkawitz R: **Negative protein 1, which is required for function of the chicken lysozyme gene silencer in conjunction with hormone receptors, is identical to the multivalent zinc finger repressor CTCF.** *Mol Cell Biol* 1997, **17**:1281-1288.