

Health Risks of Some Meat Additives on Male Rats

¹Rasha, A. Elsabagh, ¹Reham, A. Amin and ²Aziza Amin

¹Department of Food Control, Faculty of Veterinary Medicine, Benha University, Egypt

²Department of Pathology, Faculty of Veterinary Medicine, Benha University, Egypt

Abstract: Natural and synthetic meat additives approved by Food and Drug Administration "FDA" are commonly used to maintain or improve safety, nutrient value, taste and texture. Although many of the 3,000 these additives enhance meat supply, others are the subject of fierce controversy. The present study investigated the hazardous effects of sodium nitrite and monosodium glutamate "MSG" used in manufacturing of meat on some hematological, serological parameters and histopathological changes in male albino rats. Manually processed meat supplemented with doses of sodium nitrite and Monosodium glutamate "MSG", below, within and over the permissible limits, was fed to male rats for 30 days. All the used doses sodium nitrite and MSG, below, within and over the permissible limits, induced significant decrease in hemoglobin content and some hematological parameters, significant increase in serum levels of aspartate aminotransferase "AST" and alanine aminotransferase "ALT", urea and creatinine as well as various pathological changes in liver, kidney, brain and tests. The severity of these changes increased from mild to severe by increasing the concentrations of both sodium nitrite and MSG. In conclusion, the obtained data indicated that prolonged use of sodium nitrite and MSG, even within the permissible limits according to EOS (2005), was unsafe for consumer because they cause anemia, liver and kidney dysfunctions, damage to brain cells in association with deleterious effect on male fertility. Consequently, it is not recommended to use these meat additives in meat even within the permissible limits.

Key words: Sodium nitrite • Monosodium glutamate (MSG) • Hematological changes • Liver and kidney functions • Histopathological changes.

INTRODUCTION

Meat preservation has become an increasingly important practice in modern food technology with the increased production of processed and convenient food [1]. Natural and synthetic food additives approved by Food and Drug Administration "FDA" are commonly used to maintain or improve safety, the nutrient value and the taste and texture of food [2]. Although many of the 3,000 these additives enhance our food supply, others are the subject of fierce controversy. The discovery that children at the age of nursery consume food containing great amounts of additives prompted the scientific community to oversee this issue.

Nitrate and nitrite have been used as food additives in many types of meat products for many years primarily to fix color, contribute to the flavor of the final product,

help in the inhibition of growth and toxin production of *Clostridium botulinum* which causes botulism [3], effectively control rancidity by inhibiting lipid oxidation leading to excellent storage stability [4] and improves the microbiological safety of these foods and extends their safe shelf-life [5]. This offers significant benefits to consumers in terms of the availability of a variety of different foods that are safe, convenient and cost effective.

The toxic effects of nitrates and nitrites are well documented in mammals, including impairment of reproductive function [6], hepatotoxicity and methaemoglobinemia [7]. Instance, highly carcinogenic N-nitroso-compounds are produced when nitrite reacts with secondary amines and N-alkyl amides under acidic conditions in vitro [8]. An alternative to sodium nitrite for production of cured meats has not been identified despite significant research effort [9].

Monosodium glutamate "MSG" is the sodium salt of naturally occurring non-essential amino acid L-form of glutamic acid [10]. It is one of the main popular flavor enhancers used as a food additive in various food products in modern time [11]. It is widely used in many commercial packed food, restaurant and household cooking. The unique flavor and taste of this compound has been categorized and established as a separate taste sensation UMAMI taste [12] which is recognized as the fifth basic taste, very similar to "meat aroma" or "broth aroma" so enhances food palatability and encourages flavor acceptance [13]. MSG can be added "pure" or as a "hidden ingredient" of yeast extracts or hydrolyzed proteins, both containing high percentages of glutamic acid [14]. It is a component of many proteins and peptides. When bound to proteins, glutamate is tasteless. The sweet umami taste and flavor becomes perceptible only when free glutamate dissociates from proteins during the processes of fermentation, ripening and cooking [15]. It increases the perception of sweetness and saltiness and diminishes the sourness and bitterness of food. MSG serves as an energy source for certain tissues and as a substrate for glutathione syntheses [16]. MSG is a naturally present excitatory neurotransmitter in brain, mediating fast synaptic transmission in one third of all CNS synapses.

A report from the Federation of American Societies for Experimental Biology "FASEB" (1995) identified two groups of people susceptible to high MSG doses: a group intolerant to high quantities of MSG and asthmatics.

There is no complete agreement about the safety of MSG, even though FDA includes it among the substances generally recognized as safe (GRAS) [17, 16]. Despite its taste stimulation and improved appetite enhancement, reports indicated that MSG is toxic to human and experimental animals [18] and develops severe abnormalities as short stature, small endocrine organs (pituitary gland, adrenal gland, thyroid gland, ovaries, testes and pancreas), high risk of seizures and impaired learning [19]. On the contrary, several government institutions and international organizations as JECFA, FAO and WHO, have declared that MSG is safe for human consumption [20].

Although, different pathological changes in liver following MSG were reported previously by several authors [21,22], still not many literatures have described detailed histological features as effect of MSG in liver tissue. On the other hand, there was shortage in the literatures that had been recorded the pathological changes occurring in various organs such as kidney, brain and testes following MSG administration [21].

Therefore, the aim of this study was to determine the alterations in hematological, serological and pathological parameters of liver and kidneys in combination with pathological changes in brain and testes of male rats that had been fed on manually processed meat supplemented with different doses of sodium nitrite and MSG, including the permissible limits recommended by EOS [23] for 30-successive days.

MATERIAL AND METHODS

Experimental Animals: The present investigation was carried out on a total number of 35 apparently healthy adult male white albino rats (one month age and 130-150 g body weight). These animals were housed for two weeks at constant environmental and nutritional conditions similar to those under which the experiment was performed for accommodation. Rats were housed in separate cages a way from any stressful stimuli and supplied with diet and water ad libitum.

Meat Additives:

Sodium Nitrite: Food grade sodium nitrite was incorporated in fresh minced beef at concentrations of 100 ppm, 125 ppm (permissible limit according to EOS [23] and 150 ppm.

Monosodium Glutamate "MSG": Food grade MSG was incorporated in fresh minced beef at concentrations of 3 ppm, 5 ppm [24] and 7 ppm.

Experimental Design: The male rats were divided randomly into 7 main groups (n=5) feeding on manually processed meat supplemented with different doses of nitrite and MSG as follows:

Group	Dose
C	Control group received manually processed kofta without any treatment.
N1	Group fed on meat supplemented with 100 ppm sodium nitrite for 30 days.
N2	Group fed on meat supplemented with 125 ppm sodium nitrite for 30 days.
N3	Group fed on meat supplemented with 150 ppm sodium nitrite for 30 days.
MSG1	Group fed on meat supplemented with MSG 3 mg/Kg for 30 days.
MSG2	Group fed on meat supplemented with MSG 5 mg/Kg for 30 days.
MSG3	Group fed on meat supplemented with MSG 7 mg/Kg for 30 days.

At the end of the experimental period, overnight fasted animals were sacrificed by cervical dislocation and blood samples were collected in centrifuge tubes. Serum was separated from coagulant blood by centrifugation at 860 g for 20min and then quickly frozen at -20°C for biochemical analysis. Moreover, Specimens for histopathological examination were taken from different organs including liver, kidney, brain and testes. Tests were performed in triplicate.

Hemogram Parameters: Complete blood picture was performed using HA-Vet Automatic Hematology analyzer (CLINDIAG SYSTEM).

Chemical Analysis of Serum:

Aspartate Aminotransferase "AST" and Alanine Aminotransferase "ALT": were determined spectrophotometrically according to Schumann and Klauke [25].

Urea: Was determined spectrophotometrically according to Patton and Crouch [26].

Creatinin: Was determined spectrophotometrically according to Henry *et al.* [27].

Histopathological Examination: Five rats were sacrificed successfully from each group after 30 days post feeding of diet supplemented with different concentrations of nitrite and MSG. Specimens for histopathological examination were taken from different organs including liver, kidney, brain and testes. These samples were fixed in 10% buffered neutral formalin solution. Then after proper fixation, the samples were dehydrated in ascending grades of ethyl alcohol, then cleared in xylol, embedded in paraffin and finely blocking occurred. These samples were

sectioned at 5µm in thickness and stained with hematoxylin and eosin (H & E) for microscopical examination [28]. According to the severity of the microscopical lesions, all specimens were examined for histological alterations and classified on a modified semi-quantitative scale of Bagheri *et al.* [29] into none (-), mild (+), moderate (++) and severe (+++). Each lesion was graded as follow: (-) normal histology, (+) with up 1/3 of the examined section exhibiting the evaluated lesion, (++) in which 1/3 to 2/3 of the inspected section revealed the estimated pathological change, while (+++) means greater than 2/3 of the assessed section showed the estimated pathological alteration.

Statistical Analysis: The data was statistically treated by one-way ANOVA using SPSS program for windows (Version 16) (SPSS Inc. Chicago, IL and USA) and Duncan's post hoc test with $P < 0.05$ considered to be statistically significant. The results were expressed as means + SD.

RESULTS

Hematological Examination: With respect to the hematological changes of sodium nitrite and MSG treated rats, there were significant decrease in RBCs count, HGB, WBCs, MCV, MCH, MCHC, lymphocytes and monocytes in treated rats with the three dose levels as compared to those of control animals (Tables 1 and 3) revealing anemia in all sodium nitrite treated groups during all the experimental duration.

Serological Examination: Serum biochemical constituents revealed that sodium nitrite and MSG treated groups showed significant increase in the levels of AST, ALT, urea and creatinin levels (Tables 2 and 4).

Table 1: The effect of different doses of sodium nitrite on some hematological changes in male rats

Groups	Lymphocytes		Monocytes		HGB*	RBCs*	WBCs	MCV*	MCH	MCHC
	No.	%*	No.	%*						
Control	0.96±0.44 ^a	9.73±0.31 ^a	0.27±0.14 ^a	2.50±0.14 ^a	30.86±1.60 ^a	8.27±0.43 ^a	11.32±5.72 ^a	59.6±2.88 ^a	39.20±1.43 ^a	68.18±3.15 ^a
N1	0.83±0.07 ^a	9.07±0.43 ^{ab}	0.22±0.02 ^a	2.48±.13 ^a	29.12±1.47 ^a	7.76±0.24 ^{ab}	8.94±0.80 ^a	59.4±2.88 ^a	39.10±1.94 ^a	67.88±1.42 ^a
N2	0.81±0.06 ^a	8.96±0.24 ^b	0.21±0.02 ^a	2.42±0.15 ^a	26.96±1.22 ^{ab}	6.90±0.44 ^{bc}	8.90±0.90 ^a	55.0±1.87 ^b	37.59±0.78 ^a	65.18±1.82 ^a
N3	0.75±0.07 ^a	8.65±0.33 ^b	0.21±0.01 ^a	2.41±0.12 ^a	24.62±5.03 ^b	6.35±1.42 ^c	8.34±0.68 ^a	54.4±4.10 ^b	37.40±1.26 ^a	65.12±3.86 ^a

Table 2: The effect of different doses of sodium nitrite on some liver and kidney function tests in male rats

Groups	AST	ALT	Urea	Creatinine
Control	310.10±70.74 ^a	58.32±3.61 ^b	39.44±5.77 ^b	0.44±0.05 ^a
N1	354.68±59.48 ^a	70.52±19.67 ^{ab}	49.58±12.48 ^{ab}	0.46±0.05 ^a
N2	365.26±27.03 ^a	71.60±14.22 ^{ab}	56.42±10.50 ^a	0.50±0.00 ^a
N3	373.30±37.16 ^a	89.42±17.63 ^a	60.92±5.78 ^a	0.52±0.11 ^a

Table 3: The effect of different doses of monosodium glutamate (MSG) on some hematological changes in male rats

Groups	Lymphocytes		Monocytes		HGB	RBCs*	WBCs*	MCV*	MCH*	MCHC*
	No.*	%*	No.*	%						
Control	9.73±0.31a	0.96±0.44 ^a	2.50±0.14 ^a	0.27±0.14 ^a	30.86±1.60 ^a	8.27±0.43 ^a	11.32±5.72 ^a	59.6±2.88 ^a	39.20±1.43 ^b	68.18±3.15 ^a
N1	9.35±0.61a	0.79±0.05 ^a	2.48±0.13 ^a	0.21±0.01 ^a	30.06±0.10 ^a	8.02±.40 ^a	8.90±0.90 ^a	57.80±3.70 ^b	38.57±1.23 ^a	67.52±1.74 ^a
N2	9.16±0.40b	0.76±0.09 ^a	2.46±0.19 ^a	0.20±0.02 ^a	29.16±3.36 ^a	7.58±0.82 ^b	7.82±1.11 ^{ab}	55.20±2.77 ^b	37.57±1.46 ^b	66.34±2.38 ^b
N3	8.77±0.23b	0.64±0.10b	2.35±0.18 ^a	0.17±0.03 ^b	27.86±1.94 ^a	6.85±0.44 ^b	6.94±1.03 ^b	54.40±4.10 ^b	37.40±1.26 ^b	63.36±1.51 ^b

Table 4: The effect of different doses of monosodium glutamate (MSG) on Some liver and kidney function tests in male rats

Groups	AST	ALT	Urea	Creatinine
Control	310.10±70.74 ^a	58.32±3.61 ^a	39.44±5.77 ^b	0.44±0.05 ^a
MSG1	321.26±46.51 ^{ab}	80.68±12.28 ^a	54.44±12.50 ^a	0.46±0.05 ^a
MSG2	334.54±38.76 ^{ab}	84.54±34.05 ^a	56.16±3.79 ^a	0.50±0.00 ^a
MSG3	365.26±27.03 ^a	104.44±64.87 ^a	60.92±5.78 ^a	0.52±0.11 ^a

The values represent Mean ± SD of three experiments.

Means within a column followed by different letters are significantly different ($P < 0.05$).

Table 5: Lesions scores of various histopathological changes in different organs of rats fed manually processed kofta supplemented with various concentrations of sodium nitrite for 30- successive days

Lesion score	Treatment		
	Sodium Nitrite		
	Lower than permissible	Permissible	Higher than permissible
Liver			
Congestion of hepatic blood vessels	-	+	++
Perivascular hemorrhage	+	+	++
Leukocytic infiltration	+	++	++
Disturbance in hepatic cords	-	-	-
Degenerative changes	+	++	+++
Edematous fluid in hepatic parenchyma	-	-	-
Nuclear changes	-	+	++
Necrosis of hepatic cells	-	+	++
Hyperplasia of bile duct	+	++	++
Portal cirrhosis	+	++	++
Kidney			
Congestion	+	+	++
Cellular vacuolation	+	++	+++
Tubular epithelial necrosis	-	+	++
Hyaline casts	+	++	+++
Leukocytic infiltration	-	+	+
Periglomerular edema	-	+	++
Segmentation of glomerular tuft	-	+	++
Shrinkage of glomerular tuft	-	+	++
Brain			
Neural degeneration	+	+	++
Hemorrhage	-	++	+++
Encephalomalacia	-	+	++
Perivascular lymphocytic cuffing	-	-	-

Table 5: Continued

Lesion score	Treatment		
	Sodium Nitrite		
	Lower than permissible	Permissible	Higher than permissible
Testes			
Degeneration of lining epithelial cells of SNT	+	++	+++
Vacuolation in the SNT lumen	-	+	++
Destruction of basement membrane of SNT	-	+	++
Calcification of degenerated cells	-	-	-
Accumulation of edematous fluid either in SNT or in the interstitial tissue		-	+ ++

(-): normal histology,

(+): up to 1/3 of the examined section exhibiting the evaluated lesion,

(++): up to 1/3 to 2/3 of the inspected section revealed the estimated pathological change,

(+++): greater than 2/3 of the assessed section showed the estimated pathological alteration

Table 6: Lesions scores of various histopathological changes of different organs of rats fed manually processed kofta supplemented with various concentrations of MSG for 30- successive days

Lesion score	Treatment		
	MSG		
	Lower than permissible	Permissible	Higher than permissible
liver			
Congestion of hepatic blood vessels	-	+	++
Perivascular hemorrhage	+	+	++
Leukocytic infiltration	+	++	++
Disturbance in hepatic cords	+	+	+
Degenerative changes	+	++	+++
Nuclear changes	-	+	++
Edematous fluid in hepatic parenchyma	+	+	+
Necrosis of hepatic cells	-	+	++
Hyperplasia of bile duct	+	+	+
Portal cirrhosis	-	-	+
Kidney			
Congestion	+	+	++
Cellular vacuolation	-	++	+++
Tubular epithelial necrosis	-	-	++
Hyaline casts	+	++	++
Leukocytic infiltration	-	+	+
Periglomerular edema	-	-	-
Shrinkage of glomerular tuft	+	++	++
Segmentation of glomerular tuft	-	+	+
Brain			
Neural degeneration	+	++	++
Hemorrhage	+	++	++
Encephalomalacia	+	++	+++
Perivascular lymphocytic cuffing	+	+	++
Testes			
Vacuolation in the SNT lumen	+	+	++
Destruction of basement membrane of SNT	-	+	++
Calcification of degenerated cells	-	-	+
Accumulation of edematous fluid either in SNT or in the interstitial tissue		-	+ ++

(-): normal histology,

(+): up to 1/3 of the examined section exhibiting the evaluated lesion,

(++): up to 1/3 to 2/3 of the inspected section revealed the estimated pathological change,

(+++): greater than 2/3 of the assessed section showed the estimated pathological alteration




Fig. 1: Liver of rats fed diet supplemented with different concentrations of sodium nitrite for 30-successive days,(A) At a level lower than the permissible one showed few periportal mononuclear leukocytic cellular infiltration (black arrow),(B) At a level lower than the permissible one showed portal fibrosis infiltrated with leukocytes (black arrow) in combination with newly formed bile ductules (spotted arrow),(C) At the permissible level showed focal area of coagulative necrosis in the hepatic parenchyma with leukocytic infiltration (black arrow). Notice also diffuse hydropic degeneration of hepatocytes,(D) At a level higher than the permissible one showed multiple areas of lytic necrosis characterized by disappearance of hepatocytes and replaced by erythrocytes and few mononuclear leukocytic cells mainly lymphocytes (black arrow). H&E stain x400.




Fig. 2: Pathological alterations in various organs (kidney, brain and testes) of rats fed diet supplemented with different concentrations of sodium nitrite for 30-successive days,(A) At the permissible level, the kidney showed degeneration of the lining epithelium of renal tubules in association with pyknotic nucleus of the renal tubules epithelium (black arrow),(B) At a level higher than the permissible one, the kidney showed necrosis of the lining epithelium of renal tubules (spotted arrow) with the presence of eosinophilic hyaline casts in the lumen of some tubules, note also shrinkage of the glomerular tuft (black arrow),(C) At a level lower than the permissible one, the brain showed neural degeneration with swollen, rounded cell bodies and indistinct or eccentric pyknotic nucleus (black arrow),(D) At a level higher than the permissible one, the brain showed focal area of encephalomalacia in cerebellum admixed with erythrocytes (black arrow),(E) At a level lower than the permissible one, the testes showed degeneration of some seminiferous tubules. Note also vacuolation in the lumen some tubules.(F) At a level higher than the permissible one, the testes showed severe degeneration of spermatogenic cells of some seminiferous tubules in combination with calcification of the degenerated cells in some tubules (black arrow). H&E stain x400.




Fig. 3: Liver of rats fed diet supplemented with different concentrations of MSG for 30-successivedays,(A) At a level lower than the permissible one showed few perivascular mononuclear leukocytic cellular infiltration (black arrow), note also degeneration of hepatocytes with activation of vonkupffer cells.(B) At a level lower than the permissible one showed mild hyperplasia of the billiary epithelium (spotted arrow) with mononuclear leukocytic cellular infiltration in the portal area (black arrow),(C) At the permissible level showed portal and periportal leucocytic cellular infiltration mainly lymphocytes and few macrophage (black arrow), notice also pyknosis and karyolysis of hepatic cell nuclei,(D) At a level higher than the permissible one showed focal area of coagulative necrosis in the hepatic parenchyma (black arrow) with pyknosis and karyolysis of hepatic cell nuclei. H&E stain x400




Fig. 4: Kidney of rats fed diet supplemented with different concentrations of MSG for 30-successivedays,(A) At a level lower than the permissible one showed periglomerular edema admixed with few mononuclear leukocytic cellular infiltration, note also inter-tubular mononuclear leukocytic cellular infiltration (black arrow),(B) At the permissible level showed degeneration of the lining epithelium of some proximal and distal convoluted tubules as well as disintegration of the glomerular tuft (spotted arrow).(C) At a level higher than the permissible one showed necrosis of the lining epithelium of renal tubules as well as presence of hyaline casts in the tubular lumen (black arrow),(D) At a level higher than the permissible one showed shrinkage of the glomerular tuft with necrosis of its endothelial lining (black arrow). H&E stain x400




Fig. 5: Brain and testes of rats fed diet supplemented with different concentrations of MSG for 30-successive days,(A) Brain of rat fed diet supplemented with MSG at a level lower than the permissible one for 30 days showed perivascular hemorrhage (black arrow),(B) Brain of rat fed diet supplemented with the permissible level of MSG for 30 days showed showing focal areas of encephalomalacia in cerebellum admixed with spaces infiltrated by few numbers of glial cells and lymphocytes (black arrow),(C) Brain of rat fed diet supplemented with MSG at a level higher than the permissible level for 30 days showed focal areas of encephalomalacia in cerebellum and mixed with erythrocytes and mononuclear leukocytes (black arrow),(D) Testes of rat fed diet supplemented with MSG at a level lower than the permissible level for 30 days showed sperm giant cell in the lumen of some seminiferous tubules (black arrow),(E) Testes of rat fed diet supplemented with the permissible level of MSG for 30 days showed degeneration of spermatogenic cells of some seminiferous tubules (black arrow). Note also vacuolation in the lumen some tubules.(F) Testes of rat fed diet supplemented with MSG at a level higher than the permissible one for 30 days showed vacuolation of the lining cells of some seminiferous tubules in combination with accumulation of edematous fluid in the interstitial cells (black arrow). H&E stain x400.

Histopathological Examination:

Histopathological Changes in Rats Fed Diet Containing Different Concentrations of Sodium Nitrite and MSG:

Variable pathological alterations in liver, kidney, brain and testes were observed in treated animals. The severity of the pathological changes was increased by increasing the concentrations of sodium nitrite and MSG. Pathological lesion scores in various organs of rats fed diet supplemented with different concentrations of sodium nitrite and MSG for 30 successive days were summarized in Tables (5 and 6) and explained in figure legends.

DISCUSSION

Nitrate and nitrite have been used as meat additives in many types of meat products for many years primarily to fix color, contribute to the flavor of the final product. On the other side, the reaction of nitrite with haemoglobin

to form methaemoglobin can reduce oxygen transport in the blood and a theoretical possibility of the potential reaction with secondary amines under the acidic conditions of the stomach to form carcinogenic N-nitrosamines and free radicals in foods or in humans in vivo [30]. Such products may increase lipid peroxidation, which can be harmful to different organs including liver and kidney [31].

In the present study, sodium nitrite treated rats showed significant decrease in RBCs count, HGB, WBCs, MCV, MCH, MCHC, lymphocytes and monocytes. Such results agree with those of Smith [32], Reutov *et al.* [33], Vlaskina *et al.* [34] and Eman - Helal and Mervat Abdel- Rahman [35].

The hematological analysis of treated rats revealed anemia that could be attributed increased fragility of erythrocytes due to metabolic impairment [36] or due to slowing down or obstructing the

NAD and NADP reduction processes in erythrocytes by nitrite ions resulting in disturbing their cell respiration, lysis of RBCs and breakdown of HGB to bilirubin in the liver [37] or due to hyperpolarization of erythrocyte membranes and increase in membrane rigidity as result of RBCs oxidation by sodium nitrite [38].

Additionally, nitrates and nitrites trigger destruction of vitamin A and B-group vitamins [39], thereby promoting anemia. They additionally enhance free-radical reactions that may affect the shape and permeability of a cellular membrane of erythrocytes. Oxidation of heme iron, occurring during methemoglobin synthesis from HGB under the influence of nitrites, is a source of reactive oxygen species that initiate the chain of oxidative processes in RBCs. The oxidative stress results in enhanced peroxidation of lipids and aggregation of proteins of erythrocyte membrane. These processes affect changes in the permeability of the cellular membrane and diminish erythrocyte deformability, leading in this way to enhanced degradation of erythrocytes [40] and earlier removal of erythrocytes from circulation, thus shortening their life cycle [41].

The significant decrease in HGB concentration in blood of rats administered sodium nitrite has been shown to be a result of nitrites induced disturbances in heme biosynthesis, transformation of HGB into ethemoglobin [42]. Adverse changes in values of the erythrocytic system markers, occurring under the influence of sodium nitrite, may be possibly through disorders of erythrocyte synthesis, enhanced hemolysis and water shifting to the extracellular compartment [43].

Biochemical analysis could help to identify the target organs of toxicity and the general health status. It may also provide an early warning signal in stressed organism [44]. The serological results in relation to liver function of sodium nitrite treated groups in the present study agree with those of Atef *et al.* [36] and Hanaa-Hassan *et al.* [45].

The increase in the activity of AST and ALT enzymes in the serum of NaNO₂-treated rats could be attributed to the formation of over 300 cytotoxic N-nitrosocompounds, in the acidic environment of the stomach as a result of combination of sodium nitrite with secondary amines in the food or the body, causing severe hepatic and renal necrosis [46] or may be due to anemia and methaemoglobinemia which induced hypoxic injury to centrilobular hepatocytes that consequently cause enzyme leakage [47]. Also, elevated ALT can be used to diagnose myocardial infarction, arrhythmias and severe angina of heart.

Moreover, the serological results in relation to kidney function of sodium nitrite treated groups in the present study agree with those of Atef *et al.* [36], Til *et al.* [48] and Hanaa-Hassan *et al.* [45].

Furthermore, in response to NaNO₂ treatment, urea and creatinine increased in the serum, suggesting an impairment of kidney functions. These effects could also be attributed to the cytotoxic effect of N-nitrosocompounds in renal tubular cells and the changes in the threshold of tubular reabsorption, renal blood flow and glomerular filtration rate [49].

The changes in the liver and kidneys function parameters could be attributed to damage in the liver and kidneys that proved by the results of the histopathological alterations demonstrated in these organs. Furthermore, several pathological changes were demonstrated in the brain tissue of treated rats with different doses of sodium nitrite. These changes could be attributed to the role of nitrite in the induction of oxidative stress in brain of rat resulting in inhibition of neuron activities as it produced some neurobiochemical alterations in the brain tissues such as inhibition of Acetylcholine esterase (AChE) and elevation of lactate dehydrogenase (LDH) activity [50]. These enzymes have been used as standard biomarkers of toxic stress [51]. AChE is responsible to promote hydrolysis of the neurotransmitter "acetylcholine" released at the nerve endings to mediate transmission of the neural impulse across the synapse [52]. Interestingly, LDH is involved in energy production and the elevation of LDH activity might play a role in brain damage after treatment with NaNO₂ [53]. This elevation could be attributed to a generalized increase in membrane activity due to the increase of one of isoenzymes of LDH [54]. Furthermore, degeneration and necrosis of some of the spermatogenic cells in association with other pathological changes were detected in the testicular tissues of rats have a diet supplemented with variable concentrations of sodium nitrite. These findings could be due to defect in the function of Sertoli cell as reported previously by Grant and Butler [55]. Additionally, production of reactive nitrogen species by nitrate plays an important role in its carcinogenic effect through its reaction with body tissues and triggering lipid peroxidation, DNA lesions, enzyme inactivation and damage of different organs [56] that could clarify the detection of different pathological changes in various examined organs in the present work.

Meanwhile, MSG has been recognized in first half of twentieth century as taste enhancer but at the same time doubts were raised about MSG as a causative agent of Chinese restaurant syndrome which is characterized by headache, flushing, numbness, muscle tightness, generalized weakness and bronchoconstriction in asthmatics [57] that revealed there is no complete agreement about the safety of MSG.

Likewise, the results of the present study were nearly similar to those of reports suggesting that MSG could be toxic to erythrocytes and also cause deleterious changes in hematological and biochemical parameters [58] and [59]. However, Kolawole [60] found that treatment of adult rats with MSG (5-15 mg/kg body weight) for 28 days, showed no significant changes in HGB, RBC and WBC counts. No significant change was also observed in all the studied biochemical parameters, including indices of hepatic and renal functions. The results of this study suggest that MSG at the doses administered is not hazardous to health.

In the current study, the pathological changes in the liver parenchyma observed in MSG treated rats were nearly similar to those reported by Ortiz *et al.* [61] and Bhattacharya *et al.* [21]. MSG induced pathological alterations in the hepatic parenchyma even in the concentration at a level lower than the permissible limit. Chronic administration of MSG (4 mg/Kg and above) induced oxidative stress in experimental animals. Moreover, it produces oxygen derived free radicals [62] and causes changes in the liver parenchyma of mice around central vein, hepatocellular damage dilated sinusoids, inflammatory cells and pyknotic nuclei [21]. Previously, it was proved that hepatocytes especially in the central portion of the hepatic lobules are typically the primary site of toxic injury; as they have more surface receptors for toxins and less oxygen [63]. Additionally, the hydropic degeneration of hepatocytes could be considered as a kind of cellular defensive mechanism against injurious substances [64]. At the same time these vacuoles could preserve the biological activities of hepatocytes by collecting the injurious elements and preventing them from interfering with the biological activities of these cells [63].

In this work, different pathological changes were demonstrated in the renal tissues such as degeneration and necrosis of the lining epithelium of renal tubules as well as periglomerular and intertubular leukocytic infiltrations which are in agreement with the findings of Bopanna *et al.* [65] who described patchy tubular necrosis and interstitial infiltration. However, the

alterations in the hematological and biochemical parameters of the liver and kidney of rats treated with MSG are in correlation with the hepatic and renal histopathological changes in the present study. On the other hand, it was described that MSG is metabolized in liver and kidney which play an important role in its removal that could clarify the occurrence of pathological changes in these organs.

In the present work, different doses of MSG in rats produced variable pathological changes in the brain tissues. These changes returned to the neurotoxic effect of MSG especially when implicated in high doses and contribute to development of neurodegenerative disease [66]. However, the alterations in the brain tissues produced by MSG could be also due to either increase in the free radicals generation in the body [62] or due to the ability of MSG to induce oxidative damage in the brain, liver and kidney in experimental animals [67] which considered another clarification for the deleterious effect of MSG in these organs. The excitotoxic effect of glutamate could be attributed to its ability for induction of unlimited elevation of intracellular calcium resulting in activation of various enzymes that responsible for cell death by various mechanisms [66]. These excitotoxic lesions in the lateral hypothalamus are neuronal loss and disruption of blood brain barrier [68]. Additionally, MSG is a neurotoxic agent i.e. causing damage to brain cells, retinal degeneration, many endocrine disorders, renal damage [69], addiction, stroke, epilepsy, brain trauma, neuropathic pain, schizophrenia, anxiety, depression, Parkinson's disease, Alzheimer's disease, Huntington's disease, amyotrophic lateral sclerosis [18] and lesions in certain parts of the brain of neonates resulting in obesity because the blood-brain barrier in neonates is immature [70].

On the other hand, different concentration of MSG in manually processed kofta induced various pathological changes in the testicular tissues such as testicular degeneration, atrophy of the seminiferous tubules as well as destruction of its basement membrane. These findings confirm the implication of MSG in cases of male infertility which in agreement with the results of Eweka and Adjene, [18]. However, these alterations may be due to the toxic effect of MSG to the testicular cells [71].

Finally, we can conclude that the prolonged use of sodium nitrite and MSG even within the permissible limits leads to anemia, liver and kidney dysfunctions, damage to brain cells in association with deleterious effect on male fertility.

REFERENCES

1. Saad, B., M.F. Bari, M.I. Saleh, K. Ahmad and M.K.M. Talib, 2005. Simultaneous determination of preservatives (benzoic acid, sorbic acid, methylparaben and propylparaben) in foodstuffs using high performance liquid chromatography. *Journal of Chromatography A*, 1073: 393-397.
2. National Toxicology Program "NTP" 2001. "NTP" toxicology and carcinogenesis studies of sodium nitrite (CAS No. 7632-00-0) drinking water studies in F344/N rats and B6C3F1 mice. National Toxicology Program Technical Report, 495: 1-274.
3. Davidson, M.P., V.K. Juneja and J.K. Brannen, 2002. Antimicrobial agents. In: *Food Additives Second Edition Revised and Expanded*. Marcel Dekker, Inc.
4. Pearson, A.M. and T.A. Gillett, 1996. *Processed Meats*, 3rd Ed. Chapman and Hall, New York.
5. Pegg, R.B. and F. Shahidi, 2000. Nitrite Curing of Meat. *The N-Nitrosamine Problem and Nitrite alternatives*, Food and Nutrition Press, Inc. Trumbull, CT.
6. Sleight, S.D., D.P. Sinha and M. Uzoukwu, 1972. Effect of sodium nitrate on reproductive performance of pregnant sows. *Journal of American Veterinary Medical Association*, 161: 819-823.
7. Swann, P.F., 1975. The toxicology of nitrate, nitrite and N-nitroso compounds. *Journal of the Science of Food and Agriculture*, 26: 1761-1770.
8. Agarwal, M.K., M. Iqbal and M. Athar, 2007. Garlic oil ameliorates ferric nitrilotriacetate (Fe-NTA)-induced damage and tumor promotion: implications for cancer prevention. *Journal of Food Chemistry and Toxicology*, 45: 1634-1640.
9. European Food Safety Authority "EFSA", 2003. Opinion of the scientific panel on biological hazards on the request from the Commission related to the effects of nitrites/nitrates on the microbiological safety of meat products. *The European Food Safety Authority "EFSA" Journal*, 14: 1-31.
10. Vinodini, N.A., A.K. Nayanatara, K.M. Damodaragawda, B. Ahamad and S. Shabarinath, 2008. Effect of monosodium glutamate-induced oxidative damage on rat testis. *Journal of Chinese Clinical Medicine*, 3: 370-373.
11. Alao, O.A., J.O. Ashaolu, O.K. Ghazal and V.O. Ukwenya, 2010. Histological and biochemical effects of monosodium glutamate on the frontal lobe of adult Wistar rats. *International Journal of Biomedical and Health Sciences*, 6: 197-203.
12. Schiffman, S.S., 2000. Intensification of sensory properties of foods for the elderly. *Journal of Nutrition*, 130: 927-930.
13. Bellisle, F., 2008. Experimental studies of food choices and palatability responses in European subjects exposed to the umami taste. *Asia Pacific Journal of Clinical Nutrition*, 17: 376-379.
14. Nagodawithana, T., 1992. Yeast-derived flavors and flavor enhancers and their probable mode of action. *Journal of Food Technology*, 11: 138-144.
15. Yoshida, Y., 1998. Umami taste and traditional seasoning. *Food Reviews International*, 14: 213-246.
16. Jinap, S. and P. Hajeb, 2010. Glutamate: its applications in food and contribution to health. *Appetite*, 55: 1-10.
17. Joint FAO/WHO Expert Committee on Food Additives "JECFA", 1988. L-glutamic acid and its ammonium, calcium, monosodium and potassium salts. *Toxicological Evaluation of Certain Food Additives and Contaminants 1988:97-161*. New York Cambridge University Press.
18. Eweka, A.O. and J.O. Adjene, 2007. Histological studies of the effects of monosodium glutamate on the medial geniculate body of adult Wistar rat. *Electronic Journal of Biomedicine*, 2: 9-13.
19. Blaylock, R., 2007. Food Additives: What you eat can kill you. *The Blaylock Wellness Report*, 4: 3-4.
20. Michelle Meadows, 2000. MSG: A Common Flavor Enhancer. *Consumer Magazine*, 2: 1-5.
21. Bhattacharya, T., A. Bhakta and S.K. Ghosh, 2011. Long term effect of monosodium glutamate in liver of albino mice after neonatal exposure. *Nepal Medical College Journal*, 13: 11-16.
22. Al-Mosaibih, M.A., 2013. Effects of monosodium glutamate and acrylamide on the liver tissue of adult Wistar rats. *Life Science Journal*, 10: 35-42.
23. Egyptian Organization for Standardization and Quality Control "EOS", 2005. *Egyptian Organization for Standardization and Quality Control for meat products*.
24. Nayel, M.S., 2013. Studies on chemical preservatives in meat products, Ph.D. Thesis (Meat Hygiene), Faculty of Veterinary Medicine, Benha University, Egypt.
25. Schumann, G. and R. Klauke, 2003. New IFCC reference procedures for the determination of catalytic activity concentrations of five enzymes in serum: preliminary upper reference limits obtained in hospitalized subjects. *Clinical Chimica Acta*, 327: 69-79.

26. Patton, C.J. and S.R. Crouch, 1977. Enzymatic determination of urea. *Analytical Chemistry*, 49: 464-469.
27. Henry, R.J., D.C. Cannon and J.W. Winkelman, 1974. *Clinical Chemistry, Principles & Techniques*, 2nd Ed. Harper & Row, pp: 943 - 949.
28. Bancroft, J.D. and A. Stevens, 1990. *Theory and practice of histopathological techniques*. Churchill Livingstone, London.
29. Bagheri, F., A. Gol, S. Dabiri and A. Javadi, 2011. Preventive effect of garlic juice on renal reperfusion injury. *Transplantation*, 5: 194-200.
30. Honikel, K.O., 2008. The use and control of nitrate and nitrite for the processing of meat products. *Journal of Meat Sciences*, 78: 68-76.
31. Esterbauer, H. and K. Cheeseman, 1990. Determination of aldehydic lipid peroxidation products: Malonaldehyde and 4-hydroxyonenal. *Journal of Enzymology*, 186: 407-421.
32. Smith, R.P., 1991. Chemicals reacting with various forms of hemoglobin: biological significance, mechanisms and determination. *Journal of Forensic Sciences*, 36: 662-672.
33. Reutov, V.P., E.G. Sorokina, V.G. Pinelis, T.S. Korshunova, A.A. Rodionov, V.B. Koshelev, S.M. Strukova, L.P. Kaiushin, P. Braquet and Komissarova, 1993. The compensatory - adaptive mechanisms in nitrite - induced hypoxia in rats. *Biulleten Eksperimentalnoi Biologii i Meditsiny*, 116: 506-508.
34. Vlaskina, S.G., G.F. Zhukova, I.A. Alekseeva, Y. Aleshko Ozhevskii and S.A. Khotimchenko, 1996. Effect of iron deficiency on the toxic action of nitrite and on the endogenous synthesis of N-nitrosamine. *Voprosy Pitaniya*, 6: 27-30.
35. Eman, G.E. Helal and Mervat Abdel-Rahman, 2005. Effect of food preservative and food coloring agent on some physiological and hematological parameters in albino rats and the protective role of garlic. *Egyptian Journal of Hospital Medicine*, 18: 116-123.
36. Atef, M., M.A.M. Abo-Noragr, M.S.M. Hanafy and A.E. Agag, 1991. Pharmacotoxicological aspects of nitrate and nitrite in domestic fowls. *British Poultry Science*, 32: 399-404.
37. Agata Wawrzyniak, 2000. Effect of diet with added pectin on rats given sodium nitrite. *Polish Journal of Food and Nutrition Sciences*, 9/50: 57-60.
38. Zovodnik, I.B., E.A. Iapshina, G. Barotsz and M. Bryszewska, 1999. Membrane effect of nitrite - induced oxidation of human red blood cells. *Biochimica et Biophysica Acta*, 1421: 306-316.
39. Saint-Blanquat, G., 1980. Aspects toxicologiques et nutritionnels des nitrates et des nitrites. *Ann. Nutr. Alim*, 34: 827-864.
40. Phillips, W.E.J., 1966. Effect of dietary nitrite on the liver storage of vitamin A in the rat. *Canadian Journal of Biochemistry*, 44: 1-7.
41. Bartosz, G., 2004. *Druga twarz tlenu. Wolne rodniki w przyrodzie* [The second face of oxygen. Free radicals in the nature]. Wyd. Nauk. PWN Warszawa.
42. Dudka, J., S. Szczepaniak, K. Dawidek-Pietryka and E. Kusmierzak, 1998. Wplyw ołowiu i azotynów na aktywność niektórych enzymów uczestniczących pośrednio w redukcji methemoglobiny u szczurów [Effect of lead and nitrites on the activity of some enzymes indirectly affecting methemoglobin reduction processes in rats]. *Bromat. Chem. Toksykol*, 31: 233-236.
43. Sadowska, J. and M. Kuchlewska, 2011. Effect of diet composition and mixture of selected food additives on the erythrocytic system and iron metabolism in peripheral blood of male rats. *Acta Scientiarum Polonorum Technologia Alimentaria*, 10: 497-506.
44. Folamar, L.C., 1993. Effects of chemical contaminants on blood chemistry of teleost fish: A bibliography and synopsis of selected effects. *Journal of Environmental Toxicology and Chemistry*, 12: 337-375.
45. Hanaa-Hassan, A., S. Hani, H.S. Hafez and E. Zeghebar-Fawzia, 2010. Garlic oil as a modulating agent for oxidative stress and neurotoxicity induced by sodium nitrite in male albino rats. *Journal of Food and Chemical Toxicology*, 48: 1980-1985.
46. Kalantari, H. and M. Salehi, 2001. The protective effect of garlic oil on hepatotoxicity induced by acetaminophen in mice and comparison with N-acetylcysteine. *Saudi Medical Journal*, 22: 1080-1084.
47. Duncan, J.R., K.W. Prasse and E.A. Mahaffey, 1994. *Veterinary Laboratory Medicine (Clinical pathology)*. 3rd Ed. Iowa State University Press.
48. Til, H.P., C.F. Kuper and H.E. Falke, 1997. Nitrite induced adrenal effects in rats and the consequences for the no-observed-effect level. *Food and Chemical Toxicology*, 35: 349-355.

49. Zurovsky, Y. and C. Haber, 1995. Antioxidants attenuate endotoxin-generation induced acute renal failure in rats. *Scandinavian Journal of Urology and Nephrology*, 29: 147-154.
50. Rybezynska, M., S. Hoffman and J. Goslar, 1996. Molecular changes in erythrocyte membranes induced by nitroimidazoles and radiation. *Polish Journal of Pharmacology*, 48: 269-280.
51. EL-Demerdash, F.M. and E.I. Elagamy, 1999. Biological effects in *Tilapia nilotica* fish as indicators of pollution by cadmium and mercury. *International Journal of Environmental Health Research*, 9: 143-156.
52. Suresh, A., B. Sivaramakishna, P.O. Victoriamma and K. Radhakrishnah, 1992. Comparative study on the inhibition of acetylcholinesterase activity in the fresh water fish *Cyprinus carpio* by mercury and zink. *Biochemistry International*, 26: 367-375.
53. Jevtovic-Todorovic, V., J. Beals, N. Benshoff and J. Olney, 2003. Prolonged exposure to inhalational anesthetic nitrous oxide kills neurons in adult rat brain. *Neuroscience*, 122: 609-616.
54. Perry, C., H. Peretz, O. Ben-Tal and A. Eldor, 1997. Highly elevated lactate dehydrogenase level in a healthy individual: a case of macro-LDH. *American Journal of Hematology*, 55: 39-40.
55. Grant, D. and W.H. Butler, 1989. Chronic toxicity of sodium nitrite in the male F344 rat. *Food Chemistry and Toxicology*, 27: 565-571.
56. El-Wakf, A.M., H.A. Hassan, F.G. El-Said and A. El-Said, 2009. The association between nitrite contamination of drinking water and changes in methemoglobin level other hematological parameters in young and adult male rats. *Journal of Egyptian Society of Toxicology*, 49: 91-96.
57. Freeman, M., 2006. Reconsidering the effects of monosodium glutamate. A literature review. *Journal American Academy of Nurse Practitioners*, 18: 482-486.
58. Ashaolu, J.O., V.O. Ukwenya, A.B. Okonoboh, O.K. Ghazal and A.A.G. Jimoh, 2011. Effect of monosodium glutamate on hematological parameters in Wistar rats. *International Journal of Medicine and Medical Sciences*, 3: 219-222.
59. Meraiyebu, A., C.O. Akintayo, A.C. Uzoechi and S. Okere, 2012. The effects of orally administered monosodium glutamate (MSG) on blood thrombocyte, blood coagulation and bleeding in rats. *IOSR Journal of Pharmacy and Biological Sciences*, 4: 4-8.
60. Kolawole, O.T., 2013. Assessment of the effects of monosodium glutamate on some biochemical and hematological parameters in adult wistar rats. *American Journal of Bio. Science*, 1: 11-15.
61. Ortiz, G.G., O.K. Bitzer-Quintero, C. Beas Zárate, C. Rodríguez-Reynoso, F. Larios-Arceo, I.E. Velázquez-Brizuela, F. Pacheco-Moisés and S.A. Rosales-Corral, 2006. Monosodium glutamate-induced damage in liver and kidney: a morphological and biochemical approach. *Biomedicine and Pharmacotherapy*, 60: 86-91.
62. Singh, K. and P. Ahluwalia, 2003. Studies on the effect of Monosodium Glutamate (MSG) administration on some antioxidant enzymes in the arterial tissue of adult male mice. *Journal of Nutritional Science and Vitaminology (Tokyo)*, 49: 145.
63. Cheville, N.F., 2009. Ultrastructural pathology: The comparative cellular basis of disease. 2nd Ed. Wiley-Blackwell. A John Wiley of Sons, Inc. USA.
64. Abdel Hameed, T.F., 2004. Light and electron microscopic studies on the effect of orally administered formalin on liver and kidney of guinea pig. *Journal of the Egyptian German Society of Zoology C. Histology and Histochemistry*, 45: 203-224.
65. Bopanna, K., R. Balaraman and R. Noding, 1998. Antioxidant status of S-allyl cysteinesulphoxide on monosodium glutamate potentiated atherosclerosis. *Indian Journal of Pharmacology*, 30: 73-81.
66. Narayanan, S., R. Kumar, J. Paval and S. Nayak, 2010. Effect of ascorbic acid on the monosodium glutamate-induced neurobehavioural changes in periadolescent rats. *Bratisl Lek Listy*, 111: 247-252.
67. Farombi, M. and O. Onyema, 2006. Monosodium glutamate-induced oxidative damage and genotoxicity in the rat: modulatory role of vitamin C, vitamin E and quercetin. *Human and Experimental Toxicology*, 25: 251-259.
68. Brace, H., M. Latimer and P. Winn, 1997. Neurotoxicity, blood-brain barrier breakdown, demyelination and remyelination associated with NMDA-induced lesion of the rat lateral hypothalamus. *Brain Research Bulletin*, 43: 447-455.

69. Samuels, A., 1999. The Toxicity/Safety of MSG, A study in suppression of information. *Accountability in Research*, 6: 259-310.
70. Walker, R. and J. Lupien, 2000. The safety evaluation of monosodium glutamate. *Journal of Nutrition*, 130(4): 1049-105).
71. Moore, K.L., 2003. Congenital malformations due to environment in developing humans. 2nd Ed. Philadelphia, W.B. Saunders Co. Ltd. pp: 173-183.