



# Trial for Induced Spawning with Histological Evaluation of Gonads in Forskal Catfish *Bagrus bayad*

Mahamoud Abdelghaffar Emam<sup>1\*</sup>, Ashraf Mohamed Abd El-latif<sup>1</sup>, Ehab El-Zoghby<sup>1</sup>, Badia Abugherin<sup>2</sup> and Ramy Shourbela<sup>3</sup>

<sup>1</sup>Histology and Cytology Department, Faculty of Veterinary Medicine, Benha University, Egypt

<sup>2</sup>Department of Anatomy and Histology, Faculty of Veterinary Medicine, Tripoli University, Libya

<sup>3</sup>Department of Animal Husbandry and Animal Wealth Development, (Fish breeding and production), Faculty of Veterinary Medicine, Alex University, Egypt

## Abstract

Sixteen mature *Bagrus bayad* of both sexes (8 males and 8 females) were used to evaluate a trial for spawning induction using GnRH<sub>a</sub> combined with domperidone during spawning season. The experiment was carried out during May 2014. Fish were divided into 4 groups. Group 1 was control males and females *Bagrus bayad*, group 2 was females *Bagrus bayad* treated with dopamine antagonist (domperidone) alone, group 3 was females *Bagrus bayad* treated with domperidone and gonadotropin releasing hormone analogue (GnRH<sub>a</sub>), and group 4 was males *Bagrus bayad* treated with GnRH<sub>a</sub> and domperidone. GnRH<sub>a</sub> was used in the present study by dose of 20 ug/kg and dose of domperidone was 5 mg/kg. Results showed that no evidence for spawning induction. Histological examination showed structural changes in ovary of group 2 and group 3 in comparison to control ovary but there was no obvious change between structure of ovary of group 2 and group 3. In addition to, histological examination showed structural changes in the testis of group 4 in comparison to control testis. In both sex, histological examination did not identify any evidence of spawning. In conclusion, domperidone and/or GnRH<sub>a</sub> induced histological changes in the ovary and testis of *Bagrus bayad* but did not induce spawning. Further studies are needed with higher doses of GnRH<sub>a</sub> and/or carrying out the experiment during mid or late spawning season of *Bagrus bayad*.

**Keywords:** Forskal catfish; *Bagrus bayad*; Gonadotropin releasing hormone analogue (GnRH<sub>a</sub>); Domperidone; Ovary; Testis

## Introduction

*Bagrus bayad* is an important food fish, with flesh of good eating and of economic importance in Egypt. It has high growth rate and attains maximum weight of 12.5 kg [1] but it has not been cultured due to limited information on their breeding under captivity as they, like other catfishes, are not able to exhibit their natural spawning behavior in artificial ponds [2].

Recently, gonadotropin releasing hormone analogue (GnRH<sub>a</sub>) has been used in the induced spawning of several fish species [3,4]. In some fish, dopamine prevents the release of gonadotropin through inhibitory actions on the pituitary gland, thereby making it impossible for fish to ovulate and spawn. Pre-spawn fish exhibiting dopaminergic tone were induced to spawn using GnRH<sub>a</sub> in combination with dopamine antagonist like domperidone. Successful spawning of some fish species were achieved using GnRH<sub>a</sub> and domperidone [5,6].

Therefore, the aim of this study was to investigate the effects of using gonadotropin releasing hormone analogues (GnRH<sub>a</sub>) with dopamine antagonists (domperidone) on the spawning performance of *Bagrus bayad* and to evaluate the histological changes occurred in the gonads in relation to the hormonal treatment.

## Materials and Methods

### Experimental location

The present study was conducted in the Fish Hatchery, Central Laboratory for Aquaculture Research, Abbassa, Sharkia, Egypt during natural early spawning season (May 2014) but at that time, there was long winter.

### Experimental design

16 apparently healthy *Bagrus bayad* were divided into 4 randomly groups as shown in Table 1. The fish were selected from over-wintered

*Bagrus bayad* maintained in 6.5 faddan (4200 m) earthen ponds that had 1.5 meter depth. In late days of March, the spawners were caught, selected, sexed and transferred to a quarter faddan ponds with 1 meter depth. At the early spawning season, female *Bagrus bayad* which showed spawning signs were randomly divided and distributed in tanks of 3.5 cubic meter volume filled with water supplied with aerator. After acclimatization of fish for 24 hrs. The water quality was measured daily according to Dewis and Freiles [7]. The fish were maintained in the laboratory under normal photoperiod (12.5 h L: 11.5 h D) and temperature (20 ± 4°C). The fish were subjected to the treatments as in Table 1. In general, injection was intra-peritoneal and the second dose injected after 8 hours from the first dose.

Group	Number of fish	Drugs	1st Dose	2nd Dose after 8 hrs	Specimens for histology
Group 1	2 females 2 males	-----	-----	-----	Testes and ovaries at zero time
Group 2	3 females	DOM	5 mg/kg		Ovaries after 8 hrs
Group 3	3 females	GnRH <sub>a</sub> DOM	2 ug/kg 5 mg/kg	18 ug/kg 2.5 mg/kg	Ovaries after 11 hrs from 2 <sup>nd</sup> dose
Group 4	6 males	GnRH <sub>a</sub> DOM	20 ug/kg 5 mg/kg	-----	Testes after 18 hrs

DOM: Domperidone; GnRH<sub>a</sub>: Cystorelin

Table 1: Showing GnRH<sub>a</sub> and domperidone doses used in different groups.

\*Corresponding author: Mahamoud Abdelghaffar Emam, Histology and Cytology Department, Faculty of Veterinary Medicine, Benha University, Egypt, E-mail: [mahmoud.hussein@fvfm.bu.edu.eg](mailto:mahmoud.hussein@fvfm.bu.edu.eg)

Received January 31, 2015; Accepted March 31, 2015; Published April 01, 2015

Citation: Emam MA, El-latif AMA, El-Zoghby E, Abugherin B, Shourbela R (2015) Trial for Induced Spawning with Histological Evaluation of Gonads in Forskal Calfish *Bagrus bayad*. Fish Aquac J 6: 120. doi:10.4172/2150-3508.1000120

Copyright: © 2015 Emam MA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Preparation of GnRHa and dopamine antagonists

**Gonadotropin releasing hormone analogue GnRHa:** Cystorelin® (vail 20 ml) each 1 ml contains 50 ug synthetic Gonadorelin (GnRH) diacetate tetrahydrate manufactured by (CEVA SANTLE ANIMAL – France).

**Domperidone tablets:** Each tablet contains 10 mg from the active principle (under trade name Gastromotil® manufactured by EIPICO Company, Cairo, A. R. E.) were powdered and then re-suspended in physiological saline 0.9% when needed according to Alok [8].

## Histological examination

Specimens from ovaries and testis of all groups of *Bagrus bayad* were taken and fixed in Bouin's fluid for 24 hr. Specimens were dehydrated in alcohols, cleared in xylene and blocked in paraffin. Sections of 5 µm were cut and stained with hematoxylin and eosin stain, Masson's trichrome stain, Alcian blue stain and periodic acid Schiff technique. Fixation and staining methods and techniques were carried out according to Bancroft [9].


## Results

Both sex of *Bagrus bayad* in the current study do not show noticeable spawning after injection of GnRHa (20 ug/kg) and domperidone (5 mg/kg).

Histologically, the ovary of control *Bagrus bayad* (in the early spawning season) was surrounded by connective tissue capsule known as tunica albuginea which is measured 209 µm thickness (Figure 1A). The tunica albuginea is fibromuscular layer as it consists of dense collagenous connective tissue which contains outer longitudinal and inner circular smooth muscle layers (Figure 1B). From tunica albuginea, collagenous ovigerous lamellae are extended into the ovary and enclose different developmental follicular stages; previtellogenic (immature), vitellogenic, and postvitellogenic (mature), with predominance of previtellogenic ones (Figure 1C). Vitellogenic follicles show alcianophilic yolk granules (Figure 1D) while postvitellogenic follicles show non alcianophilic yolk globules (Figure 1E) but yolk globules show PAS positive reaction (Figure 1F). The follicular epithelium shows PAS and AB positive (Figures 1E and F).

In group 2 which treated with domperidone alone, the ovarian structure of *Bagrus bayad* has clearly changed. Tunica albuginea becomes thicker and is measured 294 µm. The vitellogenic and postvitellogenic follicles are relatively increased in number, with predominance of postvitellogenic (mature) follicles (Figure 1G). Moreover, all vitellogenic and postvitellogenic follicles shows PAS positive yolk granules, yolk globules and follicular epithelium (Figure 1H). In group 3 which treated with combination of domperidone and GnRHa, ovaries show no obvious changes in comparison to ovaries of group 2 except slightly increase in the number of postvitellogenic follicles per field (Figure 1I).

The testis of control *Bagrus bayad* is covered with connective tissue capsule known as tunica albuginea that is measured 155 µm and it is divided into several testicular lobules (Figure 2A). Tunica albuginea consists of dense collagenous connective tissues with (Figure 2B). From tunica albuginea, interlobular connective tissues are arisen and collected in between seminiferous lobules (Figure 2C). The basement membranes of the seminiferous lobules show strong PAS positive (Figure 2D). There are all different developmental stages from spermatogonia to spermatozoa (Figures 2A-D). In group 4 which treated with combination of domperidone and GnRHa, tunica



**Figures 1 and 2:** (1A) Photomicrograph of control ovary of *Bagrus bayad* showing tunica albuginea (arrow) and different developmental stages of eggs. H&E stain. (1B) Photomicrograph of ovary of *Bagrus bayad* showing muscular layers in tunica albuginea. Outer longitudinal (arrow) and inner circular smooth muscle layers (arrowhead). Masson's trichrome stain. (1C) Photomicrograph of ovary of *Bagrus bayad* showing collagen fibers in the ovigerous lamellae (arrows). Masson's trichrome stain. (1D) Photomicrograph of vitellogenic follicle in ovary of *Bagrus bayad* showing alcianophilic yolk granules. Alcian blue stain. (1E) Photomicrograph of postvitellogenic follicle in ovary of *Bagrus bayad* showing alcian blue negative yolk globules (arrowhead) but alcianophilic follicular epithelium (arrow). Alcian blue stain. (1F) Photomicrograph of postvitellogenic follicle in ovary of *Bagrus bayad* showing PAS positive yolk globules (arrow) and follicular epithelium. PAS technique. (1G) Photomicrograph of ovary of *Bagrus bayad* treated with domperidone showing increase number of vitellogenic (short arrow) and postvitellogenic follicles (long arrow) in comparison to control ovary. Compare to (1A) H&E. (1H) Photomicrograph of ovary of *Bagrus bayad* treated with domperidone showing PAS positive yolk granules and globules in the vitellogenic and postvitellogenic follicles respectively. PAS technique. (1I) Photomicrograph of ovary of *Bagrus bayad* treated with domperidone and GnRHa showing thicker tunica albuginea in comparison to (1A) and (1H). PAS technique. (2A) Photomicrograph of control testis of *Bagrus bayad* showing tunica albuginea (short arrow) and many seminiferous lobules (arrows). H&E stain. (2B) Photomicrograph of control testis of *Bagrus bayad* showing that tunica albuginea consists of dense collagenous connective tissue (arrow). Masson's trichrome stain. (2C) Photomicrograph of control testis of *Bagrus bayad* showing collagen fibers in the interlobular connective tissue (arrow). Masson's trichrome stain. (2D) Photomicrograph of control testis of *Bagrus bayad* showing PAS positive basement membranes (arrows) of the seminiferous lobules. PAS technique. (2E) Photomicrograph of testis of *Bagrus bayad* treated with domperidone and GnRHa showing spermatid cyst (arrow) and spermatozoa (arrowhead). Masson's trichrome stain. (2F) Photomicrograph of testis of *Bagrus bayad* treated with domperidone and GnRHa showing increase of interlobular connective tissue. Masson's trichrome stain.

albuginea becomes thicker and is measured 165  $\mu\text{m}$  in addition to, cysts of spermatid and spermatozoa are increased in number in comparison to control testis (Figure 2E). Moreover, the interlobular connective tissues are increased (Figure 2F).

## Discussion

The effect of GnRH in induction of fish spawning is controversial. Our results demonstrated that GnRHa (20  $\mu\text{g}/\text{kg}$ ) in combination with domperidone (5  $\text{mg}/\text{kg}$ ) was not sufficient to induce spawning during early spawning season of *Bagrus bayad*. However, Sahoo et al. [6] reported that ovaprim (sGnRHa (20  $\mu\text{g}/\text{kg}$ ) and domperidone (10  $\text{mg}/\text{kg}$ ) was ideal for induction of spawning in the Asian catfish. On the contrary, Tan-Fermin et al. [10] stated that 50  $\mu\text{g}/\text{kg}$  BW GnRHa could not induce ovulation in the Asian catfish. On the other hand, Silverstein et al. [11] recorded that 100  $\mu\text{g}/\text{kg}$  GnRHa was used in channel catfish *Ictalurus punctatus*, more hatching rate was recorded.

Histologically, the ovary of *Bagrus bayad* was consisted of tunica albuginea and different developmental follicular stages (previtellogenic, vitellogenic, and postvitellogenic stages) that was similar to ovarian structure in most of teleost fish as tilapia [12-16]. The ovarian tunica albuginea of *Bagrus bayad* was supported by muscular layer which differentiated into an outer circular layer and inner longitudinal layer as reported in *Bagrus bayad* by Khallaf et al. [17] and Gaber [18], while Gaber [18] did not differentiate two muscular layers in the ovarian wall of *Bagrus docmac*.

The testis of *Bagrus bayad*, like most of teleost fish, was of the lobular type as it was consisted of anastomosing seminiferous lobules which were separated by connective tissue stroma as stated by (Abd El-Aziz [19]; Mousa [12]; El-Gohary [13]) in tilapia and (Oteme et al. [20]; Emam and Abugherin [16]) in catfish. Spermatogenic cells were in synchronous development inside cysts in the testicular lobules of *Bagrus bayad* that was similar to testicular structure of teleost fish (Abd El-Aziz [19]; Schulz et al. [21]; El-Zoghby et al. [15]; Emam and Abugherin [16]). In the present study, the spermatozoa begun to appear within lumen of testicular lobules in group 4 after injections of domperidone and GnRHa in comparison to testis of group 1. The testis of group 4 did not show any signs for spawning where the spent testis or post-spawning testis should show some empty testicular lobules indicating spawned testes as they discharge their content during spawning as reported by Resink et al. [22] in *Clarias gariepinus*, Gaber [18] in *Bagrus bayad* and El-Zoghby et al. [15] and Emam and Abugherin [16] in *Clarias lazera*. In addition to, our results revealed relatively decrease the number of previtellogenic (immature) follicles and increase number of postvitellogenic (mature) follicles in groups 2 and 3 after injection of domperidone and/or GnRHa in comparison to ovary of group 1 that supports the findings of Chowdhury et al. [23] in the Indian catfish indicating potential role of GnRHa in induction of spawning but it may be of dose dependent. But there were no signs for spawning where the spawned ovary should show postovulatory and atretic follicles as mentioned by Gaber [18] in *Bagrus bayad*, and El-Zoghby et al. [15] and Emam and Abugherin [16] in *Clarias lazera*.

## Conclusion

From all previous findings, we concluded that GnRHa (20  $\mu\text{g}/\text{kg}$ ) in combination with domperidone (5  $\text{mg}/\text{kg}$ ) is not sufficient to induce spawning during early spawning season of *Bagrus bayad*. Therefore, a further study is needed for the use of higher doses of GnRHa and/or doing the experiment during mid or late spawning season of *Bagrus bayad*.

## References

1. Abdel-Latif AF (1974) Fisheries of Lake Nasser. Aswan regional planning, Lake Nasser Development Centre, Aswan: 235.
2. Lamai SL (1993) Aspects of the applied biology of *Clarias gariepinus*, aquacultural techniques and Deldrin toxicity. Ph.D. Thesis. Dept. of Pure and Applied Zoology, University of Reading.
3. Zohar Y, Mylonas CC (2001) Endocrine manipulation in cultured fish: From hormones to genes. *Aquaculture* 197: 99-136.
4. Rainis S, Ballestrazzi R (2005) The control of reproduction in finfish species through GnRH treatments. *Italian Journal of Animal Science* 4: 345-353.
5. Manosroi J, Meng-Umpfan K, Manosroi A (2004) Maturation induction of *Pangasius hypophthalmus* using gonadotropin releasing hormone analogue (GnRHa) in combination with domperidone, in oil suspension dosage forms. *Asian Fish Sci* 17: 39-49.
6. Sahoo SK, Giri SS, Sahu AK (2005) Induced spawning of Asian catfish *Clarias batrachus* (Linn.): Effects of various latency periods and sGnRHa and domperidone doses on spawning performance and egg quality. *Aquac Res* 36: 1273-1278.
7. Dewis J, Freiles T (1970) Physical and chemical methods of soil and water analysis. FAO soils. Bulletin 10: Rome.
8. Alok D, Pillai D, Garg LC (1997) Effect of d-Lys6 salmon sGnRH alone and in combination with domperidone on the spawning of common carp during the late spawning season. *Aquaculture International* 5: 369-374.
9. Bancroft JD, Stevens A, Turner DR (1996) Theory and practice of histological techniques. 4th Ed. Churchill Living stone, New York, London, San Francisco, Tokyo.
10. Tan-Fermin JD, Pagador RR, Chavez RC (1997) LHRHa and pimozide-induced spawning of Asian catfish *Clarias macrocephalus* at different times during an annual reproductive cycle. *Aquaculture* 148: 323-331.
11. Silverstein JT, Bosworth BG, Wolters WR (1999) Evaluation of dual injection of LHRHa and the dopamine receptor antagonist pimozide in cage spawning of channel catfish *Ictalurus punctatus*. *Journal of the World Aquaculture Society*, 30: 263-268.
12. Mousa AM (1998) Immunocytochemical and Histological studies on the reproductive endocrine glands of the Nile tilapia, *Oreochromis niloticus* (Teleostei, cichlidae). *J Egypt Ger Soc Zool* 27: 109-134.
13. El-Gohary NMA (2001) The effect of water quality on the reproductive biology of the Nile tilapia, *Oreochromis niloticus* in Lake Manzalah. Ph.D. Thesis. Faculty of Science, Ain Shams University.
14. El-Saba A, Abd Rabou MI, El-Sakhawy MI, El-Shammaa MA, Hussein SM (2013) Seasonal changes in the histology of the ovaries of Nile tilapia (*Oreochromis niloticus*). *J Vet Anat* 6: 1-21.
15. El-Zoghby, IMA, Bakry HH, Ghallab AM, Emam MA (2009) Histological studies on the gonads of the catfish during different seasons. *Lucrari Stiintifice, Medicina Veterinara* 52: 352-362.
16. Emam MA, Abugherin B (2014) Seasonal histological changes in gonads of the catfish *Clarias lazera*. *Fish Aquac J* 5: 87-91.
17. Khallaf EA, El-Saadany MM, Authman M (1991) Oogenesis of *Bagrus bayad* (Forsk.). *J Egypt Ger Soc Zool* 4: 1-4.
18. Gaber SAO (2000) Biological, Histological and Histochemical studies on the Reproductive Organs and Pituitary gland of *Bagrus docmac* and *Bagrus bayad* in the Nile water, with special reference to the Ultrastructure of supporting tissues. Ph.D. Thesis. Faculty of Science, Zagazig University.
19. Abd El-Aziz M (1992) An ultrastructure study on the spermatogenesis of *Oreochromis niloticus* (Tilapia) fish during winter season. *Egypt J Histol* 15: 269-281.
20. Oteme ZJ, Rodriguez JN, Kouassi CK, Hen S, Agnese JF (1996) Testicular structure, spermatogenesis and sperm cryopreservation in the African clariid catfish, *Heterobranchus longifiliis* (Vaenciennes, 1840). *Aquac Res* 31: 805-813.
21. Schulz RW, Menting S, Bogerd J, Franca LR, Vilela DA, et al. (2005) Sertoli cell proliferation in the adult testis - evidence from two fish species belonging to different orders. *Biol Reprod* 73: 891-898.

22. Resink JW, Schonen WG, Van Den Hurk R, Viveen WJ, Lambert JG (1987) Seasonal changes in steroid metabolism in the male reproductive system of the African catfish, *Clarias gariepinus*. Aquaculture 63: 59-76.
23. Chowdhury M, Chatterjee A, Mondal A, Chatterji U (2010) Ovaprim Abrogates Expression of GnRH Receptor-II in the Indian Catfish. Inter J Biol 2: 189-198.

**Citation:** Emam MA, El-latif AMA, El-Zoghby E, Abugherin B, Shourbela R (2015) Trial for Induced Spawning with Histological Evaluation of Gonads in Forskal Catfish *Bagrus bayad*. Fish Aquac J 6: 120. doi:[10.4172/2150-3508.1000120](https://doi.org/10.4172/2150-3508.1000120)

### Submit your next manuscript and get advantages of OMICS Group submissions

#### Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

#### Special features:

- 400 Open Access Journals
- 30,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submission>

