



Effects of an intravaginal GnRH analogue administration on rabbit reproductive parameters and welfare



C. Munari ^{a,*}, P. Ponzio ^a, A.R. Alkhwawah ^b, A. Schiavone ^a, C. Mugnai ^a

^a Department of Veterinary Sciences, University of Turin, Largo Braccini 2, 10095, Grugliasco, TO, Italy

^b Theriogenology Department, Faculty of Veterinary Medicine, Benha University, Egypt

ARTICLE INFO

Article history:

Received 1 June 2018

Received in revised form

26 October 2018

Accepted 27 October 2018

Available online 31 October 2018

Keywords:

Rabbit

Fertility

Welfare

Hormonal administration method

Semen characteristics

ABSTRACT

On commercial farms, rabbit does are subjected to a reproductive rhythm that does not account for their welfare or physiology, leading to reduced longevity and consequently high annual replacement. The European Food Safety Authority (EFSA) recommends limited and infrequent use of hormone treatments, and suggests replacement with alternative methods that do not threaten animal welfare when possible. In the present study, we aimed to determine whether the GnRH analogue lecorelin acetate could be administered by inclusion in the seminal dose during insemination. Twenty 9-month-old does (Grigio del Monferrato, autochthonous Italian breed), each having two previous deliveries, were individually housed and divided into two groups at artificial insemination. The control group received 0.2 mL of intramuscular lecorelin (Dalmarelin, Fatro[®], Italy) prior to insemination. The intravaginal group was inseminated with a seminal dose that included 0.3 mL Dalmarelin. The experiment was performed for six consecutive reproductive cycles at 42-day intervals, and included a total of 120 inseminations. Prior to each insemination, the heterospermic pooled semen samples were assessed for sperm motility and morphology. Each ejaculate was divided into two samples, with and without lecorelin addition. Compared to the control group, the does with intravaginal Dalmarelin administration showed equal or greater sexual receptivity, which resulted in a higher fertility rate over increasing cycles. The seminal dose volume was very low, possibly explaining the better results in the intravaginal group, which received a similar amount of hormone (0.3 mL/doe) as the control group (0.2 mL/doe). The negative performance of the control group may have also been due to anti-GnRH formation, and the more stressful method of ovulation induction. The number of live-born kits did not significantly differ between groups. Progressive motility was significantly positively correlated with motility characteristics, including VAP, VSL, ALH, BCF, STR, and LIN. Overall, our present findings supported that the incorporation of GnRH in a seminal dose could be used for ovulation induction in rabbit does. Further studies should identify the optimal dose of GnRH for intravaginal administration, taking into account that the intravaginal absorption capacity is about 10 times smaller than the intramuscular absorption capacity.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

On European farms, artificial insemination (AI) of rabbits has become a consolidated practice due to its ability to optimize human resources and increase the animals' reproductive performance [1]. Unlike in other species, ovulation in rabbits does not occur spontaneously but rather must be induced via a neuro-hormonal reflex, which is produced by mating under natural conditions [2]. Since males are absent during the practice of AI, ovulation must be

artificially induced. However, some of the exogenous hormonal substances used for this purpose may negatively affect rabbit welfare and cause fertility disorders in does [3]. For example, routine use of equine chorionic gonadotrophin (eCG) is associated with reduced fertility, represented by a decreased conception rate, high number of animals per litter, large number of stillbirths, and increased occurrence of hemorrhagic follicles [4–8] due to immunogenicity [9].

In field practice, the most frequently used method is the intramuscular administration of gonadotropin-releasing hormone (GnRH) or its synthetic analogues. Previous studies demonstrate that some of these hormones, such as gonadorelin or busarelin,

* Corresponding author.

E-mail address: chiara.munari@unito.it (C. Munari).

induce ovulation to the same level as a natural mating [10]. Unfortunately, GnRH and its synthetic analogues directly affect ovarian functions, influencing does' oocytes maturation both *in vivo* and *in vitro*, and altering the intra-follicular environment [11,12]. Additionally, higher doses of GnRH negatively affect rabbit welfare, as represented by a significantly increased number of rabbits per litter [3,13]. Larger litters are associated with decreased birth weights, and greater risk of doe exhaustion during pregnancy, which adversely affects both lactating and primiparous does. High doses of intramuscular GnRH are also associated with a high incidence of abortions due to the greater number of developing fetuses in the uterus [3]. Moreover, hormonal treatments, particularly when repeatedly used in rabbit does, are generally followed by decreased fertility due to the appearance of plasmatic anti-GnRH antibodies [14]. Canali et al. [4] concluded that in nulliparous does, the immune response to eCG begins to significantly decrease fertility after the third treatment, with a strong increase of anti-eCG antibody from the fourth treatment onward.

Commercial farms also subject rabbit does to a reproductive rhythm that does not take account for their welfare and physiology, thus reducing longevity [15], and necessitating high annual replacement. Rabbit does are routinely administered intramuscular treatment, but this process is invasive and best practice guidelines are not well developed [16]. The European Food Safety Authority (EFSA) [17] currently recommends that hormone treatments be used in a limited manner, as infrequently as possible, and that they be replaced, if possible, with alternative methods having no animal welfare consequences. The available evidence suggests a need to improve the welfare and fertility parameters of rabbit does [16] through the development of new reproductive strategies to induce doe ovulation, avoiding the traumatic event of intramuscular administration.

Several authors recently demonstrated ovulation induction by including different GnRH analogues in the seminal dose, and administering them through vaginal absorption [13,19]. Zhang and Qin [20] confirmed that inclusion of the GnRH analogue leuprorelin in the seminal dose induced doe ovulation, and led to the same reproductive performance obtained with the intramuscular method. This technique reduces stress for the animal, improving the welfare condition [10,21]. Moreover, GnRH intravaginal administration could be beneficial for farmers, avoiding potential mistakes derived from incorrect hormone administration and reducing the time spent on each AI [13].

In the present study, we aimed to investigate possible alternative methods for ovulation induction in rabbit does. Our goals were to reduce animal stress, avoid the negative effects of intramuscular treatment on ovary function, and minimize the annual replacement of does by improving reproductive performance and welfare.

2. Materials and methods

2.1. Animals, housing conditions, experimental groups, and feeding

Our study was conducted at the Cascina Campora commercial farm in Buttigliera d' Asti, Italy. Rabbits were raised with daily control of the environmental temperature (15–28 °C) and relative humidity (60–75%). The building was artificially ventilated (0.3 m/sec). The animals received a commercial diet containing 16.9% crude protein, 14.2% crude fiber, and 3.5% fat. Food and water were provided *ad libitum* throughout the experimental period. All animal experiments were managed in accordance with the Turin University Bioethics Committee recommendations (Prot. N° 256053 of 4/07/2017).

The experimental animals included 20 9-month-old rabbit does, each having two previous deliveries (Grigio del Monferrato,

autochthonous Italian breed). These animals were single-cage housed and divided into two experimental groups. The control group (C; n = 10) received 0.2 mL of intramuscular leirelin acetate (Dalmarelin, Fatro®, Italy), and was then inseminated with normal extended semen. The intra-vaginal group (IV; n = 10) was inseminated with an extended seminal dose including 0.3 mL leirelin acetate [19]. All does underwent AI for six consecutive cycles, using a seminal dose containing 10 ± 1 million spermatozoa in 0.5 mL of diluent [22]. No estrus synchronization was performed.

2.2. Reproductive performance

The does were managed according to a cycled production system with a 42-day interval and a 16-h light/8-h dark/light program according to Mousa-Balabel et al. [23]. The experimental phase was carried out from October 2017 to May 2018. We recorded the following reproductive traits at A.L.: sexual receptivity (vulva color and turgescency; a doe was deemed receptive when its vulva was red or purple and turgid), fertility rate (kindling/inseminations \times 100), and number of live-born kits [24].

2.3. Semen collection and evaluation

For each reproductive cycle, one ejaculate per male (n = 3) was collected early in the morning on a single day using an artificial vagina [25]. Only ejaculates exhibiting a white color were used, and if the gel was present, it was removed. Ejaculates with over 70% motile sperm were pooled (54 heterospermic pooled semen) and used for AI [26]. Each ejaculate pool was divided into two samples, which were diluted 1:5 using Galap (IMV Technologies, L'Aigle, France) [27], with addition of Dalmarelin (IV group) or without any additional hormone (C group). The solvent of Dalmarelin (physiological solution 0.9% NaCl) did not affect sperm quality [28]. Samples were assessed for motility characteristics, sperm viability, and acrosome status (Table 1).

2.4. Assessment of semen motility characteristics, sperm livability, and acrosome status

Sperm motility and motility characteristics at 37 °C were evaluated using a computer-assisted sperm analyzer (CASA; Hamilton Thorne, Inc., Beverly, MA, USA) with a 10 \times objective. A 10- μ L specimen of diluted semen was put on a pre-warmed Mackler slide and evaluated. Motility values were recorded as the percentages of

Table 1
Sperm characteristics in two different experimental groups of rabbit semen.

Morphological characteristics of sperms	Group	
	Control (C)	Intravaginal (IV)
Live sperm (%)	74.55 \pm 2.31	72.00 \pm 1.95
Sperm with intact acrosome (%)	78.65 \pm 1.92	79.00 \pm 1.81
Computer-assisted semen analysis for motility (CASA)		
VAP (μ m/s)	106.51 \pm 0.90	99.55 \pm 0.85*
VSL (μ m/s)	86.71 \pm 1.23	76.81 \pm 1.22*
VCL (μ m/s)	167.96 \pm 0.79	171.37 \pm 1.08*
ALH (μ m/s)	5.95 \pm 0.09	6.63 \pm 0.10*
BCF (Hz)	38.75 \pm 0.39	36.11 \pm 0.44*
STR (%)	79.76 \pm 0.62	75.65 \pm 0.75*
LIN (%)	52.99 \pm 0.85	46.30 \pm 0.86*
Total motility (%)	60.83 \pm 1.24	60.01 \pm 1.32
Progressive motility (%)	43.31 \pm 1.24	38.34 \pm 1.29*

Based on means \pm SE of 54 evaluated heterospermic pooled semen samples. VAP = average path velocity (μ m/s); VSL = straight-line velocity (μ m/s); VCL = curvilinear velocity (μ m/s); ALH = amplitude of lateral head displacement (μ m/s); BCF = beat cross-frequency (Hz); STR = straightness (%); LIN = linearity (%).*: significance at P < 0.01.

progressive motility (P. MOTIL, percentage of sperm exhibiting an actual space gain motility) and total motile sperm cells (TOTAL MOTIL). Additionally, based on the frequency with which the sperm track crossed the cell path in either direction and in changeable tracks [29,30], we calculated the average path velocity (VAP, $\mu\text{m/s}$), straight linear velocity (VSL, $\mu\text{m/s}$), curvilinear velocity (VCL, $\mu\text{m/s}$), amplitude of lateral head displacement (ALH, $\mu\text{m/s}$), linearity index (LIN, average value of the VSL/VCL ratio, %), straightness index (STR, average value of the VSL/VAP ratio, %), and beat-cross frequency (BCF, Hz).

Sperm viability and acrosome status (RSA) were evaluated using the Trypan blue/Giemsa dual staining technique, as previously described [31,32]. Trypan blue was used first to differentiate live from dead spermatozoa. Then the dried smears were fixed in 37% formaldehyde and stained with Giemsa for acrosome evaluation using an Advanced Automated Research Microscope System (Nikon Eclipse E200, phase contrast at 40 and 100 magnifications). We counted at least 200 sperm cells for each group. Acrosome-intact live (AIL) spermatozoa were differentiated based on staining characteristics. Only sperm displaying both head and tail were recorded as viable, while those with only either the head or the tail were considered unviable.

2.5. Statistical analysis

Statistical analyses were performed using SPSS statistical package version 16 (SPSS, Chicago, Illinois, USA) with one-way analysis of variance (ANOVA): descriptive statistics were used to prove the significant differences in seminal parameters including sperm motility, livability and acrosomal status (expressed as mean \pm SE) and reproductive parameters including receptivity, fertility and live-born kits (expressed as mean \pm SD) between the 2 experimental groups (C and IV group) during the different insemination cycles. Multiple comparisons of the means were done with Duncan test and P value was set at < 0.05 .

Multiple regression analysis was performed to develop a model for evaluating the correlation coefficients between sperm parameters and doe reproductive characteristics; Pearson's coefficients were calculated to assess the correlation between sperm and reproductive parameters; only fertility showed significant differences. P value was set at < 0.05 to indicate statistical significance.

3. Results

3.1. Sperm morphology and motility characteristics

The C group semen showed improved motility parameters compared to the IV group ($P < 0.01$) (Table 1), including a greater % progressive motility (43.31 vs. 38.34), VAP (106.51 vs. 99.55), VSL (86.71 vs. 76.81), BCF (38.75 vs. 36.11), STR (79.76 vs. 75.65), and LIN (52.99 vs. 46.30). On the other hand, the IV group semen showed an increased VCL (171.37 vs. 167.96) and ALH (6.63 vs. 5.95) compared to the C group semen ($P < 0.01$).

3.2. Reproductive parameters

Comparing the reproductive performances of does revealed effects of group and cycle, and their interaction ($P < 0.001$) (Table 2). The IV group showed greater or equal sexual receptivity compared to the C group ($P < 0.001$), along cycles. Within cycles, sexual receptivity was greater in cycles 3, 5, and 6 (100, 100 and 95, respectively) compared to in cycles 1, 2, and 4 (0, 75 and 50, respectively) ($P < 0.001$).

We found a significant group effect ($P < 0.001$) with regards to

Table 2
Reproductive parameters of does.

Cycle	1		2		3		4		5		6		Effects			
	C	IV	C	IV	C	IV	C	IV	C	IV	C	IV		Group	Cycle	GxC
Receptivity (%)	0.00 \pm 0.00	0.00 \pm 0.00	50.00 ^a \pm 8.33	100.00 ^b \pm 0.00	100.00 \pm 0.00	100.00 \pm 0.00	0.00 ^a \pm 0.00	100.00 ^b \pm 0.00	100.00 ^b \pm 0.00	100.00 ^b \pm 0.00	100.00 \pm 0.00	90.00 ^a \pm 9.09	100.00 ^b \pm 0.00	**	**	**
Fertility (%)	60.00 \pm 0.00	60.00 \pm 0.00	60.00A \pm 0.00	100.00B \pm 0.00	60.00A \pm 0.00	50.00B \pm 0.00	60.00A \pm 0.00	100.00B \pm 0.00	100.00B \pm 0.00	40.00A \pm 0.00	80.00B \pm 0.00	80.00 \pm 0.00	80.00 \pm 0.00	**	**	**
Live-born kits (N ^o)	3.30 ^a \pm 0.47	4.60 ^b \pm 1.30	5.60 ^b \pm 1.55	7.30 ^b \pm 0.42	5.20 ^a \pm 0.42	5.20 ^a \pm 1.20	4.70 ^b \pm 1.32	8.30 ^b \pm 0.52	8.00 ^b \pm 0.42	2.60 ^a \pm 1.33	6.60 ^b \pm 0.85	5.70 ^b \pm 1.05	6.60 ^b \pm 1.16	NS	*	NS

20 does/group/cycle (20 x 6 = 120 does/group). Values are expressed as mean \pm SD. C = control group; IV = intravaginal group; Group = effect of Group; Cycle = effect of Cycle; GxC = interaction between Cycle and Group; * : $P < 0.01$; ** : $P < 0.001$.

A, B: significant differences between groups ($P < 0.001$) for receptivity and fertility, a, b: significant differences between cycles ($P < 0.01$) for live born kits.

fertility rate during cycles 2, 4 and 5 between the does in the C group (60, 60 and 40%) and in group IV (100, 100 and 80%, respectively). While in cycle 3, the fertility rate was significantly ($P < 0.001$) increased with the C group compared to the IV group (60% vs. 50%). An equal fertility rate was recorded within cycle 1 (60–60%) and cycle 6 (80–80%) with the C and IV groups respectively. Regarding the number of live-born kits (results originated from both groups data), only cycle showed a significant effect ($P < 0.01$), with differences recorded within cycles 2, 4 and 6 (6.45, 8.15 vs. 6.15, respectively) (Fig. 1). Although there was no significant effect of group, the IV group showed an equal or greater number of live-born kits compared to the C group (4.60 vs. 3.30 in cycle 1; 6.60 vs. 5.70 in cycle 6).

3.3. Correlation between sperm and reproductive parameters

We identified several significant correlations between sperm motility, morphological parameters, and doe reproductive aspects (Table 3) (data was originated from both groups, C and IV). The % of live sperm showed a positive correlation (0.40, $P < 0.01$) with the % of sperm with intact acrosome. Moreover, progressive motility was positively correlated ($P < 0.001$) with several motility characteristics, including VAP (0.77), VSL (0.87), ALH (0.76), BCF (0.72), STR (0.79), and LIN (0.84). On the other hand, progressive motility was negatively correlated with fertility rate (-0.19 , $P < 0.001$).

3.4. Regression model for fertility

The sperm parameters VAP, VSL, VCL, ALH, BCF, STR, LIN, total motility, progressive motility, and % of live sperm with intact acrosome explained only 4% of the variation in fertility rate (Table 4) (data was originated from both groups, C and IV). Regression analysis revealed that fertility was negatively impacted ($P < 0.05$) by VAP (-0.56) and progressive motility (-0.24), and was positively impacted by VSL (0.36).

4. Discussion

Our present results indicated that the addition of GnRH to rabbit sperm did not significantly impair motility characteristics, similar to prior findings [33]. Numerous proteolytic enzymes are present in the seminal plasma and spermatozoa of mammals and avian species [34,35], and GnRH analogues could be susceptible to peptidase degradation. Vicente et al. [36] suggested that decreased hormonal activity is largely due to seminal plasma—finding that a low dilution rate of seminal plasma is associated with high amino peptidase activity, and that GnRH analogues can be hydrolyzed like many other proteins and peptides. In contrast, a high dilution rate is associated with low amino-peptidase activity, allowing the attainment of high ovulation frequency.

The intravaginal absorption of GnRH added to diluted semen may be influenced by both the mucosal state (secretions induced by




Fig. 1. Effect of insemination cycle on live-born kits number. Values are expressed as mean \pm SD; a, b: significant difference between different letters ($P < 0.01$).

Table 3
Correlation coefficients between sperm parameters and doe reproductive characteristics.

	VAP	VSL	VCL	ALH	BCF	STR	LIN	Total motility	Progressive motility	Live sperm	Sperm with intact acrosome	Fertility	Receptivity
VSL	0.908**												
VCL	0.082	-0.264**											
ALH	-0.628**	-0.863**	0.605**										
BCF	0.589**	0.841**	-0.483**	-0.934**									
STR	0.655**	0.907**	-0.521**	-0.943**	0.950**								
LIN	0.780**	0.955**	-0.522**	-0.944**	0.878**	0.950**							
Total motility	0.557**	0.510**	-0.044	-0.343**	0.260**	0.325**	0.433**						
Progressive motility	0.776**	0.871**	-0.291**	0.756**	0.718**	0.788**	0.838**	0.816**					
Live sperm	-0.127	-0.184	0.164	0.293	-0.278	-0.237	-0.231	-0.173	-0.174				
Sperm with intact acrosome	0.174	0.131	0.198	-0.032	-0.008	0.061	0.050	0.224	0.171	0.401*			
Fertility	-0.226**	-0.228**	-0.032	0.157*	-0.200**	-0.216**	-0.192*	-0.112	-0.195**	0.031	0.071		
Receptivity	-0.055	-0.021	-0.041	-0.035	0.034	0.006	-0.003	-0.010	-0.047	0.001	0.000	0.138	
Live kits	-0.072	-0.053	-0.067	0.023	0.035	-0.031	-0.043	-0.139	-0.094	-0.013	-0.220	0.282**	-0.051

Based on means of 54 evaluated heterospermic pooled semen samples, and results are expressed for both the C and IV groups. VAP = average path velocity ($\mu\text{m/s}$); VSL = straight-line velocity ($\mu\text{m/s}$); VCL = curvilinear velocity ($\mu\text{m/s}$); ALH = amplitude of lateral head displacement ($\mu\text{m/s}$); BCF = beat cross-frequency (Hz); STR = straightness (%); LIN = linearity (%); Total motility = total motile sperm (%); Progressive motility (%); Sperm with intact acrosome (%). * = $P < 0.01$. ** = $P < 0.001$.

the receptivity status) and the sperm concentration [18]. Sexual receptivity is an important factor in the pregnancy rate, as negative results have been obtained after insemination of non-receptive does [37]. Good assessment of sexual receptivity in rabbit does is fundamental to successful AI, but is subjective and affected by technician experience [18]. In our study, does inseminated with diluted semen plus Dalmarelin showed equal or greater sexual receptivity compared to the control group. In contrast, the control group showed an improved fertility rate only in cycle 3. In contrast, Dal Bosco et al. [38] reported that the intramuscular GnRH administration was associated with a higher fertility rate (80%) compared to intravaginal GnRH administration (20%).

It has been suggested that a fraction of GnRH analogue can be lost due to seminal backflow, and that a reduced seminal dose can help reduce the required quantity of added hormone [13]. In our present study, the seminal dose volume was very low, possibly explaining the better results in the intravaginal group, which received a similar amount of hormone (0.3 mL/doe) as the control group (0.2 mL/doe). The negative performance of the control group may have also been influenced by anti-GnRH antibody formation due to the repeated intramuscular hormone application, as previously described by Canali et al. [4]. On the other hand, the level of anti-eCG antibodies is reportedly eCG dose dependent [14]. Notably, the immune reaction appeared after cycle 6, and was not

significantly correlated with the reproductive parameters [14]. It is also possible that the difference between groups was amplified by the fact that intramuscular administration is probably the more stressful method of ovulation induction [16].

The two groups in our study did not significantly differ with regards to the number of live-born kits, which is in agreement with the findings of a previous study by Quintela et al. [13]. Notably, the IV group showed an equal or greater number of live-born kits compared to the C group across all cycles.

When assessing correlations between sperm parameters and fertility, receptivity, and number of live-born kits, several factors should be standardized, including the semen collection and dilution, and the doe's reproductive status. In our present study, we identified several significant correlations between sperm motility, morphological parameters, and reproductive aspects. We found a significant positive correlation between the two sperm morphological parameters: % of live sperm and % of sperm with intact acrosome. These results are in agreement with the report by Saacke and White [39], who analyzed bull semen and showed a significant correlation between sperm fertility and the percentage of spermatozoa with intact acrosome after 2 h of incubation at 37 °C. On the other hand, another study showed a higher percentage of sperm with intact acrosome among fertile stallions (74%) than infertile ones [40]; however, these percentages are not often

Table 4
Summary of selected multiple regression model for fertility (CONSTANT).

Parameters (CASA and RSA)	Coefficient ($R^2 = 4.3\%$)	S.E. (11.66)	R^2 (%)	P
CONSTANT	127.583	91.125	–	–
VAP	-0.561	0.802	2.6	0.03
VSL	0.364	0.804	2.7	0.03
VCL	0.127	0.326	0	–
ALH	-2.628	3.081	1	–
BCF	-0.270	0.594	2	–
STR	-0.605	0.987	2	–
LIN	0.376	1.003	2	–
Total motility	0.127	0.200	0.7	–
Progressive motility	-0.244	0.301	2.3	0.04
Live sperm	-0.009	0.225	0	–
Sperm with intact acrosome	0.039	0.261	0	–

Significance of statistical model = $P < 0.05$. S.E. = standard error; R^2 = the partial R^2 for each parameter; P = significance of individual parameters; CASA = computer-assisted sperm analysis; RSA = morphological analysis; VAP = average path velocity ($\mu\text{m/s}$); VSL = straight-line velocity ($\mu\text{m/s}$); VCL = curvilinear velocity ($\mu\text{m/s}$); ALH = amplitude of lateral head displacement ($\mu\text{m/s}$); BCF = beat cross-frequency (Hz); STR = straightness (%); LIN = linearity (%); Total motility = total motile sperm (%); Progressive motility (%); Sperm with intact acrosome (%).

The results presented come from both the C and IV groups data.

correlated with *in vivo* fertility. The percentage of sperm cells with normal morphological traits is an important indicator and is highly correlated with fertility rate [40]. Regarding motility, our present results revealed that progressive motility was significantly positively correlated with motility characteristics, including VAP, VSL, ALH, BCF, STR, and LIN. These results are in agreement with the findings of Nagy et al. [41] who evaluate the different kinematic (velocity) parameters of frozen/thawed bull semen and found that VAP is the most useful semen motility characteristic that has clinical relevance in fertility prediction. Unfortunately, we observed a significant negative correlation between progressive motility and the does' fertility rate. In the literature, there is no clear evidence demonstrating that a positive correlation between sperm motility parameters and sperm fertility is good, and that it can truly reflect the reproductive parameters of the doe. This lack of information is commonly due to the variation between individual animals and the use of insemination doses with spermatozoa numbers that are too high or too low [41].

The inclusion of morphological and motility parameters in a multiple regression model to evaluate fertility explained only a very low portion of the variation. However, some sperm parameters (VAP, VSL, and total motility) had a significant impact. The low correlation obtained in the present trial is likely due to the low number of sperm parameters evaluated. It is difficult to find a perfect test for evaluating *in vivo* fertility, i.e., one that consistently shows high correlations between the parameters and the animals' fertility rate. An ideal sperm assay would evaluate several spermatozoa attributes in a large number of sperm, to determine the proportion of the cells that possess all of the characteristics necessary to fertilize the oocyte [42].

5. Conclusions

Our present findings indicated that the incorporation of GnRH in a seminal dose could be used for ovulation induction in rabbit does. This method achieved sexual receptivity, fertility, and number of live-born kits equal or greater to those with conventional GnRH administration. Intravaginal administration may also carry lower risks of hemorrhagic follicles and thereby enable a higher conception rate and a longer reproductive career for the doe.

Further studies are needed to determine the ideal GnRH level for intravaginal administration based on the doe's physiological status. Compared to the intramuscular absorption capacity, the intravaginal absorption capacity is about 10 times smaller, such that this administration route requires a greater dosage. Identifying the optimal dose will enable avoidance of the conventional invasive treatment with GnRH analogues, along with the associated antibody formation and traumatic action, thus improving doe physiological welfare and longevity.

Name of the project

Sistema di Allevamento Alternativo del coniglio: benessere e produttività - Resp. Scientifico MUGNAI.

Acknowledgements

This work was supported by Turin University Research Project: "Rabbit Alternative Breeding System: well-being and productivity" (MUGC_RILO_17_01).

References

[1] Dal Bosco A, Rebollar PG, Boiti C, Zerani M, Castellini C. Ovulation induction in rabbit does: current knowledge and perspectives. *Anim Reprod Sci* 2011;129:

- 106–117.
- [2] Hafez ESE. Reproducción, hormonas y factores de crecimiento. In: Hafez ESE, editor. *Reproducción e Inseminación Artificial en Animales*. sixth ed. Mexico, DF: Interameca-McGraw-Hill; 1993.
- [3] Zapletal D, Pavlik A. The effect of leclirelin (GnRH) dosage on the reproductive performance of nulliparous and lactating rabbit does. *Anim Reprod Sci* 2008;104:306–15.
- [4] Canali C, Boiti C, Castellini C, Zampini D. Risposta anticorpale delle coniglie trattate ripetutamente con PMSG nella pratica della sincronizzazione degli estri. In: 2° meeting nazionale studio della Efficienza riproduttiva degli animali di Interesse zootecnico; 1991 24 Nov. p. 103–8. Italia.
- [5] Maertens L, Luzi F. Note concerning the effect of eCG stimulating on the mortality rate at parturition and the distribution of litter size in artificially inseminated does. *World Rabbit Sci* 1995;3:57–61.
- [6] Maertens L, Luzi F, Grilli G. Effects of PMSG induced oestrus on the performances of rabbit does: a review. *World Rabbit Sci* 1995;3:191–9.
- [7] Castellini C. Recent advances in rabbit artificial insemination. In: *Proceeding of the 6th world rabbit congress*, toulouse, France; 1996. p. 13–26.
- [8] Boiti C, Castellini C, Canali C, Zampini D, Monaci M. Long term effect of eCG on rabbit does reproductive performance. *World Rabbit Sci* 1995;3:51–6.
- [9] Rebollar PG, Milanès A, Pereda N, Millan P, Cano P, Esquifino AI, Villarreal M, Silvan G, Lorenzo PL. Oestrus synchronisation of rabbit does at early postpartum by doe–litter separation or ECG injection: reproductive parameters and endocrine profiles. *Anim Reprod Sci* 2006;93:218–30.
- [10] Theau-Clément M, Bolet G, Roustan A, Mercier P. Comparaison de différents modes d'induction de l'ovulation chez les lapines multipares en relation avec leur stade de physiologie et la réceptivité au moment de la mise à la reproduction. *5es Journées de la Recherche Cunicole*. 1990. Déc 12–13; Tome I, Communication 6. Paris.
- [11] Yoshimura Y, Nakamura Ando YM, Jinno M, Nanno T, Oda T. Protein kinase C mediates gonadotropin-releasing hormone agonist induced meiotic maturation of follicle-enclosed rabbit oocytes. *Biol Reprod* 1992;47:118–25.
- [12] Zagagnolo V, Dharmarajan AM, Hesla J, Wallach EE. Effects of a gonadotropin-releasing hormone analog on rabbit ovarian function. *Endocrinology* 1996;137:5400–6.
- [13] Quintela LA, Peña AI, Vega MD, Gullón J, Prieto MC, Barrio M, et al. Ovulation induction in rabbit does submitted to artificial insemination by adding buserelin to the seminal dose. *Reprod Nutr Dev* 2004;44:79–88.
- [14] Theau-Clément M, Lebas F, Beckers JF, Drion PV. Evolution of anti-eCG antibodies in response to eCG doses and number of injections. Relationship with productivity of rabbit does. *Animal* 2008;2:746–51.
- [15] Sanchez J, Baselga M, Peiro R, Silvestre MA. Analysis of factors influencing longevity of rabbit does. *Livest Prod Sci* 2004;90:227–34.
- [16] Castellini C. Reproductive activity and welfare of rabbit does. *Ital J Anim Sci* 2007;6:743–7.
- [17] EFSA. The impact of the current housing and husbandry systems on the health and welfare of farmed domestic rabbit. *EFSA J* 2005;267:1–31.
- [18] Viudes de Castro MP, Lavara R, Marco Jiménez F, Cortell C, Vicente JS. Ovulation induced by mucosa vaginal absorption of buserelin and triptorelin in rabbit. *Theriogenology* 2007;68:1031–6.
- [19] Dal Bosco A, Cardinali R, Brecchia G, Rebollar PG, Fatnassi M, Millan P, Mattioli S, Castellini C. Induction of ovulation in rabbits by adding Leclirelin to the seminal dose: in vitro and in vivo effects of different excipients. *Anim Reprod Sci* 2014;150:44–9.
- [20] Zhang YQ, Qin Y. Reproductive performance of rabbit does by adding leuporelin in semen to induce ovulation. In: *Proceedings of the 10th World rabbit congress*; 2012. p. 419–23.
- [21] Battagliani M, Constantini F, Baldissera CN, Castrovilli CF. Induction of ovulation and artificial insemination in the rabbit. *Concigliocultura* 1982;19:45–54.
- [22] Castellini C, Lattaioli P. Effect of number of motile sperms inseminated on reproductive performance of rabbit does. *Anim Reprod Sci* 1999;57:111–20.
- [23] Mousa-Balabel TM, Mohamed RA. Effect of different photoperiods and melatonin treatment on rabbit reproductive performance. *Vet Q* 2011;31:165–71.
- [24] Castellini C, Dal Bosco A, Cardinali R. Effect of post-weaning rhythm on the body fat and performance of rabbit does. *Reprod Nutr Dev* 2005;46:195–204.
- [25] Boiti C. Guidelines for the handling of rabbit bucks semen. *World Rabbit Sci* 2005;13:71–91.
- [26] Lavara R, Moce E, Lavara F, Viudes de Castro MP, Vicente JS. Do parameters of seminal quality correlate with the results of on-farm inseminations in rabbits? *Theriogenology* 2005;64:1130–41.
- [27] Brun JM, Theau-Clément M, Bolet G. The relationship between rabbit semen characteristics and reproductive performance after insemination. *Anim Reprod Sci* 2002;70:139–49.
- [28] Błaszczyk M, Slanina T, Massanyi P, Stawarz R. Semen quality assessment of New Zealand white rabbit bucks. *J Microbiol Biotechnol Food Sci* 2013;2:1365–76.
- [29] Lukac N, Massanyi P, Krokova J, Nad P, Slamecka J, Ondruska L, et al. Relationship between trace element concentrations and spermatozoa quality in rabbit semen. *Czech J Anim Sci* 2009;42:46–50.
- [30] Lukac N, Bardos L, Stawarz R, Roychoudhury S, Makarevich AV, Chrenek P, et al. In vitro effect on nickel on bovine spermatozoa motility and annexin V-labeled membrane changes. *J Appl Toxicol* 2011;31:144–9.
- [31] Kovacs A, Foote RH. Viability and acrosome staining of bull, boar and rabbit spermatozoa. *Biotech Histochem* 1992;67:119–24.
- [32] Boccia L, Di Palo R, De Rosa A, Attanasio L, Mariotti E, Gasparrini B. Evaluation

- of buffalo semen by Trypan blue/Giemsa staining and related fertility in vitro. *Ital J Anim Sci* 2007;6:739–42.
- [33] Fernández Serrano P, Casares Crespo L, Viudes de Castro MP. Chitosan–dextran sulphate nanoparticles for GnRH release in rabbit insemination extenders. *Reprod Domest Anim* 2017;52:72–7.
- [34] Métayer S, Dacheux F, Dacheux JL, Gatti JL. Comparison, characterization, and identification of proteases and protease inhibitors in epididymal fluids of domestic mammals Matrix metalloproteinases are major fluid gelatinases. *Biol Reprod* 2002;66:1219–29.
- [35] Kotiowska M, Kowalski R, Glogowski J, Jankowski J, Ciereszko A. Gelatinases and serine proteinase inhibitors of seminal plasma and the reproductive tract of Turkey (Meleagris gallopavo). *Theriogenology* 2005;63:1667–81.
- [36] Vicente JS, Lavara R, Marco Jimenez RF, Viudes de Castro MP. Detrimental effect on availability of buserelin acetate administered in seminal doses in rabbits. *Theriogenology* 2011;76:1120–5.
- [37] Armero E, Garcia-Ximenez F, Vicente JS, Baselga M. Cycle synchronization of rabbit does naturally mated or artificially inseminated. *World Rabbit Sci* 1994;2:107–13.
- [38] Dal Bosco A, Cardinali R, Rebollar PG, Millan P, Brecchia G, Castellini C. Ovulation induction in rabbit does submitted to artificial insemination by adding lecorelin to the seminal dose. Preliminary results. In: Proceedings of the 10th world rabbit congress, vol. 6; 2012. p. 389–93.
- [39] Saacke RG, White LM. Semen quality and their relationship to fertility. In: Proceedings of the 4th NAAB technical conference on animal reproduction AI; 1972. p. 23–7.
- [40] Zhang J, Boyle MS, Smith CA, Moore HDM. Acrosome reaction of stallion spermatozoa evaluated with monoclonal antibody and zona-free hamster eggs. *Mol Reprod Dev* 1990;27:152–8.
- [41] Nagy Á, Polichronopoulos T, Gáspárdy A, Solti L, Cseh S. Correlation between bull fertility and sperm cell velocity parameters generated by computer-assisted semen analysis. *Acta Vet Hung* 2015;63:370–81.
- [42] Graham JK. Analysis of stallion semen and its relation to fertility. *Vet Clin N Am Equine Pract* 1996;12:119–30.