



Available online freely at www.isisn.org

Bioscience Research

Print ISSN: 1811-9506 Online ISSN: 2218-3973

Journal by Innovative Scientific Information & Services Network



RESEARCH ARTICLE

BIOSCIENCE RESEARCH, 2019 16(2): 1258-1266.

OPEN ACCESS

Genetic diversity among wheat genotypes using RAPD markers and its implication on genetic variability of diallel crosses

A.A.El-Hosary¹; M. El. M. El-Badawy¹; S.A.S Mehasen¹, A.A.A.; El-Hosary¹, T.A. El-Akkad² and A. El-Fahdawy¹

¹Agronomy Dep., Fac. of Agric., Benha Univ., Egypt

²Genetic Dep., Fac. of Agric., Benha Univ., Egypt

*Correspondence: a.elhousary@fagr.bu.edu.eg Accepted: 09 Apr. 2019 Published online: 06 May. 2019

Hybrids involving diverse parental genotypes generally produce high specific combining ability effects and hybrid vigor. Therefore, the choice of the suitable breeding program depends on the inheritance of the quantitative traits. This research aims at detecting the extent of the genetic diversity (GD) among 8 wheat parental genotypes, and its implication on F₁ crosses for biological and grain yield plant⁻¹. Thus the 8 parents were screened using ten RAPD primers. These primers produced 89 fragments, 58 of them were polymorphic. A GD value as high as 0.474 was found just one cross (P₁xP₄). Half diallel mating, design involving 8 parents, was made to initiate a set of 28 F₁ crosses. Parents and their crosses were tested under normal and drought stress conditions for biological and grain yield per plant. Genotypes, general (GCA) and specific (SCA) combining ability were significant for the two traits. The non-additive gene action is substantial in controlling these traits. Five crosses showed all showed relatively high mean performance and SCA effects for both traits in both and across normal and drought environment. For biological and per plant grain yield, Pearson's correlation coefficients with GD were 0.34 and 0.69. Hence, RAPD marker can be used as a tool for determining the extent of genetic diversity among wheat genotypes and to precisely predict the yield performance value for F₁ hybrids.

Keywords: combining ability, drought stress, GCA, RAPD marker, SCA, Wheat.

INTRODUCTION

Wheat (*Triticum aestivum* L) yield is exposed to a wide range of damaging drought stress in many production areas around the globe due to climate change conditions. One way to guard against these harmful changes is to develop wheat crosses that exhibit relatively more water stress resistance.

Molecular techniques are now the most reliable, easy, cheap and valuable tool for assessing the genetic diversity in genome research (Tonk et al., 2014 and EL Saadoon et al., 2018). Random amplified polymorphic DNAs tool (RAPDs) has been proved to be independent

of the confounding effects of environmental factors.

Drought, among abiotic stress factors, is considered the most wide spreading and limiting to crop productivity. (EL Saadoon et al., 2017) illustrated that many factors like, potential evapotranspiration, evapotranspiration, precipitation, temperature, humidity individually or in combination leads to drought stress. Water stresses can happen at any stage of plant growth and development; therefore, studying the genetic nature of resistance and productivity is needed.

Diallel crosses schemes are extremely used in plant breeding research to get information

about genetic properties of parental lines and/or to estimate both general combining ability (GCA) and specific combining ability (SCA) (Shrief et al., 2017). In addition, this technique provide prior information on the genetic behavior of these attributes in F_1 (Chowdhry et al., 1992 and Topal et al., 2004). Heterosis is a complex phenomenon, which depends on the balance of different combinations of gene effects, as well as on the distribution of plus and minus alleles in the parents of a mating system. In self-pollinated crops, like wheat, the scope for utilizing heterosis depends mainly upon the direction and magnitude of heterosis. Heterosis over better parent may be useful in identifying the best crosses might show immense practical value if they involve the best cultivars of the area (El Hosary and Nour El Deen 2015).

Therefore, this investigation aims at applying the RAPD-PCR marker technique to detect DNA polymorphism, to identify parents and to estimate genetic diversity among wheat genotypes. In addition to using a half diallel crosses among eight genotypes to assess genetic variability under drought and normal irrigation treatments and to correlate F_1 performance to genetic divergence of the parental genotypes.

MATERIALS AND METHODS

Plant material

Eight wheat genotypes, P_1 - P_8 (Table 1) were crossed in 8x8 diallel cross excluding reciprocals giving a total of 28 crosses in the first season of 2015/2016.

Laboratory Experiment

DNA extraction

Ten seeds of parental genotypes were planted in pots. Then, leaf tissue was collected from 5-7 days old germinated seedlings of each parent. The samples were bulked, lyophilized and ground with a mortar and pestle. Genomic DNA was isolated and extracted according to the manufacturer's protocol carefully using DNeasy® Plant mini Kit for DNA isolation. The purity, quantification and qualification of the extracted DNA were tested.

RAPD-PCR

Random amplified polymorphic DNA (RAPD reactions) were conducted using a set of ten 10-

mer oligonucleotide (Primer from Operon Technologies, Inc.) i.e. OP A2, A9, B5, B7, C6, C20, D2, D10, D16 and E7 were screened for the ability to provide a suitable band pattern with various parents. These primers give polymorphic results for parents under study. All PCR reactions were performed as reported by Williams et al., (1990). Products of amplification were separated by electrophoresis on 1.2% agarose gels, stained with ethidium bromide, then photographed under UV-trans illuminator by digital camera with UV filter adaptor.

Field layout and management, and experimental design

Two field trials were conducted during 2016/2017 seasons at the Moshtohor Experiment Research Station (30° 21' 07" N and 31° 13' 34" E), Al-Khalubiah, Egypt. In each trail, seeds of 8 parents and its resultant 28 F_1 crosses (thereafter named $P_i \times P_j$), were planted on 4 December 2016 in three completely randomized blocks. The first trial was irrigated only once after planting irrigation and the second one was normally irrigated 5 times. Both parent and F_1 seeds were put in 0.20-m hills in 3.0 m x 0.30 m single-row plots. All field and crop management practices were followed as needed.

Meteorological data in season 2016/ 2017 were obtained from the Agro-meteorological Station at Moshtohor, Benha Univ. from December to May, the Maximum temperatures were 19.7, 17.7, 20.4, 25.8, 29.1 and 34.5°C, and the minimum temperatures were 9.2, 6.1, 7.8, 11.4, 14.4, and 19.0 °C, relative humidity were 51.3, 55.9, 47.2, 37.3, 38.9 and 32.1% and the mean precipitation were 0.5, 1.6, 0.8, 0.4, 0.3 and 0.00 mm respectively.

Sampling and studied traits

A 10-plant random sample was selected from each individual plot to calculate mean biological yield and per plant grain yield.

Data analysis

Separated bands for each RAPD marker were filed as binary data matrices, where 0 for absence and 1 presence of band. Similarity coefficients between a pair of parental genotypes were calculated according to Jaccard (1908). A dendrogram tree was constructed by the UPGMA clustering algorithm from the SAHN option of NTSYS-PC version 2.1 (Rohlf, 2000).

Table (1): The name, pedigree and source of the studied parental genotypes

| .NO | Entry name | Pedigree | Source |
|-----|-------------|--|--------|
| 1 | Yakora Rojo | Ciano 67/Sonora 6411 Klien Rendidor/3/1L815626Y-2M-1Y-0M-302M | CIMMYT |
| 2 | Sakha 93 | S 92/TR 810328 S8871-1S-2S-1S-0S | Egypt |
| 3 | Masr 1 | OASIS/SKAUZ//4*BCN/3/2*PASTOR | Egypt |
| 4 | Drought 4 | Landraces developed by Prof. Dr. M. | Egypt |
| 5 | Shandawel 1 | SITE//MO/4/NAC/TH.AC//3*PVN/3/MIRLO/B | Egypt |
| 6 | Gemiza 9 | ALD"S"/HUAC"S"/CMH74A.630/5X | Egypt |
| 7 | Giza 171 | SAKHA 93 / GEMMEIZA 9 S.6-1GZ-4GZ- | Egypt |
| 8 | Sides 13 | KAUZ"S"/TSI/SNB"S" | Egypt |

NOVA was conducted as outlined by Steel and Torrie (1980) for all characters. Both GCA and SCA mean squares were calculated according to Griffing's (1956) Method II, Model I. A combined analysis of the two experiments was carried out whenever homogeneity of mean squares was detected (Gomez and Gomez 1984).

RESULTS AND DISCUSSION

RAPD Polymorphism

The polymorphic amplification products set of ten RAPDs primers were presented in table (2). Per primer fragment mean was 8.9 that was generated in a range of 5-13 fragments, among a total of 89 generated fragments.

Among of them 58 were polymorphic within the eight genotypes (Table 2). Other studies indicated different results (Cao et al., 2002, Guoyue and Hui, 2007, Tahir, 2008, Siddiqui et al., 2010, and Kumar et al., 2017).

Genetic similarity

The genetic similarity matrix was produced from the RAPD data using Nei and Li's formula (1979) (Table 3). Genetic similarity ranged from 0.526 for P1 and P4 to reach 0.812 for P1 and P2, with and overall mean of 0.669 among all genotypes.

Cluster analysis:

Based on RAPD data, the dendrogram constructed from cluster analysis (Fig. 1) shows two main clusters. Each of P4, P5, and P7 are in Cluster 1, and the rest of genotypes is in Cluster 2. The latter one shows two sub clusters: one contains P8, and in the other one, P1 and P2 were closely related.

Field experiment

Table 4 shows the ANOVA for both per plant biological and grain yields for each of and across irrigations. For the two yield characters, differences between the two irrigations (environments) were significant ($p < 0.01$). Genotypes and its components (parents, crosses and parent vs crosses) mean squares were significant for studied yield traits indicating wide diversity among all genotypes used in this work. Moreover, significant mean squares between each of genotypes, parents and crosses by environment interaction were detected for biological and grain weight/ plant. For both yield characters, the interaction effects of irrigation regime with each of genotypes, parents, and crosses greatly varied ($p < 0.01$) (Table 4). In addition, the same trend existed for all main effects of genotype, and for parent, cross, and parent vs. cross.

Mean trait performance

Across environment (irrigation regime), per plant biological yield widely ranged from about 111 to 212 g and from about 29 to 52 g for per plant grain weight among all parents and the 28 crosses (Table 5). This 2-fold range for both yield traits implied how genotypes unevenly responded to greatly varied irrigational environments. Percentage performance improvement, for the parent and F_1 hybrids, showed significant differences ($p < 0.05$) relative to average genotypes (Table 5). Concerning Across environment (irrigation regime), per plant biological yield widely ranged from about 111 to 212 g and from about 29 to 52 g for per plant grain weight among all parents and the 28 crosses (Table 5).

Table (2): Primer used in RAPD analysis of eight wheat genotypes and total number of fragments detected by each primer and number of polymorphic fragments.

| Primers | Sequence | TSB | TF | NPF | PPF |
|---------|------------|------|-----|-----|-------|
| OPA-02 | TGCCGAGCTG | 75 | 11 | 7 | 63.64 |
| OPA-09 | GGGTAACGCC | 76 | 13 | 7 | 53.85 |
| OPB-05 | TGCGCCCTTC | 49 | 10 | 8 | 80.00 |
| OPB-07 | GTCCACACGG | 53 | 11 | 9 | 81.82 |
| OPC-06 | GAACGGACTC | 49 | 9 | 6 | 66.67 |
| OPC-20 | ACTTCGCCAC | 46 | 9 | 6 | 66.67 |
| OPD-02 | GGACCCAACC | 29 | 5 | 4 | 80.00 |
| OPD-10 | GGTCTACACC | 33 | 6 | 3 | 50.00 |
| OPD-16 | AGGGCGTAAG | 43 | 6 | 3 | 50.00 |
| OPE-07 | TCAGGGAGGT | 55 | 9 | 5 | 55.56 |
| Total | | 508 | 89 | 58 | - |
| Mean | | 50.8 | 8.9 | 5.8 | 65.17 |

Where: TSB = Total number of scorable bands, TF= Total number of fragments, NPF = Number of polymorphic fragments, PPF = Percentage of polymorphic fragments.

Table (3): Similarity matrix based on Nei and Li's coefficient for eight genotypes in wheat revealed by RAPD.

| Rows/ cols | p1 | p2 | p3 | p4 | p5 | p6 | p7 | p8 |
|------------|-------|-------|-------|-------|-------|-------|-------|----|
| p1 | 1 | | | | | | | |
| p2 | 0.812 | 1 | | | | | | |
| p3 | 0.792 | 0.806 | 1 | | | | | |
| p4 | 0.526 | 0.600 | 0.569 | 1 | | | | |
| p5 | 0.615 | 0.567 | 0.617 | 0.779 | 1 | | | |
| p6 | 0.788 | 0.753 | 0.760 | 0.544 | 0.573 | 1 | | |
| p7 | 0.608 | 0.620 | 0.609 | 0.794 | 0.792 | 0.605 | 1 | |
| p8 | 0.685 | 0.724 | 0.618 | 0.637 | 0.600 | 0.657 | 0.704 | 1 |

Where, P1 = Yakora Rojo, P2 = Sakha 93, P3 = Masr 1, P4 = Drought 4, P5 = Shandawel 1, P6 = Gemiza 9, P7 = Giza 171, P8 = Sides 13

Table (4): Mean squares for yield traits under drought stress condition and normal irrigation as well as the combined over them.

| S.O.V | df | | Biological yield/ plant | | | Grain weight / plant | | |
|----------------|----|----|-------------------------|-----------|-------------|----------------------|----------|-----------|
| | S | C | Drought | Normal | Combined | Drought | Normal | Combined |
| Irrigation (I) | | 1 | | | 136604.74** | | | 1628611** |
| Rep/ I | 2 | 4 | 64.33* | 30.7 | 47.519** | 17.76** | 2.12 | 9.94 |
| Genotypes (G) | 35 | 35 | 3213.69** | 1751.33** | 3555.66** | 176.72** | 149.35** | 233.03** |
| Parent (P) | 7 | 7 | 1992.57** | 1454.07** | 1914.43** | 102.05** | 146.03** | 185.69** |
| Cross (C) | 27 | 27 | 3639.48** | 1843.45** | 4060.93** | 195.69** | 149.42** | 227.73** |
| P vs C. | 1 | 1 | 265.01** | 672.45** | 1401.94** | 187.20** | 170.52** | 357.53** |
| G x I | | 35 | | | 1409.36** | | | 103.04** |
| p x I | | 7 | | | 1532.21** | | | 62.40** |

| | | | | | | | | |
|-------------|----|-----|-----------|----------|-----------|---------|---------|----------|
| C x I | | 27 | | | 1422.01** | | | 117.39** |
| P.vs.C x I | | 1 | | | 207.95** | | | 0.19 |
| Error | 70 | 140 | 13.29 | 12.54 | 12.91 | 5.43 | 8.17 | 6.8 |
| GCA | 7 | 7 | 1007.41** | 701.48** | 701.08** | 12.98** | 38.84** | 28.68** |
| SCA | 28 | 28 | 1087.18** | 554.35** | 1306.25** | 70.39** | 52.52** | 85.76** |
| GCA x L | | 7 | | | 1007.80** | | | 23.14** |
| SCA x L | | 28 | | | 335.28** | | | 37.15** |
| Error | 70 | 140 | 4.43 | 4.18 | 4.3 | 1.81 | 2.72 | 2.27 |
| GCA/SCA | | | 0.93 | 1.27 | 0.54 | 0.18 | 0.74 | 0.33 |
| GCA x L/GCA | | | | | 1.44 | | | 0.81 |
| SCA x L/SCA | | | | | 0.26 | | | 0.43 |

* and ** significant at 0.05 and 0.01 levels of probability, respectively.

Table (5): Mean performance for the studied traits under normal irrigation and drought stress condition as well as the combined over them.

| | Biological yield/ plant (g) | | | Grain yield/ plant (g) | | |
|----------------------|-----------------------------|--------|----------|------------------------|--------|----------|
| | Drought | Normal | Combined | Drought | Normal | Combined |
| P1 | 159.67 | 174.00 | 166.83 | 24.73 | 46.30 | 35.52 |
| P2 | 121.67 | 165.67 | 143.67 | 25.87 | 33.20 | 29.53 |
| P3 | 94.33 | 191.33 | 142.83 | 21.20 | 44.43 | 32.82 |
| P4 | 151.67 | 216.33 | 184.00 | 34.43 | 44.27 | 39.35 |
| P5 | 154.33 | 226.00 | 190.17 | 35.03 | 51.47 | 43.25 |
| P6 | 123.00 | 180.33 | 151.67 | 26.53 | 52.37 | 39.45 |
| P7 | 163.67 | 168.67 | 166.17 | 35.30 | 51.00 | 43.15 |
| P8 | 166.00 | 185.00 | 175.50 | 35.57 | 55.47 | 45.52 |
| 1x2 | 124.00 | 212.67 | 168.33 | 28.27 | 36.97 | 32.62 |
| 1x3 | 193.33 | 224.33 | 208.83 | 41.20 | 47.83 | 44.52 |
| 1x4 | 134.00 | 162.00 | 148.00 | 31.57 | 60.37 | 45.97 |
| 1x5 | 148.67 | 188.00 | 168.33 | 33.73 | 56.17 | 44.95 |
| 1x6 | 160.67 | 201.67 | 181.17 | 35.20 | 45.33 | 40.27 |
| 1x7 | 178.67 | 187.67 | 183.17 | 37.93 | 54.73 | 46.33 |
| 1x8 | 192.67 | 198.00 | 195.33 | 39.13 | 48.97 | 44.05 |
| 2x3 | 123.33 | 211.67 | 167.50 | 27.63 | 42.47 | 35.05 |
| 2x4 | 113.67 | 207.67 | 160.67 | 41.50 | 57.40 | 49.45 |
| 2x5 | 212.00 | 213.00 | 212.50 | 45.33 | 58.03 | 51.68 |
| 2x6 | 121.00 | 177.67 | 149.33 | 23.33 | 56.87 | 40.10 |
| 2x7 | 188.33 | 236.00 | 212.17 | 39.03 | 48.63 | 43.83 |
| 2x8 | 148.67 | 170.67 | 159.67 | 33.53 | 53.47 | 43.50 |
| 3x4 | 170.00 | 216.67 | 193.33 | 46.67 | 57.43 | 52.05 |
| 3x5 | 124.33 | 216.67 | 170.50 | 28.80 | 39.20 | 34.00 |
| 3x6 | 119.33 | 171.00 | 145.17 | 22.80 | 43.97 | 33.38 |
| 3x7 | 147.33 | 223.33 | 185.33 | 33.47 | 50.13 | 41.80 |
| 3x8 | 132.00 | 222.00 | 177.00 | 36.33 | 63.07 | 49.70 |
| 4x5 | 119.67 | 221.00 | 170.33 | 23.30 | 46.40 | 34.85 |
| 4x6 | 126.67 | 198.33 | 162.50 | 30.50 | 55.30 | 42.90 |
| 4x7 | 81.33 | 141.67 | 111.50 | 21.10 | 44.70 | 32.90 |
| 4x8 | 117.33 | 179.33 | 148.33 | 22.53 | 50.80 | 36.67 |
| 5x6 | 200.00 | 218.33 | 209.17 | 45.47 | 55.07 | 50.27 |
| 5x7 | 81.33 | 149.00 | 115.17 | 19.00 | 45.23 | 32.12 |
| 5x8 | 171.00 | 216.00 | 193.50 | 37.40 | 55.87 | 46.63 |
| 6x7 | 144.67 | 164.00 | 154.33 | 38.03 | 40.90 | 39.47 |
| 6x8 | 117.00 | 197.00 | 157.00 | 22.23 | 54.57 | 38.40 |
| 7x8 | 184.67 | 188.00 | 186.33 | 38.97 | 39.50 | 39.23 |
| Average of genotypes | 144.72 | 195.02 | 169.87 | 32.30 | 49.66 | 40.98 |
| L.S.D 5% | 5.92 | 5.75 | 5.75 | 5.92 | 5.75 | 5.75 |

Table 6. Estimates of combining ability effects for yield traits in both and across environments.

| Parent | Biological yield/ plant | | | Grain weight/ plant | | |
|-------------------|-------------------------|----------|----------|---------------------|----------|----------|
| | Drought | Normal | Combined | Drought | Normal | Combined |
| P1 | 14.88** | -3.28** | 5.80** | 0.58 | -0.40 | 0.09 |
| P2 | -2.82** | 0.55 | -1.13** | -0.03 | -2.67** | -1.35** |
| P3 | -10.42** | 11.32** | 0.45 | -1.14** | -1.40** | -1.27** |
| P4 | -13.65** | 0.42 | -6.62** | -0.46 | 1.40** | 0.47** |
| P5 | 6.32** | 11.88** | 9.10** | 1.24** | 1.19* | 1.22** |
| P6 | -6.72** | -6.65** | -6.68** | -2.00** | 0.98* | -0.51** |
| P7 | 3.12** | -12.82** | -4.85** | 0.75 | -2.11** | -0.68** |
| P8 | 9.28** | -1.42** | 3.93** | 1.06** | 3.02** | 2.04** |
| L.S.D(0.05) gi | 1.24 | 1.2 | 0.48 | 0.79 | 0.97 | 0.35 |
| L.S.D(0.05) gi-gj | 1.87 | 1.82 | 0.91 | 1.20 | 1.47 | 0.66 |
| P1xP2 | -32.79** | 20.38** | -6.20** | -4.58** | -9.62** | -7.10** |
| P1xP3 | 44.14** | 21.28** | 32.71** | 9.46** | -0.03 | 4.71** |
| P1xP4 | -11.96** | -30.15** | -21.05** | -0.85 | 9.71** | 4.43** |
| P1xP5 | -17.26** | -15.62** | -16.44** | -0.39 | 5.71** | 2.66** |
| P1xP6 | 7.78** | 16.58** | 12.18** | 4.32** | -4.91** | -0.29 |
| P1xP7 | 15.94** | 8.75** | 12.35** | 4.31** | 7.58** | 5.95** |
| P1xP8 | 23.78** | 7.68** | 15.73** | 5.19** | -3.32** | 0.94 |
| P2xP3 | -8.16** | 4.78** | -1.69 | -3.50** | -3.12** | -3.31** |
| P2xP4 | -14.59** | 11.68** | -1.45 | 9.70** | 9.01** | 9.36** |
| P2xP5 | 63.78** | 5.55** | 34.66** | 11.82** | 9.85** | 10.84** |
| P2xP6 | -14.19** | -11.25** | -12.72** | -6.93** | 8.90** | 0.99 |
| P2xP7 | 43.31** | 53.25** | 48.28** | 6.02** | 3.76** | 4.89** |
| P2xP8 | -2.52 | -23.49** | -13.00** | 0.21 | 3.46** | 1.83 |
| P3xP4 | 49.34** | 9.91** | 29.63** | 15.97** | 7.77** | 11.87** |
| P3xP5 | -16.29** | -1.55 | -8.92** | -3.60** | -10.26** | -6.93** |
| P3xP6 | -8.26** | -28.69** | -18.47** | -6.36** | -5.27** | -5.81** |
| P3xP7 | 9.91** | 29.81** | 19.86** | 1.56 | 3.98** | 2.77** |
| P3xP8 | -11.59** | 17.08** | 2.75** | 4.11** | 11.78** | 7.95** |
| P4xP5 | -17.72** | 13.68** | -2.02 | -9.78** | -5.85** | -7.81** |
| P4xP6 | 2.31 | 9.55** | 5.93** | 0.67 | 3.26** | 1.97** |
| P4xP7 | -52.86** | -40.95** | -46.90** | -11.48** | -4.25** | -7.86** |
| P4xP8 | -23.02** | -14.69** | -18.85** | -10.36** | -3.28* | -6.82* |
| P5xP6 | 55.68** | 18.08** | 36.88** | 13.93** | 3.23* | 8.58** |
| P5xP7 | -72.82** | -45.09** | -58.95** | -15.29** | -3.51* | -9.40** |
| P5xP8 | 10.68** | 10.51** | 10.60** | 2.80* | 1.99 | 2.40** |
| P6xP7 | 3.54 | -11.55** | -4.00** | 6.99** | -7.63** | -0.32 |
| P6xP8 | -30.29** | 10.05** | -10.12** | -9.12** | 0.91 | -4.11** |
| P7xP8 | 27.54** | 7.21** | 17.38** | 4.86** | -11.07** | -3.10** |
| LSD5%(sij) | 3.8 | 3.69 | 2.61 | 2.43 | 2.98 | 1.89 |
| LSD5%(sij-sik) | 5.62 | 5.46 | 3.86 | 3.59 | 4.41 | 2.80 |
| LSD5%(sij-skL) | 5.3 | 5.15 | 1.29 | 3.39 | 4.15 | 0.93 |

** p < 0.05




Figure 1. Phonogram generated by UPGMA cluster analysis based on Nei and Li coefficients showing clustering of eight genotypes.

Where, P1 = Yakora Rojo, P2 = Sakha 93, P3 = Masr 1, P4 = Drought 4, P5 = Shandawel 1, P6 = Gemiza 9, P7 = Giza 171, P8 = Sides 13

This 2-fold range for both yield traits implied how genotypes unevenly responded to greatly varied irrigational environments.

The parents 4 and 5 and crosses: P1xP3, P1xP6, P2xP5, P2xP7, P3xP4, P5xP6 and P5xP8, out yielded the average genotypes in in both and across irrigation treatments. As for grain yield/ plant the parent No. 8 and the crosses P2xP4, P3xP4 and P5xP6 differed significantly relative to mean genotype.

Combining ability

General (GCA) and specific (SCA) combining ability were highly significant for all studied traits in both and across environments (Table 4). Moreover, the ratios between GCA and SCA were less than unity for the two studied traits at both and across environments except, for biological yield plant⁻¹ at normal environment. Hence, non-additive types of gene action are more relatively distinct than additive and additive x additive gene action in controlling these traits. For the exceptional case, the additive and additive x additive types of gene action are more likely pronounced than non-additive gene action in controlling this trait when subjected to such conditions. The genetic variance was previously reported (Abd El-Aty and Katta, 2002, El Hosary et al., 2012, and Gomaa et al., 2014) to be mostly

due to additive effects for grain yield/ plant.

The GCA x irrigation and SCA x irrigation mean squares were significant differed ($p < 0.05$) for the studied traits. Such result indicated that the additive and non-additive types of gene action differed significantly from one environment to another for these traits. Similar results were reported by Abd El-Aty and Katta, (2002), El Hosary et al., (2012), and Gomaa et al., (2014). The additive type of gene action was much more influenced by environments than non-additive genetic one because the ratio GCA x environment/ GCA was much higher that of SCA x irrigation/ SCA treatments for the two studied traits. Such results are in harmony with those obtained by EL Saadoon et al., 2018.

General combining ability (GCA) effects

A parent's (P_i) relative average performance (Table 6) is assessed to be advanced to further breeding programs for drought resistance selection. For per plant biological yield, P₁ ranked first for a positive \hat{g}_i effect when grown under stress environment, and it responded similarly when averaged over both environments. Also, for the same trait, P₇ had a significant positive \hat{g}_i effect under stress conditions. However, P₃ had a desirable \hat{g}_i effect yet under favorable conditions. For per plant grain yield, P₄ had significant positive \hat{g}_i effect when tested under normal

irrigation conditions and over both environments. It also took a relative top rank over both environments. For both yield traits, P₅ considered a good combiner when grown in any of the two current environments and over both of them. This response of P₅ extended to P₈ except for per plant biological yield at normal environment.

Specific combining ability \hat{s}_{ij} effects

For per plant biological yield, crosses varied in their significant positive \hat{s}_{ij} effects under each of the two irrigation regimes and averaged over both (Table 6). Among the 28 crosses, 36% (10 crosses) were under drought, 64% (18 crosses) under normal, and 46% (13 crosses) when averaged over both environments. In case of per plant grain yield, 46% (13 crosses) of all crosses had significant positive \hat{s}_{ij} effects for each environment and over both. Only 25%, seven crosses, had in common significant positive \hat{s}_{ij} effects at each environment and when averaged over the two growing conditions (Table 6). Based on the above, these crosses might be of interest in breeding programs in favor of developing pure line varieties for relatively higher biological, and grain yields/ plant under drought conditions. If a cross shows a high specific combining ability and involves only one good combiner, such combinations would result in desirable transgressive segregates. Providing that an additive genetic system is present in a good combiner and a complementary and epistatic effects are present in a cross. All act in the same direction to reduce undesirable plant characteristics and maximize the character in view.

Correlation between genetic distance and each of mean performance and SCA

For the 28 hybrids, Pearson's coefficients of correlation, r , between GD and mean performance for per plant biological yield, were 0.353, 0.205, and 0.334 at drought, normal, and over both environments. Its values were 0.417 ($p < 0.05$), 0.201, and 0.366 between GD and SCA, which did not differ much from GD's r values with per plant biological yield. On the other hand, the r values between GD and per plant grain yield were quite higher ($p < 0.05$): 0.520, 0.615, and 0.695. These results throw light on how GD is related to heterosis for both yield characters. The reason for high morbidity may be due to the number of genes that control the inheritance of grain yield compared to biological yield. This study showed that GD can be used to precisely predict the yield

performance and heterosis value for F₁ hybrids. This suggests that the more primers/markers be applied, the better detection of differences between parents.

CONCLUSION

To sum up, specific combining ability effects, high mean performance and hybrid vigor produce from hybridization between diverse parental. Genetic diversity (GD) can be estimated from RAPD-PCR marker. Thus, its easy to predict the hybrid performance from screening parents by RAPD primers.

CONFLICT OF INTEREST

All authors contributed in collecting and analyzing data. All authors participated in writing every part of this study. All authors read and approved the final version.

ACKNOWLEDGEMENT

The authors would like to thank Benha University for its contribution to providing equipments through project D / 1/2 entitled "Improvement of wheat productivity under lower inputs", funded by the fund for scientific research account (Phase IV). Also, authors greatly indebted by all means to Prof. Dr. M.M. Iraqi (dean) and Prof. Dr. N.K. Elgizawy (vice dean) at Fac. Agric. for their supporting any facilities to conduct this research.

AUTHOR CONTRIBUTIONS

This work was carried out in collaboration between all authors. Author A.A. El-Hosary designed the study, wrote the protocol and wrote the first draft of the manuscript. Authors A.A. El-Hosary, A.A.A. El-Hosary and M. El. M. El-Badawy supervised the study and managed the literature searches. Author M. El. M. El-Badawy, A.A.A. El-Hosary, Tamer El Akaad and A. El-Fahdawy managed the experimental process and performed data analyses. All authors read and approved the final manuscript. The Authors have read and are fully aware of the journal policy.

Copyrights: © 2019 @ author (s).

This is an open access article distributed under the terms of the [Creative Commons Attribution License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or

reproduction is permitted which does not comply with these terms.

REFERENCES

- Abd El-Aty MA , Katta YS, 2002. Genetic analysis and heterosis of grain yield and related traits in bread wheat (*Triticum aestivum* L.). J. Agric. Res. Tanta Univ., 28 (2): 287-300.
- Cao W, Hucl P, Scoles G, Chibbar RN, Fox PN, Skovmand B, 2002. Cultivar identification and pedigree assessment of common wheat based on RAPD analysis. Wheat Information Service Number 95: 29-35.
- Chowdhry MA, Rafiq M, Alam K, 1992. Genetic architecture of grain yield and certain other traits in bread wheat. Pakistan J Agric Res 13: 216-220.
- EL Saadoon AW, EL Hosary AA, Sedhom SA, ELBadawy MELM and El Hosary AAA, 2018. Determination of Combining Ability and Genetic Diversity Using ISSR Markers to Evaluate the Genetic Variability in Wheat. The 7th Scientific and 1st International Conference of Agricultural Researches, 10-11 April 2018. 18: 67-80
- EL Saadoon AW, EL Hosary AA, Sedhom SA, ELBadawy MELM, El Hosary AAA, 2017. Genetic analysis of diallel crosses in wheat under stress and normal irrigation treatments. Egypt. J. Plant Breed. 21 (5): 279 – 292.
- EL-Hosary AA, EL-Badawy MEM, Mustafa AK, EL-Shal MH, 2012. Evaluation of diallel wheat crosses under drought tolerance. Egypt. J. Plant Breed. 16 (1): 19-40.
- El-Hosary AAA, Nour-El-Deen GA, 2015. Genetic analysis in the F₁ and F₂ wheat generations of diallel crosses. Egypt. J. Plant Breed. 19 (2): 355-373.
- Gomaa MA, El-Banna MNM, Gadalla AM, Kandil EE, Ibrahim ARH, 2014. Heterosis, combining ability and drought susceptibility index in some crosses of bread wheat (*Triticumaestivum* L.) under water stress conditions . Middle- East j. Agric. Res., 3(2): 338-345.
- Gomez KN, Gomez AA, 1984. Statistical Procedures for Agricultural Research. John. Wiley and Sons. Inc., New York, 2nd ed.
- Griffing B, 1956. Concept of general and specific combining ability in relation to diallel crossing system. Aust. J. Biol. Sci.9: 463-493.
- Guo-yue C, Hui LL, 2007. Detection of genetic diversity in synthetic hexaploid wheats using microsatellite markers. Agri. Sci. in China, 6(12): 1403-1410.
- Jaccard P, 1908. Nouvelles recherches sur la distribution florale. Bull. Soc. Vaudoise Sci. Nat., 44: 223-270.
- Kumar A, Sengar RS, Rao VP, Shukla G, Dixit R, Singh R, (2017) Assessment of genetic diversity in bread wheat (*Triticum aestivum* L.) using RAPD markers. Journal of Applied and Natural Science 9 (3): 1751 -1755.
- Nei M, Lis WEN-H, 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases Proc. Natl. Acad. Sci. USA ., 76 (10) 5269-5273.
- Rohlf FJ, 2000. NTSYS-pc: Numerical taxonomy and multivariate analysis system Version 2.1. Exeter Publishing Setauket, New York.
- Shrief SA, Abd El-Shafi MA, El-Sadi SA, 2017. Heterosis, gene action and combining ability of grain yield and its components in six bread wheat crosses. Bioscience Research, 14(4): 1204-1215
- Siddiqui MF, Iqbal S, Erum S, Khan S, 2010 DNA landmarks for genetic relatedness and diversity assessment in Pakistani wheat genotypes using RAPD markers. Pak. J. Bot. 42(2):1013-1020.
- Steel RGD, Torrie JH, 1980. Principles and Procedures of Statistics: A Biometrical Approach. McGraw Hill Book Co., Inc., New York, USA.
- Tahir NA, 2008. Assesment of genetic diversity among wheat varieties in sulaimanyah using random amplified polymorphic DNA (RAPD) analysis. Jordan Journal of Biological Sciences. 1(4): 159-164.
- Tonk FA, Tosun M, Ilker E, Istipliler D, Tatar O, 2014. Evaluation and comparison of ISSR and RAPD markers for assessment of genetic diversity in triticales genotypes. Bulgarian J. of Agric., 20 (6) :1413-1420 .
- Topal A, Aydin C, Akgiin N, Babaoglu M, 2004. Diallel cross analysis in durum wheat (*Triticum durum* Desf.): identification of best parents for some kernel physical features. Field Crop Res 87(1): 1-12.
- Williams, JGK, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV, 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. Nucleic Acid Res. 18: 6531-6535.