

*Annals of Agric. Sc., Moshtohor,*  
*Vol. 32(1):217-228, 1994*

**GENERAL AND SPECIFIC COMBINING ABILITY INTERACTIONS  
WITH YEAR IN MAIZE  
BY**

**El-Hosary, A.A.; Mohamed, M.K.; Sedhom, S.A.  
and Abo El-Hassan, G.K.A.**

Agronomy Dept., Fac. of Agric. Moshtohor, Zagazig Univ., Egypt.

**ABSTRACT**

General (GCA) and specific (SCA) combining abilities and their interaction with year were investigated in one set of F's among eight inbreeds lines. Significant GCA was detected in the combined analysis for most of the traits except for the date of mid-silking. Also, significant SCA was detected for all traits with the exception of ear height, ear length and mid-tasselling date. Both GCA\*year and SCA\*year were significant only for mid-silking date, plant height and number of kernels/row. Calculated ratios of GCA\*year/GCA was higher than the ratio GCA\*year/SCA for mid-silking date indicating that the additive and additive by additive effects were highly influenced by year for this trait. As for plant height and number of kernels/row, the ratio SCA\*year/SCA was higher than GCA\*year/GCA indicating that the non-additive effects were more influenced by year. Significant GCA by year interaction was detected for mid-tasselling date. On the contrary, significant SCA by year interaction was detected for ear diameter and grain yield/plant.

**INTRODUCTION**

Interaction of general (GCA) and specific (SCA) combining ability with year is one of the aspects that requires special attention. Diallel cross analysis has been used effectively to estimate the interaction of GCA and SCA by environmental factors. In retrospect, Matzinger *et al.* (1959) reported that the additive genetic variance was much more influenced by environment than the non-additive portion. Singh *et al.* (1977), however, reported that the additive genetic variance was stable over environment for at least 14 characters; only the characters of mid-silking date, ear height and tryptophan content showed the presence of additive by location interaction. Stuber *et al.* (1977) reported that estimates of GCA and SCA showed significant interaction with plant density for ear and plant heights. Likewise, El-Zeir (1984) found that both GCA and SCA were stable over plant density with the exception of plant and ear height. Nawar (1985) found that GCA and SCA by year interaction were significant with the latter being of higher magnitude than the former. Galal *et al.* (1987) indicated that GCA interaction with location was of higher magnitude than SCA by location. Galal *et al.* (1977) and El-Hosary *et al.* (1988a) found significant interaction of planting date with both types of combining

ability with year. El-Hosary *et al.* (1988) reported significant interaction of both types of combining ability and location for mid-tasselling date, plant height, number of rows/ear, 100-kernel weight and grain yield/plant. Badr (1989) and El-Hosary and Sedhom (1990) reported that the additive genetic variance was more biased up by the interaction with the environment than the non-additive effects. Finally, El-Zeir (1990) reported that the effects of location, year and plant density were more prominent on GCA and SCA than nitrogen fertilization.

Here, we report estimates of GCA and SCA in a set of 28 F<sub>1</sub> crosses, derived from 8 inbreds with the aim to elucidate the effect of year on both estimates.

### MATERIALS AND METHODS

28-F<sub>1</sub> crosses were used in this study. These were the progeny of 8 inbred parents. These include the six inbreds designated as Mosht<sub>1</sub>, Mosht<sub>2</sub>, Mosht<sub>3</sub>, Mosht<sub>18</sub>, Mosht<sub>19</sub> and Mosht<sub>17</sub>, respectively. These are referred to herein as P<sub>1</sub>, P<sub>2</sub>, P<sub>3</sub>, P<sub>4</sub>, P<sub>5</sub> and P<sub>6</sub>, respectively. The latter two were G303A and Rg10. Pedigree and credentials of inbreds are given elsewhere by Abo-El-Hassan (1994). In 1991 and 1992 seasons the single crosses were grown at the Experimental station of the Faculty of Agriculture at Moshtohor, Kalubia on the 20<sup>th</sup> of June. The experimental design was complete randomized blocks with three replications. Plots were two ridges each of 6 m long and 70 cm wide. Intra hills spacing was 20 cm. Hills were thinned to one plant/hill. Plots were treated alike as to fertilization, irrigation and other cultural practices. Data collected included mid-tasselling date, mid-silking date, plant height (cm), ear height (cm), ear length (cm), ear diameter (cm), number of rows/ear, number of kernels/row, weight of ear/plant and grain yield/plant.

Data obtained were taken on individual plant basis except for mid-tasselling and mid-silking date which were taken on whole plot basis. The ordinary anova was firstly performed on separate seasons assuming a fixed model. A combined analysis of both seasons was afterwards carried out after checking the homogeneity of the error variance. A one tail F test was used to test the significance of various sources of variations. General and specific combining abilities were computed according to Griffing's (1956) Model 1 (fixed effects) Method 4.

### RESULTS AND DISCUSSION

Combined sum of squares for GCA and SCA are shown for various traits in Table (1) together with the calculated ratio GCA/SCA. GCA sum of squares were highly significant for the ten studied characters. SCA sum of squares were highly significant for plant height, mid-dates of tasselling and silking, ear diameter, number of grains/row, ear weight and grain yield/plant and only significant for ear height and ear length and number of rows/ear. Such results emphasized the

Table (1): Combined mean squares from ordinary analysis for ten agronomic characters of maize.

| S.O.V.             | d.f. | Plant height (cm) | Ear height (cm) | Mid silking date | Mid tasseling date | Ear length (cm) | Ear diameter (cm) | No. of rows/ear | No. of grains/row | Ear weight (g) | Grain yield (g)/plant |
|--------------------|------|-------------------|-----------------|------------------|--------------------|-----------------|-------------------|-----------------|-------------------|----------------|-----------------------|
| Year (Y)           | 1    | 7791.32**         | 1461.01**       | 70.72            | 1.53               | 9.33            | 0.23              | 16.46**         | 202.61**          | 984.00         | 9320.25**             |
| Blocks within      | 4    | 124.46            | 320.20          | 2.69             | 4.74               | 12.85           | 0.19*             | 0.53            | 4.70              | 905.83*        | 725.52*               |
| Year Genotypes (G) | 27   | 1817.93**         | 1075.59**       | 11.20**          | 19.36**            | 10.46**         | 0.39**            | 2.50**          | 58.57**           | 3554.72**      | 2146.89**             |
| Genotype X year    | 27   | 411.06**          | 171.31          | 6.49**           | 8.54**             | 2.70            | 0.13**            | 3.49**          | 26.74**           | 345.52         | 266.54                |
| GCA                | 7    | 6303.84**         | 3433.47**       | 12.76**          | 34.89**            | 31.29**         | 1.02**            | 5.18**          | 138.88**          | 9925.08**      | 5731.12**             |
| SCA                | 20   | 247.86**          | 250.47*         | 10.65**          | 13.92**            | 3.16*           | 0.17**            | 1.56*           | 30.86**           | 1325.09**      | 888.36**              |
| GCA X year         | 7    | 453.38**          | 143.05          | 10.58**          | 14.36**            | 3.02            | 0.10              | 1.24            | 33.32**           | 127.42         | 111.47                |
| SCA X year         | 20   | 396.25**          | 181.20          | 5.05**           | 3.01               | 2.59            | 0.14**            | 4.28            | 24.48**           | 421.85         | 320.83                |
| Error              | 108  | 116.23            | 122.30          | 1.67             | 2.74               | 1.70            | 0.06              | 0.82            | 5.22              | 311.69         | 189.13                |
| GCA/SCA            |      | 21.39             | 13.71           | 1.19             | 2.51               | 9.90            | 6.11              | 3.32            | 4.50              | 7.49           | 6.45                  |

\* and \*\* significant at 0.05 and 0.01 levels of probability, respectively.

importance of both estimates in the inheritance of the traits in this group of inbred lines. In addition high calculated GCA/SCA ratios exceeding by for the unity were obtained for all traits indicating the overwhelming effect of the additive genetic portion over the non-additive in the inheritance of all ten traits.

**GCA\*season an SCA\*season interaction:**

Results in Table (1) show that both GCA and SCA by season sum of squares were significant for mid-silking date, plant height and number of kernels/row. Also the ratios of SCA\*season/SCA were higher than the ratio of GCA\*season/GCA, for the plant height and number of kernels/row. This indicates that the non-additive genetic variance was influenced more by season than the additive one for these two respective traits. However, for mid-silking date the opposite was true indicating that the additive and additive by additive effects were more influenced by season for this particular trait. Results also show significant GCA\*season and insignificant SCA\*season for mid-tasselling date indicating that the additive effects only interplayed with season. However, significant SCA\*season and insignificant GCA\*season was obtained for ear diameter and grain yield/plant indicating that SCA alone interplayed with season.

Data on GCA effects are shown in Table (2). Evidently, P<sub>1</sub> was a good combiner for grain yield/plant, most of yield components and mid silking and mid-tasselling date. P<sub>3</sub> ranked second expressing favorable GCA for grain yield and some yield components. Significant and positive SCA effects are manifested in the five crosses (P<sub>2</sub>xP<sub>5</sub>), (P<sub>2</sub>xP<sub>8</sub>), (P<sub>3</sub>xP<sub>8</sub>), (P<sub>5</sub>xP<sub>6</sub>) and (P<sub>6</sub>xP<sub>7</sub>) for grain yield/plant, Table (3). Also, results show that the single crosses (P<sub>1</sub>xP<sub>6</sub>), (P<sub>2</sub>xP<sub>5</sub>) and (P<sub>2</sub>xP<sub>8</sub>) were the best combinations.

Favorable SCA is manifested in seven crosses for mid-silking date, of which P<sub>3</sub>xP<sub>4</sub> had the highest negative value. Favorable SCA effects for mid-tasselling is manifested in eleven crosses of which P<sub>3</sub>xP<sub>5</sub> had the highest value.

Seven favorable SCA effects for shortening plant height are encountered; the highest negative value is manifested by the cross P<sub>1</sub>xP<sub>4</sub>.

Five favorable SCA effects in the direction for lowering ear height are encountered of which P<sub>1</sub>xP<sub>7</sub> expressed the highest negative value.

For ear characteristics the most favorable SCA effects are shown in the crosses (P<sub>5</sub>xP<sub>8</sub>) for ear length, (P<sub>1</sub>xP<sub>2</sub>), (P<sub>2</sub>xP<sub>8</sub>) and (P<sub>3</sub>xP<sub>8</sub>) for ear diameter; (P<sub>7</sub>xP<sub>8</sub>) for number of rows/ear (P<sub>3</sub>xP<sub>7</sub>), (P<sub>2</sub>xP<sub>6</sub>) and (P<sub>3</sub>xP<sub>8</sub>) for number of kernels/row; (P<sub>2</sub>xP<sub>8</sub>) for ear weight; and (P<sub>1</sub>xP<sub>6</sub>), (P<sub>3</sub>xP<sub>5</sub>) and (P<sub>2</sub>xP<sub>8</sub>) for grain yield/plant.

The results conclusions of this study on the basis of pooled data analysis of variance of combining ability over the two seasons show that both SCA and SCA mean squares were significant for all characters. Although SCA mean squares

Table (2): Estimates of general (GCA) combining ability effects for ten agronomic and yield traits for the eight parents in 1991 and 1992 seasons.

| characters    | Parent | year | plant height (cm) | Ear height (cm) | Mid silking date | Mid tasseling date | Ear length (cm) | Ear diameter (cm) | No. rows/ear | No. kernels/row | Ear weight (g) | Grain yield (g/plant) |
|---------------|--------|------|-------------------|-----------------|------------------|--------------------|-----------------|-------------------|--------------|-----------------|----------------|-----------------------|
| P1            | Y1     |      | 21.18             | 17.32           | -1.69            | -0.85              | 0.93            | 0.31              | 0.66         | 1.77            | 27.29          | 19.53                 |
|               | Y2     |      | 30.98             | 20.45           | N.S.             | -1.09              | 1.51            | 0.23              | 0.66         | 4.50            | 26.50          | 23.05                 |
|               | Comb.  |      | 25.97             | 18.88           | -0.90            | -1.51              | 1.22            | 0.27              | 0.66         | 3.08            | 21.29          | 15.41                 |
| P2            | Y1     |      | 9.99              | 5.93            | -1.19            | 0.38               | 1.15            | 0.26              | 0.34         | 2.11            | 19.73          | 15.41                 |
|               | Y2     |      | 14.00             | 6.05            | N.S.             | -0.48              | 1.45            | 0.11              | -0.69        | 3.63            | 18.96          | 12.14                 |
|               | Comb.  |      | 11.92             | 5.99            | -0.56            | -0.29              | 1.30            | 0.18              | 0.51         | 3.08            | 20.04          | 13.78                 |
| P3            | Y1     |      | -8.46             | 8.18            | 0.42             | 0.26               | 0.68            | -0.07             | -0.03        | 0.75            | -12.72         | -6.66                 |
|               | Y2     |      | -13.29            | 10.08           | N.S.             | 0.49               | -0.72           | -0.01             | 0.15         | -0.91           | -11.06         | -6.64                 |
|               | Comb.  |      | -10.96            | 9.13            | 0.27             | 0.65               | -0.70           | -0.04             | -0.37        | -0.86           | -6.35          | -2.02                 |
| P4            | Y1     |      | -10.13            | 6.07            | -0.36            | 0.65               | -0.61           | -0.11             | -0.30        | -1.70           | -4.92          | -3.18                 |
|               | Y2     |      | -7.84             | 6.01            | N.S.             | -0.77              | 0.05            | 0.05              | -0.10        | -1.48           | 2.06           | 0.29                  |
|               | Comb.  |      | -9.05             | 6.04            | -0.06            | -0.24              | -0.69           | -0.07             | -0.30        | -1.70           | -4.92          | -3.18                 |
| P5            | Y1     |      | -2.57             | 0.54            | 0.97             | 0.32               | -0.37           | 0.05              | -0.10        | -1.48           | 2.06           | 0.29                  |
|               | Y2     |      | 2.83              | 2.76            | N.S.             | 0.65               | -0.20           | -0.08             | -0.20        | -1.68           | 5.48           | 3.02                  |
|               | Comb.  |      | 0.06              | 1.68            | 0.63             | -1.79              | -0.09           | 0.06              | -0.17        | 0.18            | -12.34         | -8.67                 |
| P6            | Y1     |      | -4.90             | 8.79            | -0.31            | 0.65               | -0.46           | -0.13             | -0.20        | -1.68           | 5.48           | 3.02                  |
|               | Y2     |      | -16.83            | 7.98            | N.S.             | -1.18              | -1.60           | -0.21             | -0.17        | 0.18            | -12.34         | -8.67                 |
|               | Comb.  |      | -10.93            | 8.42            | -0.48            | 0.15               | -1.03           | -0.17             | -0.31        | -0.76           | -14.64         | -11.24                |
| P7            | Y1     |      | -9.79             | 9.84            | 0.64             | 0.15               | 0.81            | -0.19             | -0.01        | -2.35           | 21.51          | -17.50                |
|               | Y2     |      | -8.50             | 5.55            | N.S.             | 0.79               | -0.49           | -0.25             | -0.25        | -1.02           | -22.12         | -16.78                |
|               | Comb.  |      | -9.21             | 7.70            | 0.38             | 0.88               | -0.65           | -0.23             | -0.32        | -0.11           | 3.83           | -0.35                 |
| P8            | Y1     |      | -4.76             | 9.10            | 1.53             | 0.79               | 0.86            | -0.09             | -0.32        | -0.11           | 3.83           | -0.35                 |
|               | Y2     |      | -1.31             | 0.37            | 0.43             | 0.88               | 0.43            | 0.08              | -0.01        | -0.08           | -0.37          | -0.24                 |
|               | Comb.  |      | 1.66              | 4.74            | 0.72             | 1.29               | 0.65            | -0.01             | -0.01        | 0.98            | 8.25           | 6.63                  |
| L.S.D. (g)    | Y1     | 5%   | 4.66              | 4.68            | 0.68             | 0.78               | 0.59            | 0.12              | 0.40         | 1.30            | 11.00          | 8.84                  |
|               | Y2     | 5%   | 6.21              | 6.42            | 0.90             | 1.04               | 0.79            | 0.16              | 0.54         | 1.30            | 11.00          | 8.84                  |
|               | Comb.  | 5%   | 4.89              | 5.11            | 1.36             | 1.57               | 1.20            | 0.24              | 0.81         | 1.97            | 16.63          | 13.37                 |
| L.S.D. (g-qt) | Y1     | 1%   | 6.51              | 6.82            | 0.40             | 0.51               | 0.40            | 0.08              | 0.28         | 0.70            | 5.45           | 9.10                  |
|               | Y2     | 1%   | 3.41              | 3.41            | 0.53             | 0.68               | 0.53            | 0.10              | 0.37         | 0.94            | 7.24           | 12.09                 |
|               | Comb.  | 5%   | 4.42              | 4.53            | 1.02             | 1.18               | 0.90            | 0.18              | 0.61         | 1.47            | 12.47          | 10.03                 |
| L.S.D. (g-qt) | Y1     | 5%   | 7.05              | 7.08            | 1.36             | 1.57               | 1.20            | 0.24              | 0.81         | 1.97            | 16.63          | 13.37                 |
|               | Y2     | 5%   | 7.39              | 7.73            | 1.36             | 1.57               | 1.20            | 0.24              | 0.81         | 1.97            | 16.63          | 13.37                 |
|               | Comb.  | 5%   | 9.86              | 10.30           | 0.88             | 1.13               | 0.89            | 0.17              | 0.62         | 1.56            | 12.06          | 9.40                  |
| L.S.D. (g-qt) | Y1     | 1%   | 7.37              | 7.56            | 1.17             | 1.50               | 1.18            | 0.22              | 0.82         | 2.07            | 16.02          | 12.48                 |
|               | Comb.  | 1%   | 9.78              | 10.04           | 1.17             | 1.50               | 1.18            | 0.22              | 0.82         | 2.07            | 16.02          | 12.48                 |













were significant for all characters, still most of the total genetic variance was associated with GCA as shown by its greater magnitude. GCA and SCA were found to interact with season for plant height, mid-silking date and number of kernels/row. This indicates that both additive and non-additive effects of parents were bound to change by season. No interaction of season with GCA or SCA was detected for ear height, ear length, number of rows/ear, ear weight and grain yield/plant indicating the stability of GCA effects over seasons for these traits. For ear diameter only SCA showed sensitivity to change over season.

These results support earlier work by Singh *et al.* (1977), Stuber *et al.* (1977), Nawar (1985) and El-Hosary (1985, 1988 and 1989). Hence, additive and additive by additive epistasis, if present could be exploited in breeding for various characters of corn. Thus, selection on the basis of GCA effects, as suggested by Jensen (1970) in cereal crops is by far of great importance. Interestingly enough, the parents P<sub>1</sub> and P<sub>3</sub> showing favorable GCA for yield are also blessed with high number of favorable SCA effects.

#### REFERENCES

- Abo El-Hassan, G.K.A. (1994): Heterosis and combining ability for some quantitative characters in maize. M.Sc. Thesis, Fac. of Agric. Moshtohor, Zagazig Univ., Benha Branch, Egypt.
- Badr, M.M. (1989): Breeding studies on some divergent maize populations. M.Sc. Thesis, Fac. Agric. Moshtohor, Zagazig Univ., Egypt.
- El-Hosary, A.A. (1985): Study of combining ability in some top crosses in maize. Egypt. J. Agron. 10(1-2):39-47.
- El-Hosary, A.A. (1988): Heterosis and combining ability of ten maize inbred lines as determined by diallel crossing over two planting dates. Egypt. J. Agron. 13(1-2):13-25.
- El-Hosary, A.A. (1988): An analysis of combining ability of inbred lines of maize (*Zea mays* L.) in diallel cross system. Egypt. J. Agron., 13(1-2):27-39.
- El-Hosary, A.A. (1989): Heterosis and combining ability of six inbred lines of maize in diallel crosses over two years. Egypt. J. Agron., 14(1-2):47-58.
- El-Hosary, A.A. and Sedhom, S.A. (1990): Diallel analysis of yield and other agronomic characters in maize (*Zea mays* L.). Ann. of Agric. Sc., Moshtohor, 28(4):1987-
- El-Zeir, F.A.A. (1984): Combining ability and heterosis of some inbred lines of maize. Thesis, Fac. Agric., Zagazig Univ., Egypt.
- El-Zeir, F.A.A. (1990): Genetic studies on some inbred lines of maize and their single crosses. Ph.D. Thesis, Fac. of Agric., Al-Azher Univ., Egypt.
- Galal, A.A., El-Zeir, F.A. and Younis, M.A. (1987): Estimation of general and specific combining ability in three sets of new inbreds of maize. J. Agric. res. Tanta Univ. Egypt.
- Griffing, N.E.G. (1958): Concept of general and specific combining ability in relation to diallel crossing systems. Aus. J. Biol. Sci., 9:463-493
- Jensen, N.F. (1970): Diallel selective mating system for cereal breeding. Crop Sci. 10:629-650

- Matzinger, D.F.; Sprague, G.F. and Cockerham, C.C. (1959): Diallel crosses of maize in experiments reported over locations and years. *Agron. J.* 51:346-350.
- Nawar, A.A. (1985): Hybrid vigor and combining ability and their interaction with years in maize (*Zea mays* L.) Minufiya J. Res. 10:115-132.
- Singh, S.V.; Singh, J. and Kaht, D.K. (1977): Genetic environmental interaction in a diallel cross of opaque-2 maize. *Maydica* 22:89-95.
- Stuber, C.W. and Moll, R.H. (1977): Genetic variances and hybrid production of maize at two plant densities. *Crop Sci.* 17:503-506.

دراسة للقدرة العامة والخاصة على التآلف وتفاعلها بالبيئة  
في بعض سلالات الذرة الشامية

على الحصرى ، محمد قاسم محمد ، سيدهم أسعد سيدهم ، جمال محمد قرنى  
قسم المحاصيل - كلية الزراعة بمشتهر - جامعة الزقازيق - مصر .

أجرى هذا العمل بهدف دراسة القدرة العامة والخاصة على التآلف وتفاعلها مع السنوات لبعض سلالات الذرة الشامية. أستخدم لهذا الغرض ثمانية سلالات تمثل مدى الأختلاقات الوراثية لعشرة من الصفات وفي ١٩٩٠ تمت زراعة السلالات الأبوية وأجرى بينهم جميع التهجينات الفردية الممكنة عدا الهجن العكسية وتم الحصول على ثمانية وعشرون هجينا فرديا. وفي سنة ١٩٩١، ١٩٩٢ تم تقييم هذه الهجن الفردية. أجريت التجريبتان في التصميم القطاعات الكاملة العشوائية في ثلاث مكررات بالنسبة لعامي الدراسة.

أجرى التحليل التجميى لنباتات التجريبتان تمت تقديرات القدرة العامة والخاصة على التآلف طبقا للطريقة الرابعة في النموذج الأول لجريفتنج. وكانت أهم النتائج كما يلي :

- ١- كان التباين الراجع للسنوات معنويا لجميع الصفات المدروسة: وهي: طول النبات - ارتفاع الكوز - قطر الكوز - عدد صفوف الكوز - عدد حبوب الصف - وزن الكوز - وزن الحبوب/نبات - موعد خروج ٥٠% للحريرة والنورة المذكورة.
- ٢- كان تباين الهجن عالى المعنوية لجميع الصفات المدروسة.
- ٣- كان التباين الراجع لتفاعل كلا من القدرة العامة والخاصة على التآلف مع السنوات معنويا لميعاد خروج الحريرة وارتفاع النبات وعدد حبوب الصف وكان تفاعل القدرة الخاصة للتآلف في السنوات معنويا ومصحوبا بعدم معنوية القدرة العامة على التآلف بالنسبة لصفى قطر الكوز ومحصول الحبوب/نبات.
- ٤- أستأثر مجموع المربعات الخاص بالقدرة العامة على جزء كبير من مجموع مربعات التباين الوراثى مما يدل على أهمية هذا الجانب في توريث الصفات المدروسة.
- ٥- أظهرت النتائج وجود تفاعل عالى المعنوية بين القدرة العامة للأتلاف والسنوات مشيرا الى حساسية هذا التقدير لظروف السنوات بالنسبة لصفات ارتفاع النبات، عدد الحبوب/صف - خروج ٥٠% حريرة وخروج ٥٠% شوشة.
- ٦- أظهرت النتائج وجود تفاعل عالى المعنوية بين القدرة الخاصة للأتلاف والسنوات مشيرا الى حساسية هذا التقدير لظروف السنوات بالنسبة لصفات ارتفاع النبات قطر الكوز، عدد حبوب الصف.
- ٧- لم يتأثر كل من القدرة العامة والخاصة للأتلاف بالنسبة للصفات الأتية: ارتفاع الكوز، طول الكوز، عدد صفوف الكوز، وزن الكوز ووزن الحبوب/نبات مما يدل على ثبات هذه التقديرات تحت ظروف البيئة.