



Egyptian Journal of Rabbit Science

Published by : **Egyptian Rabbit Science Association**

ESTIMATION OF CROSSBREEDING EFFECTS FOR SOME LITTER TRAITS IN CROSSING OF NEW ZEALAND WHITE WITH EGYPTIAN GABALI RABBITS

M.H. Khalil¹, E.A. Afifi², A. Farid³, Y.K. Youssef³

¹Department of Animal Production & Breeding, College of Agriculture & Veterinary Medicine, King Saud University, Buriedah P.O.Box 1482, Saudi Arabia

²Department of Animal Production, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt

³Animal Production Research Institute, Ministry of Agriculture, Dokki, Cairo, Egypt.

A crossbreeding experiment was carried out involving New Zealand White (N) and Egyptian Gabali (G) rabbits to produce six genetic groups of NN, GG, $\frac{1}{2}G\frac{1}{2}N$, $\frac{1}{2}N\frac{1}{2}G$, $\frac{3}{4}G\frac{1}{4}N$ and $\frac{3}{4}N\frac{1}{4}G$. Estimates of direct (G^d) and maternal (G^m) additive effects, direct (H^d) and maternal (H^m) heterosis and direct recombination effects (R^d) were genetically evaluated for litter size at birth (LSB) and weaning (LSW) and litter weight at weaning (LWW).

N rabbits had superior performance in litter size and weight compared to G rabbits. Crossbred litters obtained from mating N bucks with G does were generally associated with superiority compared to those litters obtained from the reverse mating. In most cases, genetic group of $\frac{1}{2}N\frac{1}{2}G$ gave larger litter size and heavier litter weight compared to the other genetic groups. The estimates of crossbreeding effects for LSB, LSW and LWW were 2.62, 7.52 and 12.37 % for G^d , -2.34, 9.56 and -7.50 % for G^m , 9.5, 19.2 and 15.8 % for H^d , -4.31, 3.43 and -3.77 % for H^m , and 45.5, 17.8 and 23.8 % for R^d , respectively. The estimates of direct additive effects were moderate or high and in favour of N rabbits and consequently N bucks could be used as sires in crossbreeding stratification systems under hot climatic conditions. The estimates of G^m for all traits were somewhat low and in favour of G rabbits. Crossbred dams recorded favourable maternal heterotic effects on litter size at weaning and unfavourable maternal heterotic effects on litter size at weaning, i.e. crossbred dams recorded larger litter size at weaning associated with a reduction in litter weight at weaning than their crossbred daughters. Estimates of direct recombination effects for litter size and weights at weaning were insignificant and indicate that epistatic recombination effects for crossbred litters during the suckling period were negligible.

Information for contributors

Membership of the Society of Rabbit Sciences:

Egyptian Journal of rabbit science is published two time a year. Annual dues in the society which include a subscription to the journal are 12 L.E., in Egypt and 50 US Dollars in other countries. Dues are payable in advance, each year.

Subscription prices for non members are 15 Egyptian pounds and 15 US Dollars, respectively for a single part net.

Form and style of Manuscripts:

- Papers submitted should tackle rabbit production problems in an original way. Review papers on subjects of general interest and a "News and Notes" section may be published.
- Manuscripts of 3 copies should be typed double-spaced on quarto size with a wide margin. The manuscript should not be longer than 10 pages including tables, figures and references.
- Following the title(s), author(s) name(s) (with initials preceding) and address(es), articles are often divided into the following sections: Summary (not more than 20 lines), Introduction, Materials and Methods, Results and Discussion, Conclusions, Acknowledgement(s) and References.
- Figures: Values of x and axes and title of the figures(s) for each figure to be developed by computer when printing the Journal. Photos should be high gloss-prints of good contrast, size 13 by 18c.

- Line drawing with Chines ink, should be on white or transparent paper. All figures should be numbered consecutively. A separate list of clear captions for illustrations has to be added.
- The approximate location of figures and tables in the text should be indicated in the margins.
- References on. The text should be made by name(s) of the author(s) and the years and the References should have an alphabetical order giving: Surname and abbreviated first name(s) of the author(s), year, title of the paper, name of the Journal complete (not abbreviated), number of the volume, number of first and last pages., and for books title and place of publication.
- Underline the Latin names and (in the References) title of books.
- All pages including those of tables and figures must be numbered.
- On publication of his paper, each first author receives 20 off prints of his paper.
- The costs for publication are 100 L.E. for each paper of (in maximum) 10 L.E. The costs for non-Egyptians are 50 and 5 US include refereeing costs.
- The definitive draft, typed and recorded in a diskette should be submitted in word star, word perfect or as a ASCII file.
- Manuscripts should be addressed to: Prof. Dr. A.H. Daader, Editor in Chief, Department of Agriculture, Zagazig University, Zagazig, Egypt

Key words: Egyptian Gabali rabbits, crossbreeding, litter size and weight, direct and maternal additive, heterosis, recombination effects.

Direct and maternal heterosis, direct and maternal additive effects and direct and maternal recombination losses for reproductive efficiency from crossbreeding experiments including Gabali rabbits were of considerable importance (Ali, 1998; Abdel-Aziz, 1998). The New Zealand White was found to exhibit an outstanding maternal ability as related to behavior, fecundity and lactation. Results of most crossbreeding experiments carried out in Egypt reported that crossing of New Zealand White breed with Gabali or other local breeds were generally associated with considerable heterotic effects on most litter and growth traits (Oudah, 1990; El-Desoki, 1991; Khalil *et al*, 1995; Ali, 1998; Abdel-Aziz, 1998).

The objective of the present paper was to estimate direct (G^I) and maternal (G^M) additive effects and direct (H^I) and maternal (H^M) heterosis as well as direct recombination loss (R^I) for litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW) in crossbreeding experiment involving New Zealand White (N) and Egyptian Gabali (G) rabbits.

MATERIALS AND METHODS

A crossbreeding project involving New Zealand White (N) and Egyptian Gabali (G) rabbits was started in September 1994 in the experimental rabbitry, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt.

Breeding plan and management:

Rabbits involved in this crossbreeding experiment represented desert Egyptian Gabali rabbits (came from north Sinai and raised in the rabbitry of Maryout Experimental Station, Desert Research Center since 1991) and New Zealand White (bought from Bank El-Nil rabbitry). The breeding plan permitted the simultaneous production of NN, GG, $\frac{1}{2}G\frac{1}{2}N$, $\frac{1}{2}N\frac{1}{2}G$, $\frac{3}{4}G\frac{1}{4}N$ and $\frac{3}{4}N\frac{1}{4}G$ litters. Matings started in September 1995. Distribution of breeding does and their sires, dams and bucks as well as number of litters of the six genetic groups are presented in Table 1. The bucks were randomly assigned to mate the does with a restriction to avoid the matings of animals with common grandparents. Each buck

Table 1. Mating groups and number of sires, dams, does, bucks and litters used in different genetic groups of the study

Mating group ⁺			Sires	Dams	Does	Bucks	Litters
Sires	Dams	Does					
N	N	N	43	78	159	52	373
G	G	G	8	9	18	9	53
G	N	½G½N	37	50	73	21	252
N	G	½N½G	10	12	24	9	64
G	½N½G	¾G¼N	3	7	9	6	27
N	½G½N	¾N¼G	14	25	39	11	171
Total			115	181	322	108	940

⁺ N = New Zealand White; G = Egyptian Gabali; Breed of buck listed first.

was allowed to sire all his litters from the same does, i.e. separate bucks in each of the six genetic groups were used. Rabbits were raised in semi-closed rabbitry. Breeding does and bucks were housed separately in individual wired-cages arranged in flat-deck batteries allocated in three rows along the rabbitry. All cages are equipped with feeding hoppers and drinking nipples. Each doe was palpated 12 days thereafter to determine pregnancy and those failed to conceive were returned to the same mating-buck. After kindling, litters were checked and recorded. Young rabbits were weaned at four weeks, ear tagged, sexed and transferred to standard progeny wire cages. Rabbits were fed on a commercial grower pelleted diet during the whole experimental period. On dry matter (DM) basis, diet contained 18.5% crude protein (CP), 8.0% crude fiber (CF), 3.0% ether extract (EE) and 6.5% ash. Feed and water were available *ad libitum*.

Data collected and model of analysis:

Litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW) were analyzed using the following mixed model (Harvey, 1990):

$$Y_{ijklm} = \mu + G_i + D_{ij} + YS_k + P_l + e_{ijklm}$$

Where Y_{ijklm} is the observation on the $ijklm^{th}$ trait; μ is the overall mean; G_i is the fixed effect of i^{th} genetic group; D_{ij} is the random effect of j^{th} doe nested within the fixed effect of i^{th} genetic group; YS_k is the fixed effect of the k^{th} year-season of kindling combination; P_l is the fixed effect of l^{th} parity; and e_{ijklm} is the random deviation particular to the m^{th} litter, NID (0, σ^2_e).

Genetic model and estimation of crossbreeding effects:

Coefficients presented in Table 2 for the expected contribution of genetic effects (in G or N and their crosses) were computed according to Dickerson (1992). To

Table 2. Coefficients of expected contribution for genetic effects in different groups of purebreds and crossbreds

Sire genotype	Dam Genotype	Doe Genotype ⁺	Direct Additive (G^I_{N-G})	Maternal Additive (G^M_{N-G})	Direct heterosis (H^I)	Maternal heterosis (H^M)	Recombination effect (R^I)
N	N	N	1	0	0	-0.5	0.5
G	G	G	-1	0	-1	-0.5	0.5
G	N	$\frac{1}{2}G\frac{1}{2}N$	0	1	1	-0.5	0.5
N	G	$\frac{1}{2}N\frac{1}{2}G$	0	-1	0	-0.5	0.5
G	$\frac{1}{2}N\frac{1}{2}G$	$\frac{3}{4}G\frac{1}{4}N$	-0.5	0	0.5	1	-1
N	$\frac{1}{2}G\frac{1}{2}N$	$\frac{3}{4}N\frac{1}{4}G$	0.5	0	-0.5	1	-1

⁺ Sire-breed listed first.

find the appropriate linear functions of breed group class means to use in order to estimate the genetic effects of interest, expression of the mean performance of breeds and crosses in terms of their genetic expectations were calculated as shown in Table 2. The general approach used to estimate the crossbreeding parameters (Komender, 1988) was:

$$M = X P$$

Where M is a vector of least-square means of genetic groups in the experiment, P is a vector of crossbreeding estimates and X is a design matrix including the expected contribution of genetic effects to the mean of each genetic group. The genetic effects were derived as:

$$P = X^I M$$

For data of the two generations, the following genetic components were derived:

G^I = Individual (direct) additive effect of the doe:

$$G^I_N - G^I_G = NN - GG - GN + NG$$

G^M = Maternal additive effect of the dam of doe:

$$G^M_N - G^M_G = GN - NG$$

H^I = Direct heterosis in the crossbred doe:

$$H^I = (GN + \frac{3}{4}G\frac{1}{4}N) - (GG + \frac{3}{4}N\frac{1}{4}G)$$

H^M = Maternal heterosis in the crossbred dam:

$$H^M = (\frac{3}{4}N\frac{1}{4}G + \frac{3}{4}G\frac{1}{4}N) - \frac{1}{2}(NN + GG + GN + NG)$$

R^I = Direct recombination effect in the individual doe:

$$R^I = \frac{1}{2}[(NN + GG + GN + NG) - 2(\frac{3}{4}N \frac{1}{4}G + \frac{3}{4}G \frac{1}{4}N)]$$

The standard errors were computed for all the genetic effects using the inverse matrix of the sub-class numbers and the error standard deviation.

RESULTS AND DISCUSSION

Genetic-groups comparisons:

Least square means of genetic groups and comparisons among purebreds for different traits are given in Table 3. Genetic group of $\frac{1}{2}N\frac{1}{2}G$ resulted in a larger litter size and heavier litter weight compared to the other genetic groups. **G** rabbits were similar to **N** rabbits in litter size at birth and at weaning, while **N** rabbits had relatively heavier litter weight at weaning than **G**. These results were expected and reflected the superiority of **N** rabbits in milking ability.

Genetic-group significantly affected litter size and weight at weaning (Table 4). Similarly, Khalil *et al* (1995) found that mating group constituted a significant source of variation in litter size and weight at weaning. On the contrary, Afifi *et al* (1976ab), Lukefahr *et al* (1983a&b), Afifi and Emara (1984), Oudah (1990) and Abd El-Aziz (1998), all working on crossbreeding experiments found that genetic group showed insignificant effects on most litter traits studied.

Direct breed additive effect (G^I)

The linear contrasts of G^I for litter size and weight at weaning were significant and in favour of **N** rabbits (Table 5). Such differences in G^I between the two breeds lead to state that **N** buck-breed produced litters with larger size and heavier weight at weaning than the **G** buck-breed and therefore **N** rabbits could be used as a buck-breed in crossbreeding programs in Egypt. The percentages of G^I for litter size and weight at weaning were moderate (Table 5), being 7.52 and 12.37 % for **LSW** and **LWW**, respectively. Abd El-Aziz (1998) found that **G**-sired litters had larger **LSB** and heavier **LWW** as compared to **N**-sired litters, while **N**-sired litters had larger **LSW**. Khalil and Afifi (2000) reported that New Zealand White rabbits had higher estimates of G^I than Gabali rabbits for litter weight at birth and weaning ($P < 0.01$ or $P < 0.001$) in crossbreeding experiment involving the two breeds.

Maternal breed additive effect (G^M):

The percentages of G^M for pre-weaning litter traits were low or moderate (Table 5); being -2.34, -9.56 and -7.5 % for **LSB**, **LSW** and **LWW** ($P < 0.05$), respectively. The estimates of G^M were mainly in favour of **G** dams. This unexpected reverse trend was observed also by Abd El-Aziz (1998) who

reported that litters of dams gave larger LSB and LSW compared to N dams. This might be true since N rabbits raised in Egypt for a long period and had a reduction in pre-weaning litter performance due to the decrease in milk production compared to G rabbits. Opposite to the present results, there was a general trend indicating that

Table 3. Purebred and crossbred least-squares means ($\pm$ SE) for litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW)

Trait ⁺	NN	GG	$\frac{1}{2}G\frac{1}{2}N$	$\frac{1}{2}N\frac{1}{2}G$	$\frac{3}{4}G\frac{1}{4}N$	$\frac{3}{4}N\frac{1}{4}G$
	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE	Mean $\pm$ SE
LSB	6.81 $\pm$ 0.13	6.80 $\pm$ 0.32	7.02 $\pm$ 0.16	7.18 $\pm$ 0.32	7.02 $\pm$ 0.44	6.60 $\pm$ 0.20
LSW	5.18 $\pm$ 0.14	5.30 $\pm$ 0.33	5.52 $\pm$ 0.17	6.05 $\pm$ 0.33	6.00 $\pm$ 0.45	5.21 $\pm$ 0.20
LWW	3252.5 $\pm$ 85	3096.7 $\pm$ 203	3175.4 $\pm$ 101	3416.7 $\pm$ 198	3387.4 $\pm$ 274	2963.5 $\pm$ 125

Table 4. F ratios (F) of least-squares ANOVA of factors affecting litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW)

Source	DF	F-ratio		
		LSB	LSW	LWW
Genetic group	5	0.959 ^{NS}	2.036*	1.465*
Does within genetic group	316	1.358***	1.301**	1.344***
Parity	4	3.997**	2.921*	2.641*
Year-season	5	1.376	3.776**	2.958**
Remainder	609			
Error mean squares		3.491	3.875	1391808

^{NS} = Non-significant, * = $P < 0.05$, ** = $P < 0.01$ and *** = $P < 0.001$.

Table 5. Estimates of direct (G^I) and maternal (G^M) additive effects for litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW)

Trait	Direct additive			Maternal additive		
	Units	SE	$G^I\%$	Units	SE	$G^M\%$
LSB	0.1799	0.04	2.62 ^{NS}	-0.1626	0.03	-2.34 ^{NS}
LSW	0.4122	0.04	7.52*	-0.5363	0.03	-9.56*
LWW	397.1	24.8	12.36**	-241.4	17.6	-7.50*

$G^I\% = [G^I \text{ in units} / (\text{average of N} + \text{N-sired crosses})] \times 100$

$$G^M\% = [G^M \text{ in units} / (\text{average of } G + G\text{-dammed crosses})] \times 100$$

NS = Non-significant, * = $P < 0.05$, ** = $P < 0.01$.

litters mothered by exotic breeds recorded better performance than litters mothered by native breeds. Crossbreeding experiments carried out in Egypt (Afifi and Khalil, 1989; Oudah, 1990; Khalil *et al.*, 1995; Khalil and Afifi, 2000) indicated desirable additive maternal breed effect in New Zealand White rabbits for pre-weaning litter traits compared to other breeds.

Direct heterosis (H^I):

Estimates of H^I calculated in actual units and as percentages for different traits are given in Table 6. The estimates of H^I were 9.5, 19.2 and 15.8 % for LSB, LSW and LWW, respectively. These estimates were positive and significant for all traits of the study; indicating that crossing between N and G rabbits was associated with an existence of direct heterotic effects in crossbred preweaning litters of individual does of the present study. However, results of the present study are in agreement with Abd El-Aziz (1998) with the same crossbreeds and for the same traits. Results of the different crossbreeding experiments carried out in Egypt (Afifi and Emara, 1984; Afifi and Khalil, 1989; El-Desoki, 1991; Khalil *et al.*, 1995; Khalil and Afifi, 2000) revealed also that heterotic effects were evidenced in most of the possible crossbreds for litter size, litter weight, and milk yield.

Maternal heterosis (H^M):

Estimates of H^M calculated in actual units for different traits are given in Table 6. The estimates were negative for LSB and LWW, but it was positive and significant for LSW. The parental epistasis in the second-generation (F_2) may be responsible for these low estimates of maternal heterosis. The positive estimate of maternal heterosis for litter size at weaning is favourable and indicating that crossbred dams could produce larger litter size associated with lighter litter weight at weaning than their crossbred daughters. Results of the different crossbreeding experiments carried out in Egypt (i.e. Afifi *et al.*, 1976ab; Afifi and Emara, 1984) revealed that maternal heterotic effects were evidenced in most of the crossbred dams for litter size and litter weight.

Direct recombination effect (R^I):

At kindling, estimate of direct recombination effects (R^I) for litter size at birth was significant (Table 7) and was quite different to that estimate of direct heterosis. This notation implies that dominance effects on this trait recorded at kindling were of considerable importance. At weaning, negative direct recombination loss for litter size at weaning

(Table 7) reveal that crossbred does with N genes could mother does with lower litter size at weaning than purebred N does when both groups of does were mated to bucks from the same N

Table 6. Estimates of direct (H^I) and maternal (H^M) heterosis for litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW)

Trait	Direct heterosis			Maternal heterosis		
	Units	SE	$H^I\%$	Units $H^M\%$		SE
LSB	0.6439	0.048	9.5*	-0.2932	0.04	-4.31*
LSW	1.0052	0.051	19.2**	0.1798	0.05	3.43*
LWW	502.6	17.6	15.8**	-119.7	27.7	- 3.77*

$H^I\% = [H^I \text{ in units} / (\text{average of NN} + \text{GG})] \times 100$

$H^M\% = [H^M \text{ in units} / (\text{average of NN} + \text{GG})] \times 100$

^{NS} = Non-significant, * = $P < 0.05$, ** = $P < 0.01$.

Table 7. Estimates of direct recombination effect (R^I) for litter size at birth (LSB), litter size at weaning (LSW) and litter weight at weaning (LWW)

Trait	Direct recombination effect		
	Units	SE	$(R^I/H^I) \times 100$
LSB	0.2932*	0.044	45.5
LSW	-0.1797 ^{NS}	0.046	-17.8
LWW	119.7 ^{NS}	27.7	23.8

^{NS} = Non-significant, * = $P < 0.05$.

purebred. The insignificant effects for direct recombination loss for litter size and weight at weaning (Table 7) indicate that epistatic recombination losses for these traits recorded at weaning in crossbred rabbits were negligible. Information in the literature concerning estimates of direct recombination effects of crossbreeding experiments in rabbits is scarce. Most of these available results are contradicted.

CONCLUSIONS

- (1) Since post-kindling litter performance in New Zealand White and Egyptian Gabali rabbits were not significantly different in their breed performance, one may use New Zealand White as sires. For commercial rabbits industry in Egypt, Gabali does could be used in terminal crossbreeding system.
- (2) Crossing Egyptian Gabali rabbits with New Zealand White was associated with direct heterotic effects in litters of individual does.

- (3) Single cross resulted from mating New Zealand White sires with Gabali dams is recommended and producers and processors could potentially benefit economically through commercial production by this synthetic simple cross.
- (4) The low or negative estimates of maternal heterosis recorded for pre-weaning litter traits indicate that genetic differences in these traits in crossbred does produced by crossbred or purebred dams are very limited.

REFERENCES

- Abdel-Aziz, M. M. (1998). Crossbreeding between Al-Gabali and New Zealand White rabbits in the North Coast-Belt of the Egyptian Western Desert. *Ph. D. Thesis*, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt.
- Afifi, E.A. and Emara, M.E. (1984). Litter weight in local Egyptian and exotic breeds of rabbits and their crosses. *3rd World Rabbit Congress*, 4-8 April 1984, Rome, Italy, Vol. QG1:126-135.
- Afifi, E.A. and Khalil, M.H. (1989). Observation on purebred and crossbred litters of Bouscat and Grey Giant Flander rabbits in Egypt. *Journal of Applied Rabbit Research*, 12 (4): 273-277.
- Afifi, E.A., Galal, E.S.E., El-Tawil, E.A., El-Khishin, S.S. (1976a). Litter weight in three breeds of rabbits and their crosses. *Egyptian Journal of Animal Production*, 16 (2): 99- 108.
- Afifi, E.A., Galal, E.S.E., El-Tawil, E.A., El-Khishin, S.S. (1976b). Litter size at birth and at weaning in three breeds of rabbits and their crosses. *Egyptian Journal of Animal Production*, 16 (2):109-119.
- Ali, S. M. (1998). Evaluation of growth and production performance of Al-Gabali rabbits and their crosses under semi-arid conditions. M. Sc. Thesis, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt.
- Dickerson, G.E. (1992). Manual for evaluation of breeds and crosses of domestic animals. *Food and Agriculture Organization of the United Nations*, Rome, PP 47.
- El-Desoki, A.E.M. (1991). Study of the effect of some genetic and environmental factors affecting meat yield from some foreign and local breeds of rabbits and their crosses. M.Sc. Thesis, Faculty of Agriculture, Mansoura University, Egypt.
- Harvey, W.R. (1990). User's Guide for LSMLMW. Mixed model least-squares and Maximum Likelihood Computer Program. PC-Version 2, Ohio State University, Columbus, USA, (Mimeograph).

- Khalil, M.H.; Afifi, E.A. (2000).** Heterosis, maternal and direct additive effects for litter performance and post-weaning growth in Gabali rabbits and their F_1 crosses with New Zealand White. *7th World Rabbit Congress*, Valencia, Spain, 4-7 July 2000, Volume A:431-437.
- Khalil, M.H., Afifi, E.A., Youssef, Y.M.K. and Khadr, A.F. (1995).** Heterosis, maternal and direct genetic effects for litter performance and reproductive intervals in rabbit crosses. *World Rabbit Science*, 3(3):99-105.
- Komender, P. (1988).** Crossbreeding in farm animals . III. A general method of comparing models to estimate crossbreeding parameters with an application to diallel crossbreeding experiments. *J. Animal Breeding and Genetics*, 105: 191-2.
- Lukefahr, S., Hohenboken, W.D., Cheeke, P.R., Patton, N.M. (1983a).** Doe reproduction and pre-weaning litter performance of straightbred and crossbred rabbits. *Journal of Animal Science*, 57(5):1090-1099.
- Lukefahr, S., Hohenboken, W.D., Cheeke, P.R., Patton, N.M. (1983b).** Characterization of straightbred and crossbred rabbits for milk production and associative traits. *Journal of Animal Science*, 57(5):1100-1107.
- Oudah, S.M. (1990).** Studies on some rabbit breeds and their crosses. M. Sci. Thesis, Faculty of Agriculture, Mansoura University, Egypt.

تقدير تأثير الخلط على بعض صفات خلفه البطن في خلطة أرانب النيوزيلندي الأبيض مع الجبلي المصري

ماهر حسب النبي خليل^١ ، عزت عطا عفيفي^٢ ، احمد فريد محمود^٣ ، يوسف كمال^٣

^١ كلية الزراعة والطب البيطري - جامعة الملك سعود - القصيم - بريدة - المملكة العربية السعودية

^٢ قسم الإنتاج الحيواني - كلية الزراعة بمشنتهر - جامعة الزقازيق - مصر

^٣ معهد بحوث الإنتاج الحيواني - وزارة الزراعة - الدقي - القاهرة - مصر.

أجريت تجربة خلط بين أرانب النيوزيلندي الأبيض (N) وأرانب الجبلي المصري (G) لإنتاج ٦ مجموعات تربية NN، GG، $G\frac{1}{2}N\frac{1}{2}$ ، $G\frac{3}{4}N\frac{1}{4}$ ، $N\frac{3}{4}G\frac{1}{4}$ ، وذلك لتقدير التأثير التجمعي الفردي (G^I) والأمي (G^M) وكذلك قوة المهجين في الفرد (H^I) وفي الأم (H^I) وأيضاً تأثير إعادة توليف الجينات في الفرد (R^I) لصفات عدد خلفه البطن عند الميلاد والقطام ووزن البطن عند القطام.

وقد أوضحت النتائج تفوق أرانب النيوزيلندي الأبيض على أرانب الجبلي في الصفات المدروسة، وأعطت خلطة الأرانب الناتجة من ذكور نيوزيلندي وإناث جبلي أفضل النتائج بصفة عامة عن الخليط العكسي.

وقد كان التأثير التجمعي الفردي (G^I) لصفات عدد خلفه البطن عند الميلاد ووزن خلفه البطن عند القطام ٢،٦٢ و ٧،٥٢ و ١٢،٣٧ % على التوالي بينما كانت نتائج التأثير التجمعي الأمي (G^M) -٢،٤٣ و ٩،٥٦ و -٧،٥ % على التوالي، كما كانت نتائج قوة المهجين الفردية (H^I) ٩،٥ و ١٩،٢ و ١٥،٨ % على التوالي وكانت قوة المهجين في الأم (H^I) -٤،٣١ و ٣،٤٣ و -٣،٣٧ % على التوالي، أما قيم التأثير الاندماجي فقد كانت ٤٥،٥ و -١٧،٨ و ٢٣،٨ % على التوالي.

هذا وقد أوضحت نتائج تقديرات التأثير التجمعي الفردي أنما ما بين متوسطة وعالية وفي صالح الأرانب النيوزيلندي الأبيض وبالتالي فإنه يمكن استخدام ذكور أرانب النيوزيلندي الأبيض في استراتيجيات نظم الخلط تحت ظروف المناطق الحارة.

كما كانت تقديرات قيم تأثير إعادة توليف الجينات (R^I) لصفات عدد ووزن الخلفة عند القطام غير معنوية مما يشير إلى أن تأثير إعادة توليف الجينات في خلال فترة الرضاعة يمكن تجاهله.