

# Accepted Manuscript

Geological heritage of the Bahariya and Farafra oases, the central Western Desert of Egypt

Ekaterina E. Plyusnina, Emad S. Sallam, Dmitry A. Ruban



PII: S1464-343X(16)30002-4

DOI: [10.1016/j.jafrearsci.2016.01.002](https://doi.org/10.1016/j.jafrearsci.2016.01.002)

Reference: AES 2462

To appear in: *Journal of African Earth Sciences*

Received Date: 23 November 2015

Revised Date: 3 January 2016

Accepted Date: 4 January 2016

Please cite this article as: Plyusnina, E.E., Sallam, E.S., Ruban, D.A., Geological heritage of the Bahariya and Farafra oases, the central Western Desert of Egypt, *Journal of African Earth Sciences* (2016), doi: 10.1016/j.jafrearsci.2016.01.002.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

***The Paleogene reefal structure -  
example of paleogeographical type of geological heritage  
from the Bahariya Oasis (central Western Desert of Egypt)***



ACCEPTED MANUSCRIPT

1 **Geological heritage of the Bahariya and Farafra oases, the central Western Desert of**  
2 **Egypt**

3

4 Ekaterina E. Plyusnina<sup>a</sup>, Emad S. Sallam<sup>b</sup>, Dmitry A. Ruban<sup>a,c,\*</sup>

5

6 <sup>a</sup> *Department of Tourism, Higher School of Business, Southern Federal University, 23-janina*  
7 *Street 43, Rostov-na-Donu, 344019, Russia*

8 <sup>b</sup> *Department of Geology, Faculty of Science, Benha University, Farid Nada Street 15,*  
9 *Benha 13518, Egypt*

10 <sup>c</sup> *P.O. Box 7333, Rostov-na-Donu, 344056, Russia (for postal communication)*

11

12 *E-mail addresses: ruban-d@mail.ru (D.A. Ruban); emad.salam@fsc.bu.edu.eg (E.S. Sallam)*

13

14 \*Corresponding author

15

16 **ABSTRACT**

17 Archaeological and cultural heritage of Egypt is world-known, but its geological  
18 heritage is yet to be revealed. Investigations in the central Western Desert of Egypt permitted  
19 finding a lot of unique features that can be assigned to this heritage. In the Bahariya Oasis,  
20 10 geological heritage types are established, namely stratigraphical, paleontological,  
21 sedimentary, igneous, mineralogical, economical, paleogeographical, geomorphological,  
22 hydrological and hydrogeological, and pedological types. In the Farafra Oasis and vicinities,  
23 only geomorphological and hydrological and hydrogeological types are found. On the area  
24 between these oases, sedimentary, mineralogical, paleogeographical, and geomorphological  
25 features are established. Chalk and nummulitic limestones, invertebrate and dinosaur

26 localities, paleoreefs and paleokarst, iron ore deposit, and peculiar landforms occur on the  
27 study territory. Taken together, these features constitute a highly diverse geological heritage  
28 that can be judged global (even if the rank of individual objects is often relatively low). This  
29 heritage is well suitable for the purpose of geotourism; for instance, thematic excursions  
30 explaining the geological evolution during the last 100 Ma are possible. Participants of such  
31 excursions can also see different facies. A geopark in the central Western Desert of Egypt  
32 would facilitate geoconservation and geotourism activities.

33

34 *Keywords:* Geological heritage; Geotourism; Geomorphology; Upper Cretaceous; Cenozoic;  
35 Western Desert; Egypt.

36

## 37 **1. Introduction**

38 Geological heritage has attracted a lot of attention since the beginning of the 1990s.  
39 Its inventory deems essential for conservation (geoconservation) and use for the purposes of  
40 tourism development (geotourism) (Wimbledon, 1996, 1999; Wimbledon et al., 1998;  
41 Prosser et al., 2006, 2011; Gray, 2008, 2013; Dowling and Newsome, 2010; Ruban, 2010;  
42 Ruban and Kuo, 2010; Dowling, 2011; Henriques et al., 2011; Hose and Vasiljević, 2012;  
43 Wimbledon and Smith-Meyer, 2012; Erikstad, 2013; Prosser, 2013; Bradbury, 2014; Ruban,  
44 2015). Although most research is concentrated in Europe, geological heritage of other parts  
45 of the world is of equal importance. For example, Henriques et al. (2013), Enniouar et al.  
46 (2014), Zangmo Tefogoum et al. (2014), Errami et al. (2015), and Henriques and Neto  
47 (2015) have shown its unprecedented richness in Africa. Geological heritage of Egypt is of  
48 special interest because of three reasons. First, the diversity of geological features known  
49 from this country and the complex nature of its geological evolution (Issawi, 2002, 2005;  
50 Guiraud et al., 2005; Tawadros et al., 2006; Issawi et al., 2009; Tawadros, 2011) make it

51 very promising for inventory, conservation, and tourism use. Second, geological research in  
52 Egypt has been intense for many decades, and the available information is rigorously  
53 systematized. For instance, stratotypes of many units are sufficiently established and  
54 described (e.g., El Kelani et al., 2003) making them ready to be evaluated as geological  
55 heritage sites. Third, the cultural heritage of Egypt (e.g., pyramids) is well-known, and it is  
56 clear how effectively cultural heritage facilitates the promotion and tourism utility of  
57 geological heritage (Migon, 2009; Last et al., 2013; Woo et al., 2013; Gontareva et al., 2015;  
58 Moroni et al., 2015).

59 Unfortunately, very few sites have been declared as parts of the geological heritage of  
60 Egypt. One example is Wadi Al-Hitan ("Whale Valley"), which is located in the Fayium  
61 Depression (~90 km southwest of Cairo). It is included in the list of the UNESCO World  
62 Heritage Sites (see on-line at <http://whc.unesco.org/en/list/1186>). Some geological heritage  
63 sites along the Mediterranean coast of Egypt are described by El-Asmar et al. (2012). In the  
64 present paper, we attempt to fill the noted gap. Field investigations in the central Western  
65 Desert of Egypt and, particularly, in the Bahariya and Farafra oases permitted the recognition  
66 of numerous unique features that when taken together can account for an important  
67 geological heritage. Their characteristics are presented in this paper.

## 69 **2. Geologic setting**

70 The Bahariya and Farafra depressions lie on the Uweinat-Bahariya-Port Said arch  
71 (Issawi et al., 2009). The Bahariya Oasis is located in the central part of the Western Desert  
72 of Egypt (between 27°48'-28°30' N and 28°35'-29°10' E) (Fig. 1). It is oval in shape  
73 stretching NE-SW by ~94 km (the width of the depression is ~42 km). The Bahariya Oasis  
74 was a subject of many geological studies aimed at structural geology, stratigraphy, iron ore  
75 deposits, sedimentology, paleontology, geoarchaeology, etc. (Ball and Beadnell, 1903;

76 Stromer 1914, 1936; Lebling, 1919; Weiler, 1935; El-Akkad and Issawi, 1963; Said and  
77 Issawi, 1964; Soliman et al., 1970; Slaughter and Thurmond, 1974; Khalifa, 1977; El Aref et  
78 al., 2006; Catuneanu et al., 2006; Khalifa and Catuneanu, 2008; Tanner and Khalifa, 2010;  
79 Salama et al., 2012, 2013, 2014; Afify et al., 2015a, b). The Bahariya Formation  
80 (Cenomanian), which is composed of ferruginous sandstones and shales, outcrops on the  
81 bottom of the depression. The siliciclastic beds of the Bahariya Formation are weathered into  
82 black conical-like hills, mesas, and buttes. The black conical hills, which are distributed on  
83 the southern part of the Bahariya Depression, are known as “the Black Desert”. Most of  
84 these hills are capped by basalt sills, giving them a characteristic black color. The  
85 Campanian, Maastrichtian, and lower Eocene rocks outcrop on the flanks of the Bahariya  
86 Depression; the Paleocene is totally missing there (Issawi, 2009) (Fig. 1). The Bahariya  
87 depression superposes the Bahariya anticline that stretches from Gebel Ghorabi in the north,  
88 passing through the central hills of the depression to the southern part of the oasis, and  
89 extends southward to include the Farafra Oasis.

90 The Farafra Oasis is located ~140 km southwest of the Bahariya Oasis in the central  
91 part of the Western Desert (between 26°45'-27°40' N and 27°00'-28°50' E) (Fig. 1). Similarly  
92 to the Bahariya Oasis, the Farafra Oasis occupies an oval-shaped depression with an area of  
93 ~10,000 km<sup>2</sup>. On the bottom of the Farafra Depression, the Dakhla Shale (Maastrichtian) is  
94 outcropped, and it is intertonguing laterally into the Maastrichtian Khoman Chalk in the  
95 central and northern parts of the oasis (Hermina, 1990; Issawi et al., 2009). The Farafra  
96 Depression is surrounded by high escarpments, and its bottom rises gradually to the general  
97 level of the surrounding desert southwards (Beadnell, 1901; Said, 1962; Issawi et al., 2009).  
98 The scarps of the Farafra Depression are composed of the Tarawan Formation (Paleocene)  
99 overlain by the Esna Shale (Paleocene-lower Eocene) and the Farafra Limestones (lower  
100 Eocene) (Issawi et al., 2009; Orabi and Zaky, 2016). The eastern part of the depression is

101 covered by sand sheets, and the depression is bounded to the west by the Great Sand Sea  
102 (Fig. 1). The Farafra Depression forms a dome structure, which represents the southern  
103 extension of the Syrian Arc System (Omara et al., 1970). Its axis stretches in the NE-SW  
104 direction.

105 Since the beginning of the Late Cretaceous, the evolution the Bahariya-Farafra  
106 platform was influenced greatly by eustatic fluctuations and the tectonic activity along the  
107 Syrian Arc fold system (Aram, 1990; El Emam et al., 1990; Sehim, 1993; Moustafa et al.,  
108 2003). During the Cenomanian, the Bahariya-Farafra platform was located on a passive  
109 continental margin (El Emam et al., 1990). This was followed by a phase of tectonic folding  
110 (Said, 1962). Throughout the Turonian-Santonian, the northern part was tectonically uplifted,  
111 while the southern part was subsiding (El Emam et al., 1990). Folding took place later, and  
112 the anticline became even more pronounced (Sehim, 1993; Moustafa et al., 2003). In the  
113 Maastrichtian, the western and southern parts of the Bahariya Oasis and the northern part of  
114 the Farafra Oasis were covered by sea (El Emam et al., 1990). During the Paleocene, the  
115 Bahariya structure was uplifted to become an island, and the Tarawan chalky limestones and  
116 the overlying Esna Shale were deposited on the area of the present-day Farafra Oasis (Issawi  
117 et al., 2009). In the early Eocene, the whole territory submerged (Said and Issawi, 1964;  
118 Issawi et al., 2009). The last phase of the Eocene deposition took place in the Bartonian-  
119 Priabonian, when thick-bedded mollusc-rich siliciclastic-carbonate sediments of the Hamra  
120 Formation were deposited (Said and Issawi, 1964; Issawi et al., 2009). Seaward shoreline  
121 shift continued in the Oligocene, and many river systems were developed: a fluvialite cross-  
122 bedded sandstone and grit sediments were accumulated (the Radwan Formation) (Said and  
123 Issawi, 1964). Later, extensive volcanic eruptions (Mandisha's basalt) and hydrothermal  
124 activity took place on the study territory (Meneisy and El Kalioubi, 1975; El-Etr and  
125 Moustafa, 1978; Morsy, 1987; Meneisy, 1990; El Aref et al., 1999).

126           The formation of the many oases in the Western Desert is related to the thickness of  
127 the hard limestones above the clastic beds and also to the level of the underground water  
128 table. Geological structure is very important in the formation of depressions. In all known  
129 oases in the Western Desert, the successions include relatively thin limestones above a thick  
130 succession of clastics. An uplift of the area of the Bahariya double-plunging anticline  
131 "cracked" the thin limestone, and wind deflation coupled with rainfall completed "hewing"  
132 of the clastics beneath the limestone. Erosion continued to the level of the underground  
133 water. The same occurred with the Farafra Oasis, which is located on a high basement arch  
134 known as the Uweinat-Bahariya-Port Said Arch (Issawi, 2009), where sediments are thin  
135 above the basement complex.

136

### 137 **3. Methods**

138           The geological heritage of the central Western Desert of Egypt was established on  
139 three areas, namely the Bahariya Oasis, the Farafra Oasis and vicinities, and the area  
140 between the noted oases (chiefly along the road connecting these oases). Field investigations  
141 (by the second author) supported by the analysis of literature sources permitted the  
142 delineation of unique geological features. Numerous photos of the latter were taken. The  
143 uniqueness of each feature is determined by its rarity or, in contrast, very typical appearance  
144 that can be established on the local, regional, national, or global scale. All heritage objects  
145 can be further employed for scientific, educational, and tourism purposes. This information  
146 permits the delineation of the geological heritage significance of the above-mentioned three  
147 areas (each taken entirely). It was not necessary to emphasize on particular geological  
148 heritage sites (geosites) within these areas, because of three reasons: 1) these areas are  
149 relatively small in size, 2) some features occupy large areas and intersect spatially with one  
150 another, and 3) the "density" of unique features is sufficient in all areas. It is better to



151 consider each area as a large geosite.

152 The analysis of the geological heritage requires distinction of unique features by their  
153 essence. For this purpose, various classifications (Wimbledon et al., 1998; Prosser et al.,  
154 2006; Ruban, 2010; Ruban and Kuo, 2010; Bradbury et al., 2014) have been proposed. In  
155 this work, the classification proposed by Ruban (2010) and Ruban and Kuo (2010) is used. It  
156 necessitates the presence of about two dozens of types of geological features (stratigraphical,  
157 paleontological, sedimentary, igneous, metamorphic, mineralogical, economical,  
158 geochemical, seismical, structural, paleogeographical, cosmogenic, geothermal,  
159 geocryological, geomorphological, hydrological and hydrogeological, engineering,  
160 radiogeological, neotectonical, pedological, and geohistorical). Ruban (2010) underlined that  
161 many geosites are essentially complex with many types co-occurring in each given object.  
162 For instance, in an outcrop of sedimentary rocks, not only sedimentary, but also  
163 stratigraphical, mineralogical, paleogeographical, and, probably, paleontological features can  
164 be identified. To solve this problem, Ruban (2010) suggested to pay attention to the  
165 dominant types that determine the uniqueness of any given object. It should be also added  
166 that the types may differ by their rank in a given geosite or on a given area (local, regional,  
167 national, or global) (Ruban, 2010). For the three areas of the study territory, the dominant  
168 geological heritage types were established, and the rank was assigned to them tentatively.

169

## 170 **4. Results**

### 171 *4.1. Bahariya Oasis*

172 In the Bahariya Oasis, numerous peculiar geological features were established (Fig.  
173 2). These belong to 10 dominant geological heritage types (Table 1), and some representative  
174 examples are given below.

175 *Stratigraphical type.* Gebel El Dist (Fig. 2a) exhibits a section of the Bahariya

176 Formation (Cenomanian, Upper Cretaceous) and the Naqb Formation (Ypresian, lower  
177 Eocene) that can be used for the stratigraphical correlation purposes. An angular  
178 unconformity between these formations is well visible (Fig. 2h, i). This is an important, but  
179 "ordinary" stratigraphical section, and, therefore, this is a kind of local geological heritage.

180 *Paleontological type.* The same Gebel El Dist (Fig. 2a) is a famous fossil locality,  
181 from which Cretaceous and Eocene bivalves, silicified wood, leaf imprints, remains of  
182 sharks, and bones of dinosaurs were reported (Ball and Beadnell, 1903; Stromer, 1914; Said,  
183 1962; El Akkad and Issawi, 1963; Smith et al., 2001; Schweitzer et al., 2003). With regard to  
184 the noted paleobiodiversity, this can be judged as national geological heritage.

185 *Sedimentary type.* The snow-white nummulitic chalky limestones of the Qazzun  
186 Formation (Ypresian, lower Eocene) reported from several places, including Gebel El Gar El  
187 Hamra (Fig. 2l), are of special interest because such rocks are rare in the geological record.  
188 Of course, the Bahariya Oasis is not the only place to observe them, and the rank of this  
189 feature is regional at maximum.

190 *Igneous type.* Although the geology of the Bahariya Depression is dominated by  
191 sedimentary rocks, some magmatic (volcanic) formations are also known there. The so-  
192 called "Mandisha's basalt" (Oligocene) is represented by a combination of columnar and  
193 pillow-like basalts that comprise a sill that caps some hills nowadays (Fig. 2d, e). The noted  
194 lava structures are well-known, but their very typical appearance is established in this oasis,  
195 where outcrops are also large and well accessible. This implies a national rank of this  
196 geological heritage type.

197 *Mineralogical type.* Iron minerals are known from the Bahariya oasis (Fig. 2r). The  
198 relevant minerals are represented mainly by goethite, hematite, and siderite with manganese  
199 oxides intercalations (Said and Issawi, 1964; Salama et al., 2012, 2013; Afify et al., 2015 a,  
200 b; Baioumy et al., 2014, Baioumy, 2015). This is a kind of local geological heritage.

201 *Economical type.* Iron ore is mined directly in the oasis (Fig. 2r), which is an  
202 important mining site in Egypt. The iron ore lies at the basal part of the Naqb Formation  
203 (Ypresian, lower Eocene) replacing carbonates. The origin of the Bahariya iron ore has been  
204 discussed by several workers since the early work of Ball and Beadnell (1903) who  
205 considered the iron ore as Oligocene lacustrine deposits. Gheith (1955) suggested that the  
206 origin of the Bahariya iron ore is due to replacement processes in the post-Eocene time.  
207 Nakhla (1961) attributed the iron ore of the Ghorabi mine to metasomatic replacement of the  
208 lower Eocene carbonates, whereas El Shazly (1962) proved that the ore was formed as a  
209 primary sedimentary deposit in lagoons during late Eocene-Oligocene times. El Akkad and  
210 Issawi (1963) attributed the origin of the iron ore to the replacement of the carbonate rocks  
211 after their direct deposition in shallow depressions in early-middle Eocene. Said and Issawi  
212 (1964) suggested that the ore is of diagenetic origin. They believed that the iron ore was  
213 deposited together with the early Eocene Naqb carbonates. The small basins, where this  
214 deposition took place, were converted into shallow lagoons during the gradual regression. In  
215 these lagoons, iron minerals concentrated due to leaching of the ferruginous layers of the  
216 Bahariya high (Said and Issawi, 1964). Irrespective of what point of view is correct, the  
217 disputed origin of this mineral deposit (Ball and Beadnell, 1903; Alling, 1947; Gheith, 1955,  
218 1959; Nakhla, 1961; El Shazly, 1962; El Akkad and Issawi, 1963; Said and Issawi, 1964; El  
219 Bassyouny, 2004; Salama et al., 2012, 2013; Afify et al., 2015a, b; Baioumy et al., 2014,  
220 Baioumy, 2015) makes it very interesting. The rank of this heritage feature is regional.

221 *Paleogeographical type.* Different facies are known from this area. Probably, the  
222 most interesting amongst them is linked to the paleoreefs of the Hamra Formation (Lutetian-  
223 Priabonian) (Fig. 2m-j). These carbonate buildups were produced by bivalves and  
224 gastropods, communities of which flourished on the sea bottom (Said and Issawi, 1964;  
225 Issawi et al., 2009). The reefal constructions dip inward, i.e., concentrically toward the center

226 of the structure with angles of  $40^\circ$ , and form saucer-like structures. It appears that such  
227 paleoreefs are rare on the global scale, and, therefore, global rank should be assigned to this  
228 type of the geological heritage.

229 *Geomorphological type.* A lot of peculiar landforms occur in the Bahariya Oasis. The  
230 most interesting are black conical hills in the south of the oasis (Fig. 2f), ball-like  
231 concretions (average diameter 50-60 cm) of hard siliceous dolomitic limestones (so-called  
232 "melon fields") (Fig. 2j, k), and spectacular sand dunes (Fig. 2t, u). These forms are so  
233 unusual and impressive that their rank should be judged as national (at least) or global (most  
234 probably).

235 *Hydrological and hydrogeological type.* Thermal mineral springs are known in the  
236 oasis, and these are used already for the purposes of recreation and spa therapy (Fig. 2v). The  
237 rank of these heritage objects is local.

238 *Pedological type.* Middle-late Eocene paleosols have been found above the iron ore  
239 in the Bahariya iron ore mine (Fig. 2s). However, these are rather "ordinary" features that  
240 can be judged as only local.

241

#### 242 4.2. Farafra Oasis and vicinities

243 In the Farafra Oasis and in its vicinities, peculiar geological features are also  
244 numerous (Fig. 3). However, these belong to the only two dominant geological heritage  
245 types (Table 1). Examples of unique features are given below.

246 *Geomorphological type.* Diverse geomorphological features are reported from this  
247 area. These include "melon fields" (like those in the Bahariya Oasis), peculiar landforms,  
248 and, paleokarst phenomena (Fig. 3). The latter are of special interest. The Maastrichtian  
249 (Late Cretaceous) chalky limestones karstified intensively in the Miocene (Pickford et al.,  
250 2010), and the results of these processes created a very unique landscape (Fig. 3b-e). The

251 number of the co-occurrence of all these geomorphological features permits judging the rank  
252 of the relevant geological heritage as national.

253 *Hydrological and hydrogeological type.* Thermal mineral springs are known in the  
254 oasis, and these are used already for the purposes of recreation and spa therapy. The rank of  
255 these heritage objects is local, similarly to the Bahariya Oasis.

256 It should be added that some micropaleontological finds that permit interesting  
257 taphonomic judgements has been made in the Farafra Oasis recently (Orabi and Zaky, 2016).  
258 However, it is questionable whether this indicates any paleontological and paleogeographical  
259 types of geological heritage on this area.

260

#### 261 4.3. Area between the Bahariya and Farafra oases

262 On the area between the oases, some peculiar geological features also occur (Fig. 4).  
263 These belong to 5 dominant geological heritage types (Table 1) that are illustrated with some  
264 examples below.

265 *Sedimentary type.* The Khoman Chalk Formation (Maastrichtian, Late Cretaceous) is  
266 represented on this area by distinctive snow-white chalk and chalky limestones with  
267 abundant chert bands and thin shale beds at top. This formation is 25-30 m in thickness, and  
268 it overlies conformably the Hefhuf Formation (Campanian, Late Cretaceous) and underlies  
269 the Tarawan Formation (Paleocene). Although this chalk is peculiar, it is very well  
270 distributed both in Egypt and other places of the world, and, therefore, this is the only local  
271 geological heritage.

272 *Mineralogical type.* Large calcite crystals (also in the form of speleothems) are  
273 known from the so-called "Crystal Hill" (Fig. 4e). These would attract potentially the  
274 attention of visitors, although this is the only local geological heritage.

275 *Paleogeographical type.* The Khoman Chalk was deposited under open marine, outer

276 shelf environmental conditions (Issawi et al., 2009). These rocks are rather specific, but very  
277 common in many regions of the world. If to treat the outcrops of the Khoman Chalk in terms  
278 of paleogeographical heritage, its rank would be only local.

279 *Geomorphological type.* On the study area, there are paleokarst features similar to  
280 those described above in the Farafra Oasis. However, there is a feature that deserves close  
281 attention. The well-arranged crystals of calcite form large, medusa-shaped body (Fig. 4f) that  
282 is so unusual and spectacular that can be assigned to the national geological heritage.

283

## 284 **5. Discussion**

### 285 *5.1. Diversity of the geological heritage and its rank*

286 The three study areas differ by the diversity of geological heritage features, and the  
287 number of the dominant types is the largest in the case of the Bahariya Oasis (Table 1).  
288 However, in all cases, the most common are geomorphological features. These include the  
289 peculiar landforms mentioned above ("natural sculptures", "melon fields", and paleokarst  
290 features), as well as the "Black Desert" and the "White Desert" taken entirely. The White  
291 Desert (El-Sahara El-Beida in Arabic) is located ~45 km to the north of the Farafra Oasis,  
292 and it has a total area of 3010 km<sup>2</sup>. It has been declared as a natural protectorate in 2002. The  
293 White Desert is called so because of the white color dominating its whole landscape. The  
294 Black Desert (El-Sahara El-Soda in Arabic) lies between the White Desert in the south and  
295 the Bahariya Oasis in the north. It is located ~50 km southwest of the Al-Bawetii City. It is  
296 called "the Black Desert" because of many widely-spaced black conical hills and the black  
297 sheets of gravels and rock fragments strewn on the floor of this desert giving it a  
298 characteristic rough physiography. These hills vary in their forms and height.

299 A central idea in the modern understanding of geological heritage is geodiversity that  
300 can be characterized either quantitatively or qualitatively (Gray, 2008, 2013; Ruban, 2010;

301 Knight, 2011; Crawford and Black, 2012). As demonstrated above, the geological features of  
302 the study territory are chiefly of intermediate to low ranks (local or regional). However, the  
303 co-occurrence of diverse features by itself increases the overall rank of their entity (such a  
304 situation was discussed by Ruban (2010)). Rare places of the world can boast by such a  
305 diversity of peculiar geomorphological and geological features as the Bahariya and Farafra  
306 oases and the area in between them. If so, the entire geological heritage of the study territory,  
307 i.e., the central Western Desert, can be ranked as global.

308

### 309 *5.2. Geotourism perspectives*

310 The geological heritage of the central Western Desert of Egypt seems to be very  
311 suitable for the purposes of geotourism development because of its diversity and uniqueness  
312 (Table 1). Individual tourists and tourist groups can reach there easily via luxury tourist  
313 buses or four-wheel safari cars owned by tourism enterprises. Travellers to the Bahariya and  
314 Farafra oases may visit particular geological features to learn about the general geology and  
315 geomorphology. Moreover, thematic excursions, as explained by Plyusnina et al. (2015),  
316 provide another opportunity to learn about the geological history of a given area. In the case  
317 of the study territory, visitors can learn about the last 100 Ma of the geological history (Late  
318 Cretaceous-Present) seeing different facies, fossils, and other relevant features (Table 2).  
319 Importantly, some of these features are not only typical for the Western Desert, but they also  
320 characterize the general scheme of geological evolution. For instance, there are specific  
321 formations like chalk and nummulitic limestones, paleoreefs, invertebrate and dinosaur  
322 localities. As the discussed geological heritage is concentrated on rather small area restricted  
323 to two oases and their vicinities (Fig. 1), the potential for such a thematic  
324 (paleogeographical) excursion seems to be large.

325 The geological heritage, especially when it is ranked globally, should be employed

326 for tourism purposes in good balance with conservation (e.g., Prosser et al., 2006, 2011;  
327 Gray, 2008, 2013). Two possibilities for this exist in the central Western Desert. First, the  
328 Natural Protectorate of the White Desert has been designated in 2002 to include the hills of  
329 El-Dist and El-Maghrafa and the Black Desert, and it is managed now by the Egyptian  
330 Ministry of the Environment. The most activities offered at this protectorate include Safari  
331 trips, camping inside the desert, and sand boarding on the sand dunes. This kind of protected  
332 area can itself work for the purposes of geotourism. Second, a geopark that will become  
333 further a member of the global geopark network ([http://www.unesco.org/en/natural-  
334 sciences/environment/earth-sciences/geoparks/some-questions-about-geoparks/where-are-  
335 the-global-geoparks/](http://www.unesco.org/en/natural-sciences/environment/earth-sciences/geoparks/some-questions-about-geoparks/where-are-the-global-geoparks/)) can be established on the basis of the diverse and abundant geological  
336 heritage features of the Bahariya and Farafra oases and the area between them. Experience  
337 from many other places of the world (Farsani et al., 2012; Henriques et al., 2012; Fukami,  
338 2013; Palacio Prieto, 2013; Lazzari and Aloia, 2014; Sun, 2014; Wang et al., 2015) suggests  
339 that such a geopark may facilitate the joint geoconservation and geotourism activities.  
340 However, establishment of geopark requires significant efforts from the national geological  
341 community in Egypt.

342

## 343 **6. Conclusions**

344 The present study underlines the importance of geological heritage inventory in  
345 Egypt. The diverse geological features of the Bahariya and Farafra oases are essential part of  
346 the geological heritage. The latter is represented by 10 dominant types, of which the  
347 geomorphological features are the most common. This geological heritage is of global  
348 importance because of the diversity of features co-occurring on a rather restricted territory.  
349 The Bahariya and Farafra oases and the area between them are well suitable for geotourism  
350 development. The already established natural protectorate and the possible geopark can



351 contribute substantially to both geoconservation and geotourism.

352

### 353 **Acknowledgements**

354 The authors gratefully thank P.G. Eriksson (South Africa) for editorial support, the  
355 two anonymous referees for useful suggestions, M.K. Zobaa (USA) for linguistic polishing,  
356 and D.E. Bruno (Italy), E. Errami (Morocco), M.H. Henriques (Portugal), B. Issawi (Egypt),  
357 I.-M. Necheş (Romania), W. Riegraf (Germany), E. Tawadros (Canada), W.A.P. Wimbledon  
358 (UK), and the other colleagues for discussions and/or help with literature.

359

### 360 **References**

361

- 362 Afify, A.M., Sanz-Montero, M.E., Clavo, J.P., Wanas, H.A., 2015a. Diagenetic origin of  
363 ironstone crusts in the Lower Cenomanian Bahariya Formation, Bahariya Depression,  
364 Western Desert, Egypt. *Journal of African Earth Sciences* 101, 333-349.
- 365 Afify, A.M., Sanz-Montero, M.E., Calvo, J.P., 2015b. Ironstone deposits hosted in Eocene  
366 carbonates from Bahariya (Egypt) - new perspective on cherty ironstone occurrences.  
367 *Sedimentary Geology* 329, 81-97.
- 368 Alling, H.L., 1947. Diagenesis of the Clinton hematite ores of New York. *Bulletin of the*  
369 *Geological Society of America* 58, 991-1018.
- 370 Aram, R.B., 1990. Seismic stratigraphic study of the Cretaceous Bahariya Formation, south  
371 Um Baraka concession, Western Desert, Egypt. In: *Proceedings of the 9<sup>th</sup> Petroleum*  
372 *Exploration Production Conference 2, Egypt*, pp. 460–514.
- 373 Baioumy, H.M., 2015. Rare earth elements, S and Sr isotopes and origin of barite from  
374 Bahariya Oasis, Egypt: Implication for the origin of host iron ores. *Journal of African*  
375 *Earth Sciences* 106, 99-107.

- 376 Baioumy, H.M., Ahmed H. H., Mohamed Z. K., 2014. A mixed hydrogeneous and  
377 hydrothermal origin of the Bahariya iron ores, Egypt: Evidences from the trace and rare  
378 earth elements geochemistry. *Journal of Geochemical Exploration* 146, 149-162.
- 379 Ball, J., Beadnell, H.J.L., 1903. Baharyia Oasis: its topography and geology. Geological  
380 Survey of Egypt, Cairo, 84 pp.
- 381 Beadnell, H.J.L., 1901. Farafra Oasis, its topography and geology. *Geol. Surv. Egypt, Cairo,*  
382 39 pp.
- 383 Bradbury, J., 2014. A keyed classification of natural geodiversity for land management and  
384 nature conservation purposes. *Proceedings of the Geologists Association* 125, 329-349.
- 385 Bruno, D.E., Crowley, B.E., Gutak, Ja.M., Moroni, A., Nazarenko, O.V., Oheim, K.B.,  
386 Ruban, D.A., Tiess, G., Zorina, S.O., 2014. Paleogeography as geological heritage:  
387 Developing geosite classification. *Earth-Science Reviews* 138, 300-312.
- 388 Catuneanu, O., Khalifa, M.A., Wanas, H.A., 2006. Sequence stratigraphy of the Lower  
389 Cenomanian Bahariya Formation, Bahariya Oasis, Western Desert, Egypt. *Sedimentary*  
390 *Geology* 190, 121-137.
- 391 Crawford, K.R., Black, R., 2012. Visitor Understanding of the Geodiversity and the  
392 Geoconservation Value of the Giant's Causeway World Heritage Site, Northern Ireland.  
393 *Geoheritage* 4, 115-126.
- 394 Dowling, R., 2011. Geotourism's Global Growth. *Geoheritage* 3, 1-13.
- 395 Dowling, R., Newsome, D., 2010. Geotourism: a Global Activity. In: Dowling, R.,  
396 Newsome, D. (Eds.), *Global Geotourism Perspectives*. Goodfellow Publishers,  
397 Wood Eaton, pp. 1-17.
- 398 El-Akkad, S., Issawi, B., 1963. Geology and iron ore deposits of the Bahariya Oasis. *Geol.*  
399 *Surv. Egypt, Cairo, paper 18.* 300 pp.
- 400 El Aref, M.M., El Sharkawi, M.A., Khalil, M.A., 1999. Geology and genesis of the

- 401 stratabound and Stratiform Cretaceous-Eocene iron ore deposits of the Bahariya region,  
402 Western Desert, Egypt. GAW 4<sup>th</sup> International Conference, Cairo University, Egypt,  
403 450-475.
- 404 El Aref, M.M., Mesaed, A.A., Khalil, M.A., Salama, W.S., 2006. Stratigraphic setting, facies  
405 analyses and depositional environments of the Eocene ironstones of Gabal Ghorabi mine  
406 area, El Bahariya Depression, Western Desert, Egypt. *Egyptian Journal of Geology* 50,  
407 29-57.
- 408 El Asmar, H.M., Ahmed, M.H., Taha, M.M.N., Assal, E.M., 2012. Human Impacts on  
409 Geological and Cultural Heritage in the Coastal Zone West of Alexandria to Al-  
410 Alamein, Egypt. *Geoheritage* 4, 263-274.
- 411 El Bassyouny, A.A., 2004. Stratigraphy of El-Harra area, Bahariya Oasis, Western Desert,  
412 Egypt. *Sedimentology of Egypt* 12, 207-232.
- 413 El Emam, A., Dishopp, D., Dunderdale, I., 1990. The structural setting of the central  
414 Western Desert, Egypt. In: 10<sup>th</sup> Egyptian General Petroleum Corporation Seminar 2, 30-  
415 70.
- 416 El Etr, H.A., Moustafa, A.R., 1978. Field relations of the main basalt occurrences of the  
417 Bahariya Region, central Western Desert, Egypt. *Proceedings of the Egyptian Academy*  
418 *of Science* 31, 191-201.
- 419 El Kelani, A., El Hag, I.A., Bakry, H., Sheira, M., 2003. Type and stratotype sections of the  
420 Mesozoic in Sinai. Geological Survey of Egypt, Cairo, 178 pp.
- 421 El Shazly, E.M., 1962. The results of drilling in the iron ore deposits of Gebel Ghorabi,  
422 Bahariya Oasis, Western Desert, and report on the mineralogy of the low grade iron pres  
423 of El-Heiz area, Bahariya Oasis, Western Desert, Egypt. Geological Survey of Egypt,  
424 Cairo.
- 425 Enniouar, A., Lagnaoui, A., Habib, A., 2014. A Middle Jurassic sauropod tracksite in the

- 426 Argana Basin, Western High Atlas, Morocco: An example of paleoichnological heritage  
427 for sustainable geotourism. *Proceedings of the Geologists' Association* 125, 114-119.
- 428 Erikstad, L., 2013. Geoheritage and geodiversity management - the questions for tomorrow.  
429 *Proceedings of the Geologists Association* 124, 713-719.
- 430 Errami, E., Brocx, M., Semeniuk, V. (Eds.), 2015. *From Geoheritage to Geoparks: Case  
431 Studies from Africa and Beyond*. Springer, Cham, 269 pp.
- 432 Farsani, N.T., Coelho, C., Costa, C., 2012. Geotourism and Geoparks as Gateways to Socio-  
433 cultural Sustainability in Qeshm Rural Areas, Iran. *Asia Pacific Journal of Tourism  
434 Research* 17, 30-48.
- 435 Fukami, S., 2013. Perspective on geoparks and geotourism based on case studies of geoparks  
436 in Japan and China. *Japanese Journal of Human Geography* 65, 58-70.
- 437 Geological Survey of Egypt, 1981. *Geologic map of Egypt*, scale 1: 2,000,000. Cairo.
- 438 Gheith, M.A., 1955. Classification and review of Egyptian iron ore deposits. In: *Symposium  
439 on Applied Geology in the Near East*, UNESCO, Ankara, 106-113.
- 440 Gheith, M.A., 1959. Mineralogy, thermal analysis and origin of the Bahariya iron ores of  
441 Egypt. In: *International Geological Congress, Mexico*, 7, 195-206.
- 442 Gontareva, E.F., Ansari, M.K., Ruban, D.A., Ahmad, M., Singh, T.N., 2015. Geological  
443 dimension of the cultural heritage: A case example of the Ajanta Caves (Maharashtra,  
444 India). *Cadernos do Laboratorio Xeolóxico de Laxe* 38, 67-78.
- 445 Gray, M., 2008. Geodiversity: developing the paradigm. *Proceedings of the Geologists'  
446 Association* 119, 287-298.
- 447 Gray, M., 2013. *Geodiversity. Valuing and Conserving Abiotic Nature*. Wiley-Blackwell,  
448 Chichester, 495 pp.
- 449 Guiraud, R., Bosworth, W., Thierry, J., Delplanque, A., 2005. Phanerozoic geological  
450 evolution of Northern and Central Africa: An overview. *Journal of African Earth*

- 451 Sciences 43, 83–143.
- 452 Henriques, M.H., Neto K., 2015. Geoheritage at the Equator: Selected Geosites of São Tomé  
453 Island (Cameron Line, Central Africa). *Sustainability* 7, 648-667.
- 454 Henriques, M.H., Pena dos Reis, R., Brilha, J., Mota, T., 2011. Geoconservation as an  
455 Emerging Geoscience. *Geoheritage* 3, 117-128.
- 456 Henriques, M.H., Tomaz, C., Sa, A.A., 2012. The Arouca Geopark (Portugal) as an  
457 educational resource: A case study. *Episodes* 35, 481-488.
- 458 Henriques, M.H., Tavares, A.O., Bala, A.L.M., 2013. The geological heritage of Tundavala  
459 (Angola): An integrated approach to its characterisation. *Journal of African Earth*  
460 *Sciences* 88, 62-71.
- 461 Hermina, M., 1990. The surroundings of Kharga, Dakhla and Farafra oases. In: Said, R.  
462 (Ed.), *The Geology of Egypt*. Egyptian General Petroleum Corporation, 259-292.
- 463 Hose, T.A., Vasiljević, D.A., 2012. Defining the nature and purpose of modern geotourism  
464 with particular reference to the United Kingdom and south-east Europe. *Geoheritage* 4,  
465 25-43.
- 466 Issawi, B., 2002. Egypt during the Phanerozoic. In: Youssef, E.A.A. (Ed.), *Geology of the*  
467 *Arab World. Proceedings of the Sixth International Conference on Geology of the Arab*  
468 *World*, Cairo University, 2002. V. 2. Cairo University, Cairo, pp. 401-450.
- 469 Issawi, B., 2005. Archean-Phanerozoic birth and development of the Egyptian land. In:  
470 Youssef, E.A.A. (Ed.), *Geology of the Tethys. Proceedings of the First International*  
471 *Conference on the Geology of the Tethys*, Cairo University, November 2005. V. 2.  
472 Cairo University, Cairo, 339-380.
- 473 Issawi, B., Francis, M., Youssef, A., Osman, R., 2009. *The Phanerozoic of Egypt: A*  
474 *geodynamic approach*. Geological Survey of Egypt, Cairo, 589 pp.
- 475 Khalifa, M.A., 1977. Geological and sedimentological studies of the El Hefhuf area,

- 476 Bahariya Oases, Western Desert, Egypt. Unpublished M.Sc. Thesis, Cairo University,  
477 181 pp.
- 478 Khalifa, M.A., Catuneanu, O., 2008. Sedimentology of the fluvial and fluvio-marine facies  
479 of the Bahariya Formation (Early Cenomanian), Bahariya Oasis, Western Desert, Egypt.  
480 *Journal of African Earth Sciences* 51, 89-103.
- 481 Knight, J., 2011. Evaluating geological heritage: correspondence on Ruban, D.A.  
482 'Quantification of geodiversity and its loss' (*PGA*, 2010, 121(3): 326-333). *Proceedings*  
483 *of the Geologists' Association* 122, 508-510.
- 484 Last, J., Brown, E.J., Bridgland, D.R., Harding, P., 2013. Quaternary geoconservation and  
485 Palaeolithic heritage protection in the 21st century: developing a collaborative approach.  
486 *Proceedings of the Geologists Association* 124, 625-637.
- 487 Lazzari, M., Aloia, A., 2014. Geoparks, geoheritage and geotourism: Opportunities and tools  
488 in sustainable development of the territory. *GeoJournal of Tourism and Geosites* 13, 8-9.
- 489 Lebling, C., 1919. Ergebnisse der Forschungsreisen Prof. E. Stroners in den Wusten  
490 Aegyptens: Teil III, Forschungen in der Bahariya Oase und andern Gegenden  
491 Aegyptens. *Abhandl. Konig. Bayerische Akademie der Wissenschafte Math. Phus.*  
492 *Klasse.* 29.
- 493 Meneisy, M.Y., 1990. Vulcanicity. In: R. Said (Ed.), *The geology of Egypt*. Balkema,  
494 Rotterdam, 157-172.
- 495 Meneisy, M.Y., El Kalioubi, B., 1975. Isotopic ages of the volcanic rocks of the Bahariya  
496 Oasis. *Ann. Geological Survey of Egypt* 5, 119-122.
- 497 Migon, P., 2009. Geomorphosites and the World Heritage List of UNESCO. In: Reynard, E.,  
498 Coratza, P., Regolini-Bissig, G. (Eds.). *Geomorphosites*. F. Pfeil, München, pp. 119-  
499 130.
- 500 Moroni, A., Gnezdilova, V.V., Ruban, D.A., 2015. Geological heritage in archaeological

- 501 sites: case examples from Italy and Russia. Proceedings of the Geologists' Association  
502 126, 244-251.
- 503 Morsy, M.A., 1987. Geology and Radioactivity of Late Cretaceous-Tertiary Sediments in the  
504 Northern Western Desert, Egypt. Unpublished Ph.D. Thesis, Faculty of Science,  
505 Mansoura University, 175 pp.
- 506 Moustafa, A.R., Saoudi, A., Moubasher, A., Ibrahim, I.M., Molokhia, H., Schwartz, B.,  
507 2003. Structural setting and tectonic evolution of the Bahariya Depression, Western  
508 Desert, Egypt. *GeoArabia* 8, 91-124.
- 509 Nakhla, F.M., 1961. The origin ore deposits of El-Bahariya Oasis, Egypt. *Economic*  
510 *Geology*, 56, 1103-1111.
- 511 Orabi, O.H., Zaky, A.S., 2016. Differential dissolution susceptibility of Paleocene  
512 foraminiferal assemblage from Farafra Oasis, Egypt. *Journal of African Earth Sciences*  
513 113, 181-193.
- 514 Omara, S., Hemida, I., Sanad, S., 1970. Structure and hydrogeology of Farafra Oasis,  
515 Western Desert, U.A.R. In: 7th Arab Petroleum Congress, Kuwait, paper 65.
- 516 Palacio Prieto, J.L., 2013. Geosites, geomorphosites and geoparks: Importance, actual  
517 situation and perspectives in Mexico. *Investigaciones Geograficas* 82, 24-37.
- 518 Pickford, M., Wanas, H.A., Mein, P., Ségalen, L., Soliman, H., 2010. The extent of  
519 paleokarst and fluvio-lacustrine features in the Western Desert, Egypt; Late Miocene  
520 subaerial and subterranean paleohydrology of the Bahariya-Farafra area. *The Tethys*  
521 *Geological Society, Cairo, Egypt* 5, 35-42.
- 522 Plyusnina, E.E., Ruban, D.A., Zayats, P.P., 2015. Thematic dimension of geological  
523 heritage: an evidence from the Western Caucasus. *Journal of the Geographical Institute*  
524 "Jovan Cvijić" SASA 65, 59-76.
- 525 Prosser, C.D., 2013. Our rich and varied geoconservation portfolio: the foundation for the

- 526 future. *Proceedings of the Geologists' Association* 124, 568-580.
- 527 Prosser, C., Murphy, M., Larwood, J., 2006. *Geological conservation: a guide to good*  
528 *practice*. English Nature, Peterborough, 145 pp.
- 529 Prosser, C.D., Bridgland, D.R., Brown, E.J., Larwood, J.G., 2011. *Geoconservation for*  
530 *science and society: challenges and opportunities*. *Proceedings of the Geologists'*  
531 *Association* 122, 337-342.
- 532 Ruban, D.A., 2010. *Quantification of geodiversity and its loss*. *Proceedings of the*  
533 *Geologists' Association* 121, 326-333.
- 534 Ruban, D.A., 2015. *Geotourism - A geographical review of the literature*. *Tourism*  
535 *Management Perspectives* 15, 1-15.
- 536 Ruban, D.A., Kuo, I., 2010. *Essentials of geological heritage site (geosite) management: a*  
537 *conceptual assessment of interests and conflicts*. *Natura Nascosta* 41, 16-31.
- 538 Said, R., 1962. *The geology of Egypt*. Elsevier, Amsterdam, 377 pp.
- 539 Said, R., Issawi, B., 1964. *Geology of Northern Plateau, Bahariya Oasis, Egypt*. *Geological*  
540 *Survey of Egypt, Cairo*, 41 pp.
- 541 Salama, W., El Aref, M.M., Gaupp, R. 2012. *Mineralogical and geochemical investigations*  
542 *of the Middle Eocene ironstones, El Bahariya Depression, Western Desert, Egypt*.  
543 *Gondwana Research* 22, 717-736.
- 544 Salama, W., El Aref, M.M., Gaupp, R. 2013. *Mineral evolution and processes of ferruginous*  
545 *microbialite accretion – an example from the Middle Eocene stromatolitic and ooidal*  
546 *ironstones of the Bahariya Depression, Western Desert, Egypt*. *Geobiology* 11, 15–28.
- 547 Salama, W., El Aref, M.M., Gaupp, R. 2014. *Facies analysis and palaeoclimatic significance*  
548 *of ironstones formed during the Eocene greenhouse*. *Sedimentology* 61, 1594-1624.
- 549 Schweitzer, C.E., Lacovara, K.J., Smith, J.B., Lamanna, M.C., Mandela, M.A., Attia, Y.,  
550 2003. *Mangrove-dwelling crabs (Decapoda: Brachyura: Necrocarcinidae) associated*



- 551 with dinosaurs from the Upper Cretaceous (Cenomanian) of Egypt. *Journal of*  
552 *Paleontology* 77, 888-894.
- 553 Sehim, A.A., 1993. Cretaceous tectonics in Egypt. *Egyptian Journal of Geology* 371, 335-  
554 372.
- 555 Slaughter, B.H., Thurmond, J.T., 1974. A Lower Cenomanian (Cretaceous) Ichthyofauna  
556 from the Bahariya Formation of Egypt. *Annals of the Geological Survey of Egypt* 4, p.  
557 25-40.
- 558 Smith, J.B., Lamanna, M.C., Lacovara, K.J., Dodson, P., Smith, J.R., Poole, J.C.,  
559 Giegengack, R., Attia, Y., 2001. A giant sauropod dinosaur from an Upper Cretaceous  
560 mangrove deposit in Egypt. *Science* 292, 1704-1706.
- 561 Soliman, S.M., Faris, M.I., Badry, O.E., 1970. Lithostratigraphy of the Cretaceous  
562 Formations in the Bahariya Oasis, Western Desert, Egypt. In: 7<sup>th</sup> Arab Petroleum  
563 Congress, Kuwait.
- 564 Stromer, E., 1914. Die topographie und geologie der strecks Gharag Baharije nebst  
565 Ausfahrungen, uber die geologische Geschichte Agyptens. *Abhandl. Konig. Bayerische*  
566 *Akademie der Wissenschafte Math.-Phys. Klasse 24, München, 1-78.*
- 567 Stromer, E., 1936. *Ergebnisse der Forschungsreisen Prof. E. Stromers in den Wusten*  
568 *Agyptens. VII. Bahariya Kessel und Stufe mit deren Fauna und flora. Eine ergänzende*  
569 *usammenfassung. Abhandl. Konig. Bayerische Akademie der Wissenschafte Math.-Phys.*  
570 *Klasse 26 (3).*
- 571 Sun, J.H., 2014. The integration and development of heritage resources based on RMP  
572 analysis a case study of Songshan world Geopark. *Advanced Materials Research* 889-  
573 890, 1653-1659.
- 574 Tanner, L.H., Khalifa, M.A., 2010. Origin of ferricretes in fluvial-marine deposits of the  
575 Lower Cenomanian Bahariya Formation, Bahariya Oasis, Western Desert, Egypt.

- 576 Journal of African Earth Sciences 56, 179-189.
- 577 Tawadros, E., 2011. Geology of North Africa. CRC Press, London, 930 pp.
- 578 Tawadros, E., Ruban, D., Efendiyeva, M., 2006. Evolution of NE Africa and the Greater  
579 Caucasus: Common Patterns and Petroleum Potential. In: The Canadian Society of  
580 Petroleum Geologists, the Canadian Society of Exploration Geophysicists, the Canadian  
581 Well Logging Society Joint Convention. May 15-18, 2006. Calgary, pp. 531-538.
- 582 Wang, L., Tian, M., Wang, L., 2015. Geodiversity, geoconservation and geotourism in Hong  
583 Kong Global Geopark of China. Proceedings of the Geologists' Association 126, 426-  
584 437.
- 585 Weiler, W., 1935. Fischreste aus dem nubischen Sandstein von Mohamid und Edfu und aus  
586 den Phosphaten oberoegyptens und der Oase Bahariye. Abhandl. Konig. Bayerische  
587 Akademie der Wissenschafte Math.-Phys. Klasse 7, 12-42.
- 588 Wimbledon, W.A.P., 1996. Geosites - a new conservation initiative. Episodes 19, 87-88.
- 589 Wimbledon, W.A.P., 1999. GEOSITES - an International Union of Geological Sciences  
590 initiative to conserve our geological heritage. Polish Geological Institute Special Papers  
591 2, 5-8.
- 592 Wimbledon, W.A.P., Smith-Meyer, S. (Eds.), 2012. Geoheritage in Europe and its  
593 conservation. ProGEO, Oslo, 405 pp.
- 594 Wimbledon, W., Ishchenko, A., Gerasimenko, N., Alexandrowicz, Z., Vinokurov, V.,  
595 Liscak, P., Vozar, J., Bezak, V., Kohut, M., Polak, M., Mello, J., Potfaj, M., Gross, P.,  
596 Elecko, M., Nagy, A., Barath, I., Lapo, A., Vdovets, M., Klincharov, S., Marjanac, L.,  
597 Mijovic, D., Dimitrijevic, M., Gavrolovic, D., Theodossiou-Drandaki, I., Serjani, A.,  
598 Todorov, T., Nakov, R., Zagorchev, I., Perez-Gonzalez, A., Benvenuti, M., Boni, M.,  
599 Bracucci, G., Bortolani, G., Burlando, M., Costantini, E., D'Andrea, M., Gisotti, G.,  
600 Guado, G., Marchetti, M., Massolli-Novelli, R., Panizza, M., Pavia, G., Poli, G.,

- 601 Zarlenga, F., Satkunas, J., Mikulenas, V., Suominen, V., Kananajo, T., Lehtinen, M.,  
602 Gonggrijp, G., Look, E., Grube, A., Johansson, C., Karis, L., Parkes, M., Paudsep, R.,  
603 Andersen, S., Cleal, C., Bevins, R., 1998. A first attempt at a GEOSITES framework for  
604 Europe - an IUGS initiative to support recognition of world heritage and European  
605 geodiversity. *Geologica Balcanica* 28, 5-32.
- 606 Woo, K.S., Sohn, Y.K., Ahn, U.S., Yoon, S.H., Spate, A., 2013. Jeju Island Geopark - A  
607 Volcanic Wonder of Korea. Heidelberg, Springer, 88 pp.
- 608 Zangmo Tefogoum, G., Kagou Dongmo, A., Nkouathio, D.G., Wandji, P., Gountie Dedzo,  
609 M., 2014. Geomorphological features of the Manengouba Volcano (Cameroon Line):  
610 Assets for potential geopark development. *Geoheritage* 6, 225-239.

611

## 612 FIGURE CAPTIONS

613

- 614 Fig. 1. Geological map of the study territory (modified from Geological Survey of Egypt,  
615 1981).

616

- 617 Fig. 2. Geological heritage of the Bahariya Oasis: a-c - Gebel El Dist (Cenomanian  
618 siliciclastics overlain by Lower Eocene nummulitic limestones), d, e - the Mandisha's  
619 basalt (Oligocene), f - the Black Desert (black hills consisted of Cenomanian  
620 siliciclastics topped by basalts), g - Quaternary paleolake deposits, h, i - Lower Eocene  
621 dolomitic limestones overlying Cenomanian strata with angular unconformity, j, k - ball-  
622 like concretions (average diameter 50-60 cm) of hard siliceous dolomitic limestones  
623 ("melon field"), l - Gebel El Gar El Hamra (landmark conical-hill buildup, type section  
624 of the Qazzun and Hamra formations, paleoreefs and other deposits), m - Lutetian-  
625 Priabonian semi-circular reefal structure), n, o - Middle Eocene saucer-like reefal

626 structures, p, q - pedestal rocks, r - Bahariya iron ore mine (origin of ore is debatable), s  
627 - cross-bedded paleosols highly pierced by root casts, t - asymmetrical ripple marks, u -  
628 sand dunes of Ghard Ghorabi, v- thermal mineral spring equipped with a kind of bath.

629

630 Fig. 3. Geological heritage of the Farafra Oasis and vicinities: a-i - paleokarst (Miocene?)  
631 features, j - "melon field" of Quaternary sediments, k-n - peculiar landforms.

632

633 Fig. 4. Geological heritage of the area between the Bahariya and Farafra oases: a-d - Crystal  
634 Hill (brecciated structure on c, d reflects the collapsed cave roof), e - calcite crystals  
635 from the Crystal Hill, f - medusa-shaped (umbrella-shaped) body consisting of well-  
636 arranged crystals of calcite.


Table 1. Geological heritage types in the study territory. Typology is based on Ruban (2010) and Ruban and Kuo (2010).


Area	Dominant types																					
	ST	PL	SE	IG	MT	MI	EC	GC	SI	SR	PG	CO	GT	GY	GM	HY	EN	RA	NE	PE	GH	
Bahariya Oasis	X	X	X	X		X	X				X				X	X					X	
Farafra Oasis and vicinities															X	X						
area in between of oases			X			X					X				X							

Types abbreviations: ST – stratigraphical, PL- paleontological, SE – sedimentary, IG – igneous, MT – metamorphic, MI – mineralogical, EC – economical, GC – geochemical, SI – seismical, SR – structural, PG – paleogeographical, CO – cosmogenic, GT – geothermal, GY – geocryological, GM – geomorphological, HY - hydrological and hydrogeological, EN – engineering, RA – radiogeological, NE – neotectonical, PE – pedological, GH – geohistorical.

Table 2. Paleogeographical features in the geological heritage of the study territory. Typology is based on Bruno et al. (2014).

Geological age	Dominant subtypes of paleogeographical type					
	Facies	Paleoecosystems	Ichnology	Taphonomy	Events	Geoarchaeology
Quaternary	paleolake		burrows and traces			artifacts
Miocene	karstic					
Oligocene	fluvial					
Eocene	shallow-marine	nummulites			gradual basin deepening	
	shallow-marine	reefs (bivalves and gastropods)				
Upper Cretaceous	chalk-dominated outer shelf	bivalves				
	deltaic, estuarine	bivalves, dinosaurs, flora		fossil wood, other fossilized remains		





ACCEPTED MANUSCRIPT






ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT

Global-ranked geological heritage in central Western Desert of Egypt

Ten geological heritage types (geomorphological, igneous, paleogeographical, etc.)

Possibility for thematic geotourist excursions linked to geological history

Potential for geopark establishment in Bahariya and Farafra oases

ACCEPTED MANUSCRIPT