

References

- Adams JM, Cockburn MI, Difazio LT, (2006). Spinal clearance in the difficult trauma patient: a role for screening MRI of the spine. *J.Am.Surg.* January;72(1):101-105.
- Amador, M. J. & Guest, J. D. (2005) An appraisal of ongoing experimental procedures in human spinal cord injury. *J. Neurol. Phys. Ther.* 29, 70–86.
- Anderson PA, Steinman JC. (1997) Internal fixation of the cervical spine. In: Frymoyer JW (ed). *The Adult Spine. Principles and Practice*. 2nd ed.: Lippincott-Raven;: Philadelphia .Pps:1119-1147.
- Anderson, D. K . (2005) Recommended guidelines for studies of human subjects with spinal cord injury. *Spinal Cord* 43, 453–458.
- Bareyre FM, Schwab ME. 2003. Inflammation, degeneration and regeneration in the injured spinal cord: Insights from DNA microarrays. *Trends in Neurosciences* 26(10): 555-563.
- Baydur A, Adkins RH, Milic-Emili J. Lung mechanics in individuals with spinal cord injury: effects of injury level and posture. *J. Appl. Physiol.* 2001;90:405-411,
- Benevento BT, Sipski ML(2002). Neurogenic bladder, neurogenic bowel, and sexual dysfunction in people with spinal cord injury. *J .Physical Therapy* 82(6): 601-612.
- Benzel EC, Hart B, Ball P, Baldwin N, Orrison W, Espinosa M. Magnetic resonance imaging for the evaluation of patients with occult cervical spine injury. *J Neurosurg*, 85:824-829, 1996.
- Blam OG, Ehrler DM, Rauschnig W (2003): Anatomy and Pathophysiology of traumatic spinal cord injury. In: Vaccaro AR ed: *Fractures of the cervical, thoracic, and Lumbar Spine*. New York. Marcel Dekker, Inc.Ch 2: 9-21.

- Brazis PW, Masdeu JC, Biller J(2003): The localization of lesions affecting the spinal cord. In: "Localization in Clinical Neurology". 3rd ed. Philadelphia, Pa: Lippincott Williams & Wilkins: 79-108.
- Brummett R, Rushton S, Vaccaro AR (2003): Imaging in spinal trauma. In: Vaccaro AR ed: fractures of the cervical, Thoracic, and lumbar spine. New York. Marcel Dekker, Inc. 7: 81-102
- Bunge MB, Wood PM (2004): Transplantation of Schwann cells and olfactory ensheathing cells to promote regeneration in the CNS. In: Selzer ME, Clarke S,(eds). Textbook of Neural Repair and Rehabilitation. Cambridge University Press. United Kingdom.
- Burney RE, Maio RF, Maynard F(2006): Incidence, characteristics, and outcome of spinal cord injury at trauma centers in North America. Arch Surg May; 128(5): 596-9.
- Canadian Institute for Health Information CIHI (2006): Life after traumatic spinal cord injury: From inpatient rehabilitation back to the community. Analysis in Brief.
- Carrino JA, Manton GL, Morrison WB, et al(2006).Posterior longitudinal ligament status in cervical spine bilateral facet dislocations. *Skeletal Radiol.*;35(7):510-514.
- Catharyn T Liverman, Bruce M. Altevogt, and Richard T. Johnson, Editors,,: (2005) Spinal Cord Injury: Progress, Promise, and Priorities National Academy of Sciences press Washington D.C.ch 2 pps. 34-38.
- Chen Y, DeVivo MJ and Jackson AB:(2005)Pressure ulcer prevalence in people with spinal cord injury: age-period-duration effects. Arch Phys Med Rehabil.;86(6):1208-1213.
- Chen J, Dihne M & Schachner M (2005): Cell adhesion molecule L1-transfected embryonic stem cells with enhanced survival support regrowth of oritospinal tract axons in mice after spinal cord njury. J. Neurotrauma 22, 896–906

- Cheng, H, Liao, K K , & Shih, Y. H. (2004): Spinal cord repair with acidic fibroblast growth factor as a treatment for a patient with chronic paraplegia. *Spine* 29, E284–E288.
- Chiu WC, Haan JM, Cushing BM, et al (2001). Ligamentous injuries of the cervical spine in unreliable blunt trauma patients: incidence, evaluation, and outcome. *J Trauma*.;50:457-464.
- Dennis A (2005) Spine Injuries *In* “Neurosurgery Principles and Practice. Anne J. Moore and David W. Newell (Eds)” Velez and David W. Newell Springer-Verlag Limited London. Pps. 210-243
- Dennis, GC, Dehkordi, O, Millis, RM , Said B., and Baganz D.(2000)., Somatosensory evoked potentials, neurological examination and magnetic resonance imaging for assessment of cervical spinal cord decompression, *J. Life Sci.*, 66(5):389–397,
- DeVivo MJ(2002): Epidemiology of Traumatic Spinal Cord Injury. In: Kirshblum S et al, (eds). *Spinal Cord Medicine*. Vol 1. Lippincott Williams & Wilkins Baltimore, Ch3, pps 69-81.
- Drake's R (2006): Anatomy of Medulla spinalis .*In* "Gray's Anatomy Of The Human Body" 38th ed. Lea and Faberger. Philadelphia. Ch.35: Pp 722-748.
- Faden AL, Ellison JA, Noble LJ (1999): Effects of competitive and non competitive NMDA receptor antagonists in spinal cord injury. *Eur. Pharmacol.*, 175: 165.
- Fehlings MG, Perrin RG(2006): The role and timing of early decompression for cervical spinal cord injury: update with a review of recent clinical evidence. *Injury*; 36(Suppl 2): S13-S26.
- Fehlings MG, Tator CH (1999) An evidence-based review of decompressive surgery in acute spinal cord injury: rationale, indications, and timing based on experimental and clinical studies. *J Neurosurg* 91:1–11
- Flanders AE, Croul SE. (2002) Spinal trauma. *In* "Atlas SW. Magnetic Resonance Imaging of the Brain and Spine". 3rd ed.: Lippincott Williams & Wilkins Philadelphia .Pps:154-167.

- Formal CS, Saulino MF, Ditunno JF (2003): Rehabilitation after traumatic spinal cord injury. In: Vaccaro AR ed: Fractures of the cervical, thoracic, and lumbar Spine.. Marcel Dekker, Inc New York., ch52: 717-734.
- Fouad, K. (2005) Combining Schwann cell bridges and olfactory-ensheathing glia grafts with chondroitinase promotes locomotor recovery after complete
- Freed CR, Greene PE, Breeze RE and Fahn S. (2001): Transplantation of embryonic dopamine neurons for severe Parkinson's disease. *New Engl. J Med* 344: 710–719,
- Frost FS(2000). Spinal cord injury medicine. In: Braddom RL editor. Physical medicine and rehabilitation. 2nd ed.: WB Saunders; Philadelphia. Ch 22p. 1230-82.
- Geisler FH, Coleman WP, Grieco G, Dorsey FC,(2001): The GM-1 ganglioside multi-center acute spinal cord injury study. **Spine** 26:S87-S98,.
- Gonzales EG, Myers SJ, Edelstein JE, Lieberman JS (eds.)(2000): Downey and Darling's Physiological Basis of Rehabilitation Medicine (3rd Ed.): Butterworth-Heinemann Publishers, Boston. Ch 8 Pp: 698-704.
- Grabb PA, Pang D (1999): Magnetic resonance imaging in the evaluation of spinal cord injury without radiographic abnormality in children. *Neurosurgery*, 35: 406-414.
- Greenberg M (2006): Spine injuries *In "Hand Book Of Neurosurgery"*.6th ed. Thieme medical publishers. New York . Ch 25,Pps 701-703.
- Hadley MN, Walters BC, Grabb PA(2002): Guidelines for the management of acute cervical spine and spinal cord injuries. *Clin Neurosurgery*; 49: 407-98.
- Hains BC, Black JA, Waxman SG.(2003a): Primary cortical motor neurons undergo apoptosis after axotomizing spinal cord injury. *Journal of Comparative Neurology* 462(3): 328-341.
- Hains BC, Klein JP, Saab CY.(2003b): Up regulation of sodium channel Nav1.3 and functional involvement in neuronal hyperexcitability associated with central neuropathic pain after spinal cord injury. *Journal of Neuroscience* 23(26): 8881-8892.

- Harris MB, Sethi RK(2006): The initial assessment and management of the multiple-trauma patient with an associated spine injury. *Spine J.* 15; 31(11 Suppl): S9-15.
- Heller J, Pedlow F (2002): Anatomy of the cervical spine and spinal cord . *In*: Clark, C., ed. "The cervical spine". Philadelphia: Lippincott-Raven, 12-14.
- Hickey JV: (2003)*The Clinical Practice of Neurological and Neurosurgical Nursing* [5th Ed.],. Lippincott Williams & Wilkins Philadelphia, , pp 409–412
- Hoffman JR, Mower WR, Wolfson AB(2006): Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. National Emergency X-Radiography Utilization Study Group. *N Engl J Med* Jul 13; 343(2): 94-99.
- Hosten N, Liebig T (2004): Spinal trauma. *In*: CT of the head and spine ed by Hosten N, Liebig T, Georg Thieme Verlag 12: 324-336.
- Hulsebosch CE(2002). Recent advances in pathophysiology and treatment of spinal cord injury. *Advances in Physiology Education* 26(1-4): 238-255.
- Insko EK, Gracias VH, Gupta R(2002). Utility of flexion and extension radiographs of the cervical spine in the acute evaluation of blunt trauma. *J Trauma*.;53:426-429.
- J. R. Ball, L. H. Sekhon (2006)Timing of Decompression and Fixation After Spinal Cord Injury – When is Surgery Optimal?J.Critical Care and Resuscitation; 8: 56-63
- Jay Jagannathan and John A. Jane JR. (2006), Cervical spine injuries in pediatric athletes: mechanisms and management *Neurosurg. Focus / Vol 21 /Pps* 124-128
- Jeffery, N. D., & Franklin, R. J. (2005):Autologous olfactory glial cell transplantation is reliable and safe in naturally occurring canine spinal cord injury. *J. Neurotrauma* 22, 1282–1293.
- Jones LL, Sajed D, Tuszynski MH.(2003). Axonal regeneration through regions of chondroitin sulfate proteoglycan deposition after spinal cord injury: A balance of permissiveness and inhibition. *Journal of Neuroscience* 23(28): 9276-9288.

- Jones ML, Mathewson CS, Adkins VL and Ayllon T. (2003) Use of behavioral contingencies to promote prevention of recurrent pressure ulcers. *Arch Phys Med Rehabil*;84(6):796-802.
- Keirstead HS, Nistor G, Bernal G, Totoiu M, Cloutier F, Sharp K, Steward O: Human embryonic stem cell-derived oligodendrocyte progenitor cell transplants remyelinate and restore locomotion after spinal cord injury. *J Neurosci* 2005, 25:4694-4705.
- Kerr DA, Shambloot MJ, and Rothstein JD (2003) Human embryonic germ cell derivatives facilitate motor recovery of rats with diffuse motor neuron injury. *J Neurosci* 23: 5131–5140,.
- Kilburn MPB, Smith DP, Hadley MN. (2001) The Initial Evaluation and Treatment of the Patient with Spinal Trauma. In: “Batjer HH and Loftus CL (eds.) Textbook of Neurological Surgery”. Lippincott/Williams & Wilkins, Philadelphia, PA. Pps 234-239,.
- Knoller, N. (2005): Clinical experience using incubated autologous macrophages as a treatment for complete spinal cord injury: Phase I study results. *J. Neurosurg. Spine* 3, 173–181.
- Kossmann T, Trease L, and Malham G. Damage control surgery for spine trauma. *Injury* 2004;35:661-670).
- Krause JS and Broderick L. (2004) Patterns of recurrent pressure ulcers after spinal cord injury: Identification of risk and protective factors 5 or more years after onset. *Arch Phys Med Rehabil*;85:1257-1264.
- Krause JS, Sternberg M, Lottes S, et al (2000): Mortality After Spinal Cord Injury: An 11-year Prospective Study. *Archives of Physical Medicine and Rehabilitation* 78: 815-821
- Li AE, Fishman EK. Cervical spine trauma (2003): evaluation by multidetector CT and three-dimensional volume rendering. *Emerg Radiol.*;10:34–39.
- Mamas MA, Reynard JM, Brading AF (2003). Nitric oxide and the lower urinary tract: Current concepts, future prospects. *Urology* 61(6): 1079-1085.

- Mathias CJ, Bannister R(2002): Autonomic Failure: A Textbook of Clinical Disorders of the Autonomic Nervous System. 4th ed. Oxford University Press; London, England:Ch12,Pps187-199.
- Maxwell RA, Parker DE, Durham RM, et al.(2002)Routine prophylactic vena cava filtration is not indicated after acute spinal cord injury. The Journal of Trauma; 52(5):902-906.
- McGuire RA (2001): Physical examination in spinal trauma. IN: Levine AM Eismont FJ, Garfin SR, Zigler JE d. Spinal Trauma. W.B. Saunders Company 3: 13-27.
- McMinn RM (2005): Cervical spinal cord. In “Last's anatomy: Regional and applied”. 9th ed. Oxford University Press; London, England: Ch6 Pps: 535-548.
- O'Connor PJ. (2005)Survival after spinal cord injury in Australia. Arch Phys Med Rehabil;86:37-47.
- Ogawa, Y. (2002). Transplantation of in vitro-expanded fetal neural progenitor cells results in neurogenesis and functional recovery after spinal cord contusion injury in adult rats. J. Neurosci. Res. 69, 925–933
- Pandyan AD, Gregoric M& GR Johnson. (2005) Spasticity: Clinical perceptions, neurological realities and meaningful measurement. Disability and Rehabilitation; 27: 2-6.
- Pang D. (2004)Spinal cord injury without radiographic abnormality in children, 2 decades later. *Neurosurgery*.;55:1325-1343.
- Parent A, Carpenter MB(2006): Spinal cord fiber tracts. In: Carpenter's Human Neuroanatomy. 9th ed.: Lippincott, Williams & Wilkins; Philadelphia, Pa: 325-417.
- Payer M , Cyrille Sottas(2008):Mini-open anterior approach for corpectomy in the thoracolumbar spine Surgical Neurology 69 25–32
- Peckham PH, Keith MW and Wiegner A(2001). Efficacy of an implanted neuroprosthesis for restoring hand grasp in tetraplegia: A multicenter study. Arch Phys Med Rehabil; 82: October, 1380-1388.

- Perrier AL, Tabar V, Barberi T and Studer L.(2004)Derivation of midbrain dopamine neurons from human embryonic stem cells. *Proc Natl Acad Sci USA* 101: 12543–12548,.
- Popovich PG, (2003). Manipulating neuroinflammatory reactions in the injured spinal cord: Back to basics. *Trends in Pharmacological Sciences* 24(1): 13-17
- Priebe MM, Sherwood AM, Markowski(1996).J. Clinical assessment of spasticity in spinal cord injury: a multidimensional problem. *Arch Phys Med Rehabilitation*; 77: 713–716.
- Raineteau O, Schwab ME.(2001). Plasticity of motor systems after incomplete spinal cord injury. *Nature Reviews Neuroscience* 2(4): 263-273.
- Reid DC, Henderson R, Saboe L(2002): Etiology and clinical course of missed spine fractures. *J Trauma*,Sep; 27(9): 980-6.
- Saboe LA, Reid DC, Davis LA, et al(2006): Spine trauma and associated injuries. *J Trauma* Jan; 31(1): 43-8.
- Schurch B, Rossier AB.(1996):Post-traumatic syringomyelia (cystic myelopathy): a prospective study of 449 patients with spinal cord injury. *J. Neurology Neurosurg. and Psychiatry*;60: 61–67.
- Selden NR, Quint DJ, Patel N, et al(1999). Emergency magnetic resonance imaging of cervical spinal cord injuries: clinical correlation and prognosis. *Neurosurgery*;44:785–792.
- Shehu B. B and N. J. Ismail.(2004): Practical management of spinal cord injury. *Annals of African Medicine* Vol. 3, No. 1;; 46 – 5
- Shergill IS, Hamid R, et al(2004). The importance of autonomic dysreflexia to the urologist. *British J of Urology Int*;93:923–6.
- Siddall PJ, McClelland JM, Cousins MJ. (2003). A longitudinal study of the prevalence and characteristics of pain in the first 5 years following spinal cord injury. *Pain* 103(3): 249-257.

- Sipski, M.L., Alexander, C.J. (2000). Sexual function and dysfunction after spinal cord injury. *Physical Medicine and Rehabilitation Clinics of North America* WB Saunders Company.Philadelphia: (pp. 811-828).
- Smith D, Walmsley (1999): Segmental cervical spine motion during orotracheal intubation of the intact and injured spine with and without external stabilization. *J Neurosurg. (Spine 2)*: vol.92: 201-206.
- Smith, H.C., Savic, G., (2000), Corticospinal function studied over time following incomplete spinal cord injury, *Spinal Cord*, 38(5):292–300,.
- Snell R (2004): *Clinical Neuroanatomy for medical student*. 3rd ed. Little Brown Boston,ch4.pps76-85
- Snoek GJ, Sorensen F(2004). Survey of the needs of patients with spinal cord injury: impact and priority for improvement in hand function in tetraplegics. *Spinal Cord J.*; 42; 526-532.
- Stiell IG, McKnight RD(2003): The Canadian C-spine rule versus the NEXUS low-risk criteria in patients with trauma. *N Engl J Med* Dec 25; 349(26): 2510-8
- Swischuk LE(2000): The spine and spinal cord, in *Emergency Imaging of the Acutely Injured Child*, 4th ed.: Lippincott Williams & Wilkins, Philadelphia, pp 532–587
- Tator CH(2004): Acute management of spinal cord injury. *Br J Surg* May; 77(5): 485-6.
- Tay B, Eismont FJ (2003): Physical examination: ASIA motor/ sensory examination and spinal cord injury syndrome. *In: Vaccaro AR ed: Fractures of cervical spine*. 1st ed. Lippincott/Williams & Wilkins, Philadelphia,
- Terre R, Valles M, Vidal J (2000). Post-traumatic syringomyelia following complete neurological recovery. *Spinal Cord* 38(9): 567-570. transection of the spinal cord. *J. Neurosci.* 25, 1169–1178.
- Tubbs R, Grabb p, Spooner, (2002): The apical ligament; anatomy and functional significance ,*j Neursoug.(spine2)*94;197-200.

- Vaccaro A, Ahmad S, Raushcing W(2002): Anatomy and pathophysiology of spinal cord injury with APSCI. *In*: “Spine Trauma Levine, A, ed.”. Philadelphia: WB Saunders, 75-86.
- Vitarbo L, Levi ADO. (2005) Cervical laminoplasty—open door. *In*: “Benzel EC editor. Spine surgery techniques, complication avoidance, and management”. 2nd ed. Harcourt Health Science; Chapter 166 pp. 2103-2112.
- Waxaman SG (2003): Spinal cord and spine. *In*: Clinical neuroanatomy ed. by Waxman SG 25th ed. McGraw Hill 3: 45-80.
- Weber DJ, Stein RB, Chan KM, Loeb GE, (2004)Functional electrical stimulation using microstimulators to correct foot drop: a case study. *Can J Physiol Pharmacol*;82:784-792.
- Wobus, Anna M., and Kenneth R. Boheler. (2005)Embryonic Stem Cells: Prospects for Developmental Biology and Cell Therapy. *Physiol Rev* 85: 635–678.
- Wu SS, Hwa SY, Lin LC, et al(1996).Management of rigid post traumatic kyphosis. *J.Spine*;21:2260–2267.
- Wu X, Ding S, Ding Q, Gray NS, and Schultz PG.(2004):Small molecules that induce cardiomyogenesis in embryonic stem cells. *J.Am. Chemistry Soc* 126: 1590–1591,
- Young PA, Young PH(2002): Basic Clinical Neuroanatomy.1st ed Lippincott, Williams & Wilkins; . Philadelphia, Pps13-22; 99-116; 127-152; 217-234.
- Zheng, B. (2005) Genetic deletion of the Nogo receptor does not reduce neurite inhibition in vitro or promote corticospinal tract regeneration in vivo. *Natl Academy. of Science. USA* 102, 1205–1210.