

REFERENCES

- 1) **Czajka I and Zgliczynski W. (2005):** Gynecomastia-Pathogenesis,diagnosis and treatment.Endokrynol Pol.; 56(3): 269-77.
- 2) **Braunstein GD, Friedman SB, Fisher M and Schonberg SK. (1992):** Gynecomastia.NEJM; 328: 490-494.
- 3) **Griffen JE and Wilson JD. (2003):** Disorders of the testes and the male reproductive tract. Abnormalities in estrogen metabolism. Gynecomastia. In: Larsen PR, Kronenberg HM, Melmed S, Polonsky KS, editors. Williams Textbook of Endocrinology. 10th ed. Ch.18. Philadelphia, PA: Elsevier; 2003. p. 741-746,753.
- 4) **Bembo SA and Carlson HE. (2004):** Gynecomastia:its features,and when and how to treat it.CLEVELAND CLINIC JOURNAL OF MEDICINE; 71:6.
- 5) **Hodgson EL, Fruhstorfer BH and Malata CM. (2005):** Ultrasonic liposuction in the treatment of gynecomastia. Plast Reconstruct Surg. 2005 Aug; 116(2): 646-53; discussion 654-5.
- 6) **Einav-Bachar R, Phillip M, Aurbach-Klipper Y and Lazar L.(2004):** Prepubertal gynecomastia: etiology, course and outcome. Clin Endocrinal; 61(6): 779-80.
- 7) **Mc Minn RM. (1994):** Anatomy of the Breast. In: lasts Anatomy; regional and applied, 9th edition; 2:70-73.
- 8) **Ellis H. (1992):** Female breast clinical anatomy. 8th edition, Oxford; 169-173.
- 9) **Healey. (1990):** Structure of the breast, Surgical Anatomy of Breast; 20-23.
- 10) **Burns PN, Davis JD and Halliwell M. (1984):** Doppler US in diagnosis of breast cancer. In: US in breast and endocrine disease, Leopold GR (ed); 41-56.

- 11) Hideo Nakajima, Nobuaki Imanishi and Sadaka ZU Aiso.(1996):** Arterial anatomy of the nipple and areola complex. *Plast. Recon. Surg*; 843-844.
- 12) Haagensen CD. (1971):** Carcinoma of the male breast. In: *Diseases of breast*. Philadelphia: WB. Saunders; 779-780.
- 13) Herniques CQ. (1982):** The veins of the vertebral column and their role in the spread of cancer. *Ann. R. Coll. Surg.*; 31:7-22.
- 14) Gregory S. Georgiade and Nicolas G Georgiade. (1997):** Innervation of the breast. In: *Plastic, Maxillofacial and Reconstructive surgery*; 3rd edition; 64: 752.
- 15) J. Dirk Iglehart and Carolyn M. Kaelin. (2004):** Diseases of the breast. *Sabiston textbook of surgery*, 17th edition; 32:867-944. Edited by: **Courtney M.T, R. Daniel Beauchamp, B. Mark Evers, Kenneth L. Mattox**. Published by: **Elsevier Sanuders**.
- 16) Adel HR, Reza M and Mohammed K. (2006):** Gynecomastia evaluation and management. *Recent advanced in surgery*; 29:13. Edited by: **C.D.Johnson and I.Taylor**. Published by: **the Royal Society of Medicine Press Ltd**.
- 17) Niewoehner CB and Nuttal FQ.(1984):** Gynecomastia in a hospitalized male population. *AM J Med*; 77:633-638.
- 18) Braunstein GD. (2000):** Management of gynecomastia. In:Harris JR, Lippman ME, Morrow M, Osborne CK. (eds) *Diseases of the Breast*, 2nd edn. Williams & Wilkins, 2000.
- 19) Richard A.Maldick. (2004):** Gynecomastia. *Aesthetic Surgery Journal* 2004; 24:471-479.
- 20) Ando S, De AF, Rago V, Carpino A et al. (2002):** Breast cancer: from estrogen to androgen receptor. *Mol Cell Endocrinal* 2002; 193: 121-128.

- 21) **Hershkovitz E, Leiberman E. (2002):** Gynecomastia: a review. *Endocrinologist* 2002; 12: 321-332.
- 22) **Hall A, Feldman HA, Mckinley JB et al. (1991):** Age, disease, and changing sex hormone levels in middle-aged men: results of the Massachusetts male aging study. *J Clin Endocrinal Metab* 1991; 73: 1016-1025.
- 23) **Korenman SG. (1986):** The endocrinology of the abnormal male breast. *Ann NY Acad Sci* 1986; 464:400-408.
- 24) **Zayed A, Stock JL, Liepman MK et al. (1994):** Feminization as a result of both peripheral conversion of androgens and direct estrogen production from an adrenocortical carcinoma. *J Endocrinol Invest* 1994; 17:275-278.
- 25) **Braunstein GD. (1993):** Gynecomastia. *N Engl J Med* 1993; 328:490-495.
- 26) **Carlson HE. (1992):** Gynecomastia. In: Morley JE, Korenman SG, editors. *Endocrinology and Metabolism in the Elderly*. Boston: Blackwell Scientific Publications Inc, 1992:294-307.
- 27) **Turkington RW. (1972):** Serum prolactin levels in patients with gynecomastia. *J Clin Endocrinol Metab* 1972; 34:62-66.
- 28) **Gooren LJG and Daantje CRE. (1986):** Psychological stress as a cause of intermittent gynecomastia. *Horm Metab Res* 1986; 18:424.
- 29) **Choi W-T, Mac Lean HE, Chus, Warne GL and Zajac JD. (1993):** Kennedy`s Disease: genetic diagnosis of an inherited form of motor neuron disease. *Aust NZ. J. Med*; 23:187-192.
- 30) **Loriaux DL, Menard R, Taylor A, Pita JC and Santen R. (1976):** Spironolactone and endocrine dysfunction. *Ann Intern Med* 1976; 85:630-636.
- 31) **Braunstein GD. (1999):** Aromatase and gynecomastia. *Endocrine-Related Cancer* 1999; 6:315-324.

- 32) MacDonald. (1979):** J Clin Endocrinal Metab 1979; 49:905-916.
- 33) Tavassoli Fattanch. (1999):** Male breast lesions. In: Pathology of the breast, chapter 16, 2nd edition; 829-849.
- 34) Bannayan GA, Hajdu SI. (1972):** Gynecomastia: clinicopathologic study of 351 cases. Am. J. Clin. Pathol.1972; 57:431-437.
- 35) Pages A and Ramos J. (1984):** Contribution a l'etude des gyne'comasties. A propose de 122 cas. Ann. Pathol. (Paris); 4: 137-142.
- 36) De May RM. (1996):** The art and science of cytopathology. Volume 2. Chicago, American Society of Clinical Pathologies; pp.890-891.
- 37) Malini H, Melissa PM, Paul PR and Sandra JS. (2005):** Prepubertal gynecomastia with lobular differentiation. The Breast Journal. 2005; 11:48-51.
- 38) Robert AE, Martin S, Patrick HB and Mark AS. (2000):** Gynecomastia a clinical review. Aesthetic Surgery Journal- September/October 2000; 20:381-386.
- 39) Pinedo F, Nargas J, de Agustin P, Perez-Barrios A and Ballestin C. (1991):** Epithelial atypia induced by chemotherapeutic drugs: A possible pit fall in fine aspiration biopsy. Acta. Cytol.; 35: 229-233.
- 40) Sneige N, Holder PD, Katz RL, Fanning CV, Dekmezian RI, Shabb NS and Sigletry SE. (1993):** Fine needle aspiration cytology of the male breast in a cancer center. Diagn. Cytopathol.; 9:691-697.
- 41) Fowler LJ, Smith SS, Snider T and Schultz MR. (1996):** Apocrine metaplasia in gynecomastia by fine needle aspiration as a possible indicator of anabolic steroid use. Acta. Cytol.; 40: 734-738.
- 42) Pensler JM, Silverman BL, Sanghavi J, Goolsby C, Speck G, Brizio-Molteni L and Molteni A. (2000):** Estrogen and progesterone receptors in gynecomastia. Plast. And Recon. Surg.; Vol. 6, No. 5, 1011-1013.

- 43) Webster JP. (1946):** Mastectomy for gynecomastia through a semicircular intra-areolar incision. Ann. Surg 1946;124:557.
- 44) Simon BE, Hoffman S and Kahn S. (1973):** Classification and surgical correction of gynecomastia. Plast. Recon. Surg; 51:48.
- 45) Carlson HE. (1980);** Current concepts of gynecomastia N. Engl. J. Med, 303: 795.
- 46) Deutinger M and Freilinger G. (1986):** Die gynakomastie: versuch einer klassifizierung und operative ergebnisse. Hanchirurgie; 18, 239-243.
- 47) Rohrich Rod J, Richard YH, Jeffery M, Kenkel, William P and Adam JR. (2003):** Classification and management of gynecomastia: Defining the role of ultrasound assisted liposuction. Plast. And Recon. Surg; Vol. 3, No. (2) 909-925.
- 48) Robert S.Hamas and Chalm W.Williams. (1998):** A sharp cutting liposuction cannula for gynecomastia. Aesthetic Surg J 1998; 18: 261-265.
- 49) Andersen JA, Gram JB. (1982):** Male breast at autopsy. Acta Pathol Microbiol Immunol Scand [A] 1982; 90:191-197.
- 50) Treves N. (1958):** Gynecomastia; the origins of mammary swelling in the male: an analysis of 406 patients with breast hypertrophy, 525 with testicular tumors, and 13 with adrenal neoplasms. Cancer 1958; 11: 1083-1102.
- 51) Joshi A, Kapila K and Verma K. (1999):** Fine needle aspiration cytology in the management of male breast masses. Acta. Ctol; 43: 334-338.
- 52) Chantra PK, So GJ, Wollman JS and Basset LW. (1995):** Mammography of the male breast. Am. J. Roent; 165:853-858.
- 53) Svensson WE. (2000):** The value of ultrasound scanning in breast disease. Hosp. Med; 61:233-239.

- 54) Myking OL and Thorsen T. (1981):** Conjugated and unconjugated estrogens in men with hyperestrogenism and in normal men: The day to day variation and the relation ship between different estrogens. *J. Steroid Biochem*; 14:935-940.
- 55) Ismail AA and Barth JH. (2001):** Endocrinology of gynecomastia. *Ann. Clin. Bioderm*; 38: 596-607.
- 56) Xciarra A, Casale p, Di Nicola S, Chiro C and Disilverio F. (1998):** Hormonal profile of patients with leydig cell tumors, a urologic cause of gynecomastia. *Mineria Urol. Nefrol*; 50:225-231.
- 57) The Copenhagen Study Group for Liver Diseases. (1986):** Testoserone treatment of men with alcoholic cirrhosis: adouble-blind study. *Hepatology* 1986; 6: 807.
- 58) Kuhn JM, Roca R, Laudat MH, Rieu M, Luton JP and Bricaire H. (1983):** Studies on the treatment of idiopathic gynecomastia with percutaneous dihydrotestosterone. *Clin Endocrinal (Oxf)*; 19: 513-520.
- 59) Jones DJ, Holt SD, Surtees P, Davison DJ and Coptcoat MJ. (1990):** A comparison of danazol and placebo in the treatment of adult idiopathic gynecomastia: results of a prospective study in 55 patients. *Ann R Coll Surg Engl* 1990; 72: 296-298.
- 60) Stepanas AV, Burnet RB, Harding PE and Wise PH. (1977):** Clomiphene in the treatment of pubertal-adolescent gynecomastia: a preliminary report. *J Pediatr*; 90: 651-653.
- 61) Plourde PV, Kulin HE and Santner SJ. (1983):** Clomiphene in the treatment of adolescent gynecomastia. *Clinical and endocrine studies. Am J Dis Child*; 137: 1080-1082.
- 62) LeRoith D, Sobel R and Glick SM. (1980):** The effect of clomiphene citrate on pubertal gynecomastia. *Acta Endocrinal (Copenh)*; 95: 177-180.

- 63) Zachmann M, Eiholzer U, Muritano M, Werder EA and Manella B. (1986):** Treatment of pubertal gynecomastia with testolactone. Acta Endocrinol Suppl (Copenh); 279: 218-226.
- 64) McDermott MT, Hofeldt FD and Kidd GS. (1990):** Tamoxifen therapy for painful idiopathic gynecomastia. South Med J; 83: 1283-1285.
- 65) Parker LN, Gray DR, Lai MK and Levin ER. (1986):** Treatment of gynecomastia with tamoxifen: a double-blind crossover study. Metabolism; 35: 705-708.
- 66) Alagaratnam TT. (1987):** Idiopathic gynecomastia treated with tamoxifen: a preliminary report. Clin Ther; 9: 483-487.
- 67) Ting AC, Chow LW and Leung YF. (2000):** Comparison of tamoxifen with danazol in the management of idiopathic gynecomastia. Am Surg; 66: 38-40.
- 68) Hanavadi S, Banerjee D, Monypenny IJ and Mansel RE. (2005):** The role of tamoxifen in the management of gynecomastia. Breast 2005.
- 69) Tyrrel CJ. (1999):** Gynecomastia: aetiology and treatment option. Prost. Cancer Prost. Dis; 2: 167-171.
- 70) Dicker Adam p. (2003):** The safety and tolerability of low-dose irradiation for the management of gynecomastia caused by antiandrogen monotherapy. The Lancet Oncology; Vol. 4, No.1, 1-11.
- 71) Chou JL, Easely JD, Feldmeier JJ, et al. (1988):** Effective radiotherapy in palliating mammalgia associated with gynecomastia after DES therapy. Int. J. Radiat. Oncology Biol. Phys.; 15: 749-751.
- 72) McGrath MH. (1991):** Gynecomastia. In: Lewis, ed. The Art of Aesthetic Plastic Surgery. Boston: Little, Brown; 915-925.
- 73) Maldick RA. (1991):** Gynecomastia, liposuction and excision. Clin Plast Surg; 18:815-822.

- 74) Gurcan A, Dogan T, Ahmet T and Ferruh B. (2005):** PTP Incision for Gynecomastia. *Annals of plastic surgery*; 54:130-134.
- 75) Schlenz I, Kuzbari R, G rubber H, etal. (2000):** The sensitivity of the nipple-areola complex: an anatomic study. *Plast Reconstr Surg*; 105:905-909.
- 76) Terino EO. (1990):** A transareolar circumthelial approach to augmentation mammoplasty. In:Georgiade NG, Georgiade GS, Riefkohl R, eds. *Aesthetic Surgery of the Breast*. Philadelphia: WB Saunders; 1990:101-111.
- 77) Mohammad T, Mohammad M, Jamal MH, Mohammad BH and Mohammad S. (2004):** The surgical management of high grade gynecomastia. *Annals Plastic Surgery* 2004; 53: 17-20.
- 78) Murphy TP, Ehrlichman RJ and Seckel BR. (1994):** Nipple placement in simple mastectomy with free nipple grafting for severe gynecomastia.*Plast Reconstr Surg*. 1994; 94:818-823.
- 79) Ward CM and Khalid K. (1989):** Surgical treatment of grade 3 gynecomastia. *Ann R Coll Surg Engl*;71: 226-228
- 80) Letterman G and Schurter M. (1972):** Surgical correction of massive gynecomastia. *Plast Reconstr Surg*; 49: 259-262.
- 81) Bretteville-Jensen G. (1975):** Surgical treatment of gynecomastia. *Br J Plast Surg*; 28: 177-180.
- 82) Douglas HF, Rodrigo GA and Nivaldo A. (2006):** Treatment of severe gynecomastia (grade III) by resection of periareolar skin. *Aesthetic Surg J* 2006; 26:669-673.
- 83) Metin Y, Erol K, Naile NK and Sabri A. (2006):** Light retractor-assisted transaxillary approach in gynecomastia correction. *Ann Plast Surg* 2006; 57: 370-373.
- 84) Balch CR. (1978):** A transaxillary incision for gynecomastia. *Plast Reconstr Surg*. 1978; 61: 13-16.

- 85) Gamal I. EL-Habbaa. (2005):** Modified surgical technique for treatment of high-grade gynecomastia. Benha M.J. 2005; 22: 63-74.
- 86) Matarasso MD. (2001):** Liposuction JAMA; 285 (3): 266-268.
- 87) Jennifer LW, Robert PS and Steven JB. (2004):** Cross-chest lipoplasty and surgical excision for gynecomastia. : a 10-year experience. Aesthetic Surg J 2004; 24: 216-223.
- 88) Hunstad JP. (1994):** The tumescent technique: a safe and effective method to minimize blood loss and provide consciousness anesthesia for liposuction procedures. Presented at the 27th annual meeting of ASAPS, Dallas, Texas.
- 89) De Souza Pinto, Patricia EI, Adriane da costa Muniz and Yobanka PM. (1996):** Superficial liposuction. Clinics in Plastic Surgery; 23 (4): 529-549.
- 90) Zocchi ML. (1996):** Ultrasonic assisted lipoplasty. Technical refinements and clinical evaluations. Clinics in Plastic Surgery. 3:575.
- 91) Rohrich Rod J, Richard YH, Jeffery M, Kenkel, William P and Adam JR. (2003):** Classification and management of gynecomastia: Defining the role of ultrasound assisted liposuction. Plast. And Recon. Surg; 111:909.
- 92) Elaine Hodgson, Birgit Fruhstorfer and Charles Malata. (2005):** Ultrasonic liposuction in the treatment of gynecomastia. Plast Reconstr Surg; 116:646.
- 93) Baker JL. (1999):** A practical guide to ultrasound-assisted lipoplasty. Clin Plast Surg. 26:363.
- 94) Bekenstein MS and Grotting JC. (2000):** Ultrasound-assisted lipectomy using the solid probe: Aretrospective review of 100 consecutive cases. Plast Reconstr Surg. 105:2161.

- 95) Rohrich RJ, Beran SJ, Kenkel JM, Adams WP and Dispaltro F. (1998):** Extending the role of liposuction in body contouring with ultrasound-assisted liposuction. *Plast Reconstr Surg.* 101:109.
- 96) Calobrace MB. (1999):** Ultrasound-assisted lipoplasty. *Clin Plast Surg.* 26:103.
- 97) Ablaza VJ, Gingrass MK, Perry LC, Fisher J and Maxwell GP. (1998):** Tissue temperatures during ultrasound-assisted lipoplasty. *Plast Reconstr Surg.* 102:534.
- 98) Thornton LK and Nahai F. (1999):** Equipment and instrumentation for ultrasound-assisted lipoplasty. *Clin Plast Surg.* 26:299.
- 99) Maxwell GP and Gingrass MK. (1998):** Ultrasound-assisted lipoplasty. Aclinical study of 250 consecutive patients. *Plast Reconstr Surg.*101:189.
- 100) Eaves Felmont F, John Bostwick and Foad Nahai. (1995):** Instrumentation and setup for endoscopic plastic surgery. Minimal invasive techniques in aesthetic and reconstructive surgery. *Clinics in Plastic Surgery; Vol. 22 No. (4).* 591-603, 683-695.
- 101) Kaoru K, Mayumi I, Hiromichi I, Junko K, Makoto H and Keizo S. (2002):** Early results of an endoscope-assisted subcutaneous mastectomy and reconstruction for breast cancer. *Surgery.*2002; 131:S324-329.
- 102) Ytzhack R, Lucian F, Isaac JP, Liron E, Dana E and Yehuda U. (2005):** Multimodality gynecomastia repair by cross-chest power-assisted superficial liposuction combined with endoscopic-assisted pull-through excision. *Ann Plast Surg.*2005; 55:591-594.
- 103) Arturo C. Pardo and Paulo F. Castillo. (2005):** Minimal surgical access to treat gynecomastia with the use of power-assisted arthroscopic-endoscopic cartilage shaver. *Plast Reconstr Surg.* 115:939-942.

- 104) Courtiss EH, Becker H, Rosenberg GJ, et al. (1994):** Treatment of gynecomastia. Aesthetic surgery; 4-9.
- 105) Gingrass MK and Kenkel JM. (1999):** Comparing ultrasound assisted liposuction with suction assisted liposuction. Clinics in Plastic Surgery; 26: 283.
- 106) Kenkel J, Robinson J, Berans et al. (1998):** The tissue effects of ultrasound assisted lipoplasty. Plast. Recon. Surg.; 102: 213, 1998.
- 107) Azurin D, Gingrass M, Fisher J, et al. (1998):** Seroma following ultrasound assisted lipoplasty presented at the 67th annuals ASPRS meeting, Boston.
- 108) Davis FA. (1981):** Taber's Cyclopedic medical dictionary; ed. (14): Philadelphia.
- 109) Christman K. (1986):** Death following suction lipectomy and abdominoplasty. Plast. Recon. Surg.; 78:428.
- 110) Skarzynski J, Salvin J, Spencer R, et al. (1986):** Matching ventilation perfusion images in fat embolism. Clin. Nucl. Med.; 11:40.
- 111) Zocchi M. (1993):** Clinical aspects of ultrasound liposculpture. Persp. Plast. Surg.; 7: 153.
- 112) Klein JA. (1987):** The tumescent technique for liposuction surgery: Am. J. Cosmetic surg.; 4:263.
- 113) Samdal F, Anland P and Bugge J. (1994):** Plasma lidocaine levels during suction assisted lipectomy using large doses of dilute lidocaine with epinephrine. Plast. Recon. Surg.; 93:1217.
- 114) Ostad A, Kageyama N and Moy R. (1996):** Tumescent anesthesia with lidocaine doses 55 mg/Kg is safe for liposuction. Derm. Surg.; 22:921.
- 115) Ovrebo UK, Grong K and Vindenes H. (1997):** Small intestinal perforation and peritonitis after abdominal lipoplasty. Ann. Plast. Surg.; 38: 642.