

1.8. Radiolabelling methods	27
1.8.1. Radionuclide selection for labelling	27
1.8.2. Chemistry of labelling with iodine	29
1.8.3. Electrophilic substitution mechanism	30
1.8.4. Choice of radioiodination method	31
1.8.4.1. Iodine monochloride methods	32
1.8.4.2. Chloramine-T method	32
1.8.4.3. Alternative chemical oxidation methods	34
1.8.4.3.1. Sodium hypochlorite method	34
1.8.4.3.2. N-bromosuccinimide method	34
1.8.4.4. Iodogen method	35
1.8.4.5. Electrolytic iodination method	36
1.8.4.6. Enzymatic iodination method	37
1.8.5. Conjugation labelled methods for the iodination proteins and peptides method.	38
1.8.6. Purification of radiolabelled protein from unreacted radioiodide.	40
1.8.7. Assessment of the radioiodinated tracer	41
1.8.7.1. Radiochemical purity	41
1.8.7.2. Immunoreactivity	42
1.8.7.3. specific activity (SA)	43

1.8.7.4. Stability and shelf-life (storage)	45
1.9. Performance characteristic of the optimized RIA system	46
1.9.1. Sensitivity	46
1.9.2. Precision	46
1.9.2.1. Intra assay precision	47
1.9.2.2. Inter assay precision	47
1.9.3. Accuracy	48
1.9.3.1. Recovery test	48
1.9.3.2. Dilution effect test	48
1.9.4. Validity and comparison with the available commercial kits	49

Chapter 2

2.1. Materials and methods	50
2.1.1. Chemical and reagents	50
2.1.2. Equipment	52
2.1.3. Reagent and buffer preparation	53
2.2. Radioiodination of AFP antigen	57
2.2.1. Chloramine-T method	58
2.2.2. Iodogen method	58
2.2.3. N-bromosuccinimide method	59
2.2.4. Lactoperoxidase method	59

2.2.5. Purification of ¹²⁵ I-AFP antigen	60
2.2.6. Immunoreactivity check of the prepared tracer	60
2.2.7. Comparison study between different oxidizing agent to prepared radioiodination of AFP antigen.	61
2.2.8. Storage of radioiodinated tracers	62
2.3. Preparation of AFP standards and quality controls	62
2.3.1. Human cord blood	62
2.3.2. Purification of AFP antigen from cord serum	63
2.3.3. Determination of protein concentration	64
2.4. Preparation of polyclonal antibody	64
2.4.1. Choice of host animals	64
2.4.2. preparation of the immunogen	65
2.4.3. Booster immunization	65
2.4.4. Blood sampling and harvest	65
2.4.5. Purification of rabbit anti-serum IgG obtained using ammonium sulfate method.	66
2.4.6. Estimation of protein content	67
2.4.7. Characterization and selection of antibodies or antisera	68
2.5. Formulation of RIA systems for AFP went into two ways	67
2.5.1. Double antibody liquid phase RIA assay.	68
2.5.2. Solid phase coated tube RIA system	69

2.5.2.1. Phosphate buffer approach	69
2.5.2.2. Borate buffer approach	69
2.5.2.3. Assay protocol for solid phase coated tube RIA system.	70
2.6. Optimization of the prepared RIA systems	70
2.6.1. Sample, standard or quality control volume	70
2.6.2. Incubation time	70
2.6.3. Radioactivity	71
2.6.4. Washing manner for coated tubes (RIA)	71
2.7. Performance characteristics of the optimized RIA system	71
2.7.1. Determination of the sensitivity	71
2.7.2. Determination of the precision	72
2.7.3. Effect of recovery	72
2.7.4. Effect of dilution	73
2.7.5. Method comparison	73
2.8. Statistical analysis:	73
2.8.1. Arithmetic mean value	74
2.8.2. standard deviation	74
2.8.3. Standard error of the mean	75
2.8.4. Simple linear regression and correlation coefficient	75

Chapter 3

3.1. Radioiodination of AFP antigen	77
3.1.1. Chloramine –T method	77
3.1.1.1. Effect of reaction time	77
3.1.1.2. Effect of reaction volume	81
3.1.1.3. Effect of chloramine-T content	83
3.1.1.4. Effect of reaction pH	87
3.1.2. Iodogen method	90
3.1.2.1. Effect of reaction time	90
3.1.2.2. Effect of reaction volume	93
3.1.2.3. Effect of iodogen content.	96
3.1.2.4. Effect of reaction pH	99
3.1.3. N-bromosuccinimide method	102
3.1.3.1. Effect of reaction time	102
3.1.3.2. Effect of reaction volume	105
3.1.3.3. Effect of N-bromosuccinimide content.	108
3.1.3.4. Effect of reaction pH	111
3.1.4. Lactoperoxidase method.	114
3.1.4.1. Effect of the reaction time	114
3.1.4.2. Effect of the reaction volume	117
3.1.4.3. Effect of the lactoperoxidase content.	120

3.1.4.4. Effect of the reaction pH	123
3.1.5. Effect of specific activity and storage conditions	126
3.1.6. Comparison study between different oxidizing agent.	132
3.2. Preparation of AFP standard and quality controls	134
3.3. Anti-AFP antibody evaluation	137
3.4. Formulation of RIA double antibody of PEG liquid phase system for the estimation AFP	141
3.5. Optimization of the RIA double antibody of PEG liquid phase system.	144
3.5.1. standard, sample or QC volume	144
3.5.2. Incubation time	145
3.5.3. Radioactivity counts	146
	147
3.6. Solid phase coated tubes RIA system	150
3.6.1. Optimization of a solid phase RIA coated tube system.	150
3.6.1.1. Standard, sample or QC volume	150
3.6.1.2. Incubation time	151
3.6.1.3. Radioactivity counts	152
3.6.1.4. Washing method.	153
3.6.2. Performance characteristics of coated tube AFP-RIA assay.	154