

# CONTENTS

	<i>Page</i>
<b>I- Introduction.</b>	
<b>1. Microbial enzymes in relation to their activities</b>	<b>1</b>
<b>2. Pectinases and pectic substances</b>	<b>1</b>
<b>3. Application of pectinases</b>	<b>6</b>
▪ Acidic pectinases	<b>10</b>
▪ Sparkling clear juices	<b>11</b>
▪ Apple	<b>11</b>
▪ Juice extraction	<b>11</b>
▪ Clarification	<b>12</b>
<b>4. Solid state fermentation (SSF) processes</b>	<b>7</b>
4.1 Definition of the (SSF) methods	<b>7</b>
4.2 Characteristics	<b>8</b>
4.3 Advantages of SSF	<b>9</b>
<b>II. Review of literature</b>	<b>16</b>
<b>III. Materials and methods</b>	<b>31</b>
<b>3. Construction of galacturonic acid standard curves</b>	<b>31</b>
<b>4. Protein determination</b>	<b>32</b>
4.1 Standard curve for protein determination	<b>32</b>
4.2 Determination of the total water-soluble proteins	<b>33</b>
<b>5. Growth and maintenance medium</b>	<b>34</b>
<b>6. Agro-industrial wastes</b>	<b>35</b>
6.1 Potato peels wastes (PPW)	<b>35</b>
6.2 Other agro-industrial wastes	<b>35</b>

<b>5. Production media</b>	<b>35</b>
5.1 Basal media	<b>35</b>
5.2 Potato peels basal medium (PPBM)	<b>36</b>
<b>1. Isolation of bacteria</b>	<b>31</b>
<b>2. Purification of bacterial isolates</b>	<b>31</b>
<b>8. Identification of the most potent bacterial isolates</b>	<b>36</b>
<b>8.a Media used</b>	<b>36</b>
8.a.1 Nutrient agar medium (NA)	<b>36</b>
8.a.2 King, ward and Raney's media	<b>36</b>
8.a.3 Hugh and leifson (OF) medium	<b>37</b>
8.a.4 Methyl red (MR) test medium	<b>37</b>
8.a.5 Levan production medium	<b>38</b>
8.a.6 Citrate medium	<b>38</b>
8.a.7 Urea medium	<b>39</b>
8.a.8 Gelatin agar medium	<b>39</b>
8.a.9 Czapek-Dox's tributyrin medium (CDTM)	<b>39</b>
8.a.10 Pectinase medium	<b>40</b>
8.a.11 Cellulase agar medium	<b>40</b>
<b>8.b Morphological characteristics</b>	<b>41</b>
(a) Staining	<b>41</b>
(i) The Gram stain	<b>41</b>
(a) Gram's crystal violet	<b>41</b>
(b) Gram's iodine	<b>42</b>
(c) Gram's safranin	<b>42</b>
(ii) Endospore staining	<b>42</b>
Motility medium	<b>42</b>

<b>9. Biochemical tests methods</b>	<b>43</b>
9.1 Aceton production	43
9.2 Blood haemolysis test	44
9.3 Carbohydrate fermentation	44
9.4 Catalase test	45
9.5 Cytochrome (oxidase test)	45
9.6 MacConkey agar	46
9.7 Nitrate reduction	46
9.8 Starch hydrolysis	47
9.9 Urease test	47
9.10 Oxidation of fermentation (OF) of glucose	48
9.11 Citrate utilization	48
9.12 Catalase activity	49
9.13 Gelatin hydrolysis	49
9.14 Hydrogen sulphide production	49
9.15 Potassium cyanide	49
9.16 Levan formation	49
9.17 Lipase enzyme production	50
9.18 Potassium hydroxide test	50
9.19 Methyl red (MR) reaction	50
9.20 Pectinolytic activity	50
9.21 Production of cellulase	51
<b>10. Characterization test reagent and methods</b>	<b>51</b>
10.1 Acid mercuric chloride	51
10.2 Lugol's iodine solution	51
10.3 Kovac's reagents	51

10.4 Hydrogen peroxide	52
10.5 Methyl red solution	52
10.6 $\alpha$ -Naphthol solution	52
10.7 Nitrate test reagents	52
10.8 Oxidase test reagents	52
<b>11. Qualitative screening test media, methods, and conditions (first survey)</b>	<b>53</b>
11.1 Pectinolytic enzyme production medium	53
11.2 Pectinase production and activity assay medium	53
<b>12. Qualitative screening test media, methods and conditions (second survey).</b>	<b>54</b>
12. (a) 1. Pectinolytic enzyme production medium (PEPM)	54
12. (b) Analysis of reducing sugar by Nelson's technique	55
Reagents	55
12. (b) 1. Nelson's reagent	55
12. (b) 2. Arsenomolybdate reagent	55
<b>13. Preparation of cell-free filtrate (CCF)</b>	<b>56</b>
<b>14. Parameters controlling the polygalacturonase productivities</b>	<b>56</b>
14.1 Different inoculum sizes	56
14.2 Different substrate concentration	56
14.3 Different incubation period	57
14.4 Different pH values	57
14.5 Different temperatures	57
14.6 Different nitrogen sources	58
14.7 Different carbon sources	58

14.8 Different amino acids	<b>58</b>
14.9 Different vitamin requirements	<b>59</b>
14.10 Different flask volumes	<b>59</b>
<b>15. Purification of polygalacturonase produced using potato peels under solid state fermentation (SSF) conditions</b>	<b>60</b>
15.1 Polygalacturonase production	<b>60</b>
15.2 Preparation of cell-free filtrate	<b>60</b>
15.3 Ammonium sulphate fractionation	<b>61</b>
15.4 Dialysis	<b>61</b>
15.5 Applying column chromatography technique	<b>62</b>
15.5 (a) Applying on sephadex G-200	<b>62</b>
15.6 Amino acid analytical data for the purified enzymes	<b>63</b>
<b>16. SDS-PAGE</b>	<b>63</b>
16.1 Detection of protein using silver stain	<b>64</b>
<b>17. Factors affecting the purified polygalacturonase activities</b>	<b>65</b>
17.1 Effect of incubation temperatures	<b>65</b>
17.2 Different substrate concentrations	<b>65</b>
17.3 Different pH values	<b>65</b>
17.4 Different enzyme concentrations	<b>65</b>
17.5 Heat stability	<b>66</b>
17.6 pH stability	<b>66</b>
<b>18. Factors affecting on the application of polygalacturonase in the clarification of juava juice.</b>	<b>66</b>

18.1 Substrate volumes.	<b>66</b>
18.2 Polygalacturonase concentrations.	<b>67</b>
18.3 Different temperatures.	<b>67</b>
18.4 pH values.	<b>67</b>
<b>Results (Section A)</b>	<b>68</b>
(A) Qualitative screening for selection of the most potent bacterial isolates pectinase producer (First survey by clearing zon technique)	<b>68</b>
(B) Qualitative screening for selection of the most potent bacterial isolates pectinase producer (second survey by clearing zone technique).	<b>68</b>
(C) Quantitative screening for selection of the most potent pectinase producer bacterial isolates (second survey by Nelson's technique, 1944).	<b>71</b>
(D) Identification of the three most potent bacterial isolates	<b>75</b>
<b>Section (B)</b>	
<b>Parameters controlling the polygalacturonase (PG) productivity</b>	<b>78</b>
1. Different inocula size	<b>78</b>
2. Different substrate concentrations	<b>78</b>
3. Different incubation periods	<b>81</b>
4. Different initial pH values.	<b>81</b>
5. Different incubation temperatures	<b>81</b>
6. Different nitrogen sources	<b>85</b>
7. Different carbon sources	<b>85</b>
8. Different amino acids	<b>85</b>

9. Different vitamins	90
10. Different flask volumes	90
<b>Section (B)</b>	<b>95</b>
<b>Purification of polygalacturonase produced by B. firmus-I-10104, allowed to grow on S. tuberosum (ST) peels under solid state fermentation conditions</b>	<b>95</b>
<b>Subsection (I): Purification of polygalacturonase</b>	<b>95</b>
Step (1): Enzyme production and preparation of CFF	96
Step (2): Fractional precipitation by ammonium sulphate	96
Step (3): Concentration by dialysation against sucrose	99
Step (4): Purification on sephadex G-200 gel column	99
Step (5): Amino acid analysis of the purified enzyme	99
Step (6): Gelectrophoresis analysis of the purified enzyme	99
<b>Amino acid analysis of the purified polygalacturonase</b>	<b>104</b>
<b>Subsection II</b>	<b>106</b>
Characterization of the purified enzyme produced by B. firmus-I-10104 at 37 °C	106
1. Effect of incubation temperature	106
2. Effect of different substrate concentrations on the purified enzyme activity	106
3. Effect of different pH values on the activity of purified enzyme	109
4. Different concentrations of the purified enzyme in relation to its activity	109
5. Thermostability of the purified enzyme	109
6. pH stability of the purified enzyme activity	109
<b>Section (C).</b>	<b>109</b>

<b>Biotechnological application of purified PG in clarification of Juava Juice</b>	<b>115</b>
<b>Applying of the purified PG in the clarification of Juava Juice using Nelson's technique</b>	<b>115</b>
1. Effect of different substrate (Juava Juice) concentrations in the clarification of Juava juice	<b>115</b>
2. Effect of different enzyme concentrations on the clarification of Juava Juice	<b>117</b>
3. Effect of different incubation temperatures on the clarification of Juava Juice	<b>117</b>
4. Effect of different pH values on the clarification of Juava juice	<b>117</b>
<b>Discussion</b>	<b>121</b>
<b>Summary</b>	<b>141</b>
<b>References</b>	<b>145</b>
<b>Arabic summary</b>	