

Reference

Andraus,M.H.;Bastos-De-Siqueira,MEP(1997): Determination and derivatization for the analysis of butorphanol by GC-MS in race horse urine.

Biomed-Chromatogr. (Dec.), 12(1):**P.** 34-37.

Aitkenhead-AR;Lin-ES;Achola-KJ(1988): Pharmacokinetics of oral and intravenous nalbuphine in healthy volunteers.

(British-Journal-of-Clinical Pharmacology);25(Feb); **P.** 264-268.

Badwan, A.A.; Nabusli, L.;Al-Omari, M.M.;Daraghmeh, N.and Ashour, M. (2001): Analytical profiles of drug substances and excipients,27: P. 339-377.

Bathory, G.;Sazko, E.;Magyer, K. and Deutsch, T. (1990): Properties of the melanin binding of p-bromomethyl amphetamine. Pol. J.pharmacol. pharm., **42**:P. 19-27.

Baumgartner, W.A.; Hill, V.A. and Bland, W. H. (1989): Hair analysis of drugs of abuse.

J. Forensic Sci., **34**:P. 1433-1453.

Borg,B.J.;Sitaram,B.R.;Taylor,D.A.(1993): Ion pair extraction and liquid chromatographic analysis of morphine in rat brain and plasma.

J.Chromatogr, Biomed Appl.; (Nov.), Vol. 132(2"621"); **P.165-172.**

Burtis,C.A.;Bowers,L.D.;Chattoraj,S.C.andUllman,M.D.(1987):

"Chromatography" In "Fudametals of clinical chemistry ".

3rd ed. (N.W. Tietz,ed.),W.B. Saunders company,Philadelphia,London.

p. 105-124.

Camachoa; Matthews,SC;Dimsdeli JE (2001): Invisible synthetic opiates and act psychosis.

New-England Journal of medicine; 345p.469.

Casarett,L.J.and Doull,J. (1996): "The basic science of poisons ".

5 th. ed. (Curtis D. klassen,ed.),McGraw-Hill. New York. **P. 89-186.**

Christiasen S (1987): Nalbuphine in patient controlled analgesia;comparison with morphine.

Current Theraputic-research; 41:**P. 933-945.**

Christensen, H.; Steentoft, A. and Worm, K. (1985): Muscle as an autopsy material for evaluation of fatal cases of drug overdose.

J. Forensic sci. soc., **25**: 191-206.

Clarke, E.G.C. (1986): "Isolation and identification of drugs".

London the pharmaceutical press, 2nd ed. **P.794-795.**

Clarke,E.G.C.(2004): "Analysis of drugs and poisons in pharmaceuticals,body fluids and postmortem materials".

London the pharmaceutical press, 3rd ed. **P.1312-1313.**

Chou,-JZ;Albeck,H.;Kreek,MJ(1993): Determination of nalmefene in plasma by HPLC with electrochemical detection.

J-Chromatogr. , Biomed Appl. (Apr.); Vol.:124(2"613"); **P. 359-364.**

Cone, E.J.;Darwin ,W.D. and Wang ,W. L. (1993): The occurrence of cocaine, heroin and metabolites in hair of drug abusers.

Forensic sci. Int., **63**:P. 55-68.

Davidson,A.F. ;Emm,T.A.;Pieniaszek,-H.J. ,J.R (1996):

Determination of naltrexone and its major metabolite in human plasma using liquid chromatography with electrochemical detection.

J.-Pharm-Biomed-Anal., (Sep.); 14(12); P. 1717-1725.

De-Cazanove, F.; Kinowski, J.M.; Audran, M.; Rochette, A.;

Bressolle, F. (1997): Determination of nalbuphine in human plasma by HPLC with electrochemical detection.

J.,Chromatogr.,Biomed Appl. (Mar.),Volume 690 no. (1-2); P. 203-218.

Dube – L.M.; Beaudoin, N.; Lalande, M.; Mc Gilveray,I.J. (1988) :

Determination of nalbuphine by HPLC with electrochemical detection.

J – chromatogr.; 427 (may 13); P. 113–120.

Fang , H ; Duan ,-H.;Zhou,T ; C,Chen , (1989) : Studies on the analysis of some narcotic analgesics and CNS stimulants.

Zhongguo – Yaoxue Zazhi, oct.; **24 (10):**P. 759-768.

Garriott, J.Cq. (1991): Skeletal muscle as an alternative specimen for alcohol and drugs analysis. J. Forensic sci., **36(1):**P. 60-69.

Giorgi, M.; Bertini, S.; Soldani, G. and Giusiani, M. (2000):

Comparison of HPLC and GC-MS methods for determination of embutamide (A component of tanax ® or T-61®) in biological specimens.

J. Anal. Toxicol. , **24**(5):P. 323-326.

Groenendaal,D.;Blom-Roosemalen,M.C.M.;Danhof,M.;De

Lange,E.C.M*(2005): HPLC of nalbuphine ,butorphanol and morphine in blood and brain.

J. Chromatogr.,B.,Biomed. , Aug., 822(1-2):P. 230-237.

Harreleson J.C. and Wong Y.J. (1988): Species variation in the disposition of nalbuphine and its acetyle salicylate ester.

anal.;Xenobiotica Nov.;**1(11)**:P.1239-47.

Henderson, G.L. (1993): Mechanisms of drug incorporation into hair.

Forensic Sci. Int., **63**:P. 19-29.

Ho,S.T., Wang, J.J.,Hu O.Y.P.,HO,(1991); J. Chromatogr.,Biomed

Appl.(Oct.);Vol. 108(2"570");**P. 339-350.**

Ho,S.T., Wang, J.J.,Hu O.Y.; HuT.M.(1995): Effect of aging on the pharmacokintice of nalbuphine in rabbits.

Biopharm, Drug Dispose; 16 (nov); P. 695 – 703.

HO, S.H.; Wang, J.J., Hu O.Y.; P.; Ching, P.S; Lee, S.C. (1996):
Determination of nalbuphine by HPLC with UV detector in human and rabbit.

J. Chromatogr.,Biomed Appl.:(April), 678(2), P. 289-296.

Chang-RK; Hussain M.A. (1998): Improvement of nalbuphine bioavailability through rectal administration of a prodrug.

Pharmaceutical development and technology; 3(1):p. 131-134.

Hussain MA; Aungst BJ; Maurin MB; WU Ls(1991): Injectable suspension for prolonged release nalbuphine.

Drug-development and Industrial-Pharmacy; 16(1); P.67-76

Hussain MA;Aungst-BJ;Shefter-E(1986): Buccal and oral bioavailability of nalbuphine in rats.

Journal of pharmaceutical Sciences; 75; P.218-219.

Hussar DA (1980): New drugs ("13 approved drugs from FDA").

Nurse-80; **P.24-32.**

Ito, S.;Kudo, K.; Lmamura, T.; Jitsufuchi, N. and Kimura,k.(1997): Detection of drugs and poisons in postmortem tissues –determination of paraquat in tissues of rabbits buried under ground.

Nippon Hoigaku Zasshi, **51(2):P. 83-88.**

Jacqz Aigrain E.,Debillon T.,Daound P.,Boithias C.,Hamon I.,Ray,Popon M.and Menotre F. (2003): Population pharmacokinetics of nalbuphine in neonates,paediatric and perinatal drug therapy .

J.Chromatogr. Biomed Vol. 5 No. 4(9) **P.190-198.**

Jailon p.; gardin – ME; Lecocq – B.; Richard – MO. (1989): Healthy volunteers and elderly patients.

Clin. – pharmacol. – ther.; 46 (aug);**P. 226 – 233.**

Jin young Kim, Moon kyo In,ki–jung paeng and Bong chul chung (2004): Simultaneous determination of nalbuphine and opiates in human hair by GC–MS. chromatographia vol. 591103–4– **P.219–226.**

Julien B; Demailly S; Inghels Y; Urbina MA (2005): Questioning the obstetrical use of nalbuphine.

ASHP –Midyear- Clinical meeting; 40, p-17d

Keegan, M; Kay, B. (1984): Detection of nalbuphine in plasma improved HPLC.

J. Chromatogr., biomed. 9 Nov.; 311:**P.** 223 – 226.

Kessler J M (1986): New drug profile"nalbuphine"

South African pharmaceutical journal; 53; **P.**36-37.

Kim, JY; IN, MK; Paeng; Chung, BC (2004): Simultaneous determination of nalbuphine and opiates in human hair by GC - MS.

Chromatographia; 59(3-4) : **P.**219-226

Kintz,P.;Tracqui A.;Mangin-P.(1992): Determination of nalbuphine using HPLC coupled to photodiode array detection and GC-MS Appl.

J. Chromatogr. Biomed Appl.; (Aug.7) **P.** 172-176.

Kuhnert, P; Erhard, P; Dixon.a; kuhert, b; Gross, T (1988):
determinationof ritodine in plasma using HPLC with internal standard
nalbuphine.

J-Liq.Chromatogr.; 6(14); **p.27775-2783**

Lake, CL; Difario, CA; Duckworth, EN; Moscick, JC; Engle, IS;
dusbin, CG (1982): HPLC analysis of plasma levels of nalbupine in
cardiosurgical patients.

J-Chromatographia, Biomed Appl.; 22(233): **P.410-416**

Leung, G.N.W.; Chung, E.W.; D.KK; Tang, FPW; WAN, TSM*,
YU, NH (2005).J-Chromatogr.,B. Anal. Technol. Biomed life sci ;(
15) Oct.; 825(1):P.47-56.

Liaw, WJ,HO,ST,Wang,J,J;HV,OYB;LIJA(1998): Determination
of morphine by HPLC with electrochemical detection.

J-chromatogr, Biomed; 714 (2):p.237-245.

Logan, B.K. and Stafford, D.T. (1989): Liquid /solid extraction on diatomaceous earth for drug analysis in postmortem blood.

J. Forensic Sci., 34(3):P. 553-564.

Lo, M.W., Juergens G.p and whitney JR (1984): Determination of nalbuphine in human plasma by automated HPLC with electrochemical detection.

J. Res commun chem. Pathol pharmacol. Jan; 43 (1):P. 159 – 68.

Lukas, G.; Brindle, S.D. and Greengard, P. (1971): The route of absorption of interperitoneally administered compounds.

J. Pharmacol. Exp. Ther., **178**:P. 562-566.

Martindale (1999): "The complete drug reference "32nd ed.

London the pharmaceutical press p. 60-61.

Miyazawara, N.; Uematsu, T.; Mizuno, A.; Nagashima, S. and Nakashima, M. (1991): Ofloxacin in human hair determined by higher performance liquid chromatography.

Forensic Sci. Int., **51**:P. 65-77.

Mule,S.J.(1971): Routine identification of drugs of abuse in human urine.

J. Chromatogr., **55**:P. 255-257.

Nakahara, Y. U. J.; Kikura, P. U. R. I. and Takahashi,k. (1995):
Effect of structural factors on incorporation of drugs into hair. The incorporation rates of amphetamine analogs.

Pro. Of TIAFT Int. Conference and workshop for hair analysis in Forensic Toxicology. **P.28-49.**

Nakamura, G.R.;Liu,Y. and Noguchi, T.T(1981): A method for the separation and determination of neutral compounds in postmortem tissues.

J.Anal. Toxicol., **5(4)**:P. 162-164.

Nickolls,L.C.(1956): "The scientific investigation of crime"

1st ed. London, Butterworth. **P.382-383.**

Nicolle,E;Michaut,S;SERRE-

Debeauvais,F;Bessard,G(1995):Rapid and sensitive HPLC assay for nalbuphine in plasma.

J. Chromatogr. Biomed, B. Appli. (Jan.) vol. 663(1); **P.111-117.**

Nicolle,E.;Veih,S.;Guimier,C;Bessard,G.,(1997): Modified method of nalbuphine determination in plasma ,validation and appl. to pharmacokinetics of the rectal route.

J.Chromatogr.,B :Biomed.(Mar.);690(1-2):**P.** 89-97.

Pao,LH;Hsiong,CH;HU,OYP;HO,ST(2000): HPLC method for the simultaneous determination of nalbuphine and its prodrug sebacyl dinalbuphine ester in dog plasma and application to pharmacokinetics studies in dogs.

J-Chromatographia, b: Biomed-appl.; 746(2):**P.**241-147.

Pounder, D.J.; Adams, E.; Fuke, c. and Longford, A.M. (1996): Variability of post mortem drug concentration in liver and lung.

J.Forensic Sci., **41(6)**:P. 927-932.

Püschel, K.; Thomasch,P. and Arnold, W.(1983): Opiates leveles in hair.

Forensic Sci. Int., **21**:P. 181-186.

Quarry, M.A.; Williams, RC; Sebastian, D.C. (1998); J.Liq. Chromatogr. Relat Technol. (Nov.); 21(18), **P. 2841-2852.**

Quarry,M.A.;Sebastian,D.S.;Williams,R.C.(1998): Determination of nalbuphine ,Methylparben and probyle parben.
J. Chromatogr.; (May); 47(9-10), **P.515-522.**

Quinones Torrelo, -C; Sagrado, S.; Villanueva-Camanas, RM; Medina-Hernandez, M.J. (2002): Opioid analgesics retention pharmacologic activity models using biopartioning micellular chromatography .
J., Chromatogr. B; Anal. Technol. Biomed life sci., (25Jan.) 766(2):**P.265-277.**

Smith,J.E.; Guly,H.R.(2004): Developed a nalbaphine and slow release morphine.
British medical journal ; **328** (7453);**P.1426.**

Spratt,e.(1977): Interfering compound in urine screening for drugs.
Excepra Medical, Drug Dependence, 5:2121.

Steinert,L. and Coffman, N.(1992): Techniques of drug analysis in Clinical laboratory medicine.

(Edited by Tilton, R.C.; Balows,A.; Hohnadel,D.C.and Reiss, R.F.),Mosby year Book,Inc.:p.346.

Stadnnyk A, Grossman R F (1986): Nalbuphine induced pulmonary oedema.

Chest 90:p.773:774.

Tek, D; Mellon, M (1987): The effectiveness of nalbuphine and hydroxyzine for the emergency treatment of severe headache.

Ann-Emerg.-Med. 16(3):P.308-313.

Traqui, A.; Kintz, P.and Mangin, P. (1995): Hair analysis a worthless tool for therapeutic compliance monitoring.

Forensic Sci. Int., 70: P.183-189.

Uematoso, T.; Sato, R.; Fujimori, O. and Nakashima, M. (1990): Human scalp hair as evidence of individual dosage history of haloperidol :a possible linkage of haloperidol excretion into hair with hair pigment .

Arch. Derm. Res., 282:P. 120-125.

Uematsu, t.; Sato, R.; Suzuki, K.; Yamaguchi, S. and Nakashima, M. (1989): Human scalp hair as evidence of individual dosage history of haloperidol: Method retrospective study.
Eur .J. Clin. Pharmacol., **37:P.239-244.**

Uematsu, M.; Nakano, M.; Akiyama, H. and Najkashima, M. (1992): Possible effect of pigment on the pharmacokinetics of ofloxacin and its excretion in hair.
J. Pharm. Sci., 81:P.45-48.

Wetzelsberger, N., Luckner, P.W.; Erking, W.: (1988): Internally standardized method for the determination of nalbuphine in human plasma by means of HPLC with electrochemical detection.
Arzneimittelforschung; Dec. (1988); 38 (II) (12), **P. 1768 – 1771.**

Wilson, S.J.; Errick, J.K.; Balkin, J. (1986): Pharmacokinetics of nalbuphine during parturition.
Am.J ,Obstet.–Gynecol ; 1986 ;155 (Aug):P. 340 – 344.

Wilson, J.F.; Green, s.; Dunastan, D.G.; Wicks, J.F.C.; Brewer, C. and Perfect, N. (1995): Relationship between methadonedose, hair methadone concentration and hair color in man.

Ther. Drug Monit., **17:P. 419-421.**

Wischnik A; Wetzelsberger N. and Luckner (1988): Elimination of nalbuphine in human milk.

Arzneimittelforschung Oct.; 38(10):P.1496-8.

Xu, -Y.; Fang , H, Xu, Y; Duan, H; Wu, Y(1991): Studies on the analysis anileridine, levorphanol, nalbuphine and ethamivan in urine.

Yaoxue-xuebao; 26(8); **P. 606-610.**

Yoo Y.C.,Chung H.S.,Kim I.S.,Jin W.T. and kim M.K. (1995): Determination of nalbuphine in drug abuser's urine.

J.Anal-Toxicol., (Mar.-April) (1995); Vol.19 (2); **P. 120-123.**