

CONTENTS

	Page
ACKNOWLEDGEMENT	i
I. INTRODUCTION	1
I-1- General introduction	1
I-2- High temperature superconductors	3
I-3- The bismuth strontium calcium copper oxide (BSCCO) system	5
I-4- The lanthanum copper oxide system	16
I-5- AIM OF THE WORK	27
II- Theories and methods of calculation	29
II-1- Electrical conductivity	29
II-2- Theory of superconductivity	30
II-3- Magnetic susceptibility	32
II-4- Rotating ring-disk electrode	38
III- EXPERIMENTAL	44
III-1- Preparation of samples	44
III-1-i- Materials	44
III-1-ii- Preparation of BSCCO samples	44
III-1-iii- Preparation of Lanthanum copper oxide compound	46
III-2- Methods of investigation	46
III-2-i- X-ray powder diffraction (XRD) patterns	46
III-2-ii- Thermal gravimetric analysis TGA	46
III-2-iii- Surface morphology	47
III-2-iv- Density measurements	47
III-2-v- Moisture effect	47
III-2-vi- Electrical conductivity measurements	48
III-2-vii- Magnetic measurements	50
III-2-viii- Preparation of La_2CuO_4 ring-disk electrode	51
IV- CHARACTERIZATION OF SAMPLES	54
IV-1- X-ray powder diffraction	54
(a) Sample EM1	54
(b) Sample EM2	57
(c) Sample EM3	57
(d) Sample EM4	62
(e) Sample EM5	62
(f) Sample EM6	62

To be continue

IV-2-	Thermal Gravimetric analysis (TGA)	68
IV-3-	Surface morphology	73
IV-4-	Density measurements	78
IV-5-	Moisture effect	80
V-	RESULTS AND DISCUSSION	81
	PART A	81
V-A-1-	Dc Electrical conductivity measurements	81
	(a) Sample EM1	81
	(b) Sample EM2	84
	(c) Sample EM3	84
	(d) Sample EM4	87
	(e) Sample EM5	87
	(f) Sample EM6	90
V-A-2-	DC magnetic measurements	95
	(a) Sample EM1	95
	(b) Sample EM2	98
	(c) Sample EM3	100
	(d) Sample EM4	100
	(e) Sample EM5	100
	(f) Sample EM6	100
V-A-3-	General Discussion	106
	PART (B)	110
V-B-1-	Cuprate surface reactivity to protons	110
V-B-2-	Coulometric efficiency and rate of $\text{La}_2\text{CuO}_{4+x}$ Formation	122
V-B-3-	General Discussion	131
	SUMMARY AND CONCLUSION	133
	REFERENCES	138
	ARABIC SUMMARY	

List of Figures

	Page
Figure (II-1): The effect of magnetic field on the superconductor.	37
Figure (II-2): Ring and disk current-potential curves in solution of A. (I) isolated disk; (II) isolated ring; (III) ring and disk.	42
Figure (III-1): Four-point probe measurement set-up.	49
Figure (III-2): Cross sectional view of ring-disk electrode.	53
Figure (IV-1): X-ray diffraction patterns of EM1 sample • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	55
Figure (IV-2): X-ray diffraction patterns of EM2 sample • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	58
Figure (IV-3): X-ray diffraction patterns of EM3 sample • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	60
Figure (IV-4): X-ray diffraction patterns of EM4 sample. • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	63
Figure (IV-5): X-ray diffraction patterns of EM5 sample. • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	65
Figure (IV-6): X-ray diffraction patterns of EM6 sample. • 2212 phase; ▽ 2223 phase; ■ 2201 phase.	67
Figure (IV-7): TG curves in air of samples: (a) EM1 (b) EM2 and (c) EM3	71
Figure (IV-8): TG curves in air of samples: (a) EM4 (b) EM5 and (c) EM6.....	72
Figure (IV-9): TG curves in nitrogen of samples: (a) EM1 (b) EM2 and (c) EM3	74
Figure (IV-10): TG curves in nitrogen of samples: (a) EM4 (b) EM5 and (c) EM6	75
Figure (IV-11): Surface morphology of the investigated samples....	76
Figure (V-A-1): Temperature dependence of the electrical resistance for sample EM1.....	83
Figure (V-A-2): Temperature dependence of the electrical resistance for sample EM2.....	85
Figure (V-A-3): Temperature dependence of the electrical resistance	

for sample EM3.....	86
Figure (V-A-4): Temperature dependence of the electrical resistance for sample EM4.....	89
Figure (V-A-5): Temperature dependence of the electrical resistance for sample EM5.....	91
Figure (V-A-6): Temperature dependence of the electrical resistance for sample EM6.....	92
Figure (V-A-7): The T_c -values verses the relative concentration of the superconductor phase (2212) for all the prepared samples; ■ $T_{c,H}$, • $T_{c,L}$, ▲ $T_{c,off}$	93
Figure (V-A-8): The T_c -values verses the relative concentration of the superconductor phase (2223) for all the prepared samples; ■ $T_{c,H}$, • $T_{c,L}$, ▲ $T_{c,off}$	94
Figure (V-A-9): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM1 in a 5 G field.....	97
Figure (V-A-10): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM2 in a 5 G field.....	99
Figure (V-A-11): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM3 in a 5 G field.....	101
Figure (V-A-12): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM4 in a 5 G field.....	102
Figure (V-A-13): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM5 in a 5 G field.....	103
Figure (V-A-14): The field-cooled (FC) and the zero field-cooled (ZFC) magnetic moment for sample EM6 in a 5 G field.....	104
Figure (V-B-1): Ring potential scans at 10 mV/s for a La_2CuO_4 disk-Au ring electrode rotated at 900 rpm in 0.1 mM H^+ /1.0M NaCl. (A) La_2CuO_4 disk lacquer coated, (B) La_2CuO_4 disk exposed, (C) La_2CuO_4 disk- Au ring electrode in 1.0 M NaCl.	114
Figure (V-B-2): $i_{R(\text{unsh})}-\omega^{1/2}$ plot for La_2CuO_4 at different $[\text{H}^+]$,	

marked as mM, in 1M NaCl; disk is held at open circuit and i_R measured at -0.56 V vs Ag/AgCl.	115
Figure (V-B-3): $i_{R(sh)}-\omega^{1/2}$ plot for La_2CuO_4 at different $[\text{H}^+]$, marked as mM, in 1M NaCl.	116
Figure (V-B-4): $i_{\text{RCu(II)}}-\omega^{1/2}$ plot for La_2CuO_4 at different $[\text{H}^+]$, marked as mM, in 1M NaCl; disk is held at open circuit and i_R measured at -1.1 V vs Ag/AgCl.	117
Figure (V-B-5): $\Delta i_R-i_{\text{RCu(II)}}$ plot for La_2CuO_4 at 1,2,3 mM $[\text{H}^+]$, in 1M NaCl.	119
Figure(V-B-6a): $i_R-\omega^{1/2}$ plot for La_2CuO_4 at different $[\text{H}^+]$, marked as mM, in 1M NaCl, i_R measured at -0.52 V vs Ag/AgCl.	120
Figure(V-B-6b): $i_R-\omega^{1/2} - [\text{H}^+]$ plot for La_2CuO_4	121
Figure (V-B-7): Potentiodynamic curve characterizing film formation of the superconductor in 1M KOH, scan rate 5 mV/sec. Electrode area = 0.178 cm^2	124
Figure (V-B-8): Three successive i_R-i_D curves for La_2CuO_4 disk-Au ring in 1M KOH. E_R , -1.2 V; ω , 900 rpm. Potential-controlled disk was scanned at 5 mV/sec. Arrows on curve indicate scan direction.	125
Figure (V-B-9): Galvanostatic data for La_2CuO_4 at $200 \mu\text{A}$, E_D , i_R - t plots were obtained in 1M KOH. The area under the curve is the missing O_2 reduction current (superconductor formation).	127
Figure (V-B-10): XPS spectra of La_2CuO_4 pellet (a) before oxidation and (b) after oxidation in 1M KOH solution.	130