CONTENTS

. INTRODUCTION	•••
I. REVIEW OF LITERATURE	•••
1. Seasonal abundance of predatory spiders	
2. Seasonal abundance of S. littoralis (Boisd.),	A.
craccivora (Koch.), A. gossypii (Glover) and T. urtic	cae
(Koch.)	.
3. Biological studies	
4. Toxicity of some alternative pesticides on predat	ory
spiders	
5. Effect of some alternative pesticides on total protein a	and
protein bands	
I. MATERIAL AND METHODS	
1.Seasonal abundance of the spiders and their preys dur	ing
successive seasons of cotton and broad bean plants	
A. Estimation of spider populations	• • • •
B. Estimation of <i>S. littoralis</i> populations	
C. Estimation of Aphids populations: Aphis craccive	ora
in broad bean plants and Aphis gossypii in cotton plan	nts
D. Estimation of phytophagous mite populations	
2. Biological studies	
A. Spider's rearing	
B. Spider's food	
C. Prey's rearing.	
3. Toxicological of alternative pesticides	
4. Biochemical studies	.
1.Tissue preparation	

2. Determination of total protein	
3. Separation of protein bands by electrophoresis (using	ing
SDS-PAGE)	
5. Statistical analysis	
IV. RESULTS AND DISCUSSION	••••
1.Ecological studies	.
1.1. Occurrence of spider families collected from bro	oad
bean and cotton crops	
1.1.1.Occurrence of spiders families in Qaha Station.	• • • • •
1.1.2.Occurrence of spiders families in Seds Station	
1.2. The percentage of spider families and occurrence	e of
their species their families during two successive year	S
1.2.1. In Qaha Station (2003/2004)	.
1.2.2. In Qaha Station (2004/2005)	· • • • •
1.2.3. In Seds Station (2003/2004)	
1.2.4. In Seds Station (2004/2005)	
Seasonal abundance	
1.3.1.Population density of predatory spider and the	heir
preys (Spodoptera littoralis, Aphis craccivora	and
Tetranychus urticae) which collected from broad b	ean
plants at Qaha Station during two successive ye	ears
(2003-2004) and (2004-2005)	
1.3.2. Population density of predatory spider and the	heir
preys (Spodoptera littoralis, Aphis gossypii	and
Tetranychus urticae) which collected from cotton pla	ants
at Qaha Station during two successive years (2004) a	and
(2005)	
1.3.3. Population density of predatory spider and the	heir

	preys (Spodoptera littoralis, Aphis craccivora and
	Tetranychus urticae) which collected from broad bean
	plants at Seds Station during two successive years (2003-
	2004) and (2004-2005)
	1.3.4. Population density of predatory spider and their
	preys (Spodoptera littoralis, Aphis gossypii and
	Tetranychus urticae) which collected from cotton plants
	at Seds Station during two successive years (2004) and
	(2005))
1.4.	Association of spider families and preys on broad bean
and	cotton plants during two successive years (2003-2004)
and	(2004-2005) at two stations (Qaha and Seds)
	1.4.1. Association of spider families and preys on broad
	bean and cotton plants during (2003-2004) at Qaha
	Station
	a) Broad bean
	b) Cotton
	1.4.2. Association of spider families and preys on broad
	bean and cotton plants during (2004-2005) at Qaha
	Station
	a) Broad bean
	b) Cotton
	1.4.3. Association of spider families and preys on broad
	bean and cotton plants during (2004-2005) at Kaha
	Station
	a) Broad bean
	b) Cotton
	1.4.4. Association of spider families and prevs on broad

bean and cotton plants during (2004-2005) at Seds
Station
a) Broad bean
b) Cotton
1.5. Average number of spiders' predators and their families
from broad bean and cotton plants during two successive
years (2003-2004) and (2004-2205) at Kaha and Seds Stations
2. Biological studies
3. Toxicity of some alternative pesticides on adults (females
and male)
4.Biochemical study to investigate the effect of nutrition and
alternative pesticide on total protein and protein bands in adult
spiders
4.1. Total protein content.
4.2. Refraction of protein patterns
4.2.1. The effect of treated the adult of (T. abini) with
alternative pesticides on its protein bands molecular
weight
4.2.2. Molecular weight of protein bands for adult (T.
abini) which feeding on different prey
V. SUMMARY
VI. REFERENCES
ARABIC SUMMARY