

REFERENCES

- Abbott, W. S. (1925):** A method of computing the effectiveness of an insecticide. J. Econ. Entomol., 18: 265-267.
- Abdel-Rahman, S. L.; A.A.Ibrahim; H. Abdel-Fattah and S. M. Abdel-Tawab (2001):** Biological studies on the true spider, *Thanastus albini* (Audouin) (Fam.: Philodromidae) Predator on two- different prey species. Biol. Control Conf. Cairo, 1-12 Dec.
- Abo-El-Ela, R. G.; N. M. Helmy; O. M. El-Mpnairy and H. Sahah (1995):** Effect of certain plant extracts on some biochemical aspects of the housefly larvae, *Musca domestica* (Diptera : Muscidae). Bull. Ent. Soc. Egypt, 22 : 17-25.
- Abo-El-Ghar, G. E. S.; M. S. Khalil and T. M. Eid (1995):** Some biochemical effects of plant extracts in the black cutworm, *Agrotis ipsilon* (Hufnagel). Bull. Ent. Soc. Egypt, Econ. Ser., 22: 85.
- Abo-El-Ghar, G. E. S.; M. S. Khalil and T. M. Eid (1996):** Effects of some plant extracts on development and fecundity of *Agrotis ipsilon* (Lepidoptera : Noctuidae). Bull. Ent. Soc. Egypt, Econ. Ser., 21: 171-190.
- Abou-Zeid, A. E. (1998):** Physiological and toxicological effect of certain plant extracts on *Ostrinia nubilalis* (Hubner). M. Sc. Thesis, Fac. Sci., Ain-Shams Univ., Cairo, Egypt.
- Agnew, C. W.; A. D. Dean and J. J. W. Smith (1985):** Spiders collected from peanuts and nonagricultural habitats in the Texas west crops timbers the south western Nat., 30: 1-12.
- Azab, A. K.; M. F. S. Tawfik and I. I. I. Ismail (1965):** Seasonal changes in the abundance of certain aphids and their predators in Giza. Bull. Soc. Ent. Egypt, 44: 11-24.

- Bahgawan, C. N.; K. D. Reddy and K. Sukumar (1992):** Annona-induced growth anomalies and protein depletion in red cotton bug, *Dysdercus keoenigii*. Indian J. Exp. Biol., 30: 10, 908-912.
- Cai, S. H. (1987):** Studies on the occurrence of *Tetranychus cinnabarinus* and its control target. China Cotton, 5: 39-40.
- Chatzaki, M.; A. Trichas; G. Markakis; M. Mylonas and P. A. Seldon (1998):** Seasonal activity of the ground spider fauna in a Mediterranean ecosystem (MTY Ouchtas, Grete, Greece). Proceeding of the 7th European Colloquium of Arachnol, Edinburgh, 235-243.
- Chitra, R. C. and T. S. V. Reddy (2000):** Effect of *Annoma squamosa* L. seed extract on protein metabolism of *Spodoptera litura*. Fab. Insect Environ. 6:1, 39-40.
- Cividanes, F.; M. J. Tromozine and L. H. Silvestre (1990):** Distribution of *Tetranychus urticae* (Koch). (Acari –Tetranychidae. Cientifica, 15 (2): 147-155.
- Corbitt, T. S.; A. S. J. Green and D. T. Wright (1989):** Relative potency of abamectin against larval stages of *Spodoptera littoralis* (Boisd.) (Lepidoptera : Noctuidae). Crop Protection, 8: 2, 127-132.
- Costa, F. G. (1995):** Ecologia and daily activity of the sand spiders *Allocosa* spp. (Araneae: Lycosidae) in Narinida, a Coastal Village of South Uruguay. Rev. Bras. Bio. 55: 457-466.
- Davis, B. J. (1964):** Disc electrophoresis. 2. Method and application to human serum proteins. Ann. N. Y. Acad. Sci., 121: 404-427.
- Dean, D. A.; M. Nyffeles and G. Breene (1987):** Foraging by selected spider predators on the cotton leafhopper and other prey. South Western Entomol., 12 (3): 263-270.

- Deltshev, C. D. (1997):** Spider (Araneae) from the Coastal habitats of Shabla-Extracts, Lake, Bulgaria (Black sea Coast). *Acta Zoologica Bulgarica*, 49: 58-63.
- Devasahayam, S.; M. A. Koyak and P. P. Reddy (1998):** IPM in species-challenges for the future. *Env. Implications*, India, 157-164.
- Dong, C. X. and C. E. Xu (1980):** Spiders in cotton fields and their protection and utilization. *China Cotton*, 3: 45-47.
- Dunin, P. M. and A. A. Mamedov (1992):** Spiders of cotton fields in the southern eastern part of Azerbaijan byulleten Moskovskogo Obshchestva Ispytatelei Prirody, *Biologichskii* 97 (6): 53-61.
- El-Beltagy, A. M.; A. M. Salem; M. A. El-Hamaky; R. Abu-Aiana and M. E. Omar (1993):** Cotton aphid, jassid and whitefly population dynamic under pheromone and insecticide applications against pink bollworm. *Egypt, J. Biol. Pest. Con.*, 13 (1): 63-70.
- El-Bermawy, S. M. and H. M. Abdel Fattah (2000):** Changes in protein electrophoretic pattern of *Tribolium confusum* 4th instar larvae after treatment with volatile plant oil (VETIVER). *J. Egypt Ger. Soc. Zool.*, 31 (E): 167-182.
- El-Bokl, M. M.; M. S. Hamed and H. M. Moawad (1998):** Effect of sub lethal dosage of plant extracts and cypermethrin on certain biochemical aspects of the first filial generation of *Culex pipiens* larvae. *J. Egypt Ger. Soc. Zol.*, 16 (E): 45-55.
- El-Defrawi, G. M.; A. K. Emam; I. A. Marzouk and I. Rizkallah (2000):** Population dynamics and seasonal distribution of *Aphis craccivora* Koch and associated natural enemies in relation to virus disease incidence in faba bean fields. *J. Agric. Res., Egypt*, 78 (2): 627-641.

- El-Erksousy, M. H. (2000):** Studies on some ture spiders in Egypt. Ph.D. Thesis, Fac. Agric. Al-Azhar Univ., 130 Pp.
- El-Erksousy, M. H. M. and M. M.H. Fawzy (2001):** Biological studies on the true spiedr *Thanatus albini* (AuDouin) (F: Philodromidae) on wheat *Aphis schizophis graminum* (RonDani). Anal. Agric. Sci., Moshtohor, 39 (1): 645-649.
- El-Erksousy, M. H.; G. M. Mosa and W. O. Gomaa (2002):** The spider *Theridion egyptium* Fawzy and Elerksousy as a biological control agent on cotton aphid, *Aphis gossypii* (Glover). The 2nd Int.Conf. of Plant Prot. Res. Institute, Cairo, Egypt, 21-24 Dec., 2002; 24-25.
- El-Heneidy, A. H.; A. A. Ibrahim; Y. H. Fayad and G. M. Moawad (1996):** Survey and population dynamics of common true spiders in Egyptian cotton field. Annals of Agricultural Science, Moshtohor, 34 (3): 1177-1187.
- El-Hennawy, H. K. (1986):** A list of Egyptian spider genera. Serket, 1 (1): 12-14.
- El-Hennawy, H. K. (1992):** Distribution of spider genera in Egypt (Arachnidae : Araneidae). Serket, 3 (1): 1-32.
- El-Hennawy, H. K. (2000):** Spiders of Sinai (Egypt), a list of species (Arachnidae : Araneidae). Serket, 8 (1): 29-34.
- El-Hennawy, H. K. (2002):** List of Egyptian spider (revised in 2002) Serket, 8 (2): 73-83.
- El-Nagar, S.; M. M. Megahed and A. H. Amin (1978):** The aphid fauna of wild plant in Giza, Egypt. Bull. Soc. Ent. Egypt, 62 : 219.
- El-Saadany, G. B.; A. Shoukry; T. A. Abu-Hashish and S. M. M. Hassan (1987):** The population dynamics and seasonal abundance of *S. littoralis* (Boisd.). 2nd Nat. Conf. of Pests & Diseases of Vegetables and Fruits, Ismailia, October, 1978-: 77-92.

- El-Sheikh, M. A. and S. El-Nagar (1994):** Aphid fauna and seasonal incidence on potato of Egypt. 60:389-394.
- Finney, D. J. (1971):** Probit analysis. Cambridge Univ. Press Cambridge, 33.
- Flatz, U. (1988):** Abundance, seasonal dynamics and diversity of ground spider in montance meadows near Innsbruck (North Tyrol, Austria). Ber. Nathurwiss. Med. Ver. Innsbruck, 75: 125-141.
- Foelix, R. F. (1982):** Biology of spider. Harvard Univ. Press, London, England, 306 pp.
- Furr, R. E. and T. R. Pfrimmer (1968):** Effects of early, mid and late season infestation of two spotted spider mites on the yield of cotton. J. Econ. Ent., 62 (5): 1446-1447.
- Gabboiur, s. I.; M. M. Hussein and H. K. El-Hennawy (1996):** Spider populations associated with different crop in Menoufyia Governorate, Nile Delta, Egypt. J. Agric. Res.
- Galal, S. A. (1989):** Ecological studies on some insect pests infesting certain Legume crops, in Qalubiya Governorate. Ph. D. Thesis, Fac. Agric. Cairo Univ.
- Gormal, A. G.; C. Bardawill and M. M. I. Davis (1949):** Determination of serum protein by means of biuret reaction. J. Biol. Chem., 177: 751-766.
- Guo, J. R.; Y. X. Chen and Z. D. Fang (1989):** Relationship between aphids and incidence of IUTEO virus infecting *Vicia faba*. J. Nanjing Agric. Univ., 12 (4): 53-57.
- Hamid, S. M. A.; A. Shah and A. M. Anwar (1977):** Some ecological and behavioral studies on *Aphis craccivora* Koch (Hemiptera : Aphididae). Tech. Bull. Common Wealth, 18: 99-111.

- Hassan, H. (2002):** Biological and biochemical studies on the effect of some botanical extracts on cotton leafworm, *S. littoralis* (Boisd.) Lepidoptera: Noctuidae. M. Sc. Thesis, Fac. Science, Ain-Shams Univ.
- Hassanein, M. H.; A. H. El-Sebae; F. M. Khalil and E. A. Moustafh (1971):** Studies on *Aphis gossypii* and *Aphis craccivora* in the cotton field in Upper Egypt. Bull. Soc. Ent. Egypt, 55: 163-169.
- Hassanein, S. S. M. (1994):** Effect of some crop management practices on populations of certain insects infesting broad bean plants at Khattara region, Egypt. Zagazig J. Agric. Res., 21 (6): 1807-1816.
- Helal, H. A.; R. M. Salem; A. S. El-Khouly; M. M. Metwally and A. B. El-Mezaïen (1996):** Population dynamics of *Aphis craccivora* (Koch) and *Empoasca* spp. on faba bean in relation to associated predators and some climatic factors. J. Agric. Res., Egypt, 75 (2): 461-471.
- Hightower, B. G. and D. F. Martin (1965):** Ecological studies of spider mites attacking cotton in central Texas. J. Econ. Entomol. 49 (3): 424-425.
- Huffker, C. B.; R. I. Rabb and J. A. Logan (1977):** Some aspects of population dynamics relative to augmentation of natural enemy action. Pages 3-38 in R. L. Ridgway and S. B. Vinsondes. Biol. Control By Augmentation of natural enemies, Plenum press, 480 pp.
- Hussein, A. M. (1999):** Seasonal abundance and daily activity pattern of spider fauna in some vegetable crops in Menuofiya Governorate, Egypt. J. Agric. Res., 77 (2): 677-689.
- Hussein, A. M.; H. K.El-Hennawy and A. A. Sayed (1998):** Biodiversity of spiders (Araneae, in the western desert of Egypt in

relation to agriculture and reclamation. Bull. Fac. Agric. Cairo Univ., 49: 597-610.

- Ibrahim, A. M. A.; K. T. Awadala and M. D. Abdallah (1992):** Seasonal abundance of the aphid, *Myzua persica* Sulz and *Aphis gossypii* Glov. On apple trees and their aphidivorus association. Egypt, J. Biol. Pest. Control, 2 (1): 1-4.
- Joon, P. N. K. and M. C. Lee (1988):** Spiders from the southern region of MZ in Kangwondo Lorean Arachnology, 4: 15-33.
- Kabir, A. R. M. F. (1978):** Pest of grain legumes and their control in Bangladesh. Cf. Pest of Grain Legumes Ecology and Control., 31-36.
- Kaston, B. J. (1978):** How to know the spiders. WC. Brwon Co., Dubugue, Iowa, USA, 272 pp.
- Kim, H. J. (1990):** Studies on the faunal and seasonal dynamics of spiders in the crown of Korean pine, *Pinus koraiensis*. Lorian Arachnology, 5: 177-193.
- Kuroli, G.; I. Nemeth and L. Numeth (1988):** Aphid damage to field bean in relathion to population dynamics and ecological conditions. Mededlingen-Van-de-Faculteit-Landbouw-Wetenschappen, Rijksuniversiteit-Gent, 53 (3a): 1195-1201.
- Levy, G. (1977):** The Philodromidae spiders (Araneae, Philodromidae). 1st J. Zool., 26 : 193-229.
- Li, D. and R. R. Jackson (1996):** How temperature affects development and reproduction in spiders: a review. J. Thermal Biology, 21 (4): 245-274.
- Li, X. Z; X. S. Leng and H. S. Wang (1983):** Species of spiders and their population of fluctuations in peanut field. Natural Enemies of insects Kunchongtiandi, 5 (2); 112-115.

- Lockley, T. C.; J. W. Smith; W. P. Scott and C. R. Parenica (1979):** Population fluctuations of two groups of spiders from selected cotton field in Panola and Pontotoc countries, Mississippi. The Southwestern Entomol., 4: 20-24.
- Mackie, D. W. (1978):** How to begin the study of spiders. Br. Arachnol. Soc., 12 pp.
- Mansour, F. (1987):** Spiders in sprayed and unsprayed cotton field in Israel, their interactions with cotton pest and their importance as predators of the Egyptian cotton leaf worm, *Spodoptera littoralis*. Phytoparasitica, 15: 1, 31-41.
- Mansour, F. and U. Heiewbach (1993):** Evaluation of lycosid, micryphantis and linyphiid spiders as predators of *Rhopalosiphum padi* (Hom. ; Aphididae). Entomophaga, 38 (1): 79-87.
- Mansour, M. M. and W. Nentwing (1988):** Effect of agrochemical residues on four spider taxa : laboratory methods for pesticides tests with web-building spiders. Phytoparasitica, 16: 317-326. *Sharkia Governorate. Proc. 2nd Arab Pesticide Conf. Tanta Univ.*
- Mansour, M. M.; A. M. Metwally and H. E. Metwally (1977):** Seasonal fluctuation of *Aphis craccivora* Koch on cowpea plants in Sharkia Governorate. Proc. 2nd Arab Pesticide Conf. Tanta Univ.
- Metwally, A. M.; M. H. Mowafi and M. A. Mohafez (2002):** Seasonal abundance of spider Families inhabiting cultivation at Sohag Governorate. Al-Azhar J. Agric. Res. 36: 137 pp.
- Metwally, E. M.; S. S. Hassanein and A. F. E. Aafsah (1995):** Sesaonal population activity of sucking pests infesting common bean and squash plants at Gemmeza region, Gharbia Governorate, Egypt. J. Agric. Res., Egypt, 73 (3): 693-667.
- Mohafez, M. A. M. (2000):** Studies on true spiders in Sohag

- Governorate. M. Sc. Thesis, Fac. Agric., Al-Azhar Univ., 115 pp.
- Mohamed, M. I. and S. F. Hafez (2000):** Biological and biochemical effect of non-volatile oil (jojoba) against *Culex pipiens* Diptera : Culicidae. J. Union Arab Biol., Cairo, (13): 65-78.
- Mohamed, Z. M. A. (1984):** Studies on aphids in Sharkia region. Ph. D. Thesis, Zagazig, Univ.
- Mostafa, S. a. (1993):** Biochemical effects of some chemical compounds on *Spodoptera littoralis* (Boisd.). Ph. D. Thesis, fac. Agric. Al-Azhar Univ.
- Naser, E. A.; M. R. Tucker and D. G. Compion (1984):** Distribution of moths of the Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.) (Lepidoptera : Noctuidae) in the Nile Delta interpreted from catches in a pheromone trap network in relation to meteorological actors. Bull. Entomol. Res., 74 (3); 487-494.
- Nassef, M. A.; A. A. El-Feshawi and W. M. Watson (1996):** Seasonal abundance of certain predators as influenced by cotton planting dates and weather factors. J. Agric. Res. Egypt, 74 (3): 585-597.
- Negm, A. M.; M. F. Abou-Ghadir and A. G. A. Salman (1976):** Population trend and time of activity of certain true spiders collected from clover foliage at Assuit. Agric. Res. Rev., 54: 87-91.
- Nyffeler, M. and G. Benz (1988):** Feeding ecology and predatory importance of wolf spiders (*Pardosa* spp.) (Araneae, Lycosidae) in winger wheat fields. J. Appl. Entomol., d 106: 2, 123-134.
- Pang, D. Y. and X. M. Liang (1992):** Safety study of pesticides on spiders predatory natural enemies of pests, in paddy fields. Human Agric. Cole., 3: 234-239.
- Pekar, S. (1999):** Foraging mode a factor affecting the susceptibility of spiders (Araneae) to insecticide applications. Pesticide Sce., 55

(11): 1077-1082.

Plagens, M. J. (1983): Population of *Misaumenops* (Araneida : Thomisidae) in two Arizona cotton fields. Environ. Entomol., 12: 572-575.

Prentice, T. R.; J. C. Burger; W. R. Lcenogle and R. A. Redak (1998): Spiders from diegan coastal sage scrub (Arachnida : Araneae). Pan Pacific Ent., 74: 181-202.

Putman, W. L. (1967): Life histories and habits of two species *Philodromus* (Araneida : Thomisidae) in Ontario. Can. Entomol., 99: 622-631.

Qu, H. Z.; Y. L. Huang and R. X. Wu (1986): Population dynamics of spiders in cotton filed and their protection and utilization. Natural Enemies of Insects, 8 (3); 142-145.

Rahil, A.A.H.(1988): Ecological and biological studies on the spiders at Fayoum. M. Sc. Thesis, Fac. Agric., Cairo, Univ., 133 pp.

Rao, P. J.; K. M. Kumar; S. Singh and B. Aubrahmanyam (1999): Effect of *Artemisia annus* on development and reproduction of *Dysdercuss koenigii* F. (Hem., Pyrrhocoridae). J. Appl. Entomol., 123 (5): 315-318.

Reddy, K. D.; C. N. Bhagawam and K. Sukumar (1993): Depletion of amino acids and protein in the nymphal haemolymph of the red cotton bug, *Dysdercus koenigii* (Fab.) Hemiptera : Pyrrhocoridae) treated with *Annoma squamosa* extract. Phytophaga 5 (1): 45-50.

Rod, M. and M. Kenprestum (1984): Spiders of the world. Facts on life publication, New York, Oxford England, 189 pp.

Salah, H. A. (1993): Effect of some botanical extracts *Musca domestica*. M. Sc. Thesis, Fac. Science, Zagazig Univ. (Benha branch).

Saleh, M. R. A.; M. H. Hassanein and M. El-Sebae (1972): Population

dynamics of *Aphis craccivora* Koch on broad bean and cowpea in Upper Egypt (Homoptera : Aphididae). Bull. Soc. Ent., Egypt, 65 : 135-138.

Sallam, Gihan, M. (1996): Studies on true spiders in Giza Governorate. M. Sc. Thesis, Fac. Agric. Cairo Univ. 139 pp.

Sallam, Gihan M. E. (2002): Surveyed and ecological studies on spiders in four Governorates of Egypt. Ph. D. Thesis, Fac. Agric. Cairo Univ., 144 pp.

Schmidt, G. H.; H. Remblid; A. A. I. Ahmed and M. Breuer (1998): Effect of *Melia azedarach* fruit extract on juvenile hormone titer and protein content in the haemolymph of two species of noctuid : Lepidoptera larva (Insecta : Lepidoptera : Noctuidae). Phytoparasitica, 26: 4, 283-292.

Selim, A. A.; S. A. El-Refai and A. El-Gantiry (1987): Seasonal fluctuations in the population of *Aphis craccivora* Koch, *Myzus persica* (Sulz.), *Aphis gossypii* (Glov.) and their parasites. Ann. Agric. Sci., Ain-Shams Univ., 32 (3): 1837-1848.

Semeada, A. M.; H. Isamail and Shahinaz A. Abd El-Salam (1993): *Aphis gossypii* (Glover) infesting maize plants a new record to Egypt. Ent. Soc. Egypt, 71: 41-45.

Sengonca, C. and W.Klein (1988): Prey spectrum and feeding activity in the laboratory of the Orb-Web spider, *Araniella opistographa* (Kleucx.) and the crab spider, *Philodromus cespitum* (Walck.) which are frequent in apple orchards. Zeitswchrift-fur-Angewandte-Zoo., 75:1, 43-54.

Sharma, R. P. and R. P. Yadav (1994): Population dynamics of bean aphid *Aphis craccivora* Koch and its predatory coccinellid complex in relation to crop type (lentil, lathyrus and faba bean) and weather

conditions. J. Entomol. Res., 18 (1); 25-36.

Shaurub, E. H.; Z. A. Ahmed and S. E. M. El-Naggar (1998): Impacts of pyriproxyfen and extract of *Schinus terebinthifolius* Raddi on development, reproduction and reproductive organism, *Spodoptera littoralis* (Boisd.) Lepidoptera: Noctuidae. J. Egypt Ger. Soc. Zool., 27 (E): 57-82.

Soliman (2003): Studies on true spiders associated with some vegetable crops. M. Sc. Thesis, Fac. Agric. Menufiya Univ., 113 pp.

Swidan M. H. (1994): Antifeedant activity of 24 plant extracts against larvae of *Spodoptera littoralis* (Boisd.) Lepidoptera : Noctuidae. Alex. J. Agric. Res., 39: 3, 363-374.

Taha, H. A. S.; Z. R. Sawires and A. M. Mostafa (1990): Population dynamics of the two spotted spider mite, *Tetranychus arabicus* (Attiah) on two cotton varieties. Agric. Res., Cairo, 68 (1): 5-10.

Whiteford, F.; W. Showers and G. Edwards (1987): Insecticide tolerance of ground and foliage dwelling spiders (Araneae) in European corn borer (Lepidoptera: Pyralidae) action sites. Env. Entomol., 16 (3): 779-785.

Wright, D. J.; A. Loy; A. S. J. Green and R. A. Dybas (1985): The translaminar activity of Abamectin against mites and aphids. Mededeling-Van-de-Facultiet-Rijksuniversiteit-Gent, 50: 26, 595-601.

Young, O. P. (1989): Predation by *Psecusaurinamira* (Araneae : Pisauridae) on *Lygus lineolaris* (Heteroptera : Miridae) and other arthropods. J. Archol., 17: 43-48.

Zedan, Z. H.; H. A. Abdel-Rahman; W. M. El-Deeb; M. S. Salama and E. M. Radwan (2002): Effect of sub lethal concentration of Abamectin and S-Fenvalerate on protein of the spiny bollworm,

Earias insulana (Boisd.). The First Conf. of the Central Agric. Pesticide Lab., 3., 5 Sept. 2002.

Zhao, J. (1984): Species, distribution and population fluctuations predacious spiders in cotton field in China. Natural Enemies of Insects, 6 (1): 1-12.