

VI. REFERENCES

- Abbas, M. (1992).** Comparative rates of infestation by four scale insects on citrus trees with special reference to rates of parasitism on the purple scale *Lepidosaphes beckii*. Egypt, J. Agric. Res., 70 (2): 477-486.
- Abd-Rabou, S. (1997a).** Parasitoids attacking some species of scale insects (Homoptera: Coccoidea: Diaspididae) in Egypt. Proceeding of the First Scientific Conference of Agricultural Science, Vol.II : 727-736.
- Abd-Rabou, S. (1997b).** Parasitoids attacking the olive scale insect, *Parlatoria oleae*(Colvee) (Homoptera:Coccoidea:Diaspididae) in Egypt.Proceeding of the First Scientific Conference of Agricultural Science,Vol.II:719-726.
- Abd-Rabou, S. (1999).** Parasitoids attacking the Egyptian species of armored scale insects (Homoptera: Diaspididae). Egypt. J. Agric. Res., 77 (3): 1113-1129.
- Abd-Rabou,S. (2000).** *Aphytis chrysomohali*(Hymenoptera:Aphelinidae) as an effective parasitoids of armored scale insects in Egypt.(27th October),Seventh Arab Congress of Plant Protection.p.460.
- Abdel-Fattah, M. and G. El-Saadany (1979).** The role of parasitoids in the control of the purple scale, *Lepidosaphes beckii* (Newman) in Egypt. Zeitschrift für Angewandte Entomologie, 87 (2): 154-159.

- Alam, S. (1956).** The taxonomy of some British apelinid parasites (Hymenoptera) of scale insects (Coccoidea). Trans. Roy. Entomol. Soc., London, 108: 357-384.
- Alexandrakis, V. (1990).** Effect of *Dacus* control sprays, by air or ground, on the ecology of *Aspidiotus nerii* Bouché (Hom. Diaspididae). Acta Horticulturae, No. 286: 339-342.
- Alexandrakis, V. and C. Benassy (1981).** Experiment in biological control on olive in Crete using *Aphytis melinus* DeBach (Hym. Aphelinidae), a parasite of *Aspidiotus nerii* Bouché (Hom. Diaspididae). Acta Oecologica, 2 (1): 13-25.
- Applebaum, S. and D. Rosen (1964).** Ecological studies on the olive scale, *Parlatoria oleae*, in Israel. Jor. Econ. Entomol., 57 (6): 847-850.
- Asfoor, M. (1997).** Seasonal abundance and control of the plum scale insect *Parlatoria oleae* (Colvee) on some deciduous trees. Ph.D. Thesis, Fac. Of Agric., Moshtohor, Benha Branch, Zagazig Univ., 399 pp.
- Atkinson, P. (1983).** Environmental factors associated with fluctuations in numbers of natural enemies of a population of citrus red scale, *Aonidiella aurantii* (Maskell) (Hemiptera: Diaspididae). Bull.Ent. Res., 73 (3): 417-426.

- Avidov, Z. (1970).** Biology of natural enemies of citrus scale insects and the development of methods for their mass production. Rehovot, Israel, Hebrew Univ. of Jerusalem, Fac. of Agric., 8: 247 pp.
- Bartlett, B. and T. Fisher (1950).** Laboratory propagation of *Aphytis chrysomphali* for release of control California red scale. *Jor. Econ. Entomol.*, 43 (6): 802-806.
- Battaglia, D. and G. Viggiani (1982).** Observations on the distribution and phenology of *Aonidiella aurantii* (Maskell) (Homoptera-Diaspididae) and on its natural enemies in Campania. *Annali della Facolta di Scienze Agrarie della Universita degli Studi di Napoli, Portici*, 16 (2): 125-132.
- Benassy, C. (1967).** Note sur *Parlatoria oleae* au Maroc (Hom. Coccoidae). *Annals Soc. Ent. Fr. (N.S.)*, No. 3: 165-171.
- Benassy, C. and H. Bianchi (1983).** On the development of populations of *Lepidosaphes beckii* Newman in the presence of its specific parasite, *Aphytis lepidosaphes* Compere. 38 (2): 119-124.
- Benassy, C. and G. Euverte (1967).** Notes on *Chrysomphalus dictyospermi* in Morocco. *Awamia*, 24: 95-111.
- Benassy, C. and F. Soria (1964).** Ecological observations on the Diaspine Coccids injurious to citrus in Tunisia. *Annals Inst. Natn. Rech. Agron.*, 37: 193-221.

- Benassy, C., H. Bianchi, and P. Brun (1981).** Preliminary data on the definition of a treatment threshold in *Lepidosaphes beckii* New. in standardization of biotechnical methods of integrated pest control in citrus orchards .Luxembourg, Commission de Communautés Europeennes, 19-25.
- Carrero, J. (1983).** Current status of biological control of citrus scale insects in Valencia (Spain). *Fruits*, 35 (10): 625-631.
- Casas, J., R. Nisbet, S. Swarbrick, and W. Murdoch (2000).** Eggload dynamics and oviposition rate in a wild population of a parasitic wasp. *Journal of Animal Ecology*, 69 (2): 185-193.
- Clancy, D. and M. Muma (1959).** Purple scale parasites found in Florida. *J. Econ. Ent.*, 52: 1025-1026.
- Clancy, D.; A. Selhime and M. Muma. (1963).** Establishment of *Aphytis holoxanthus* as a parasite of Florida red scale in Florida. *J. Econ. Ent.*, 56: 603.
- Compere, H. (1955).** A systematic study of the genus *Aphytis* Howard (Hymenoptera: Aphelinidae) with descriptions of new species. *Univ Calif. Publ. Entomol.*, 10: 271-319.
- Crouzel, I. and E. Merluzzi. (1980).** Biological observations on the parasite *Aphytis* near *lingnanensis* Compere (Hym. Aphelinidae) and its host *Unaspis citri* (Komstock) (Hom. Diaspididae). *Revista de la Sociedad Entomologica Argentina*, 39: 89-100.

- Das, S. (1988).** Studies on *Aphytis* sp. *chrysomphali* (Mercet)-a parasite of black scale, *Chrysomphalus aonidum* (= *C. ficus*) Ashm. Two and a bud, 35 (1-2): 44-45.
- Dean, H. (1955).** Factors affecting biological control of scale insects on Texas citrus. Jor. Econ. Entomol., 48 (1): 444-447.
- Dean, H. (1961).** *Aphytis lepidosaphes* (Hymenoptera: Chalcidoidea), an introduced parasite of purple scale. Ann. Entomol. Soci. America, 54: 918-920.
- Dean, H. (1982).** Reduced pest status of the Florida red scale on Texas citrus associated with *Aphytis holoxanthus*. Jor. Econ. Entomol., 75 (1): 147-149.
- DeBach, P. (1959).** New species and strains of *Aphytis* (Hymenoptera: Eulophidae) parasitic on the California red scale, *Aonidiella aurantii* (Mask.) in the orient. Ann. Entomol. Soci. America, 52: 354-361.
- DeBach, P. (1962).** Species of *Aphytis* for possible use in the biological control of *Chrysomphalus dictyospermi*. FAO Plant, Prot. Bull., 10 (2): 29.
- DeBach, P. (1977).** A newly imported California red scale parasite from Saudi Arabia. Calif. Agric., 31 (12): 6-7.

DeBach, P. (1979). *Aphytis riyadhi* n. sp. (Hymenoptera: Aphelinidae), a parasite of *Aonidiella* sp. (Homoptera: Diaspididae). *Entomophaga*, 12 (4): 325-342.

DeBach, F. and L. C. Argyriou (1967). The colonization and success in Greece of some imported *Aphytis* sp. (Hym. Aphelinidae) parasitic on citrus scale insects (Hom. Diaspididae). *Entomophaga*, 12 (4): 325-342.

DeBach, P. and D. Rosen (1976). Twenty new species of *Aphytis* (Hymenoptera: Aphelinidae) with notes and new combinations. *Ann. Entomol. Soci. America*, 69 (3): 541-546.

Dekle, G. (1976). Florida armored insects. Florida Dep. Agric. And Consumer Serv. Div. Con. Comm., 345 pp.

De Santis, L. (1937). Un himenoptero, parasito de la cochinilla roja de los citrus, nuevo para la fauna argentina (*Aphytis chrysomphali*, G. Mercet). *Rev. Fac. Agron. La Plata*, No. 20: 262-599.

De Santis, L. (1948). Estudio monografico de los aphelinidos de la Republica Argentina (Hymenoptera: Chalcidoidea). *Rev. Mus. La Plata. Zool.* 5: 23-280.

De Santis, L. and I. Crouzel (1994). Species of *Aphytis* occurring in the Neotropical region and their role in biological control. p. 257-77 in: Rosen, D. (Editor). *Advances in the study of Aphytis*. Intercept Ltd.

Doutt, R. L. (1954). An evaluation of some natural enemies of the olive scale. J. Econ. Entomol., 47: 39-43.

Doutt, R. and R. Smith (1974). The pesticide syndrome-Diagnosis and suggested Prophylaxis. p. 3-15 in: Huffaker, C. B. (Editor). Biological Control. Proceedings of an AAAS Symposium, P.S.H.

El-Minshawy, A. and O. Osman (1974). Biological and ecological studies on the masked scale insect *Mycetaspis personata* (Comstock) in the Alexandria area (Coccoidae: Diaspididae). Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri", Portici, 31: 152-170.

Erol, T. and B. Yasar (1999). Investigations on natural enemies, some biological characteristics, and population fluctuations of *Lepidosaphes ulmi* (L.) (Homoptera, Diaspididae) and *Palaeolecanium bituberculatum* (Targ. And Tozz.) (Homoptera, Coccoidae) harmful to apple trees in Van Province. Turkish Journal of Agriculture and Forestry. 23 (2): 151-164.

Ferriere, C. (1965). Hymenoptera Aphelinidae d'Europe et du Bassin Méditerranéen. Faune de l'Europe et du Bassin Méditerranéen. 1. Masson, Paris, 206 p.

Flanders, S. (1951). Mass culture of California red scale and its golden chalcid parasites. Hilgardia, 21 (1): 1-42.

Flanders, S. (1957). Hymenopterous parasites of three species of oriental scale insects. Bull. Lab. Zool. Portici, 33: 10-28.

Gahan, A. (1928). On some Chalcidoid scale parasites from Java. Bull. Ent. Res., 18: 149-54.

Gerson, U. (1967). The natural enemies of the chaff scale *Parlatoria pergandii* Comstock in Israel. Hebrew University of Jerusalem, Rehovot., 21-53.

Gulamahamad, H. and P. DeBach (1976). Biological control of the San Jose scale *Quadraspidiotu perniciosus* (Comstock) (Homoptera: Diaspididae) in Southern California. Hilgardia, 46 (7): 204-238.

Gürkan, S. (1983). Investigations on the bioecology of *Pseudaulacaspis pentagona* Targ., a pest of peach and mulberry in the Marmara Region. Istanbul, Turkey., 22 (4) [Tr., Fr., 13 ref., 3 fig.].

Hafez, M. (1984). Systematic survey of certain parasites of scale insects and mealy in Egypt. Ph.D. Thesis, Faculty of Agriculture, University of Alexandria., 233 p.

Hafez, M. (1988). Population fluctuations on parasites of California red scale, *Aonidiella aurantii* (Mask) (Homo. Diaspididae) in Alexandria. J. App. Ent., 106 (2): 183-187.

Hafez, M., A. El-Minshawy, and A. Donia (1987a). Population fluctuations of parasites of *Lepidosaphes beckii* Newm. and

Ceroplastes floridensis Comst. Anz. Schadlingskde,
Pflanzenschutz, Ummeltschutz, 60: 6-9.

Hafez, M., A. El-Minshawy, and A. Donia (1987b). Parasitic efficiency of some hymenopterous *Aphytis* sp. on the purple scale insect *Lepidosaphes beckii* Newm. Journal of Applied Entomology, 103 (2): 135-138.

Hare, J. and R. Luck (1991). Indirect effects of citrus cultivars on life history parameters of a parasitic wasp. Ecology, 72 (5): 1576-1585.

Hayat, M. (1983). The genera of Aphelinidae of the world. Systematic Entomology, 8: 63-102.

Hayat, M. (1994). The *Aphytis* fauna of the Oriental region. Advances in the study of *Aphytis* (Hymenoptera: Aphelinidae), 303-315.in: Rosen, D. (ed).Advances in the study of *Aphytis*. Intercept Ltd.

Heimpel, G. and J. Rosenheim (1998). Egg limitation in parasitoids: a review of the evidence and a case study. Biological control, 11 (2): 160-168.

Howard, L. (1881). Report on the parasites of the Coccoidae in the collection of this department. Report of the Entomologist, Part III, in: Ann. Rep. Comissioner Agric. Washington, D.C., p. 350-371.

Howard, L. (1900). A new genus of Aphelinidae from Chile. Canad. Entomol., 32: 167-168.

- Hsiao, S. (1983).** Natural enemies and population fluctuation of the palm scale, *Hemiberlesia cyanophilli* Signoret. Chinese Journal of Entomology, 1 (2): 69-76.
- Huffaker, C., C. Kennett, and G. Finney (1962).** Biological control of olive scale, *Parlatoria oleae* (Colvée), in California by imported *Aphytis maculicornis* (Masi) (Hymenoptera: Aphelinidae). Hilgardia, 32: 541-636.
- Imamkuliev, A. (1969).** The olive scale *Parlatoria oleae* Colvée (Homoptera, Coccoidea) in the conditions of Lenkoran (Azerbaijan). Ent. Obozr, 470-476.
- Io, C. (1938).** Ridescrizione dell' *Aspidiotus destructor* Sign. (Homoptera-Coccoidae). Boll. Lab. Zool. Portici, 30: 240-249.
- Javier, P., A. Havron, R. Morallo, and D. Rosen (1991).** Selection for pesticide resistance in *Aphytis* III. Male selection. Entomologia Experimentalis et Applicata, 61 (3): 237-245.
- Jiang, H. (1985).** Observations on the bionomics of *Pseudaulacaspis pentagona* and its natural enemies. Insect Knowledge (Kunchongzhishi), 23 (1): 19-20.
- Karaka, I. and N. Uygun (1990).** Natural enemies of *Aonidiella aurantii* (Maskell) (Homoptera, Diaspididae) in East Mediterranean citrus areas, and their population development on different citrus varieties. Ege Universitesi, 97-108.

- Kennett, C. (1967).** Biological control of olive scale *Parlatoria oleae* (Colv.) in deciduous fruit orchards in California. *Entomophaga*, 12: 461-474.
- Kfir, R. and H. Podoler (1983).** Effect of temperature and parasite density on three species of *Aphytis* (Hymenoptera: Aphelinidae), parasitising California red scale. *Researches on Population Ecology*, 25 (1): 69-80.
- Kiriukhin, G. (1946).** Les parasites du *Chrysomphalus dictyospermi* Morg. au nord de l'Iran. *Lab. Entomol. Phytopathol. Appl. Teheran*, 2: 4-6, 13-22 (in Persian, with a summary in French).
- Le Baron, W. (1870).** The chalcideous parasite of the apple-tree bark-louse (*Chalcis* [*Aphelinus*] *mytilaspidis*, n. sp.) *Amer. Entomol. Bot.*, 2: 360-362.
- Lenteren, J. and P. DeBach (1981).** Host discrimination in three ectoparasites (*Aphytis coheni*, *A. lingnarensis*, and *A. melinus*) of the oleander scale (*Aspidiotus nerii*). *Netherlands Journal of Zoology*, 31 (3): 504-532.
- LePage, H. (1942).** A "cochonilha branea" das amoreiras, *Pseudaulacaspis pentagona* (Targ. Tozz.) [The mulberry scale, *Aulacaspis pentagona*] *Biologie*, 8: 105-109.
- Liotta, G. (1983a).** Relation between *Aphytis chilensis* (How.) and *Aspidiotiphagus citrinus* (Craw.) (Hym. Aphelinidae) on the host *Aspidiotus nerii* Bouché (Hom. Diaspididae). *Instituto di*

Entomologia Agraria dell' Università di Palermo, Sicily, Italy, 35 (11): 695-699.

Liotta, G. (1983b). The role of aphelinids in the natural control of *Parlatoria pergandii* Comstock (Hom-Diaspididae) in Sicily. Istituto di Entomologia Agraria dell' Università di Palermo, Sicily, Italy, 66: 553-561.

Liotta, G., S. Burgio, and M. Montagna (1985). Parasitic biocoenosis of *Aspidiotus nerii* Bouche (Hom. Diaspididae) in Sicily, Palermo, Italy. Accademia Nazionale Italiano di Entomologia, 833-840.

Lu, Y. (1989). Studies on experimental population life table of citrus red scale (Homoptera: Diaspididae) and the role of parasites in the control of a natural population. Guangdong Entomological Institute, China, 207-217.

Luck, R. and H. Podoler (1985). Competitive exclusion of *Aphytis lingnanensis* by *A. melinus*: potential role of host size. Ecology, 66 (3): 904-913.

Malipatil, M., K. Dunn, and D. Smith (2000). An Illustrated Guide to the Parasitic Wasps Associated with Citrus Scale Insects and Mealy Bugs in Australia. Natural Resources and Environment, Agriculture Resources Conservation Land Management, Agriculture, Victoria. 152 p.

Masi, L. (1911). Contribuzioni alla conoscenza dei Calcididi Italiani. Boll. Lab. Zool. Gen. Agr. Portici, 5: 140-171.

- McClain, D., G. Rock, and J. Woolley (1990).** Influence of trap color and San Jose scale (Homoptera: Diaspididae) pheromone on sticky trap catches of 10 aphelinid parasitoids (Homoptera). *Environmental Entomology*, 19 (4): 926-931.
- Mercet, R. (1911).** Los Calcididos parasitos de Coccidos. *Bol. R. Soc. Espan. Hist. Nat.*, 11: 507-515.
- Mercet, R. (1912).** Un parasito del "poll-roig". *Bol. R. Soc. Espan. Hist. Nat.*, 12: 135-140.
- Mercet, R. (1928).** La fauna afelinina espanola. *Asoc. Españ. Prog. Cienc., Congr. (Cadiz, 1927)*, 4: 235-243.
- Miller, D., and M. Kosztarab (1979).** Recent advances in the study of scale insects. *Annual Review of Entomology*, 24: 1-27.
- Moreno, D., and R. Luck (1992).** Augmentative releases of *Aphytis melinus* (Hymenoptera: Aphelinidae) to suppress California red scale (Homoptera: Diaspididae) in southern California lemon orchards. *Jor. Econ. Entomol.* 85 (4): 1112-1119.
- Moursi, K., and E. Hegazi (1983).** The olive-tree scale, *Leucaspis riccae* Targ. (Hom.; Diaspididae) as a key pest of olive tree in dry farm system in the Egyptian western desert. *Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri"*, 40: 119-124.

- Moursi, K., and H. Mesbah (1985).** Olive pests of irrigated farm systems in the Egyptian western desert with special references to armored scale insects. *Annals of Agricultural Science, Moshtohor*, 23 (2) 901-911.
- Muma, H. (1971).** Parasitism of armored scale insects in commercial Florida citrus groves. *Fla. Entomol.*, 54 (2): 139-150.
- Murdoch, W., R. Luck, S. Swarbrick, S. Walde, D. Yu, and J. Reeve. (1995).** Regulation of an insect population under biological control. *Ecology*, 76 (1): 206-217.
- Onillon, J. (1988).** Biological and integrated control in citrus groves in the Mediterranean region. *Sta. Zool. et Lutte Biologique*, 33 (4): 481-494 [Fr, en, 59 ref.].
- Orphanides, G. (1984).** Competitive displacement between *Aphytis* sp. (Hym. Aphelinidae) parasites of the California red scale in Cyprus. *Entomophaga*, 29 (3): 275-281.
- Pickett, A. (1965).** The influence of spray programmes on the fauna of apple orchards in Nova Scotia XIV supplement to II oystershell scale, *Lepidosaphes ulmi* (L.). *Can. Ent.*, 97 (8): 816-821.
- Podoler, H. (1981).** Effects of variable temperature on responses of *Aphytis melinus* and *A. lingnanensis* to host density. *Phytoparasitica*, 9 (3): 179-190.

- Priesner, H., and M. Hosny (1940).** Notes on parasitic and predators of Coccidae and Aleurodidae in Egypt. Bull. Soc. Fouad Entomol., 31: 58-70.
- Prinsloo, G., and O. Naser (1994).** The *Aphytis* fauna of the Afrotropical region. pp. 280-302 in: Rosen, D. (ed). Advances in the study of *Aphytis*. Intercept Ltd.
- Quednau, F. (1964).** A contribution on the genus *Aphytis* Howard in South Africa (Hymenoptera: Aphelinidae). J. Entomol. Soc. S. Afr., 27: 107-116.
- Rodrigo, E., P. Troncho, and F. Garcia (1996).** Parasitoids (Hym.: Aphelinidae) of three scale insects (Hom.: Diaspididae) in a citrus grove in Valencia, Spain. Entomophaga. 41 (1): 77-94.
- Rosen, D. (1988).** Parasitic Hymenoptera in biological control: The genus *Aphytis*. p. 411-416 in: Gupta (Editor). Advances in parasitic hymenoptera research. Intercept Ltd.
- Rosen, D. (1994).** Advances in the study of *Aphytis*. Intercept Ltd., 320.
- Rosen, D., and P. DeBach (1974).** Biosystematic studies on the species of *Aphytis* (Hymenoptera: Aphelinidae). Bull. SROP, 3: 21-23.
- Rosen, D., and P. DeBach (1979).** Species of *Aphytis* of the world. The Hague, Boston, and London. Dr. W. Junk Br. Publ., 801 p.

- Salama, H. (1970).** Ecological studies on the scale insect, *Chrysomphalus dictyospermi* (Morgan) in Egypt. *Zeitschrift für Angewandte Entomologie*, 65 (4): 427-430.
- Samways, M. (1988).** Comparative monitoring of red scale *Aonidiella aurantii* (Mask.) (Hom., Diaspididae) and its *Aphytis* sp. (Hym., Aphelinidae) parasitoids. *J. App. Entomolo.*, 105 (5): 483-489.
- Savastano, L. (1931).** Della biancarossa (*Chrysomphalus dictyospermi* Morg.) negli agrumie in altre specie ospitanti nell' Italia. *Studi di Frtopatologia arbora. Ann. Staz. Agrum. Frutt. Acireale.*, 1-7.
- Selhime, A., M. Muma, W. Simanton, and C. McCoy (1969).** Control of Florida red scale in Florida with the parasite *Aphytis holoxanthus*. *Jor. Econ. Entomol.*, 62 (4): 954-955.
- Shutova, N., A. Antaze, M. Murvanidze, T. Timofeeva (1985).** Biological control of pests of citrus crops. *Zashchita Rastenii*, 2: 37.
- Siscaro, G., S. Longo, and S. Lizzio (1999).** Role of *Aonidiella aurantii* (Maskell) (Homoptera, Diaspididae) natural enemies in Sicilian citrus groves. *Phytophaga (Palermo)* 9. (supplemento)., 41-52.
- Souza, S. (1988).** Population fluctuations of *Mytilococcus beckii* (Newman, 1896) (Homoptera, Diaspididae) in Pera, Valencia, and Mandarina orange orchards in Cuaiba, Rio Grande do Sul. *Arquivos da Universidade Federal Rural do Rio de Janeiro*, 11 (1-2): 15-18.

- Steinberg, S., H. Podoler, and D. Rosen (1986).** Biological control of the Florida red scale, *Chrysomphalus aonidum* in Israel by two parasite species: current status in the coastal plain. *Phytoparasitica*, 14 (3): 199-204.
- Swaillem, S. (1974).** On the seasonal occurrence of *Lepidosaphes tapleyi* Williams (Homoptera-Hemiptera: Diaspididae). *Bulletin de la Société Entomologique d'Egypte*, 57: 67-72.
- Temarak, S. (1981).** Historical records of parasitoids in Egypt. Assiut University, Technical Bulletin, 1: 50.
- Titayavan, M., and D. Davis (1988).** Studies of a uniparental form of *Aphytis vandenboshi* (Hymenoptera: Aphelinidae), a parasite of the San Jose scale in northern Utah. *Great Basin Naturalist*, 48 (3): 388-393.
- Traboulsi, R. (1969).** Contribution à l'étude des *Aphytis Howard* du Liban (Hym. Chalcidoidea, Aphelinidae). *Ann. Soc. Entomol. Fr. (N.S.)*, 5: 5-72.
- Traboulsi, R., and C. Benassy (1965).** The coccids on citrus in the Lebanon and their natural parasites. *Magon Publ. Inst. Rech. Agron. Liban*, 5: 13.
- Troncho, P., E. Rodrigo, and F. Garcia (1992).** Observations on the parasitism of the diaspid *Aonidiella aurantii* (Maskell), *Lepidosaphes beckii* (Newman), and *Parlatoria pergandii*

(Comstock) in an orange grove. Boletín de Sanidad Vegetal, Plagas, 18 (1): 11-30.

Viggiani, G. (1994). The Aphytis fauna of the West Palaearctic region, with notes on their role in biological control. pp. 335-347 in: Rosen, D. (ed). Advances in the study of Aphytis. Intercept Ltd.

Viggiani, G., and F. Iannaccone (1973). Observations on the biology and on the parasites of the Diaspini *Chrysomphalus dictyospermi* (Morg.) and *Lepidosaphes beckii* (Newm.) carried out in Campania in the three-year period 1969-1971. Bollettino del Laboratorio di Entomologia Agraria "Filippo Silvestri", Portici, 30: 104-116.

Xu, M., and Q. Min. (1982). Studies on *Aphytis chrysomphali* (Mercet) (Hym.: Aphelinidae), a parasite of *Unaspis yanonensis* Kuwana (Hom.: Diaspididae). Natural Enemies of Insects, 4 (4): 19-21.

Yasnosh, V. (1994). *Aphytis* species occurring in the former U.S.S.R. and their role in biological control. pp. 317-333 in: Rosen, D. (ed). Advances in the study of Aphytis. Intercept Ltd.

Zhang, X., and D. Gu (1995). Study on the biology of three parasites of *Aonidiella aurantii*. Chinese Journal of Biological Control, 11 (2): 60-63.