

REFERENCES

- Abd El-Aziez, E.M. (1988): "Biostratigraphy of Upper Cretaceous-Lower Tertiary boundary from wells in northern Western Desert, Egypt": M.Sc. Fac. of Sci., Tanta Univ.
- Abd El-Kireem, M.R. and Abdou, H.F. (1979): Upper Cretaceous-Lower Tertiary planktonic foraminifera from South Galala plateau, Eastern Desert, Egypt". *Revista Espanola de Micropl.*, 11, (2), p. 175-222, pl. 1-12. Madrid.
- Abd El-Malik, W.M.; Bassiouni, M.A. and Obeid, F.L. (1974): "Biostratigraphy of Upper Cretaceous-Lower Tertiary rocks from West Central Sinai". 6th Coll. African Micropal. Tunisia, p. 181-215, 9 pls.
- Abd El-Razik, T.M. (1967): "Stratigraphy of the sedimentary cover of Anz-Atshan-south Duwi district". Bull. of the Faculty of Science No. 41, Cairo University.
- Abd El Razik, T.M. (1970): "Geological and microfacies studies on the Upper Cretaceous-Lower Eocene formations in East Rena Area, Eastern Desert, U.A.R.". Ph.D. Thesis, Cairo Univ, Cairo.
- Abd El-Razik, T.M. (1972): "Comparative studies on the Upper Cretaceous-Early Paleogene sediments on the Red Sea Coast, Nile Valley and Western Desert, Egypt". 8th Arab. Petr. Congr. paper No. 71 (B-3), S, 1-23, Algiers
- Abdou, H.F.; Naim, E.M. & Abdel Kireem, M.R. (1969): Planktonic foraminiferal zonation of the Upper Cretaceous-Lower Tertiary rocks of Gebel Ghanima, Kharga Oasis, Western Desert, UAR". *Buletin of the Faculty of Science, Alexandria University* 9, 251-270, Alexandria.
- Abu Khadrah, A.M.; Darwish, M.; El Azabi, M.H. & Abdel Fattah, M.A. (1987): "Lithostratigraphy of the Upper Cretaceous-Tertiary succession in the Gulf of Suez (Southern Galala Plateau), Egypt". In: Matheis & Schandelmeier, H.

- (eds.): Current Research in African Earth Sciences, 171-176; Rotterdam (Balkema).
- Akmal, M. (1985): "Sedimentological and Stratigraphical studies on the Upper Cretaceous-Lower Tertiary succession near Qena-Egypt. M.Sc. Thesis. F. Sci. Tanta University.
- Almogi-Labin, A.; Flexer, A.; Honigstein, A.; Rosenfeld, A. & Rosenthal, E. (1990): "Biostratigraphy and tectonically controlled sedimentation of the Maastrichtian in Israel and adjacent countries". *Revista Espanola de Paleontologia* 5, 41-52.
- Almogi-Labin, A.; Eshet, Y.; Honigstein, A.; Moshkovitz, S. & Rosenfeld, A. (1991): "Detailed biostratigraphy of the Santonian/Campanian boundary interval in Northern Israel". *J. micropaleontol.*, 10(1): p. 39-50.
- Alsharhan & Kendall (1991): "Cretaceous chronostratigraphic unconformities and eustatic sea level changes in the sediments of Abu Dhabi, United Arab Emirates". *Cret. Res.* (12), 379-401.
- Anan, H.S. (1979): "Stratigraphical and micropaleontological study on some Lower Tertiary rocks in Egypt". Ph.D. Fac. of Sci. Ain Shams Univ.
- Anan, H.S. & Hewaidy, A. (1986): "Biostratigraphy and distribution of the Paleocene benthonic Foraminifera in the Nile Valley Facies of Egypt". *M.E.R.C. Ain Shams Univ., Sci. Res., Ser.*, 6, 1-32.
- Ansary, S.E. & Tewfik, N.M. (1966): "Planktonic Foraminifera and some new benthonic species from the surface Upper Cretaceous of Ezz El Orban area, Gulf of Suez". *Jour. Geol. U.A.R.*, 10(1), p. 37-76.
- Ardenbol, J. & Berggren, W.A. (1978): "A new Paleogene numerical time scale. In: Cohee, G.; Glaessner, M. and Hedberg, H.D., Eds., Contributions to the geologic time scale". *American Association of Petroleum Geologists Studies in Geology* No. 6, 213-234.
- Aref, M.A. (1976): "Micropaleontological and biostratigraphical studies on some Late Cretaceous-Early Tertiary sections in the Eastern Desert, Nile Valley and Western Desert". M.Sc. Thesis, Assiut Univ., Egypt.

- Aubry, M.P.; Berggren, W.A.; Kent, D.V.; Flynn, J.J.; Klitgord, K.D.; Obradovich, J.D. & Prothero, D.R. (1988): "Paleogene geochronology; an integrated approach". *Paleoceanography*, 3, 707-742.
- Balbiez, F. (1955): "The Genus *Globotruncana* in Tunisia". *Micropal.* 1, p. 161-171.
- Ball, J. (1900): "Kharga oasis-its topography and geology". Egypt, Surv. Dept., Cairo, pp. 19.
- Ball, J. (1916): "The geography and geology of West-Central Sinai". Egypt Surv. Dept., 219, p. Cairo.
- Ball, J. (1939): "Contributions to geography of Egypt". Cairo. Geological Survey of Egypt.
- Bandy, O.L. (1951): "Upper Cretaceous Foraminifera from the carlsbad area", San Diego county, California: *Jour. Paleontology*, V. 25, p. 488-513, 2 fig., pl. 72-75.
- Bandy, O.L. (1963): "Larger living foraminifera on the continental borderland of southern California". *Contr. Cushman Found. Foramin. Res.*, 14, pp. 121-126, New York.
- Bandy, O.L. (1964): "Cenozoic planktonic foraminiferal zonation. *Micropaleont.*" New York, 10: 1-17.
- Bandy, O.L. (1967): "Cretaceous planktonic foraminiferal zonation". *Micropaleontology*, 13(1), 1-31.
- Bandy, O.L. & Arnal, R.E. (1960): "Concepts of foraminiferal paleoecology". *Bull. Am. Assoc. Petrol.*, Vol. 44(12). p. 1921-1932.
- Banner, F.T. (1989): "The nature of *Globanomalina* Haque". *J. Foramin. Res.*, 19, 171-179.
- Banner, E.T. & Blow, W.H. (1959): "The classification and stratigraphical distribution of the Globigerinacea". *Paleont. London*, 21-27, 3 pls.

- Banner, F.T. & Blow, W.H. (1960): "Some primary types of species belonging to the superfamily Globigerinacea": *Cushman Found. Foram. Research, Contrib.*, V. 11, p. 1-41, pl. 1-8.
- Barakat, M.G. & El-Dawoody, A.S. (1973): "A microfacies study of the Upper Cretaceous-Paleocene-Lower Eocene sediments at Duwi and Gurnah sections, Southern Egypt". *Ann. Inst. Geol. Publ. Hung. Budapest*, 1973, 391-414, 6 pls.
- Barakat, M.G. & El-Dawoody, A.S. (1973): "Nannobiostratigraphy of Upper Paleocene-Lower Eocene in Duwi Range Quseir District, Egypt". 8th Arab Petrol. Cong. Vol. 3, part 2, paper No. 70, (B-3), p. 1-43, 6 pls.
- Barazi, N. (1985): "Sedimentologie und Stratigraphie des Abyad-Beckens (NW Sudan)". *Berliner Geowissenschaftliche Abhandlungen*, A 64, 1-85.
- Barazi, N. & Kuss, J. (1987): "Southernmost outcrops of marine Lower Tertiary carbonate rocks in NE-Africa (Gebel Abyad, Sudan)". *Geologische Rundschau*, 76(2), 529-537.
- Barr, F.I. (1972): "Cretaceous biostratigraphy and planktonic foraminifera of Libya". *Microplaeontology*, 18(1), 1-46.
- Barron, I. (1907): "The topography and geology of the peninsula of Sinai (western portion)". *Egypt Surv. Dept.*, 241 p., Cairo.
- Barron, T. & Hume, W. (1902): "Topography and Geology of the Eastern Desert of Egypt (central portion)" *Egypt. Surv. Dep.* 331 pp., Cairo.
- Barthel, K.W. & Hermann-Degen, W. (1981): "Late Cretaceous and Early Tertiary Stratigraphy in the Great Sand Sea and its SE Margins (Farafra and Dakhla Oasis), SW Desert, Egypt". *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie*, 21, 141-182, München.
- Bassiouni, M.A.; Abdel Malik, W.M. and Boukhary, M.A. (1974): "Eocene planktonic foraminiferal zones from Abu Zeneima area, East Coast of Gulf of Suez, Egypt".

- Bassiouni, M.A. and Luger, P. (1990): "Maastrichtian to Early Eocene Ostracoda from southern Egypt. Paleontology, Palaeoecology, Paleobiogeography and Biostratigraphy". Berliner geowiss. Abh., A, 120.2, 755-928, 16 figs. 14 tabs, 25 pls. Berlin.
- Beadnell, H.J. (1905): "The topography and geology of the Fayium province of Egypt". Surv. Dept. Cairo, 101 pp.
- Beadnell, H.J. (1924): "Report on the geology of the Red Sea coast between Quseir and Wadi Ranga". Petroleum Research Bulletin No. 13. Survey department, Egypt.
- Beadnell, H.J. (1927): The wilderness of Sinai-London: Arnol.
- Beckmann, J.P.; El Heiny, I.; Kerdany, M.T.; Said, R. & Viotti, L. (1969): "Standard planktonic zones in Egypt". Proceedings of the First International Conference on Planktonic Microfossils, Geneva (1967), Vol. 1, 92-103, Geneva.
- Bellier, J.P. (1983): "Foraminifères planctoniques du Crétacé de Tunisie septentrionale: Systematique, biozonation, utilisation stratigraphique de l'Albien au Maastrichtien". Thèse de Doctorat D'Etat, Mémoires des Sciences de la Terre, Université Curie, Paris, 82-41:250 pp., 24 pls.
- Bellier, J.P. (1983): "Foraminifères planctoniques du Crétacé de Tunisie septentrionale": systematique, biozonation, utilisation stratigraphique de l'Albien.
- Bellier, J.P.; Caron, M.; Donze, P.; Herm, D.; Maamouri, A.L. & Salaj, J. (1983): "Le Campanien sommital et le Maastrichtien de la coupe du Kef (Tunisie septentrionale): zonation sur la base de Foraminifères planctoniques". Zitteliana, München, 10, 609-611.
- Berggren, W.A. (1960): "Biostratigraphy, planktonic foraminifera and the Cretaceous-Tertiary boundary in Denmark and southern Sweden". Report of the 21st International Geological Congress, Norden 6, 181-192.
- Berggren, W.A. (1965): "Some problems of Paleocene-Lower Eocene Planktonic foraminiferal correlations". Microplaeontology, 11, (3), S. 278-300, New York.

- Berggren, W.A. (1968): "Phylogemetic an taxonomic problems of some Tertiary planktonic foraminiferal lineages". Tulane. Univ. Stud. Geol. Paleont. New Orleans, 6, 1-22.
- Berggren, W.A. (1969): "Biostratigraphy and planktonic foraminiferal zonation of the Tertiary System of the Sirte Basin of Libya, North Africa". Proc. 1st Intern. Conf. plankt. Microfossils, 1, S. 104-120, Brill, Leiden.
- Berggren, W.A. (1971): "Tertiary boundaries and correlation". In: Funnel, B.M. & Riedel, W.R. (Eds.), Micropaleontology of the oceans, Cambridge Univ. Press, S. 693-809, Cambridge.
- Berggren, W.A. (1977): "Atlas of Palaeogene planktonic forminifera. Some species of the genera Subbotina, Planorotalies, Morosovella, Acarinina and Truncorotaloides". In: Ramsay, A.T.S., Ed., Oceanic micropalaeontology. London: Academic Press, pp. 205-299.
- Berggren, W.A. (1994): "In defense of the Selandian". GFF, 116, 44-46.
- Berggren and Miller (1988): "Paleogene tropical planktonic foraminiferal biostratigraphy and magnetobichrnology-micropaleontology". Vol. 34, no. 4, pp. 362-380.
- Berggren, W.A.; Olsson, R.K. & Reyment, R.A. (1967): "Origin and develpment of the foraminiferal genus pseudohastigerina Banner and Blow". Micropaleont. New York, 13: 265-288, 1 pl.
- Berggren, W.A.; Kent, D.; Swisher, C. & Miller, K.G. (1994): "A revised Cenozoic geochronology and chronostratigraphy". SEPM, Spc. Publ.
- Blankenhorn, M. (1900): "Neues Zur Geologie und Paleontology Aegyptens, II Das Palaogen (Eocan and Oligocan)", Z. dent. Geol. Ges., Vol. 52, S. 403-479.
- Blow, W.H. (1979): "The Cainozoic Globigerinida", E.J. Bripp, Leiden, Netherlands, 1413 pp., 264 pls.
- Boersma, A. & Shackleton, N.J. (1981): "Oxygen- and carbon-isotope variations and planktonic-foraminifer depth habitats, Late Cretaceous to Paleocene, central Pacific, Deep Sea Drilling Project Sites 463 and 465". In: T.L. Vallier, J.

- Caron, M. (1983): "Taxonomie et phylogenie de la-famille des Globotruncanidae". *Zitteliana*, 10, S. 677-681, München.
- Caron, M. (1985): Cretaceous Planktonic Foraminifera. In bolli, H.M., Saunders, J.B. and Perch-Nielsen (eds) plankton stratigraphy". Cambridge Earth Science Ser., p. 17-86.
- Carsey, D.D. (1926): "Foraminifera of the Cretaceous of Central Texas". Texas Univers. Bull. 2612, S. 1-56, Taf. 1-8, Austin.
- Cherif, O.H.; Al-Rifaiy, I.A.; Al-Afifi, F.I. & Orabi, O.H. (1989): "Planktonic foraminifera and chronostratigraphy of Senonian Exposures in west-central Sinai, Egypt." *Revue de Micropaléontologie* 32, 167-184.
- Cushman, J.A. (1925): "Some new Foraminifera from the Velasco Shale of Mexico". *Cushm. Lab. Foram. Res.*, 2, (1), S. 16-24, Taf. 2, 3.
- Cushman, J.A. (1926): "Some Foraminifera from the Mendez Shale of Eastern Mexico". *Contr. Cushm. Lab. Foram. Res.*, 2, (2), S. 16-24, Taf. 2, 3, Sharon, Mass.
- Cushman, J.A. (1927): "An outline of a reclassification of the Foraminifera": *Cush. Lab. Foram. Res. Contr.*, Massachuttes, 3, 1-105.
- Cushman, J.A. (1938): "Additional new species of American Cretaceous foraminifera". *Cushm. Lab. Foram. Res. Contr.*, 14(2), p. 31-50, pls. 5-8, Sharon, Mass., USA.
- Cushman, J.A. (1946): "Upper Cretaceous Foraminifera of the Gulf Coastal Region of the United States and adjacent area". *US. Geol. Surv. Prof. paper* 206, p. 1-241, pls. 1-66, Washington.
- Cushman, J.A. & Ponton, F.L. (1932): "An Eocene foraminiferal fauna of Wilcox age from Alabama". *Cushm. Lab. Foram. Res. Contr.* 8, p. 51-72, pls. 7-9. Mass., U.S.A.
- Cushman, J.A. & Bermudez, P.J. (1949): "Some Cuban species of Globorotalia". *Contr. Cushman Lab. Foram. Res.*, 25, S. 26-45, Taf. 5-8, U.S.A.

- Cuvillier, J. (1930): "Revision du Nummulitique Egypten (Stratigraphie et Paleontologie)". Mem. Inst. Egypt, Le Caire (Cairo), n.s. 16, 1, 371 S., 25 Taf., Fig. 23, Cairo.
- Dabous, A.A. & Mohammed, M.H. (1989): "Petrography and lithofacies studies on the Lower Tertiary sediments of Esna-Qena district, Upper Egypt". M.E.R.C. Ain Shams Univ., Sc. Ser., Vol. 3, 1989, 137-149, Cairo.
- Darmoian, S. (1975): "Planktonic Foraminifera from the Upper Cretaceous of southeastern Iraq: Biostratigraphy and systematics of the Heterohelcidae". micropaleo. vol. 21, no. 2, pp. 185-214, Pls. 1-4.
- De Klasz, I. (1953a): "Einige neue oder wenig bekannte Foraminiferen aus der helvetischen Oberkreide der bayerischen Apfen, südlich Traunstein (Oberbayern)". Geologica Bavarica, 17, 223-224. pls. 4-7.
- De Klasz, I. (1953b): "On the foraminiferal genus Gublerina Kikoine". Geologica Bavarica, 17, 245-251, pl. 8.
- De Lapparent, A.F. (1939): "A propos de L'age tertiaire due "Calcaire Pisolithique" de Vigny (S.-et-O.)". Soc. géol. France, Bull, 5 sér. vol. 9, p. 249-250.
- Dercourt, J.; Zonenshain, L.P.; Kazmin, V.G.; Le Pichon, X.; Knipper, A.L.; Grandjacquet, C.; Sobrshchikov, I.M.; Boulin, J.; Sorokhtin, O.; Geyssant, J.; Levebrier, C.; Biju-Duval, B.; Sibuet, J-C.; Savostin, L.A.; Westphal, M. & Lauer, J.P. (1985): "Presentation de 9 cartes paleogeographiques au 1/20.000.000 s entendant de l'Atlantique au Pamir pour la periode de Lias a Actuel". Bull. Soc. Geol. France, 8, (5), 637-652.
- D'Orbigny, A. (1839): "Foraminiferes In: R. de la Sagra (ed.), Histoire Physique, politique et naturelles de l'île de Cuba, Bertrand, Paris, 224 p.
- D'Orbigny, A. (1840): "Memoire dur les Foraminifers de la craie Blanche du Bassin de Paris". Mem. Soc. Geol. France, 4, P. 1-51, pls. 1-4, Paris.
- Dorreen, J.M. (1974): "The western Gaj River section, Pakistan, and the Cretaceous-Tertiary boundary". Micropaleontology, vol. 20, no. 2, pp. 178-193, pl. 1, text-figs. 1-7.

- Egger, J.G. (1899): Foraminiferen und Ostrakoden aus den Kreidemergeln der oberbayerischen Alpen". K. Bayer, Akad. Wiss., Munchen, Math. Phys. Cl., Abhandl., 21, (1), p. 3-230, pls. 1-27.
- Ehrenberg, C.G. (1840): "Ueber die Bildung der Kreidefelsen und des Kreidemergels durch unsichtbare organismen". Kon. preuss. Akad. Wiss. Berlin Abd., Jahrg. 1838, p. 59-147.
- El-Akkad, S. & Issawi, B. (1963): "Geology and Iron ore deposits of the Bahariya Oasis". Geol. Res. & Mining paper No. 18.
- El-Beshtawy, M.K. (1989): "Biostratigraphic studies on the Late Cretaceous-Early Tertiary between Wadi Tayiba and Wadi Feiran, West Central Sinai, Egypt". M.Sc. Thesis, Fac. of Science, Benha University.
- El-Gammal, R.M. (1994): "Geological studies of the Late Cretaceous-Early Tertiary succession in southwestern Sinai, Egypt". Ph.D. Thesis, Faculty of Science, Al-Azhar Univ. 336 pp., 20 pls., Cairo.
- El-Naggar, Z.R. (1966): "Stratigraphy and planktonic Foraminifera of the Upper Cretaceous-Lower Tertiary succession in the Esna-Idfu region, Nile Valley, Egypt, UAR". Bull. British Museum (Nat. Hist.) Geol. Suppl. 2, 291 p. 23 pls., 18 figs., London.
- El-Naggar, Z.R. (1968): "Stratigraphy and classification of Type Esna Group of Egypt: Discussion". AAPG Bull. 52, 9, 1792-1798.
- El-Naggar, Z.R. (1970): "On a proposed lithostratigraphic subdivision for the Cretaceous-Early Paleocene succession in the Nile Valley, Egypt. Egypt UAR". 7th Arab Petrol. Congress. Kuwait, paper No. 64, (8-3). 50 S.
- El-Naggar, Z.R. (1971): "On the classification, evolution and stratigraphical distribution of the Globigerinacea". In: Farinacci, A., Ed., Proceedings of the Second Planktonic Conference, 1, 421-476.
- El-Sheikh, H.A. & El-Beshtawy, M.K. (1992): "Biostratigraphic studies on the Late Cretaceous-Early Tertiary between Wadi Tayiba and Wadi Feiran, West

- Central Sinai, Egypt". *Geology of the Arab World*, Cairo Univ., 395-405, Cairo.
- Esker, G.L. (1968): "Danian Ostracodes from Tunisia". *Micropal.*, 14,3, 319-333, 4 pls., New York.
- Faris, M.I. (1947): "The content of the Cretaceous and Eocene rocks in the Tramsa-Tukh area (Qena; Upper Egypt)". *Inst. Egypt, Bull.* vol. 28, pp. 73-85, 2 pls.
- Faris, M. (1974): "Geological and Paleontological studies on the Late Cretaceous-Early Tertiary succession in Qena region and Kharga Oasis. M.Sc. Thesis, Assiut Univ., 189, pp., 10 pls.
- Faris, M.M. (1982): "Biostratigraphy of the Upper Cretaceous-Lower Tertiary succession of Duwi range, Quseir district, Egypt". *Revue de micropaleontologie* vol. 27, no. 2, pp. 107-112.
- Faris, M.M. (1982): "Micropaleontologie et biostratigraphie du Cretace superieur et de L'Eocene inferieur de L'Egypte centrale (Region de DUWI Vallee du Nil, Oasis de Kharga et de Dakhla)". *These de Doctorat d'etat presentee a L'Univers. Pierre et Marie Curie, Paris*, P. 1-438, pls. 1-50, figs. 1-81, Tab. 1-17, Paris.
- Faris, M.M. (1984): "The Cretaceous/Tertiary boundary in central Egypt (Duwi region, Nile Valley, Kharga and Dakhla Oases)". *N. Jb. Palaont. Mh.*, (7), 385-392; Stuttgart.
- Faris, M.M. (1985): "Stratigraphy of the Late Cretaceous-Early Tertiary sediments in the Ghanima and Ain Amur sections, Kharga area, Egypt". *Newsl. Stratig.* 14(1) p. 36-47, 4 fig., 2 pl. Berlin. Stuttgart, 15.3
- Faris, M. & A.T. Abd El-Hameed (1986): "Contribution to the Late Maastrichtian-Paleocene chalk at Jiran al Ful area, Abu Roash, west of Cairo, Egypt". *Newsl. Stratigr.* 15, 3, 139-149, Berlin-Stuttgart.
- Faris, M.; Abd El Monem, T.; Abdel Hameed & Akmal Marzuk, M. (1986): "The Cretaceous/Tertiary boundary in Taramsa section west of Qena, Ni. Valley, Egypt". *Newsletters in Stratigraphy*, 16(2), 85-97.

- Faris, M.M.; Alam, A. & Khalil, H.M. (1986): "Early Eocene planktonic foraminifera and calcareous nannofossil biostratigraphy of Gebel Qabeliat, southwest Sinai, Egypt". *J. Geol.* 30, No. 1-2, pp. 71-88.
- Finlay, H.J. (1939): "New Zealand Foraminiferal key species in stratigraphy". No. 1- Royal Soc. New Zealand Trans. Wellington, 68, 504-535, 1 pl.
- Finlay, H.J. (1947): "New Zealand foraminifera: Key species in stratigraphy". No. 5. N. 2. *J. Sci. Technol. section B.* 28, 259-92.
- Fiorentino, A. (1995): "Calcareous nannofossil Stratigraphy of Maastrichtian sections in Israel". *Cretaceous Research*, 16, 327-341.
- Fleury, J.J.; Bignot, G.; Blondeau, A. & Poignat, A. (1985): "Biogeographie de Foraminifères benthiques téthysiens de Senonien à l'Eocène supérieur". *Bull. Soc. Geol. France*, 8(1), 757-770.
- Frerichs, W.E. (1971): "Evolution of planktonic foraminifera and paleotemperatures". *Journal of Paleontology*, 45, 964-968.
- Gallitelli, E.M. (1957): "A revision of the Foraminiferal Family Heterohelcidae". *U.S. Nat. Mus. Bull.* 215, pp. 133-154.
- Ghorab, M.A. (1956): "A summary of a proposed rock stratigraphic classification for the Upper Cretaceous rocks in Egypt Road before Geol". *Soc. Egypt.* (Unpublished).
- Gibson, T. & Buzas, M.A. (1973): "Species diversity: Patterns in modern and Miocene foraminifera of the eastern margin of North America". *Geol. Soc. Amer. Bull.* 84, 277-238.
- Glaessner, M.F. (1937): "Plankton foraminiferen aus der Kreide dem Eozän und der stratigraphische Bedeutung. (Studies in micropaleontology: Moscow Univ., Lab. Paleont., publ. Moscow, 1: 27-45.
- Gondolfi, R. (1955): "The genus Globotruncana in northeastern Colombia". *Bull. Am. Paleontology*, 36, (155), 118 p., 10 pl., 12 fig.

- Graham, J.J. & Church, C.C. (1963): "Campanian foraminifera from the Stanford University campus, California". Stanford University Publications in Geological Science, 8(1), 1-90.
- Haggag, M. (1986): "Late Cretaceous and Paleocene Planktonic Foraminifera from Abu Roash, West of Cairo, Egypt". Middle East Research Centre, Ain Shams University, Scientific Research Series 6, 151-179.
- Hagn, H. (1953): "Zur Kenntnis des unteren Obercampan's (Zone der *Belemnitella mucronata* senior) in Südbayern". N. Jb. Geol. Paläont., Abh. 96, 304-338.
- Haq, B.U.; Hardenbohl, J. & Vail, P.R. (1987): "Chronology of Fluctuating Sea Levels Since the Triassic". Science, 235, 1156-1167; London.
- Hart, M.B. & Bailey, H.W. (1979): "The distribution of planktonic Foraminiferida in the mid-Cretaceous of N.W. Europ". Aspekte der Kreide Europas. I.U.G.S., Ser. A, No. 6, 527-542.
- Haslett, S.K. (1994): "Planktonic foraminiferal biostratigraphy and Palaeoceanography of the Cretaceous-Tertiary boundary section at Bidart, south-west France.
- Hendriks, F. (1987): "Cretaceous to Early Tertiary evolution of the depositional environments of southeastern Egypt". Current research in African earth Sciences, Matheis & Schandelmeier (eds.).
- Hendriks, F. & Luger, P. (1987): "The Rakhayat Formation of the Gebel Qreiya area: evidence of Middle Campanian to Early Maastrichtian synsedimentary tectonism". Berliner geowiss. Abh., (A), 75, Berlin (this volume).
- Hendriks, F.; Luger, P.; Kallenbach, H. & Schroeder, J.H. (1984): "Stratigraphical and sedimentological framework of the Kharga-Sinn el-Kaddab stretch (western and southern part of the Upper Nile Basin), Western Desert, Egypt". Berl. Geowiss. Abh. Reihe A, 40, p. 117-151, g Abb., Berlin.
- Hendriks, F.; Luger, P.; Bowitz, J. & Kallenbach, H. (1987): "Evolution of the depositional environments of SE-Egypt during the Cretaceous to lower Tertiary". Berliner geowissenschaftliche Abhandlungen (A) 75(1), 49-82, Berlin.

- Herm, D.; Hileberandt, A.; von und Perch-Nielsen, K. (1981): "Die Kreide/Tertiär-Grenze im Lattengebirge (Nordliche Kalkalpen) in mikropaläontologischer Sicht.-Geologica bavarica, 319-344. München.
- Hermine, M.H. & Issawi, B. (1971): "Rock stratigraphic classification of the upper Cretaceous-Lower Tertiary exposures in southern Egypt". In: Gray, C. (ed.): Symposium on the Geology of Libya, Fac. Sci. Univ. Libya, Tripoli, Libyan Arab Rep., 147-154, Tripoli.
- Hewaidy, A. (1987): "Biostratigraphy and Paleobathymetry of the Esna Shale in El-Qusaima area, N.E. Sinai, Egypt". M.E.R.C. Ain Shams Univ., Earth Sc. Ser., Vol. 1, pp. 180-206.
- Hewaidy, A. (1994): "Biostratigraphy and Paleobathymetry of the Garra-Kurkur area, southwest Aswan, Egypt". M.E.R.C. Ain Shams Univ. Earth Sci. Ser., Vol. 8, P. 48-73.
- Hewaidy, A. & Anan, H.S. (1986): "Paleocene benthonic foraminiferal paleoecology of Central Egypt". M.E.R.C. Ain Shams Univ., Sci. Res. Ser., 6, 81-104.
- Hillebrandt, A.; Von (1965): "Foraminifern-Stratigraphie im Alttertiär, von Zumaya (Provinz Guipuzcoa, NW Spanien) und ein Vergleich mit anderen Tethys-Gebieten". Bayerische Akademie Wissenschaften, Mathematisch-Naturwissenschaftlichen Klasse, N.F., 123, 62 pp.
- Hinte, J.E. van (1963): "Zur stratigraphie und mikropaleontologie der Oberkreide und des Eozäns des Krappfeldes (Karnten). Jarb. Geol. Bund., Sond. 8, 1-147.
- Hongstein, A.; Rosenfeld, A. & Benjamini, C. (1991): "Ostracods and foraminifera from the Early-Middle Eocene of Qeren Sartaba, Jordan Valley". J. micropaleontol., 10(1), p 95-p 107.
- Hume, W.F. (1910): "The origin of the Nile Valley in Egypt". Geol. Mag., V. 7, P. 385-389.
- Hume, W.F. (1911): "The effects of secular oscillation in Egypt during the Cretaceous and Eocene periods". Quart. Journ. Geol. Soc. of London, 67(1), S. 118-148, London.

- Hume, W.F. (1927): "The phosphate deposits in Egypt". Geol. Survey Egypt, Cairo, paper 41. 20 pp.
- Ismail, A.A. (1989): "The stratigraphy of the Cretaceous-Lower Tertiary of the southwestern region of the Gulf of Suez-Egypt". Unpublished Ph.D. Thesis, Geology Department, Fac. of Sci. Ain Shams Univ.
- Ismail, M.M. & Abd Allah, A.M. (1966): "contribution to the stratigraphy of St. Paul Monastery area by microfacies Alexandria Univ". Bull., Fac. Sci., 7: 325-334.
- Issawi, B. (1971): "Geology of Darb El Arab' an Western Desert, Egypt". Ann. Geol. Surv. Egypt, 1, 53-92, Cairo.
- Issawi, B. (1972): "Review of Upper Cretaceous-Lower Tertiary Stratigraphy in central and southern Egypt". AAPG. Bull., 56, 8, 1338-1463, 6 figs., 4 tabl., Tulsa.
- Jenkins, D.G. (1971): "New Zealand Cenozoic planktonic foraminifera": N.Z. Geol. Surv. Paleont., Wellington, 42: 1-278, 23 pls.
- Jenkins, D.G. & Luterbacher, H. (1992): "Paleogene stages and their boundaries (Introductory remarks)". N. Jb Geol. Paläont. Abh. 186, 1-5.
- Kaiho, K. (1992): "A low extinction rate of intermediate water benthic foraminifera at the Cretaceous/Tertiary boundary". Mar. Micropaleontol., 18, 229-259.
- Kassab, I.I.M. (1975): "Globotruncana falsocalcarata Kerdany and Abdelsalam from northern Iraq". Micropaleontology 21(3), 346-351, 2 pls., New York.
- Kassab, I.I.M. (1976): "Some Upper Cretaceous planktonic foraminiferal genera from northern Iraq". Micropal. 22, 215-238.
- Kassab, I.I.M. (1978): "Biostratigraphy of Upper Cretaceous-Lower Tertiary of North Iraq". Actes Vie Colloq. Afric. Micropal. (Tunis), 2, pp. 259-325.
- Kassab, I.I.M. (1979): "The genus Globotruncana Cushman from the Upper Cretaceous of Northern Iraq". J. Geol. Soc. Iraq (Baghdad), 12, 1, pp. 27-127.

- Keller, G. (1988a): "Extinction, survivorship and evolution of Planktic foraminifera across the Cretaceous/Tertiary boundary at El-Kef, Tunisia". *Marine Micropaleontology*, 13, 239-263.
- Keller, G. (1989a): "Extended K/T boundary extinctions and delayed population change in planktic foraminifera of continental shelf sections: Implications for impact and volcanism theories". *Geol. Soc. Am. Bull.*, 101, 1408-1419.
- Keller, G. (1993): "The Cretaceous-Tertiary boundary transition in the Antarctic Ocean and its global implications". *Mar. Micropaleontol.*, 21, 1-45.
- Keller, G.; Barrera, E.; Schimits, B. & Mattson, E. (1993): "Gradual mass extinction, species survivorship and long-term environmental changes across the Cretaceous-Tertiary boundary in high latitudes". *Geol. Soc. Am. Bull.*, 105, 979-997.
- Kenawy, A.I. (1966): "Die Grossforaminiferen des bayerischen Ultrahelveticums". Sonderdruck. aus *Geologica Bavarica* Nr. 56, pp. 104-176.
- Kenawy, A.I. (1967): "Large foraminifera from the Thebes Formation, Taramsa section, Qene, Egypt". *Bull. Fac. Sci. Assiut Univ.* (in press).
- Kenawy, A.I. (1972): "Planktonic Foraminifera of lower Tertiary succession in the Taramsa Section, Qene, Nile Valley, Egypt". *Fragm. Min. Pal.*, 3, 25-116.
- Kerdany, M.T. & Abdel-Salam, H. (1969): "Globotruncana falsocalcarata n.sp. from the Quseir area, Eastern Desert, U.A.R.". *Proceedings, 3rd African Micropaleontological Colloquium, Cairo 1968: 261-267, The National Information and Documentation Centre, Cairo.*
- Khalilov, D.M. (1956): "The foraminiferal fauna of the Paleogene deposits of Azerbaijan". *Akad. Nauk Azerbaj. Trudy Instituta Geologii, Akademiya Nauk Azerbaydzhanskoy SSR* 17, 234-255. (in Russian).
- Kikoine, J. (1948): "Les Heterohelidae du Crétacé supérieur pyrénéen". *Bulletin de la Société Géologique de France*, ser. 5, 18(1), 15-35, pl. 1,2.
- Krashennnikov, V.A. & Abdel-Razik, T.M. (1969): "Zonal stratigraphy of the Paleocene in Quseir, "Red Sea Coast". In: R. Said, J.P. Beckmann, et al.

- (Editors), Proceedings of the third African Micropaleontological Colloquium". NIDOC, Cairo, Egypt, pp. 229-309.
- Kroon, D. & Nederbrat, A. (1990): "Ecology and Paleoecology of Triserial Planktic Foraminifera". *Marine Micropaleontology*, 16 (1990) 25-38.
- Kulbrok, F. (1996): "Stratigraphy, Facies and Sequence Stratigraphy of the Late Cretaceous-Early Tertiary successions in the surrounding of the northern Gulf of Suez". Ph.D. thesis, Bremen University.
- Kuss, J. (1987): "Interpretation of Upper Cretaceous and Lower Tertiary facies patterns from limestone-dominated sequences of northeastern Egypt". current research in African earth sciences, Matheis & Schandelmeir (eds).
- Kuss, J. (1992): "Facies and Stratigraphy of Cretaceous Limestones from Northeast Egypt, Sinai and Southern Jordan". *Proc. 1st Int. Conf. on the Geology of the Arab World, Cairo 1992* - (in press).
- Lamolda, M.B.; X. Orue-Etxebarria & F. Proto Decima (1983): "The Cretaceous/Tertiary boundary in Sopelana (Biscay, Basque country)". *Zitteliana* 10, 633-670.
- Leonov, G.P. & Alimarina, V.P. (1961): "Stratigraphy and planktonic foraminifera of Cretaceous-Paleogene "transition" beds of the central part of the north Caucasus". *Sbornik trudov geol. fak. Moskov. University. (k 21. sessi mezhdunarod. geol. kongr.)*: 29-53, Moscow University (in Russian).
- Le Roy, L.W. (1953): "Biostratigraphy of the Maqfi section, Egypt". *Geol. Soc. Amer. Mem. Nr. 54*, 73 p., 4 figs., 14 pls., Washington.
- Loeblich, A.R. (1951): "Coiling in Heterohelcidae". *Cushman. Found. Foramin. Res. Contr.*, vol. 2, pp. 106-110.
- Loeblich, A.R. & Tappan, H. (1957): "Correlation of the Gulf and Atlantic coastal plain Paleocene and Lower Eocene formations by means of planktonic foraminifera". *Journ. Paleont.* 31, (6), p. 1109-1137.
- Loeblich, A.R. & Tappan, H. (1984): "Suprageneric classification of the Foraminiferida (protozoa)". *Micropal.*, 30, (1), p. 1-70.

- Loeblich, A. & Tappan, H. (1988): "Foraminiferal genera and their classification". Library of Congress Cataloging-in-Publication Data. Van nostrand reinhold company. New York.
- Loetterle, G.J. (1937): "The micropaleontology of the Niobrara formation Kansas, Nebraska and South Dakota". Nebraska Geological Survey Bulletins, ser. 2, 12(1), 1-73, pls. 1-11.
- Longoria, J.F. & Von Feldt, A. (1991): "Taxonomy, Phylogenetics and biochronology of single-keeled globotruncanids (Genus Globotruncanita Reiss), micropaleontology, Vol. 37, no. 3. PP. 197-243, Pls. 1-11 text-figures 1-16.
- Lu & Keller (1995): "Planktonic foraminiferal faunal turnovers in the subtropical pacific during the Late Paleocene to Early Eocene". Journal of Foraminiferal Research. v. 25, no. 2, p. 97-116, April 1995.
- Luger, P. (1985): "Stratigraphie der Marinen Oberkreide und des Alttertiars in Sudwestlichen Obruil-Becken (SW-Egypt), unter besonderer Berücksichtigung der Micropalaontologie, Palökologie und Palaogeographie". Berliner Geowiss. Abh. R, 36, 151 p., 22 Abb., 11 Tab., 24 pls. Berlin.
- Luger, P. (1988): "Campanian to Paleocene agglutinated foraminifera from fresh water influenced marginal marine (deltaic) sediments of southern Egypt". Abh. Geol. B.A. Wien, April 1988, pp. 255-263.
- Luger, P. (1988): "Maastrichtian to Paleocene facies evolution and Cretaceous/Tertiary boundary in middle and southern Egypt". Revista Espanola de Micropaleontologia, no. Extraordinario, 83-90.
- Luterbacher, H.P. (1964): "Studies in some Globorotalia from the Paleocene and Lower Eocene of the central Apeninnes". Eclog. Geol. Helvet., Basel, 57, 631-670.
- Luterbacher, H.P. (1975): "Planktonic foraminifera of the Paleocene and Eocene, Passagno section". Schweizerische Paläontologische Abhandlungen, 97, 57-67.

- Luterbacher, H.P. & Premoli-Silva (1964): "Biostratigrafia del limite Cretaceo-Terziario nell' Appennino centrale". *Rivista Italiana di Paleontologia et Stratigrafia* 70, 67-128.
- Macleod, N. (1993): "The Maastrichtian-Danian radiation of triserial and biserial Planktic foraminifera : Testing Phylogenetic and adaptational hypotheses in the (micro) fossil record". *Marine Micropaleo.*, 21, 47-100.
- Malmgren, B.A. (1991): "Biogeographic patterns in terminal Cretaceous planktonic foraminifera from Tethyan and warm transitional waters". *Mar. Micropaleontol.*, 18, 73-99.
- Manivit, H. & H. Feinberg (1984): "Correlation of magnetostratigraphy and nannofossil biostratigraphy in Upper Cretaceous and Lower Paleocene sediments of the Walvis Ridge area". In: *Repts. DSDP 74*, 469-474, Washington.
- Marina, V.P. (1963): "(Some peculiarities of development of planktonic foraminifera in connection with zonation of the lower Paleogene of the northern Caucasus) in Russ.". *Moscov Univ., Voprosy Mikropal.*, 7, S. 158-95, Abb. 1-4, Ta. 1,2, Moskau.
- Martin, L.T. (1943): "Eocene foraminifera from the type Lodo Formation, Fresno Claifornia, Stanford Univ". *Publ. Geol. Sci.*, 3, 99-105.
- Martini, E. (1971): "Standard Tertiary and Quaternary calcareous nannoplankton zonation". In: *Farinacci, A. (Ed.) Proceedings II Planktonic Conference, Roma, 1970*, 2, 739-785.
- Martini, E. (1976): "Cretaceous to Recent calcareous nannoplankton from the Central Pacific Ocean (DSDP, Leg. 33)". *Init. Repts. DSDP. 33*, pp. 383-423, Washington.
- Masters, B.A. (1976): "Planktonic foraminifera from the upper Cretaceous Selma group, Alabama". *Journal of Paleontology*, 50(2), 318-330, pls. 1, 2.

- Masters, B.A. (1977): "Mesozoic Planktonic Foraminifera. A world-wide review and analysis". In Ramsey A.T.S. (ed.), *Oceanic Micropalaeontology* 1, 301-731, pls. 1-58, London, etc.
- Masters, B.A. (1984): "Comparison of planktonic foraminifers at the Cretaceous-Tertiary boundary from the Haria Shale (Tunisia) and the Esna Shale (Egypt)". In: *Proceedings of the 7th Exploration Seminar, EGPC, Cairo*. pp. 310-324.
- Masters, B.A. (1993): "Re-evaluation of the species and subspecies of the genus *Plummerita* Brönnimann and a new species of *Rugoglobigerina* Brönnimann (Foraminiferida)". *Journal of Foraminiferal Research* 23(4), 267-274.
- McGowran, B. (1964): "Foraminiferal evidence for the Paleocene age of the King's Park Shale (Perth basin, Western Australia)". *J.R. Soc.* 47, 81-8.
- Molina, E.; Canudo, J.I.; Martinez-Ruiz, F. & Ortiz, N. (1994): "Integrated stratigraphy across the Paleocene/Eocene boundary at Caravaca, southern Spain". *Eclog. Geol. Helv.*, 87, 47-61.
- Montanaro, G. (1957): "A revision of the foraminiferal family Heterohelcidae. In: Loeblich, A.B., et al., *Studies in foraminifera, part 1- Planktonic foraminifera*. U.S. National Museum Bulletins 215, 133-154, Pls. 31-34.
- Morozova, V.G. (1939): "The stratigraphy of Upper Cretaceous and Lower Tertiary deposits in the Emba oil bearing district according to the foraminiferal fauna". *Soc. Nat. Moscow, n.s. sect. geol.*, 17, 59-86 (In Russian).
- Morozova, V.G. (1957): "The foraminiferal superfamily Globigerinidea superfam. Nova and some of its representatives". *Akad. Nauk SSSR, Doklady*, 114, 1109-1111 (In Russian).
- Morsi, S. (1984): "Micropaleontology and Biostratigraphy of some Palaeogene rocks in the Gulf of Suez area, Egypt". M.Sc. Thesis, Fac. of Sci., Cairo Univ.
- Murray, J.W. (1976): "A method of determining proximity of marginal seas to an ocean". *Marine Geology* 22, 103-119.

- Murry, J.W. (1991): "Ecology and Palaeoecology of benthic Foraminifera". 397 pp. (Longman Scientific & Technical, London).
- Nakkady, S.E. (1950): "A new foraminiferal fauna from the Esna Shales and Upper Cretaceous Chalk of Egypt". Jour. Paleont., 24, (6), p. 675-692, Tulsa, Oklahoma.
- Nakkady, S.E. (1951): "Zoning the Mesozoic-Cenozoic transition of Egypt by the Globorotalidae". Fac. Sci., Alexandria Univ., Bull., 1, pp. 45-58.
- Nakkady, S.E. (1957): "Biostratigraphy and interregional correlation of the Upper Senonian and Lower Paleocene of Egypt". Jour. Pal., Vol. 31, No. 2, pp. 428-447, Tulsa, Oklahoma.
- Nakkady, S.E. (1958): "Stratigraphy and Petroleum geology of Egypt". Assiut Univ. Press., Monograph. Series, 1, 215 pp., Assiut.
- Nakkady, S.E. (1959): "Biostratigraphy of Um El-Ghanayem section, Egypt". Micropal., 5, 453-472, New York.
- Nash, S. (1981): "A neotype for the Cretaceous genus *Pseudotextularia* Rzehak, 1891". J. of Foram. Res., v. 11, no. 1, p. 70-75, pl. 1.
- Nederbragt, A.J. (1991): "Late Cretaceous biostratigraphy and development of Heterohelidae (Planktic foraminifera)". micropaleontology, Vol. 37, no. 4, pp. 329-372, plates 1-12, text-figs. 113, table 1.
- Nederbragt, A.J. (1992): "Paleoecology of Late Maastrichtian Heterohelidae (planktic foraminifera) from the Atlantic region". Palaeoecol., 92, 361-374.
- Newton, R.B. (1898): "Notes on some Lower Tertiary shells from Egypt". Geol. Mag., 5 (Decade 4), 531-541.
- Nichols, D.; Brown, J. & Orth, C. (1992): "A new Cretaceous-Tertiary locality in the western Powder River basin, Wyoming: biological and geological". Cretaceous Research 13, 3-30.
- Norris, R.D. (1991): "Parallel evolution in the keel structure of planktonic foraminifera". J. Foraminiferal Res., 21, 319-331.

- Norris, R.D. (1992): "Umbilical structures in Late Cretaceous Planktonic foraminifera". *micropaleontology*, Vol. 38, pp. 165-181.
- Nuttall, W.L. (1930): "Eocene Foraminifera from Mexico". *J. Plaeont.* Tulsa/Oklahoma, 4: 271-293, 2 pls.
- Olsson, R. (1964): "Late Cretaceous Planktonic Foraminifera from New Jersey and Delaware". *Micropal.*, 10, p. 157-188.
- Olsson, R. (1970): "Paleocene Planktonic foraminiferal, biostratigraphy and paleozoogeography of New Jersey". *Jou. of Paleontology*, Vol. 44, pp. 589-597.
- Olsson, R.K.; Hemleben, C.; Berggren, W.A. & Liu, C. (1992): "Wall texture classification of Planktonic foraminiferal genera in the lower Danian". *Journal Foraminiferal Res.*, 22, 195-213.
- Omara, S.M. (1951): "Micro-stratigraphical studies of the Cretaceous rocks of the Nezzazat area". Fouad I Univ., M.Sc. Thesis.
- Omran, A.M. (1973): "Micropaleontological studies on Gebel Umm El-Ghanayem, Kharga Oasis, Egypt". M.Sc. Thesis, Assiut Univ.
- Orabi, H.O. (1988): "Late Cretaceous stratigraphy and geology of West Central Sina". Ph.D. Thesis, Fac. of Sci., Menoufia Univ.
- Osman, A. (1954): "Upper Cretaceous Foraminifera of Western Sinai". *Bull. Fac. Eng., Cairo Univ.*, 1, p. 335-363, Cairo.
- Parr, W.J. (1938): "Upper Eocene Foraminifera from the deep borings in Kings Park, Perth, Western Australia". *Petrol. Soc. Western Australia, Journ.*, 24, p. 69-101.
- Pearson, P.N. (1993): "A Lineage Phylogeny for the Paleogene Planktonic Foraminifera". *Micropaleontology*, Vol. 39, no. 3, pp. 193-232, text-figures 1-25, table 1-2.
- Perch-Nielsen, K. (1972): "Remarks on Late Cretaceous to Pleistocene coccolith from the North Atlantic". *Init. Repts. DSDP*, 12 pp 1003-1069, Washington.

- Perch-Nielsen, K. (1979): "Calcareous nannofossils at Cretaceous/Tertiary boundary near Biarritz, France". Cretaceous/Tertiary events symposium II. Ed. Univ. of Copenhagen.
- Perch-Nielsen, K. (1980): "New Maastrichtian and Paleocene calcareous nannofossils from Africa, Denmark, the USA and Atlantic and some Paleocene lineages". *Eclog. geol. Helv*, Basel.
- Perch-Nielsen, K. (1981): "Nouvelles observations sur les nannofossiles calcaires à la limite Crétacé/Tertiaire Près de El-Kef (Tunisie)". *Cah. Micropaléont.* 1, 25-36, Paris.
- Perch-Nielsen, K. (1985): "Mesozoic calcareous nannofossils". In: bolli, H.M., Saunders, J.B. & Perch-Nielsen K. *Plankton Stratigraphy*, 329- 426, Cambridge University press.
- Perch, Nielsen, K; Sadek, A.; Barakat, M.G. & Teleb, F. (1978): "Late Cretaceous and Early Tertiary calcareous nannofossil and planktonic foraminifera zones from Egypt". *Actes du Vie Colloque Africain de Micropal. Tunis 1974. Annales de Mineralogie et de Géologie Tunis*, 28(2), 337-403.
- Pessagno, E.A. (1967): "Upper Cretaceous planktonic foraminifera from the Western Gulf coastal plain". *Paleont. Americ.*, 5, (37), p. 245-445, Pls. 48-101, Tex. Fig. 1-63, Ithaka, New York.
- Petters, S.W. (1982): "Central West African Cretaceous-Tertiary benthic foraminifera and stratigraphy". *Palaeontogr. Abt. A; Palaeozool. Stratigr.*, 179, 104 pp.
- Picard, L. (1943): "Structure and evolution of Palestine Hebrew Univ., Geol. Dept., Bull., vol. 4, nos. 2-4.
- Plummer, H.J. (1926): "Foraminifera of the Midway Formation in Texas". *Texas Univ. (Bur. Econ. Geol.) Bull.* 2644, p. 1-206, figs. 1-11, pls. 1-15, Austin, Texas.
- Plummer, H.J. (1931): "Some Cretaceous Foraminifera in Texas". *Texas Univ. Bull.* 3101, p. 109-203, pls. 8-15, Austin, Texas.
- Postuma, J.A. (1971): "Manual of planktonic foraminifera". II. Cenozoic. Elsevier pub. com. Amst., Lon. New York.

- Premoli-Silva, I. (1977): "The earliest Tertiary Globigerina eugubina zone. Paleontological significance and geographical distribution". Mem. 2nd Congress Latinoam. Geol., Vol. 3. spec. publ. 7, 1541-55.
- Premoli-Silva, I. & Boersma, A. (1977): Cretaceous planktonic foraminifers - DSDP Leg 39 (South Atlantic)". In: Supko, P.R., Perch-Nielsen, K. et al., Initial Reports of the Deep Sea Drilling Project, Vol. 39, 615-641, pl. 1, 2. Washington, D.C.: U.S. Government Printing Office.
- Premoli-Silva and Boersma (1988): "Atlantic Eocene Planktonic foraminiferal historical biogeography and Palaeohydrographic indices". Palaeogeography, Palaeoclimatology, Palaeoecology, 67, 315-356.
- Premoli-Silva, I. & Bolli, H.M. (1973): Late Cretaceous to Eocene Planktonic foraminifera and stratigraphy of LEG 1s sites in the Caribbean Sea". Init. Rep. DSDP, 15, p. 499-547, pl. 1-10. Washington.
- Reiss, Z.; Almogi-Labin, A.; Honigstein, A.; Lewy, Z.; Lipson-Benitah, S.; Moshkovitz, S. & Zaks, Y. (1985): "Late Cretaceous multiple stratigraphic framework of Israel". Israel Journal of Earth Science 34, 147-166.
- Robaszynski, F.; Caron, N.; Gonzalez Donoso, J.M. & Wonder, A.A.H. (1984): "Atlas of Upper Cretaceous Planktonic Foraminifera". Revue de Micropaleontologie, 69, (3-4), S. 145-305, pl. 1-54.
- Romein, A.T.T. & J. Smit (1981): "The Cretaceous/Tertiary boundary; Calcareous nannofossils and stable isotopes". Proc. Kon. Nederl. Acad. Wet. Ser. B., 84(3), 295-314, Netherlands.
- Romein, A.T.T. & S.T. Hussain (1983): "Calcareous nannoplankton from the Cretaceous/Tertiary boundary interval in the Western Gaj River Section (Pakistan)". Paleontology, Proceedings B 86 (4), 343-400.
- Roth, P.H. (1978): "Cretaceous nannoplankton biostratigraphy and Oceanography of the Northwestern Atlantic Ocean". Init. Repts. DSDP, 44, pp 731-759, Washington.

- Rzehak, A. (1891): Die Foraminiferen fauna der alttertiären Ablagerungen von Bruderndorf in Niederösterreich, mit Berücksichtigung des angelichen Kreidevorkommens von Leikerdorf". K.U.K. Naturk. Hofm., Anm., Bd. 6, S. 1-12, Wien.
- Said, R. (1960): "Planktonic foraminifera from the Thebes Formation, Luxor, Egypt". *Micropaleontology*, 6, (3), S. 277-286, pl. 1, Text fig. 1, Tab. 1,2, New York.
- Said, R. (1961): "Tectonic framework of Egypt and its influence on distribution of foraminifera". *Bull. Amer. Assoc. Geologists* 45, 198-218.
- Said, R. (1962): "The geology of Egypt". 377 p., Amsterdam - New York (Elsevier).
- Said, R. & Kenawy, A. (1956): "Upper Cretaceous and Lower Tertiary Foraminifera from northern Sinai, Egypt". *Micropal.*, 2, (2) p. 105-173, pl. 1-7, 6 figs., New York.
- Said, R. & Kerdany, M.T. (1961): "The Geology and Micropaleontology of Farafra Oasis". *Micropaleontology* 7(3), 317-336.
- Said, R. & Sabry, H. (1964): "Planktonic Foraminifera from the type locality of the Esna shale in Egypt". *Micropal.*, 10, (3), p. 375-395, pls. 1-3, text. figs. 1-3.
- Salaj, J. (1980): "Microbiostratigraphie du Crétacé et du Paléogène de la Tunisie septentrionale et orientale (hypostratotypes tunisiens)". Institut Geol. Dionyz Stur, Bratislava, 238 p.
- Salaj, J. (1983): "Quelques problèmes taxinomique concernant les foraminifères planktiques et al zonation du Senonien supérieur d'El Kef". *Geologica carpathica* 34, 187-212.
- Salaj, J. (1985): "L'Harien et le Mellégien, les nouveaux étages proposés pour le stratotype du Paléocène marin près D'El Kef (Tunisie septentrionale)". *Les Actes 1er Congr. nat. Sc. de la Terre. Tunis 1981*, 1, 263-270, Tunis.
- Salaj, J. (1994): "Contribution to the microbiostratigraphy of the Middle Paleocene (Harian) of El-Kef region (North Tunisia)". *Zemni Plyn a nafta*, 38, 4, pp. 261-289, Hodanin.

- Salaj, J. & Solakius, N. (1984): "The genus *Kassabiana* Salaj, 1983 from the uppermost Maastrichtian of northeastern Tunisia". *CAN. J. Earth. Sci.* Vol. 21.
- Samanta, B.K. (1970): "Middle Eocene Planktonic foraminifera from Lakhpat, Cutch, Western India". *Micropaleont.* New York 16, 185-215, 3 pls.
- Selima, A. (1980): "Stratigraphical and paleontological studies on the Lower Tertiary sediments, (Dandara section), Qnea, Nile Valley, Egypt". M.Sc. Thesis, Ains Shams Univ., Cairo, Egypt.
- Seyve, C. (1984): "Nannofossil biostratigraphy of the Cretaceous-Tertiary boundary in the Frecnh Basque country". *Bulletm des Centres de Recherches Exploration-Production Elf. Aquitaine* 14, 553-572.
- Shahar, Y. (1968): "Type section of the Campanian-Maastrichtian Ghareb Formation in the Oron syncline, northern Negev". *Geological Survey of Israel, Stratigraphic Sections*, 6, 1-8.
- Shahein, A. (1988): "Fossil fauna and stabile isotopic composition with the Late Cretaceous-Early Tertiary at Gebel Nezzazat, Sinai, Egypt". Ph.D. Thesis, Fac. of Sci., Mansoura Univ.
- Sigal, J. (1952): "Order des foraminiferes in *J. pivetaeu*". *Triate de Paleontologie*, t. I.
- Silter, W.V. (1968): "Upper Cretaceous Foraminifera from southern California and Northwestern Baja California, Mexico". *The University of Kansas Paleont. Contributions* (16). p. 1-141, Fig. 1-9, Pl. 1-24, Tables 1-15.
- Sissingh, W. (1977): "Biostratigraphy of Cretaceous calcareous nannoplankton". *Geologie en Mijnbouw*, v. 56, pp 37-65.
- Smit, J. (1977): "Discovery of a planktonic foraminiferal association between the *Abathomphalus mayaroensis* Zone and the "*Globigerina*" *eugubina* Zone at the Cretaceous/Tertiary boundary in the Barranco del Gredero (Caravaca, SE Spain): A Preliminary Report". *Proceedings Koninklijke Nederlandse Akademie van Wetenschappen, Series B* 80 (4), 280-301.

- Smit, J. (1982): "Extinction and evolution of Planktonic foraminifera after a major impact at the Cretaceous/Tertiary boundary. In Geological implications of impacts of Large asteroids and comets on the earth". Geological Society of America Special Paper 190, pp. 329-352.
- Smith, C.C. & Pessagno, E.A. (1973): "Planktonic foraminifera and stratigraphy of the Corsicana formation (Maastrichtian) North Central Texas". *Cushman Foraminiferal Research Special Publication* 1, 12, 66 p., 27 pls.
- Solakius, N. (1983): "A new planktic foraminiferal species from the Late Maastrichtian of Tunisia". *N. Jb. Geol. paläont. Mh.* 1983, H. 2 pp. 113-118, Stuttgart, Februar 1983.
- Solakius, N. (1983): "Globotruncana falsocalcarata Kerdany-Abdel Salam, 1969, from the Upper Maastrichtian of Enfidaville, Northeastern Tunisia". *Geologický Zborník-Geologica Carpathica*, 34, 5, Bratislava, pp. 617-626.
- Solakius, N. (1983a): Taxonomy of some Globotruncanidae and biozonation of the Late Maastrichtian of El-Haria Formation at Kef en Nsoura, Enfidaville region, eastern Tunisia". *Notes du Service Géologique du Tunisie* 46, 37 pp.
- Solakius, N. (1983b): "Biozonation of the Maastrichtian of section II at Kef en Nsoura, Tunisia, and taxonomy of some Globotruncanidae". *Revista Española de Micropaleontología* 15(2), 241-256.
- Solakius, N. & Salaj, J. (1985): "Foraminiferids from the Maastrichtian of Enfidaville northeastern Tunisia". *N. Jb. Geol. Paläont. Abh.* 170, 2 pp. 229-241.
- Solakius, N. & Salaj, J. (1986): "A new Kaasabiana species from the Uppermost Maastrichtian of Northeastern Tunisia". *Revista Española De Micropaleo.* Vol. XVIII, num. 1, PP. 71-77.
- Solakius, N.; Maamouri, A. & Bensalem, H. (1984): "Planktonic foraminiferal biostratigraphy of the Maastrichtian sedimentary beds at Ain Mdeker, Northeastern Tunisia". *Geobios*, no. 17, fasc. 5. pp. 583-591, 4 fig., 1 pl.
- Solakius, N.; Papaloannou, F.B. & Alexopoulos, A. (1990): "On the Paleogeographic distribution of the Late Maastrichtian Planktonic foraminiferal genus

- Kassabiana Salaj & Solakius, 1984". *Acta Geologica Hispanica*, v. (25), no. 4, PP. 289-298.
- Speijer, R.P. (1994a): "The Late Paleocene benthic foraminiferal extinction event as observed in the Middle East". *Bull. Soc. belge Geol.*: in press (This volume, Chapter IV).
- Speijer, R.P. (1994b): "Paleocene sea-level fluctuations recorded in benthic foraminiferal assemblages from Gebel Oweina (Esna, Egypt)". local, regional or global patterns. *GFF*, 116, 63-65.
- Speijer, R.P. & Samir, A.M. (1994): "Globanomalina luxorensis, a pelagic marker for the Paleocene/Eocene boundary". *Micropaleontology*: in prep. (This volume, Chapter III).
- Stainforth, R.M.; Lamb, I.L.; Luterbacher, H.H.; Beard, J.H. & Jeffords, R.M. (1975): "Cenozoic planktonic foraminiferal zonation and characteristics of index forms". *Univ. Kansas. Paleont. Contr. Lawrence*, 62, 1-425.
- Steinecek, P.L. & Fleisher, R.L. (1978): "Toward the classical evolutionary reclassification of the Cenozoic Globigerinacea (Foraminiferida)". *Journal of Paleontology*, 52(3), 618-635.
- Strougo, A. (1986): "The velascoensis event: A significant episode of tectonic activity in the Egyptian Paleogene". *N. Jr. Geol. Palaont. Abh.* 173(2), 253-269.
- Strougo, A. (1994): "The velascoensis event in Egypt: Paleogeography and biostratigraphic implications, Meeting Proceedings". *Stratigraphy of the Paleocene in GFF volume 116*, pp. 39-67.
- Strougo, A. & Abul-Nasr, R.A. (1981): "On the age of the Thebes Formation of Gebel Duwi, Quesair area, Egypt". *Noues Jahrb. Geol. Paleontol. Monatsh.*, 49-53.
- Strougo, A.; Haggag, M. & Luterbacher, H.P. (1992): "The basal Paleocene "Globigerina" eugubina Zone in the Eastern Desert (St. Paul's Monastery, South Galala, Egypt)". *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1992 (2), 97-101.

- Subbotina, N.N. (1947): "Foraminifera of the Danian and Paleogene deposits of the northern Caucasus". Geol. Razved. Inst. Oil. Fiedls UDSSR, pb. 1, 5, 39-160, figs. 1-9.
- Subbotina, N.N. (1950): "Microfauna and stratigraphy of the Elburgan Horizon and the Horizon of Goriatchii Klitch". Trudy VNIGRI, new ser., 51 (Microfauna of the USSR 4). 5-112 (in Russian).
- Subbotina, N.N. (1953): "Globigerinidae, Hantheninidae, and Globorotaliidae Vses". Neft. Nauchno Issled. geol. Razved. Inst. (VNIGRI) Mikrof. SSSR, Trudy, geol., 5 b, 76, 1-296 (In Russian).
- Thomas, E. (1990a): "Late Cretaceous-early Eocene mass extinctions in the deep sea. In: V.L. Sharpton and P.D. Ward (Editors), GSA Spec. Pap, 247: Global Catastrophes in Earth History; An Interdisciplinary Conference on Impacts, Volcanism, and Mass Mortality. GSA, Inc. Co, pp. 481-496.
- Thomas, E. (1990b): "Late Cretaceous through Neogene deep-sea benthic foraminifers (Maud Rise, Weddel Sea, Antarctica). In: Barker, J.P., Kennett, et al. (Editors), Proceedings of the Ocean Drilling Program, Scientific Results, Vol. 113. ODP, College Station, TX, pp. 571-594.
- Tilev, N. (1951): "Etude des Rosalines maestrichtiennes (geme Globotruncana) du Sud-Est de la Turquie (Sondage de Ramandage)". Bull. Lausanne Univ. Lab. Geol. Minéral., Géophys. Mus. géol. 103, 1-101.
- Todd, R. (1970): Maastrichtian (Late Cretaceous) foraminifera from a deep-sea core of southwestern Africa". Rev. espan. Micropaleont. 2/2, 131-154.
- Toulmin, L.D. (1941): "Eocene smaller Foraminifera from the salt Mountain Limestone of Alabama". Journ. Paleont. 15, p. 567-611, figs. 75-82. Tulsa, Oklahoma.
- Toumarkine, M. & Luterbacher, H.P. (1985): "Paleocene and Eocene Planktonic foraminifera. In Bolli, H.M., Saunders, J.B. and Perch Nielsen (Eds) plankton stratigraphy". Cambridge Earth Science ser., p. 87-154.
- Tromp, S.W. (1942): "The micro-fauna of the basal Eocene and Upper Cretaceous section of Romandig Well no. 2 (S.E. Turkey)". Turkey, Maden Tetkik ve

- Arama Enstitüsü, Yayınlarından (Mining Research and Exploration Institute of Turkey, Publ.), pp. 113-125.
- Van der Zwaan, G.J.; Jorissen, F.J. & de Stigter, H.C. (1990): "The depth dependency of planktonic/benthic foraminiferal ratios. Constraints and applications". *Marine Geology* 95, 1-16.
- Van Morkhoven, F.P.C.M.; Berggren, W.A. & Edwards, S.A. (1986): "Cenozoic cosmopolitan deep-water benthic foraminifera". *Bull. Centr. Rech. Expl. Prod. Elf-Aquitaine, Mem.*, 11: 421 pp.
- Viotti, G. & El-Demerdash, G. (1968): "Studies on Eocene sediments of Wadi Nukhul area, East Coast Gulf of Suez". 3rd African Micropaleont. Colloq. Cairo, 21 pp, 3 pls.
- Vogler, J. (1941): "Ober-Jura und Kreide von Misol (Niederlaendisch-Ostindien)". *Palaeontographica, Suppl.* 4, Abt. 4, p. 246-293, pl. 19-24.
- Voorwijk, G.H. (1937): "Foraminifera from the Upper Cretaceous of Habana, Cuba". *Koninkl. Nederlandse Akad. Wetensch., proc., Sec., V.* 40, p. 190-198, 3 pl.
- Wanner, J. (1902): "Die Fauna der obersten weissen Kreide der Libyschen Wüste". *Palaeontologica, Bd.* 30, 91-151, Stuttgart.
- White, M.P. (1928a): "Some index foraminifera from the Tampico Embayment of Mexico". *Journ. Paleont.*, 2, (3), p. 177-215, pls. 27-29.
- White, M.P. (1928b): "Some index foraminifera of the Tampico embayment area of Mexico, pt. 2". *Hjour. Paleontology*, 2, p. 280-317, pls. 38-42.
- Whittaker, J.E. (1980): "Revision of *Plummerita* Brönnimann (Foraminiferida) and new Maastrichtian species from Ecuador". *Bull. Br. Mus. nat. Hist. (Geol.)* 34(4), 289-297.
- Wise, S.W. & Wind, F.H. (1977): "Mesozoic and Cenozoic calcareous nannofossils recovered by D.S.D.P. Leg 36 drilling on the Falkland Plateau, Southwest Atlantic". *Initial Reports of the Deep Sea Drilling Project*, 269-491.

- Wylie Paog, C. & Commeau, A. (1995): "Plaeocene to Middle Miocene Planktic Foraminifera of the southwestern Salisbury embayment, Virginia and Maryland: Biostratigraphy, Allostratigraphy and Sequence Stratigraphy". J. Fo. Res. v. 25, no. 2, p. 134-155.
- Yassini, I. (1979): "The littoral system ostracodes from the bay of Bou Ismail, Aliers, Algeria". Rev. Esp. Micropal. 11, 3, 353-416, Madrid.
- Youssef, M.I. (1949): "Stratigraphical studies in Kossier area ". Ph.D. Thesis, Alexandria Univ. Egypt, 146 pp. Alexandria.
- Youssef, M.I. (1957): "Upper Cretaceous rocks in Kosseir area". Bull. Inst. desert. Egypt., 7(2): 35-54.
- Zittel, K. (1883): "Beitrag zur Geologie und Palaeontologie der Libyschen Wuste und angrenzenden Gebiete von Aegyptens geolog. Teil". paleontographica, 30, (3), p. 1-147, Kassel.