

References

REFERENCES

- Aboul-Hag, S.A.T.A. (1985). Studies on the helminth parasites of some fishes from Sharkiya Governorate. M.Sc. Thesis, Department of Zoology, Faculty of Science, Zagazig University, Egypt.
- Aboul-Hag, S.A.T.A. (1990). Studies on the helminth parasites of some fishes from the Red Sea. Ph.D. Thesis, Department of Zoology, Faculty of Science, Zagazig University, Egypt.
- Acena, S.P. (1941). Preliminary notes on a trematode with two yolk reservoirs and a new species of *Lecithochirium*. Philippine. J. Sci., 75:285-289.
- Agrawal, V. (1964). On some new trematodes from water fishes of Lucknow. Indian J. Helminthol., 16:82-99.
- Agrawal, V. (1966). Studies on some trematode parasites of Fresh-water fishes from Lucknow. Ann. Parasitol. Hum. Comp., 41:217-231.
- Agrawal, V. (1967). Four trematodes parasites (Plagiorchiidae, Lühe, 1901) from reptiles of Lucknow. Rev. Biol. Trop., 14:133-135.
- Agrawal, P.K. and Agrawal, S.M. (1981). Life history of *Haplorchoides vacha* sp. nov. from the intestine of *Eutropüchthyes vacha*. Riv. Parasitol., 42:185-190.
- Ahmed, J. (1980). On five new trematodes of the genus *Uterovesiculurus* Skrjabin et Guschankaja, 1954 (Trematoda, Hemiuridae) from marine fishes of the Bay of Bengal. Acta Parasitol. Polonica, 27:415-422.
- Ahmed, J. (1981). Studies on digenetic trematodes from marine fishes from the Bay of Bengal. Part XVI. Riv. Parasitol., 47:403-413.

- Ahmed, J. (1985). Studies on digenetic trematodes of fishes from the Arabian Sea, Off the Panjim Coast, Goa, India, part 48. *Rev. Iberi. Parasitol.*, 45:185-194.
- Akhmerov, A.A. (1960). New trematodes from fish in the Amur river. *Helminthologia*. 2:286-294.
- Akhmerov, A.A. (1963). Two near far Eastern Bucephalid Species (Trematoda, Bucephalidae). In [Helminthes of man, animals and plants of their control papers on helminthology], presented to Academician K.I. Moscow: Izdatelstvo Akad. Nauk. SSSR, PP. 126-129.
- Amato, J.F.R.(1983). Digenetic trematodes of percoid fishes of Floianopolis Southern Brazil-Pleorchidae, Didymozoidae and Hemiuridae, with the description of three new species. *Rev. Brazil. Biol.*, 43:99-124.
- Annerreaux, R.F.(1947). Three new trematodes from marine fishes of California. *Trans. Amer. Micro. Soc.*, 66:249-255.
- Balozet, L. and Sicart, M. (1960). *Hemiurus communis* Odhner, Parasite de l'anguille. *Bulletin de la Societe d'Histoire Naturelle de Toulouse*, 59:105-110.
- Bashirulla, A.K.M. and D'Silva, J. (1973). Two new parasites of the genus *Lecithocladium* Lühe, 1901 (Family: Hemiuridae) Japanese J. Parasitol., 22:108-110.
- Baugh, S.O. (1957). Contributions to our knowledge of digenetic trematodes II. *Proc. Nat. Acad. Sci. (India)*, 26:295-313.
- Bear, J.G. (1924). Description of a new genus of *Lepadematidae* with a systematic essay on the family. *Parasitology*, 16:22-31.
- Belding, D.L. (1965). Textbook of parasitology. Third edition, Appleton-Century, Crofts, New York, 1374 PP.

- Beverley-Burton, N. (1962). Some trematodes from *Clarias* spp. in the Rhodesia, including *Allocreadium mazoensis* n.sp and *Eumaseia bangweulensis* n.sp. and comments on the species of the genus *Orientocreadium*. Tubangui, 1931. Proc. Helminthol. Soc. Wash., 29:103-115.
- Bhalerao, G.D. (1926). On the trematode parasites of a water snake *Tropidonotus piscater*. Parasitology, 18:4-13.
- Bhalerao, G.D. (1936). Studies on the helminthes of India. Trematoda III. Helminthol., 14:163-228.
- Bhalerao, G.D. (1937). Studies on the helminthes of India Trematoda. IV. J. Helminthol., 15:97-124.
- Bhalerao, G.D.(1940). Observation on the anatomy of *Acanthostomum burminis*. Indian Vet. Sci., 10:94-97.
- Bilqees, F.M. (1976). Two trematodes of the genus *Proisorhynchus* Odhner, 1905 (Bucephalidae) including a new species *P. erumenis* from the fish *Psettodes erumenis*, B. 4 Schn) of the Karachi Coast. Norwegian J. Zool., 24:345-348.
- Bilqees, F.M. (1977). *Plagioporus heterorchis* sp.n. (Trematoda: Opecoelidae) from the fish *Pomadasys olivaceum* (Day) of Karachi Coast. Pakistan J. Sci. Ind. Res., 20:258-260.
- Bilqees, F.M. and Khan, A. (1988). *Plagioporus gonii* nov. spec. (Trematoda-Opecoelidae) from the fish *Labeo gonius* of Kalri Lake, Sind, Pakistan. Angewante Parasitol., 29:37-41.
- Bilqees, F.M. and Nighat, H.Y. (1985). A new trematode *Lecithocladium thynensis*, new species (Trematoda: Hemiuridae) from the fish *Thynus* sp., off Karachi Coast, Pakistan J. Zool. 17:211-214.

- Boraey, F.A. (1974). Biological study of the family Mugilidae (Grey Mulletts) in Lake Qarun. Ph.D. Thesis, Faculty of Science, University of Ain Shams, Egypt.
- Boulenger, G.A. (1907). Zoology of Egypt. The fishes of the Nile. Hugh Press Ltd., London.
- Bray, R.A. (1987). A revision of the family zoogonidae Odhner, 1902 (Platyhelminths: Digenea): Subfamily Lepidophyllinae and comments on some aspects of biology. Syst. Parasitol., 9:83-123.
- Bray, R.A. (1989). Some helminth parasites of marine fishes of South Africa. Family: Opecoelidae (Digenea). J. Nat. Hist., 21:1049-1075.
- Bychowsky, B.E. and Dubinina, M.N. (1954). Systematics of digenetic trematodes belonging to the family Acanthocolpidae. Lühe, 1909. Zool.Zh., 33:788-793.
- Byrd, M.A. (1963). Helminth parasites of Antarctic vertebrates. part I. Digenetic trematodes of marine fishes. Proc. Helminthol. Soc. Wash., 30:129-148.
- Caballero, C.E. and Caballero, R.G. (1970). Work on the Howard A. Winter Collection of trematodes of marine fish from the Mexican and U.S.A. Pacific Coasts. An. Esc. Nac. Cienc. Biol. Mex., 18:265-266.
- Cable, R.M. (1952). On the systematic position of the genus *Deropristis* of *Dihemistephanus sturionis* Little, 1930 and of a new digenetic trematode from a sturgeon. Parasitology, 42:85-91.
- Cable, R.M. (1955). Taxonomy of some digenetic trematodes from sturgeons. J. Parasitol., 41:441.

- Cable, R.M. and Hunninen, A.V. (1942). Studies on *Deropristis inflata* (Molin, 1859) its life history and affinities to trematode of the family Acanthocolpidae. Biol. Bull. Woods Hole, 82:292-312.
- Carvalho-Varela, M. and Cunha-Ferreira, V. (1987). Helminth parasites of common sole, *Solea solea* and the Senegalese sole, *Solea senegalensis* on the Portuguese Continental Coast. Aquaculture, 67:135-138.
- Catalano, P.A. and Etges, F.J. (1981). *Plagioporus gyirinophili* n.sp. (Trematoda: Opecoelidae) from *Gyrinophilus porphyriticus duryi* and *Pseudtriton ruber* (Caudata: Plethodontidea). Proc. Helminthol. Soc. Wash., 48:198-201.
- Chatterji, R.C. (1933). On the trematode parasites of Rangoon siluroid fish *Clarias batrachus*. Bull. Acad. Sci. Allahabad, 3:33-40.
- Chauhan, B.S. (1943). Trematodes from Indian marine fishes part II. On some trematodes of the gasterostome family Bucephalidae (Baun, 1883) Poche, 1907 with description of four new species Proc. Indian Acad. Sci., 17:97-117.
- Chauhan, B.S. (1945). On some trematodes of the family Hemiuridae Lühe, 1901 with description of six new forms. Proc. Ind. Acad. Sci., 21:160-175.
- Chauhan, A.S. and Khoche, B.Y. (1975). Studies on trematode parasites of Madhya Pradesh, Pt. V. (Digenea: family Buephalidae), report on two new species of *Bucephalopsis* Diesing, 1855. Zool. Soc. India., 369-372.
- Chen, H.T. (1949). Systematic consideration of some Heterophid trematodes in the subfamilies: Haplorchinae and Stephanochasminae. Ann. Trop. Med. Parasitol., 43:304-312.

- Ciurea, I. (1933): "Les vers parasites de l'homme, des mammifères et des Oiseaux provenant des poissons du Danube et de la Mer Noire. Premier mémoire. Trematodes, Famille Heterophyidae Odhner, avec un essai de classification des trematodes de la superfamille Heterophyoidea Faust". Arch. Rum. Pathol. Exper. Micro., VI(1/2), 1-134.
- Cobbold, T.S. (1861). Synopsis of Distomid. Proc. Linn. Soc. Land., 5, 1-56; 117-127.
- Dawes, B. (1946). The trematoda with special reference to British and other European forms. Camb. Univ. Press, 644 PP.
- Dawes, B. (1947). The trematoda of British Fishes (London, Ray Society, No. 131), 364 PP.
- Dayal, J. (1938). Studies on the trematode parasites of fishes. A new trematode *Neoganada barakankiae* (nov. gen., nov. sp.) from *Clarias batrachus*. Proc. Indian Acad. Sci. Sec. B., 7:132-137.
- Dayal, J. (1948). Trematode parasites of Indian fishes Part I. New trematodes of the family Bucephalidae Poche, 1907 Indian J. Helminthol., 1:47-62.
- Dayal, J. (1949). Trematode parasite of Indian fishes. Part II. Ind. J. Helminthol., 1:93-120.
- Dayal, J. and Gupta, S.P. (1951). A redescription of the trematode parasite, *Bucephalopsis karvei* Bhalerao, 1937. Proc. Indddd. Sci. Congress, 83th, Part III, P. 211.
- Diesing, K.M. (1855). Revision der Myzhelminthen. Wiener sitzungsberichte, math-naturw. KL. Vol. 3, No. 2.
- Dogiel, V.A. (1962). General parasitology (Transplanted by Z. Kabatata) Oliver and Boyd, London.

- Dollfus, R.P. (1950). Trematodes recoites au Congo belge par le proesseur Paul Brien (Mai, Aout 1937). Ann. Mus. R. "Congo belge, Ser. 4, Zoologie, 1:1-136.
- Dujardin, F. (1845). Histoire naturelle des helminthes ou vers. in'estinaux, XVI, 659 PP., Paris.
- Durio, W.O. and Manter, H.W. (1968). Some digenetic trematodes of marine fishes of New Caledonia Part I. Bucephalidae, Monorchiidae, and some smaller families. Proc. Helminthol. Soc. Wash., 35:143-153.
- Dwivedi, M.P. (1966). On the genus *Astiotrema* Looss, 1900. Indian J. Helminthol., 18:33-43.
- Echmann, F. (1932). Beitrage zur kenntnis trematoden familie Bucephalidae. Z. Parasitol., 5:94-11.
- Ejsmont, L. (1930). *Astiotrema emydis* n.sp. ein trematode aus Emys *orbicularis*. Bull. Int. Acad. Pol. Sci., 2:405-417.
- El-Naffar, M.K. (1970). Studies on the parasites of Nile fishes in Assiut Province of Egypt. Ph. D. Thesis, Assiut University, Egypt.
- El-Naffar, M.K. (1980). Studies on the heterophyid cercaria from Assiut Province, Egypt II. The life cycle of *Haplorchoides cahirinus* (Looss, 1896) Chen, 1949. J. Egypt. Soc. Parasitol., 10:117-125.
- El-Naffar, M.K., Saoud, M.F. and Hassan, I.M. (1983). A general survey of the helminth parasites from some fishes from Lake Nasser at Aswan. A.R. Egypt. Assiut Vet. Med. J., 11:141-148.
- El-Serafy, S.S. anmd Abdel-Baky, T.E. (1990). Fish catch characteristics in Lake Manzllah. Egypt. J. Appl. Sci., 5:505-521.

Eysenhardt, H.W. (1829). Eingiges uber Eingeweide Würmer Verh Ges. Naturf F.R. Berlin, 1:144-152.

Fantham, H.B. (1938). *Lecithostaphylus spondylisomae* n.sp. a trematode parasite of the Hottentot fish, *Spondylisoma blochii*, found in South African waters. Trans. Roy. Soc. South Africa, 26:387-393.

Farooq, M. and Khanum, Z. (1980). Two new species of the genus *Lecithocladium* Lühe, 1901 (Digenea: Hemiuridae) from marine fishes of Karachi Coast, Pakistan. Pakistan J. Zool., 12:221-224.

Fischthal, J.H. (1973). Three digenetic trematodes of *Clarias mossambicus* peters (Clariidae) from Ethiopia Proc. Helminthol. Soc. Wash., 40:166-167.

Fischthal, J.H. and Kuntz, R.E. (1963a). Trematode parasites of fishes from Egypt. Part III. Six new Hemiuridae. Proc. Helminthol. Soc. Wash., 30:28-91.

Fischthal, J.H. and Kuntz, R.E. (1963b). Trematode parasites of fishes from Egypt. Part IV. A redescription of *Monascus typicus* (Odhner, 1911) Fellodistomidae Proc. Helminthol. Soc. Wash., 30:177-182.

Fischthal, J.H. and Kuntz, R.E. (1963c). Trematode parasites of fishes from Egypt. Part V. Annotated record of some previously described forms. J. Parasitol., 49:91-98.

Fischthal, J.H. and Kuntz, R.E. (1963d). Trematode parasites of fishes from Egypt. Part VII. *Orientocreadium batrachoides* Tubangui, 1931 (Plagiorchiidae) from *Clarias lazera*, with a review of the genus and related forms. J. Parasitol., 49:451-464.

Fischthal, J.H. and Kuntz, R.E. (1964). Digenetic trematodes of fishes from Baldwin Island, Philippines. Part II. Five Opecoelidae, including three new species. Proc. Helminthol. Soc. Wash., 31:40-46.

Fischthal, J.H. and Kuntz, R.E. (1965). Digenetic trematodes of Amphibians and Reptiles from Malaysia. *Proc. Helminthol. Soc. Wash.*, 32:124-136.

Fischthal, J.H. and Nasir, P. (1974). Some digenetic trematodes from fresh-water and marine fishes of Venezuela. *Norweg. J. Zool.*, 22:71-80.

Fischthal, J.H. and Thomas, J.D. (1968). Digenetic trematodes of some fresh-water and marine fishes from Ghana. *Proc. Helminthol. Soc. Wash.*, 35:126-140.

Fischthal, J.M. and Thomas, J.D. (1970). Digenetic trematodes of marine fishes from Ghana: Family Lepocreadiidae. *J. Helminthol.*, 44:365-386.

Fischthal, J.M. and Thomas, J.D. (1972). Digenetic trematodes of marine fishes from Senegal. *Bulletin de l'Institut fondamental d'Afrique Noire, Ser. A. Sci. Nat.*, 34:292-322.

Gaevskaya, A.V. (1975). Two new species of trematodes - *Gonocera macrouri* n.sp. and *Hemiurus macrouri* n.sp. from *Macrourus rupestris* in the north eastern Atlantic. *Parasitology*, 9:475-459.

Gaevskaya, A.V. and Rdyuk, G.V. (1988). New and rare species of trematodes from benthic fish in the south-west Atlantic. *Zest. Zool.*, No. 5:11-15.

Gibson, D.I. (1986). *Podocotyle araii* sp. nov. (Digenea: Opecoelidae) from *Sebastes spp.* off the Coast of British Columbia, with a key to the species of *Podocotyle* Dujardin, 1845, occurring off the Pacific Coast of North American *J. Nat. Hist.*, 20:735-743.

Gibson, D.I. and Bray, R.A. (1984). On *Anomalotrema*. Zukov, 1957, *Pellamyzon* Montgomery, 1957 and Opecoelidae, with a description of *Anomalotrema Kolae* sp. nov. from North Atlantic Waters. *J. Nat. Hist.*, 18:949-964.

- Gibson, D.I. and Bray, R.A. (1986). The Hemiuridae: Digenea of fishes from the North-East Atlantic. Bull. Bri. Mus. (Nat. Hist.), Zool., 51:125.
- Gomes, D.C.; Fabio, S.P. and Rolas, F.J.T. (1974). Contribution to the knowledge of parasites of fish from the Guanabara littoral 3 J Memorias do Instituto Oswaldo. Cruz., 72:9-19.
- Grabda-Kazubska, B. (1959a). *Astiotrema trituri* sp.n. (Trematoda: plagiiorchiidae) from the small intestine of *Triturus vulgaris*. Bull. Acad. Polan. Sci., 11-Sex. Sci. Biol., 7:17-21.
- Grabda-Kazubska, B. (1959b). The life cycle of *Astiotrema trituri* B. Grabda, 1959 (Trematoda-Plagiiorchiidae). Acta Parasitol. Polonica, 7:489-498 (English text; Polish Summary).
- Grabda-Kazubska, B. (1961). A redescription of *Metaleptophallus gracillimus* (Lühe, 1909) Yamaguti, 1958 (Trematoda: plagiiorchiidae) Acta. Parasitol., Polonica, 9:101-107.
- Gu, C. and Shen, J. (1976). Report on some gasterostomatous trematodes (Famiy: Bucephalidae Poche, 1907) from marine fishes in Dong Hai and Nan Hai, China. Acta. Zool. Sinca, 22:371-384.
- Gu, C. and Shen, J. (1978). Some hemiruid trematodes (sub-family: Dinurinae Looss, 1907) from marine fishes of economic importance of China. Acta Zool. Sinca, 24:373-387.
- Gupta, S.P. (1951). Trematode parasites of Indian fishes. Three new trematodes of the subfamily: Leptohallinae Dayal, 1938 from fresh-water fishes of U.P. Indian J. Helminthol., 3:29-40.
- Gupta, S.P. (1955). Trematode parasites of freshwater fishes. Indian J. Parasitol., 5:42-56.

- Gupta, N.K. and Kumari, A. (1977). On one already known Bucephalid trematode from *Xenentoden cancila* (a fresh-water fish) from Ropr (Punjab, India). Res. Bull. Panjab University, New Series, 24:113-116.
- Gupta, N.K. and Mehrotra, V. (1970). A new hemiurid parasite of the genus *Lecithocladium* Lühe, 1901, with revised key to Indian species of the genus. Res. Bull. Panjab. Univ. Sci., 21:73-76.
- Gupta, N.K. and Sayal, A. (1979). Digenetic trematodes of marine fishes from Laccative Islands (Arabia Sea) and Goa. India Folia Parasitol., 26:221-229.
- Gupta, N.K. and Sehgal, S.K. (1967). Two new parasites of digenetic trematodes from some Indian marine food fishes. Indian J. Helminthol., 18, Suppl. Appendix, A, P. III.
- Gupta, N.K. and Sehgal, S.K. (1971). Some allocradiid trematodes from marine food fishes of India. Res. Bull. Panjab Univ., 21:267-275.
- Gupta, P.C. and Govind, H. (1985a). Three species of trematode parasites of the genus *Haplorchoides* Chen, 1949, from freshwater fishes of Kanpur, India. Ind. J. Parasitol., 9:35-39.
- Gupta, P.C. and Govind, H. (1985b). Studies on trematode parasites of fishes, on four new species of the genus *Lecithocladium* Lühe, 1901 from marine fishes of Puri Coast, Orissa. Kanpur Univ. Res. J., 2:31-39.
- Gupta, P.C. and Govind, H. (1987). On three new species of the genus *Plagioporus* Stafford, 1904 (Family: Opecoelidae Ozaki, 1925) from fishes of Orissa: Kanpur Univ. Res. J., 5:1-6.
- Gupta, P.C. and Gupta, S.P. (1979). On two avian trematode parasites (Family: Heterophyidae Odhner, 1914) from Kanpur Univ. Ind. J. Helminthol., 49:73-78.

- Gupta, P.C. and Gupta, V.C. (1986). Two new piscine hemiurids of the genus *Uterovesiculurus* Skrjabin et Guschanskaja, 1954 from Orissa, Indian Proc. Parasitol., No. 2:1-5.
- Gupta, P.C. and Gupta, V.C. (1987a). Two species of the genus *Proisorhynchus* Odhner, 1905 (Digenea: Bucephalidae) from marine fishes of Puri Coast, Orissa. Ind. J. Helminthol., 38:112-118.
- Gupta, P.C. and Gupta, V.C. (1987b). Three species of the genus *Lecithocladium* Lühe, 1901 (Digenea: Hemiuridae) from Bay of Bengal, Off Puri Coast, Orissa. Kanpur Univ. Res. J., (Science), 4:1-7.
- Gupta, P.C. and Singh, R.B. (1982a). Studies on digenetic trematodes of Indian fishes and Birds. Kanpur Univ. Res. J., 3:23-36.
- Gupta, P.C. and Singh, R.B. (1982b). Two new trematodes of the genus *Lecithocladium* Lühe, 1901 from a marine fish of Puri Coast, Bay of Bengal. Ind. J. Parasitol., 6:325-329.
- Gupta, P.C. and Singh, R.B. (1985). Two new species of digenetic trematodes of fish in India. Pakistan J.Zool., 17:329-334.
- Gupta, V. and Puri, M. (1981). Six new species of the genus *Lecithocladium* Lühe, 1901 (Hemuridae Lühe, 1901) from marine fishes of Puri, Orissa. Indian J. Helminthol., 3:58-78.
- Gupta, V. and Puri, M. (1986). On some trematode parasites from marine fishes of Puri, Orissa. Indian J. Helminthol., 37:49-62.
- Gupta, V. and Tiwari, M. (1985). Trematode parasites of marine fishes. Indian J. Helminthol., 35:93-111.

- Gupta, V. and Saxena, A.M. (1988). On a new trematode *Astiotrema manteri* sp. nov. from *Kachuga intermedia* (Boulenger) from Lucknow and a review of the genus *Astiotrema* (Looss, 1898) Looss, 1900. Indian J. Helminthol., 39:114-112.
- Gupta, R.C. and Gupta, S.P. (1978). Five new trematodes of the genus *Lecithocladium* Lühe, 1901 (Hemiuridae) from marine fishes. Folia parasitol., 25:223-229.
- Gupta, S.P. and Tandon, V.L. (1985). On some trematode parasites from marine fishes of Puri, Orissa. Indian J. Helminthol., 36:143-161.
- Gupta, V. and Ahmed, J. (1977). Digenetic trematodes of marine fishes of Bay of Bengal, Puri, Orissa. Riv. Parasitol., 38:181-191.
- Gupta, V. and Ahmed, J. (1979). Digenetic trematodes of marine fishes VIII. On four new species of the genus *Haplospalchnus* Looss, 1902 from marine fishes of Puri, Orissa. Helminthology, 16:185-193.
- Gupta, V. and Johana, A. (1979). On two new trematodes of the genus *Astiotrema* (Plagiorchiidae). Indian J. Helminthol., 30:35-41.
- Hafeezullah, M. (1971). Opecoelid trematodes of marine fishes of India. Parasitology, 62:321-329.
- Hafeezullah, M. (1975). Two new trematodes (Digenea: Hemiuridae) of marine fishes from East Coast of India. Zool. Soc. India., 203-210.
- Hafeezullah, M. and Siddiqi, A.H. (1970). Digenetic trematodes of marine fishes of India Part I. Bucephalidae and Cryptogonimidae. Indian J. Helminthol., 22:1-22.
- Hassan, E.A. (1980). Studies on certain parasites of some fishes from Lake Nasser. M.Sc. Thesis, Faculty of Science, Assiut University, Egypt.

- Hassan, S.H. (1976). Studies on some parasitic worms of marine fishes in Egypt, Ph.D. Thesis, Mansoura University, Egypt.
- Hendrix, S.S. (1973). *Plagioporus hypentelii* sp.n. (Trematoda: Opecolidae) from the Kogsucker, *Hypentelium nigricans* (Lesueur) Osteichchys: Casostomidae). Proc. Helminthol. Soc. Wash., 40:144-146.
- Imam, E.A.R. (1971). Morphological and biological studies on the enteric helminthes infesting some of the Egyptian Nile fishes. Ph.D. Vet. Thesis, Cairo University, Egypt.
- Issaitschikow, I.M. (1928). Zur kenntnis der parasitischen Würmer einiger Gruppen von Wirbeltieren der russischen Arktis. Trudy Morsk Nauch. Inst. Moskva, 3:5-79.
- Johnson, W.F. (1949). A new species of trematoda, *Podocotyle gibbonsia*, from tide pool fishes of Monterey Bay, California. Trans. Amer. Micro. Soc., 68:107-109.
- Johari, L.N. and Smyth, J.D. (1956). A histochemical approach to the study of helminth morphology. Parasitology, 46:107-66.
- Karyakarte, P.P. (1967). *Acanthostomum (Atrophocaecum) alii* sp.n. from the snake *Elaphe helena* in India. J. Parasitol., 53:587-588.
- Karyakarte, P.P. (1968). *Acanthostomum (Gymnatrema) pambanense* sp.n. (Trematoda: Acanthostomatidae) from the fish, *Therapon puta* in India. Riv. Parasitol., 28:195-189.
- Khalil, L.F. (1959). On a new trematode *Astiotrema sudanensis*, n.sp. from a fish water turtle in the Sudan. J. Helminthol., 33:263-266.
- Khalil, L.F. (1961). On a new trematode *Orientocreadium lazeri* sp.nov. from a fresh-water fish *Clarias lazera* in the Sudan. J. Helminthol., 35:259-262.

- Khalil, L.F. (1963). *Acanthostomum gymnarchi* Dollfus, 1950, with notes on the genera, *Acanthostomum* Looss, 1900, *Atrophocaecum*. Bhalerao, 1940, *Gymnatrema* Morozov, 1955, and *Haplocaecum* Simha, 1958. J. Helminthol., 37:207-214.
- Khalil, L.F. (1971). Check list of the helminth parasites of African fresh-water fishes. Tech. Commun. Commonw. Inst. Helminthol., No. 42:88 PP.
- Khalil, L.F., and Thurston, J.P. (1973). Studies on the helminth parasites of fresh-water fishes in Uganda including the description of two new species of digeneans. Rev. Zool. Bot. Afri., 87:210-225.
- Kohn, A. (1961). Um novo *Proisorhynchus* parasito de Garrup sp. (Trematoda: Bucephaliformes). Atas Soc. Biol. Rio de Janeiro, 5:46-49.
- Kohn, A. (1962). Nota previa sobre uma nova especie do genero *Bucephalopsis* Diesing, 1855 (Trematoda, Bucephaliformes). Atas Soc. Biol. Rio de Janeiro, 6:14-18.
- Kohn, A. and Bührnheim, P.F. (1964). "Um novo hospedeiro e nova distribuicao geografica para *Myosaccium ecaudo* Montgomery, 1957 (Trematoda, Hemiuridae). Atas Soc. Biol. Rio de Janeiro, 8:50-52.
- Kruse, G.O.W. (1977). Some digenetic trematodes from fishes of Bering Sea with the description of *Proisorhynchus mizellei* sp.n. (Bucephalidae) and *Pseudopecoelus nossamani* sp.n. (Opecoelidae). Proc. Helminthol. Soc. Wash., 44:73-76.
- Lacaze-Duthiers, F.J.H. de. (1854). Memoire sur le Bucephale Haime, parasite des hitres et des bucardes. Ann. de Sc. Nat., 1:294-302.
- Lal, A.K. and Prasad, D. (1980). On *Astiotrema Siddiqui* n.sp. (Trematoda): Plagiorchiidae, (Astiotreminae) from *Rana cyanopchtis*. Indian J. Helminthol., 32:93-95.

- Lamloom, D.A.M. (1987). A general survey on the helminth parasites of some fish from Fayoum Governorate. Arab Republic of Egypt, M.Sc. Thesis, Faculty of Science, Ain Shams University.
- Lamloom, D.A.M. (1990). Studies on the helminth parasites of some fishes from Inland Water In Egypt. Ph.D. Thesis, Faculty of Science, Cairo University, Bani Suef Branch, Egypt.
- Lamothe-Arugumedo, R. and Rodriguez, Poncia, No. G. (1985). Revision of subfamily Acanthostominae Nicoll, 1914, and the establishment of two new genera. *Anales Inst. Biol. Uni. Nac. Autonoma de Mexico, Zool.*, 56:301-322.
- LaRue, G.R. (1926). Studies on the trematode family Strigeidae (Holostomidae). No. II. Taxonomy. *Trans. Amer. Microbiol. Soc.*, 645:11-19.
- Latife, A.F.A. (1974). Fisheries of Lake Nasser Aswan Regional Planning. Lake Nasser Development Centre, Aswan, Egypt.
- Layman, E.M. (1930). Parasitic worms from the fishes of Peter the Great Bay. *Bull. Pacif. Scient. Fish. Res.*, 3:1-120.
- Lebedev, B.I. (1968). New trematodes of pelagic fish of the order perciformes in the pacific basin. [Helminths of animals of the Pacific Ocean]. Moscow: Izdat. Nauka, 4:156-167.
- Linton, E. (1900). Fish parasites collected at Woods Hole in 1898. *Bull. U.S. Fish Comm.* (1899), 19:267-304.
- Linton, E. (1910). Helminth fauna of the Dry Tortugas, II. Trematodes. *Carneg. Inst. Wash. Publ.*, No. 133, 98 PP.
- Linton, E. (1940). Trematodes from fishes mainly from the Woods Hole region, Massachusetts. *Proc. U.S. Nat. Mus.*, 88:1-172.

- Lloyd, L.C. (1938). Some digenetic trematodes from puge sound fish. J. Parasitol., 24:103-133.
- Long, S. and Lee, W.C. (1964). Worm parasites from Taihu fishes: Digenetic trematodes VI Bucephalidae Poche, 1907, with a description of two new genera and four new species and a revision of the family. Acta. Zool. Sinca, 16:567-580.
- Looss, A. (1896). Recherches sur la fauna parasitaire de l'Egypt. Premiere Partie: Mem. Inst. Egypt, 3:252 PP.
- Looss, A. (1899). Weitere Beitrage zur kenntnis der Trematoden-fauna Aegyptens, Zugleich Versuch einer naturlichen Gliederung des Genus *Distomum*. Retzius. Zool. Jahrb. Syst., 12:521-784.
- Looss, A. (1900). Nachtragliche Bemerkungen zu den Namen der von mir vorgeschlagenen Distomidengattungen. Zool. Anz., 23:601-608.
- Looss, A. (1901). Über die Fasciolidengenera *Stephanochasmus*, *Acanthochasmus* and einige and andere. Ctbl. Bakt. I, 29:595-606; 628-634; 654-661.
- Looss, A. (1902). *Monorchis* (Monticelli) und *Haplospalchnus* n.g. zur Kenntnis der Trematoden fauna des Triester Hafens II. Ctbl. Bakt., 32:115-122.
- Looss, A. (1907). Beriträge zur Systematik der Distomen. Zur Kenntnis der Familie Hemiuridae Zool. Jahrb. Syst. , 26:63-180.
- Lühe, M. (1901). Über *Monostomum Orbiculare*. Ctbl. Bakt., 29:49-60.
- Lühe, M. (1906). Report on the trematode parasites from the marine fishes of Ceylon. Pearl Oyster Fish. Rep. Pt. 5:97-108.

Lühe, M. (1909). Parasitische Plattwürmer. I. Trematoden Süßwasserfauna Deutschl., Heft, 17:215 PP.

Machida, M. (1980). Hemiurid trematodes of *Kyphosus* collected around cape shionomisaki, Kii, Peninsula. Mem. Nat. Sci. Mus., Tokyo, No. 13, 113-120.

Madhavi, R. (1974). Digenetic trematodes from marine fishes of waltair coast, Bay of Bengal. Family Bucephalidae. Riv. Parasitol., 35:189-199.

Madhavi, R. (1979a). Digenetic trematodes from marine fishes of waltair coast, Bay of Bengal, Family: Zoogonidae. Riv. Parasitol., 40:249-259.

Madhavi, R. (1979b). Digenetic trematodes from marine fishes of waltair coast, Bay of Bengal, Families Haplosporididae. Riv. Parasitol., 40:237-248.

Maillard, C., and Lambert, M. (1978). *Plagioporus novellus* n.sp. (Trematoda: Opecoelidae) a parasite of *Conger conger*. Ann. Parasitol. Huma. Com., 53:367-371.

Mamaev, I.L. (1965). Helminth of fish from Bering Sea. Parasitic worms of domestic and wild animals. Akad. Nauk. SSSR: 1 68-188.

Manter, H.W. (1925). Some marine fish trematodes of marine. J. Parasitol., 2:11-18.

Manter, H.W. (1926). Some North American fish trematodes. Biol. Monogr., 10:127-264.

Manter, H.W. (1934). Some digenetic trematodes from deep-water fish of Tortugas, Florida. Publication (435), Carnegie Inst. wash., Papers Tortugas Lab., 28:257-345.

- Manter, H.W. (1937). The status of the trematodes genus *Deradena*. Linton with a description of six species of *Haplospilanchnus* Looss, 1902 (Trematoda) Rabot. Helminth. Skrjabin. Skrjabin Jubil., 1:381-387.
- Manter, H.W. (1940a). Digenetic trematodes of fishes from the Galapagos Islands and the neighboring pacific. Rep. Allan Hancock Pacif. Exp., 2:325-497.
- Manter, H.W. (1940b). Gasterostomes (Trematoda) of Tortugas, Florida. No. 524, PP. 1-19. (Papers from the Tortugas Laboratory, Vol. 33).
- Manter, H.W. (1947). The digenetic trematodes of marine fishes of Tortugas. Florida Amer. Midl. Nat., 38:257-416.
- Manter, H.W. (1953). Two new species of Prosorhynchinae (Trematoda: Gasterostomata) from the Fiji Islands, G.S. Thapar Commem., PP. 193-200.
- Manter, H.W. (1954). Some digenetic trematodes from fishes of New Zealand. Trans. Roy. Soc. New Zealand, 82:475-568.
- Manter, H.W. (1963). Studies on digenetic trematodes of fishes of Fiji II. Families. Lepocreadiidae, Opistholebetidae and Opecoelidae. J. Parasitol., 49:99-113.
- Manter, H.W. and Pritchard, M.H. (1960a). Some hemiurid trematodes from Hawaiiin fishes. Proc. Helminthol. Soc. Wash., 28:87-102.
- Manter, H.W. and Pritchard, M.H. (1960b). Additional hemiurid trematodes from Hawaiiin fishes. Proc. Helminthol. Soc. Wash., 27:165-180.
- Manter, H.W. and Van Cleave, H.J. (1951). Some digenetic trematodes including eight new species from marine fishes of La Jolla, California. Proc. U.S. Nat. Mus., 10:315-339.

- Margolis, L. and Adams, J.R. (1956). Description of *Genolinea onchorhynchi* n.sp. (Trematoda: Hemiuridae) from *Onchorhynchus gorbuscha* in British Columbia with notes on the genus. Proc. U.S. Nat. Mus., 34:373-577.
- Mazza, J. (1963). Some trematodes of fish from the fishing-ground of Berre. Rev. Trav. L'Institut des Peches Mari., 27:441-448.
- McCauley, J.E. (1960). Some hemiurid trematodes of Oregon marine fishes. J. Parasitol., 46:84-89.
- Mikailov, T.K. (1958). Parasitic fauna of *Mugil saliens* Risso of the Caspian Sea. Zool. Zhurnal, 37:373-378.
- Mohamed, M.A. (1978). Studies on certain protozoan and trematode parasites of some Nile fishes. M.Sc. Thesis, Ain Shams Univ., Egypt.
- Molin, R. (1859). Prospectus helminthum, quae in parte secunda prodromi faunae helminthologicae venetae continenture. Sitzungsberg. Akad. Wiss. Wien. Manth. Nat., 33:287-302.
- Montgomery, W.R. (1957). Studies on digenetic trematodes from marine fishes of La Jolla, California, Trans. Amer. Microsc. Soc., 76:13-36.
- Moravec, F. (1976). On two acanthostomatid trematodes *Acantostomum spiniceps* (Looss, 1968) and *A. absconditum* (Looss, 1901) from African bagrid fishes. Folia Parasitol., 23:201-206.
- Moravec, F. (1977). Some digenetic trematodes from Egyptian fresh-water fishes. Vës. Cës. Spol. Zool., 41:52-67.
- Moravec, F. and Sey, O. (1989). Some trematodes of fresh-water fishes from North Vietnam with a list of recorded endohelminths by fish hosts. Folia Parasitol., 36:243-262.

- Morozov, F.N. (1955). Basis of trematodology, trematodes of animal and man. Skrajabin, Moscow, 10:243-335.
- Nagaty, H.F. (1937). Trematodes of fishes from the Red Sea, Part I. Studies on the Family: Bucephalidae Poche, 1907, No. 12, 172 PP. Faculty of Medicine, Egyptian University.
- Nagaty, H.F. (1973). Trematodes of fishes from the Red Sea. Parts 1-20. A recapitulation. Bull. Zool. Soc. Egypt., 25:1-13.
- Nagaty, H.F. and Abdel Aal, T.M. (1962a). Trematodes of fishes from the Red Sea. Part 19, on three new species of *Podocotyle* (Family: Allocreadiidae). J. Parasitol., 48:746-747.
- Nagaty, H.F. and Abdel Aal, T.M. (1962b). Trematodes of fishes from the Red Sea, Part 17, on three allocreadiid sp. and one Schistorchid species. J. Arab. Vet. Med. Ass., 22:307-314.
- Nagaty, H.F. and Abdel Aal, T.M. (1969). Trematodes of fishes from the Red Sea part 18, on two new and one known Allocreadiid species. J. Egypt. Vet. Med. Ass., 29:1-5.
- Nahhas, F.M., and Cable, R.M. (1964). Digenetic and Aspidogastrid trematodes from marine fishes of Curacao and Jamaica. Tulane, Stud. Zool., 11:169-228.
- Nasir, P. (1974). Revision of genera *Acanthostomum* and *Telorchis* with a redescription of *Acanthostomum* (*Acanthostomum*) *Scyphocephalum* and *Telorchis deatus*. Riv. Parasitol., 35:1-22.
- Neafie, R.C. and Haley, A.J. (1962). Sex differences in resistance of the mouse *Mus musculus* to infection with rat neomatoda *Nippostrongylus brasiliensis*. J. Parasitol., 48:151 PP.

- Negm-El-Din, M.M. (1987). Some morphological studies on the internal parasites of fish in Delta Nile. M. Vet. Sc. Thesis, Faculty of Vet. Med. Zagazig University, Benha Branch (Moshtohor).
- Nicoll, W. (1909). Studies on the structure and classification of the digenetic trematodes. Quart. J. Microbiol. Sci., 53:391-487.
- Nicoll, W. (1910). On *Gasterostomum tergestinum* stossich. Ann. del Mus. Zool. della R. Universit. d. Napoli, Vol. 3, No. 14.
- Nicoll, W. (1913). New trematode parasites from fishes of the English channel. Parasitology, 5:238-246.
- Nicoll, W. (1914). The trematode parasites of fishes from the English channel. J. Mar. Biol. Assoc. U.K., 10:466-505.
- Nicoll, W. (1915). The trematode parasites of North Queensland. III. Parasites of fishes. Parasitology, 8:22-41.
- Nicoll, W. (1935). "Vermes" Zoll. Rec., LXXI, 1-158 (W.L23823). Cited from Bhalerao, G.D. (1936). Studies on the helminthes of India. Trematoda III. J. Helminthol., 14:163-228.
- Nigrelli, R.F. (1940). Two new species of trematodes from the deep sea scorpion fish, *Scorpaena madurensis* Cuv and Val. Zool. New York, 25:263-268.
- Noor-El-Din, S.N.E.A. (1981). Studies on some parasitic helminthes in some fresh-water fish. M.Sc. Thesis, Tanta University, Egypt.
- Odhner, T. (1902). Mitteilungen zur kenntnis der Distomen. II. 1. Drei neue Distomen aus der Gallenblase von Niffischen. 2. *Deropristis*, n.g.3. *Helicometra* n.g. Allocreadiinarum. Ctbl. Bakt., 31:152-162.
- Odhner, T. (1905). Die trematoden des arktischen geibetes. Fauna Arctica, Jena., 4:291-372.

- Odhner, T. (1911). Zum natürlichen system der digenen Trematoden. III. Zool. Anz., 38:97-117.
- Odhner, T. (1914). Nordostafrikanische Trematoden, grosstenteils vom weissen Nil (von der schwedischen Zoologischen Expedition gesammelt). I. Fascioliden (Gorgoderidae, Lepodermatidae, Lecithodendriidae, Dicrocoeliidae and Echinostomidae). Results Swedish Zool. Exped. Egypt and White Nile, 1907, Pt. 4:1-166.
- Oliveira-Rodrigus, H.De.; Varela, M.C.; Rodrigues, S.S., and Cristofaro, R. (1972). Some digenetic trematodes of fish from the Atlantic Ocean. The continental Portuguese Coast and the Continental African Coast. Atas Soc. Biol. Rio de Janeiro, 15:87-93.
- Osmanov, S.Y. (1940). See Skrajabin: Trematodes of animals and man (On helminth fauna of fish), 9:48.
- Overstreet, R.M. (1969). Digenetic trematodes of marine Teleost fishes from Biscayne Bay, Florida. Tulane Stud. Zool., 15:119-176.
- Ozaki, Y. (1924). Gastrostomatous trematodes and three new genera of them. Zool. Mag., 36:173-201.
- Ozaki, Y. (1925). Preliminary notes on a trematodes with anus. Zool. Mag., 37:461-423.
- Ozaki, Y. (1928). Some Gasterostomatous trematodes of Japan. Japanese J. Zool., 2:1-33.
- Ozaki, H. and Ozaki, Y. (1952). A new gasterostome trematode, *Bucephalopsis tylosuris*, n.sp. J. Sci. Hiroshima University, Series B, Division 1 (Zoology), 13:85-90.
- Pande, B.P. (1934). On a new trematode from an Indian fresh-water fish. Proc. Indian Acad. Sci., 4:107-112.

- Pande, B.P., and Shukla, R.P.(1976). *Haplorchoides* Chen, 1949 (Haplorchinae: Heterophyidae) in fresh-water fishes. J. Helminthol., 50:181-192.
- Pandey, K.C. (1969). *Bucephalopsis hexaglandulata* sp.n., *B. oxygasteri* sp.n. and *B. multiglandulata* sp.n.: gastrostome metacercariae from Indian fishes. Trans. Amer. Microsc. Soc., 88:486-491.
- Paperna, I. (1964). Parasitic helminths of inland water fishes in Israel. Israel J. Zool., 13:1-20.
- Park, T.T. (1937). A revision of the genus *Podocotyle* (Allocreadiidae) with a description of eight new species from tide pool fishes from Dillon's Beach, California. J. Parasitol., 23:405-422.
- Park, J.T. (1939). Trematodes of fishes from Tyosen IV. A new digenetic trematode parasite, *Bucephalopsis cybii* sp. nov. (Bucephalidae: Poche, 1907) Keijo J. Med., 10:63-65.
- Pearson, J.C. (1964). A revision of the subfamily Haplorchinae Looss, 1899 (Heterophyidae) I. The *Haplorchis* group. Parasitology, 54:601-676.
- Pigulewsky, S.W. (1931). Nue Arten Von Trematoden aus Fischen es Dnjeprbassins. Zool. Anz., 96:9-18.
- Poche, T. (1907). Einige Bemerkungen zur nomenclatur der Trematoden. Zool. Anz., 31:124-126.
- Poche, T. (1925). Zur kenntnis von amphilino foliaceae Z. Wiss. Zooll., 25:585-619.
- Poche, T. (1926). Das system der platodaria. Arch. Nat., 91:458 PP.
- Polyanski, Y.I. (1952). Two new digenetic trematodes from fish in the North Atlantic. Parazitol. Sbornik, 14:266-280.

- Por, F.D. (1978). Ecological studies 23, Lessepsian Migration, 228 PP., Springer-Verlag, Berlin, Heidelberg, New York.
- Price, E.W. (1934). New digenetic trematodes from marine fishes XCI. Smiths. Misc. Coll. 91, 8 PP.
- Pritchard, M.H. and Manter, H.W. (1961). Studies of digenetic trematodes of Hawaiian fishes; Family Haplosporidia. Proc. Helminthol. Soc. Wash., 28:191-197.
- Pritchard, M.H. (1966). A revision of the genus *Podocotyle* (Trematoda: Opecolidae). Zool. Jahrb. Syst., 93:158-172.
- Qazi, M. and Qazi, M.H. (1963). *Astiotrema lissemysi* a new trematode from the intestine of fresh-water turtle, *Lissemys punctata*. Scientist, Karachi, 6:148-152.
- Ramadan, M.M. (1979). Studies on helminth parasites of some Red Sea fishes. Ph.D. Thesis, Faculty of Science, Ain Shams University, Egypt.
- Ramadan, M.M. (1985). On three species of the trematode genus *Plagioporus* Stafford, 1904 from Red Sea fishes. Qatar Univ. Sci. Bull., 5:267-278.
- Reid, W.A., Coil, W.H., and Kuntz, R.E. (1966). Hemiurid trematodes of Formosan marine fishes 1. Subfamilies: Dinurinae and stomachicolinae. J. Parasitol., 52:39-45.
- Roitman, V.A. (1963). Morphological characteristics of the trematode families. Allocreadiidae Stossich, 1903 and Opecoelidae Ozaki, 1925. Trudy Gel'm. Lab., 13:301-302.
- Rokicki, J. (1973). Helminths of certain Clupeidae mainly of the herring *Clupea harengus* L., in South Baltic. Acta. Parasitol. Polonica, 21:443-464.