

REFERENCES

- Abd EL-Sattar, M. A., F. Awny, Massoud, Samya I. and I. N. Ali (1985). Pathological studies on powdery mildew of cucurbit plants in Egypt Proc. 1st Conf. of Pests and Dis. of veg. and Fruits in Egypt, Ismailia, 1985 PP 602-613.
- Abd-EL Kareem, F., W. E. Ashour, M. M. Diab, and M. M. Aly (1993). Induction of resistance in watermelon plants against fusarium with using biotic and chemical inducers. 5th Nat. Conf. of Pests and Dis. of Veg. and Fruits in Egypt (Ismalia), 447-455.
- Abd-EL-Kareem, F. I. (1998). Induction of resistance to some diseases of cucumber plants grown under greenhouse conditions. Ph D. Thesis, Fac. of Agric. Ain Shams Univ. 117 pp.
- Abeles, F. B. and L. E. Forrence (1970). Temporal and hormonal control of B-1, 3 glucanase in *Phaseolus vulgaris* L. Plant Physiology, 45:395-400.
- Abo-Taleb, Mona, M. A. (2001). Biochemical changes associated with the application of some resistance inducing compounds for controlling powdery mildew of cucumber. Egypt. J. Appl. Sci, 16: (12) 387-403.
- Agrios, G. N. (1988). Plant pathology, 3rd ed. Academic press Inc., San Dieg USA.
- Ahmed, Saeida. S. A. (2001). Future studies on soybean anthracnose disease in Egypt. Ph D. Thesis, Fac Agric, Cairo Univ. 159 pp.
- Allam, A. L. and J. P. Hollis (1972). Sulfide inhibition of oxidase in rice roots. Phytopathology, 62: 634-639.
- Aly, M. M., S. A. Habib and S. M. Abd-Allah (1988). Induced resistance in barley plants against powdery mildew disease using ethephone treatment pro 2nd conf. Develop. Res.
- Anonymous (1974). Brockhaus Landwirtschaft. VEB F. A. rockhause Verlag, Leipzig. 636 pp.
- Arnon, D. I. (1949). Copper enzymes in isolated chloroplasts. Polyphenolase in *Beta vulgarise*. Plant Pathology, 24: 1-5.
- Attitalla, I. H. and S. Brishammar (2002). Oxalic acid elicited resistance to fusarium wilt in *lycopersicon esculentum* Mill. Plant Protection Science. 38: 1, 128-131.
- Avdiushko, S. A., X. S. Ye and J. Kuc, (1993). Detection of several enzymatic activities in leaf prints of cucumber plants. Physiological and Molecular Plant Pathology, 42: 441-454.
- Awad, Nadia S. A. (2004). New approaches for controlling powdery mildew disease of cucumber. Ph. D. Thesis, Faculty of Agricultural, Suez Canal Univ. 168 pp.

- Babu, R. M., A. Sajeena, A.V. Samundeeswari, A. Sreedhar, P. Vidhyasekeran and M. S. Reddy (2003). Induction of bacterial blight (*Xanthomonas oryzae* pv. *oryzae*) resistance in rice by treatment with acibenzolar-s-methyl. *Annals of Applied Biology*. 143 (3): 333-340.
- Baraka, M. A., M. I. Ghonim, H. M. Abdel-Nabi, and M. I. Ahmed (2004). Biochemical mechanisms of resistance in some bean cultivars against charcoal rot disease. *Bulletin of Faculty of Agricultural, Cairo Univ.* 55 (1): 97-120.
- Baydar, H. and S. Ulger (1998). Correlations between changes in amount of endogenous phytohormones and flowering in the safflower (*Carthamus tinctorius*). *Tr. J. of Biology* 22:421-425, (c. f. International Internet, CAB).
- Benhamou, M. N., H. A. J. Joosten and P. J. M. Dewit (1990). Sub cellular localization of chitinase and its potential substrate in tomato root tissues by *Fusarium oxysporum* f. sp. *radicis-lycopersici*. *Plant Physiology*, 92:1108-1120.
- Bidawell, R. G. S. (1974). *Plant Physiology*, pp. 143-147. Macmillan Publishing co., Inc. New York, 643 pp.
- Biehn, W. L., E. B. Williams and J. Kuc. (1968). Accumulation of phenols in resistant plant fungi interactions. *Phytopathology*, 58:1255-1260.
- Biles, C. L and R. D. Martyn (1989). Local and systemic resistance induced in watermelon by formae speciales of *Fusarium oxysporum*. *Phytopathology*. 79: 856-860.
- Biryukova, N. K. and N. S. Gorshkova, (1990). Production of the cucumber hybrid Topiek for the spring greenhouse. *Selektsiya, Smenovodstvo, Moskva*, 3: 17-20. (c.f. international Internet, CAB).
- Boesewinkel, N. J. (1977). Identification of Erysiphaceae by conidial characteristics. *Rev der Mycologie*, 41: 493-507.
- Booler, T., A. Gehri, F. Mauch, and U. Vogeli (1983). Chitinase in bean leaves induction by ethylene, purification, properties, and possible function *Planta*, 157: 22-31.
- Bradford, M. M. (1976). A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein dye binding. *Analytical Biochem.*, 72 : 248-254.
- Branzanti, B., and A. Brunelli, (1992). Study of aetiology of powdery mildew of cucurbits in Emilia- Romagna. *Informatore Fitopatologico*. 42 (11): 37-44.
- Brigitte, M. and M. Felix. (2005). The role of abscisic acid in plant pathogen interactions. *Plant Biology*, 8: 409-414.
- Buell, C. R. (1999). Genes involved in plant-pathogen interaction in: Agarwal, A. A., and S. Tuzum. E. Bent (eds). *Induced Plant Defenses. Against pathogens and Herbivores*, PP. 73-93, APS press. St. Paul, M.N.
- Caruso, F. L. and J. Kuc (1977a). Field protection of cucumber, watermelon and muskmelon against *Colletotrichum lagenarium*, *Phytopathology*, 67: 1290-1292.
- Caruso, F. L. and J. Kuc (1977b). Protection of cucumber, watermelon and muskmelon against *Colletotrichum lagenarium*, *Phytopathology*. 67:1285-1289.
- Cheah, L. H., B. B. Page and J. K. Cox. (1996). Epidemiology of powdery mildew (*S. fuliginiea*) of squash. *Proc. 94th N.Z. Plant Protection Con.* 147-151.
- Chen, ChunQuan, R. R. Belanger, N. Benhamou, T. C. Paulitz and C. Q. Chen (1999). Role of salicylic acid in systemic resistance induced by *Pseudomonas* spp. against *Pythium aphanidermatum* in cucumber roots. *European Journal of Plant Pathology*. 105 (5): 477- 486.

- Chen, X. J. and F. J. Meng (1990) Zearalenone radioimmunoassay. *Acta Phytophysiological Sinica*. 16: (1) 70-76.
- Chester, K.S. (1933). The problem acquired physiological immunity in plants. *Q. Biol. Rev.* 8: 275-324.
- Christin, G. (1990). HPLC Tips and Tricks. Great Britain at the Iden Press, Oxford. 608 pp.
- Chrzanowski, G., A. P. Ciepiela, I. Sprawka, C. Sempruch, H. Sytkiewicz and P. Czerniewicz (2003). Activity of polyphenoloxidase in the ears of spring wheat and triticale infested by grain aphid (*Sitobion avenae*). *Electronic J. of Polish Agricultural Universities, Biology*, Volume 6, Issue 2. (<http://ejpau. Media pl>).
- Cohen, R., G. Leibovich and D. Shtienberg (1993). Variability in the reaction of squash (*Cucurbita pepo*) to inoculation with *Sphaerotheca fuliginea* and methodology of breeding for resistance. *Plant Pathology*, 42: 510-516.
- Conrath, U., O. Thulk, V. Katz, S. Schwindling and A. Kohler (2001). Priming as a mechanism in induced systemic resistance of plants. *European Journal of Plant Pathology*, 107: 113-119.
- Conti, G. G., A. Pianezzola, A. Arnoldi, G. Violini and D. Maffi. (1996). Possible involvement of salicylic acid in systemic acquired resistance of *Cucumis sativus* against *Sphaerotheca fuliginea*. *European J. of Plant Pathology*, 102 (6): 537-544.
- Daayf, F., R. BelRhliid and R. R. Belanger (1997). Methyl ester of P-coumaric acid: a phytoalexin like compound from long English cucumber leaves. *Journal of Chemical Ecology*, 32 (6): 1517-1526.
- Descalzo, R. C, J. E. Rahe and B.C. Mauza (1990). Comparative efficacy of induced resistance for selected disease of greenhouse cucumber. *Canadian J. of Plant Pathology*. 12 (1): 16- 24.
- Dik, A. J., M. A. Rehaar and R. R. Belanger, (1998). Comparison of three biological control agents against cucumber powdery mildew (*Sphaerotheca fuliginea*) in semi-commercial- scale glasshouse trials. *European Journal of Plant Pathology*, 104 (4): 413-423.
- Doubravera, N., R. Dean, and J. Kuc (1988). Induction of systemic resistance to anthracnose caused by *Colletotrichum lagenarium* in cucumber by oxalate and extracts from Spinach and rhubarb leaves. *Physiological and Molecular Plant Pathology*, 33: 69-79.
- Elad, Y, B. Kirshner, N.Yehuda and A. Sztejnberg, (1998). Management of powdery mildew and graymould of cucumber by *Trichoderma harzianum* T₃₉ and *Ampelomyces quisqualis* AQ10. *biocontrol*. 43: (2) 241-251.
- EL-Desouky, S. M. (1989). Studies on powdery mildew disease of cucurbitaceae. M.Sc. thesis, faculty of Agriculture, Minufeya Univ. 135 pp.
- EL-Korachy, M. A. (1991). Epidemiology and management of the downey mildew disease of cucurbits under protected agriculture in Ismailia, Ph. D. Thesis, Dept. Agric. Sci, Inst. Environ. Stud. Res., Ain Shams Univ, Egypt (141).
- EL-Naggar, Doaae R. M. A. (2005). Further studies on stem Rust of Wheat. Ph. D. Thesis. Faculty of Agric. Zagazig Univ. 182 pp.
- EL-Toony A. M. E. (1992). Studies on leaf spot fungi of some Cucurbitaceae plants. Ph.D. Thesis, Fac. Agric., AL-Azhar. Univ.
- Engstrom K and A. Stromberg (1996). Changes in sugar content during induction of systemic acquired resistance to late blight caused by *Phytophthora infestans* (Mont) de

- Bray in potato. J. of Phytopathology- Phtopathologische Zeitschrift. 144: (1) 33-36. (c.f. international Internet, CAB)
- Fadl F. A., M. S. Abdel-Moneim, EL-Shami, Mona, S. Hassan, I. Nagi. and W. R Rizkallah (1981). Chemical control and varietal reaction to cucumber powdery mildew. Res. Bull. No. 289, Zagazig Univ.
- Farkas G. D. and Z. Kiraly (1962). Role of phenolic compounds in the physiology of plant disease and disease resistance. Phytopathology. Z., 44: 106-150.
- Fawe A., M. Abu Zaid, J. G. Menzies, R. R. Belanger, B. Duffy, U. Rosenherger, and G. Defago (1998). Silicon induces the accumulation of antifungal low-molecular weight metabolites in cucumber plants. Molecular approaches in biological control. Delemont, Switzerland, 15-18 September 1997. Bulletin-OLIBSROP21 (9): 133-136.
- Fisher, R. A. (1948). Statistical Methods for Research Workers. Oliver and Boyd, London, U.K.
- Fofana B., N. Benhamou, D. J. McNally, C. Labbe, A. Seguin, and R. R. Belanger (2005). Supression of induced resistance in cucumber through distribution of the flavonoid pathway. Phytopathology, 95 (1): 114-123.
- Fraser, R. S. S., S. A. R. Loughlin and R. J. Whenham (1979). Acquired systemic susceptibility to infection by tobacco mosaic virus in *Nicotiana glutinosa* L. J. Gen.Virol, 43:131-141.
- Frohlich, G (1979). Phytopathologie und prflanzenschutz VEB Gutay Fischer Verlay Jena, PP295 (C.F. International CAB)
- Galal, A. A., L. Ammal, Z. A. Botros, A. A. Shihaca and M. F. Ouf (1997). Acquired resistance to *Puccinia heliantha* in sunflower plants. Egypt. J. Phytopathology, 25 (1-2): 45-54.
- Gamil, N. A. M. (1995). Induced resistance in squash plants against powdery mildew by cobalt and phosphate sprays. Ann. Agric. Sci. Moshtohor, 33 (1): 183-194.
- Gäuman, E. (1946). Crosstalk in plant cell signaling: structure and function of the genetic network. Trends Plant Sci. 4: 503-507.
- Gessler, C. and J. Kuc (1982). Induction of resistance to *Fusarium* wilt in cucumber by root and foliar pathogen. Phytopathology, 72: 1439-1440.
- Ghoneim, K. E. A. (1985). Studies on powdery mildew diseases in Egypt. M.Sc. Thesis, Fac, Agric., Tanta Univ.
- Gilpatrick, J. D. and M. Weintraub (1952). The effects of aspirin and poly acrylic acid on soluble leaf protein and resistance to virus infection in five cultivars of tobacco. Science, 115:701-706
- Gottstein H. D and J. Kuc (1989). Induction of systemic resistance to anthracnose in cucumber by phosphates. Phytopatholog, 79: 271-275.
- Gouvea, C. M.; C. P. Ricard and A. R. Oliveira. (1995). Accumulation of a host microsomal fraction protein in virus in infected cucumber and tomato. Plnat Physiol. and Biochemistry- Paris. 33: 453-461.
- Grieve, M. (2003). Botanical. Com, Amodern Hebal. [http:// WWW. Botanical. Com/ botanical/ mgmh. html](http://WWW.Botanical.Com/botanical/mgmh.html) Cotanical.
- Grisebach, H. (1981). Lignin, in corn, E. E. (ed): the biochemistry of plants, Academic press New Yourk. PP 451-478.
- Gross, G. G. (1979). Recent advances in chemistry and biochemistry of lignin, Recent Adv. Phytochem, 12, 77.

- Gupta, S. K, A. G. Sham and K. R. Ramesh Bhardwaj (2001). Morphological characterization and effect of meteorological factors on development of cucumber powdery mildew. *Indian Phytopathology*, 54 (3): 311-315.
- Gupta, S. K., P. P. Gupta, C. D. Kaushik and H. K. L. Chawla (1992). Metabolic changes in groundnut leaf due to infection by leaf spot pathogens. *Indian Phytopathology*, 45: 434-438.
- Heijwegen, T. and M. A. Verhaar (1995). Effects of cucumber genotype on the induction of resistance to powdery mildew *Sphaerotheca fuliginea* by 2, 6-dichloroisonicotinic acid. *Plant Pathology*, 44 (4): 756-762.
- Hynes, R. K., J. Hill, M. S. R. Reddy and G. Lazarovits (1994). Phytoalexin productions by wounded white bean (*Phaseolus vulgaris*) cotyledons and hypocotyls in response to inoculation with *Rhizobacteria*. *Canadian J. of Microbiology*, 40 (7): 548-554.
- Irving, H. R. and J. Kuc, (1990). Local and systemic induction of peroxidase chitinase and resistance in cucumber plants by K_2HPO_4 . *Physiology and Molecular Plant Pathology*, 37: 355-366.
- Ishii, H., Y. Tomita, T. Horio, Y. Narusaka, Y. Nakazawa, K. Nishimura and S. Iwamoto (1999). Induced resistance of acibenzalar-s-methyl (CGA 245704) to cucumber and Japanese pear disease. *European J. of Plant Pathology*. 105 (1): 77-85.
- Jeun, Y. C., K. S. Park, C. H. Kim, W. D. Flower and J. W. Kloepper (2004). Cytological observations of cucumber plants during induced resistance elicited by *Rhizobacteria*. *Biological Control*. 29 (1): 34-42.
- Jiao, Z. and Lu, Bing Sheng (1995). The influence of the preparation of *Trichoderma Konigii* on some physiological and biochemical properties of cotton and kidney bean seedling. *Chinese J. of Biological Control*. 11: 1, 30-32.
- Jindal, S., Y. K. Arora and K. L. Bajaj (1979). Effect of infection by *Sphaerotheca fuliginea* (Powdery mildew) on amino acids and sugars in leaves of muskmelon (*Cucumis melo* L.). *Plant Biochemical J.* 6 (2): 116-123.
- Joseph, KUC (1995). Phytoalexins, stress metabolism and disease resistance in plants. *Ann Rev. Phytopathol*, 33: 275-297.
- Kabitarani A. and T. Bhagirath, (1991). Powdery mildew of cucurbits in Manipur-Indian *Phytopath.* 44 (1): 137-139.
- Khalil, M. S. (1981). Biochemical and serological studies on flax rust. Ph. D. Thesis, Fac. Agric., Cairo Univ.
- Khan, M. W. (1981). *Sphaerotheca fuliginea* causing powdery mildew of cucumber a new record for Libyan Jamahiriya. *The Libyan Jamahiriya. The Libyan J. of Agric.* 10: 145-150.
- Kombrin, M., E. M. Schrodeder and M. Hahlbrok (1988). Several pathogenesis-related proteins in potato are B-1,3- -glucanase and chitinase. *Proc Natl Acad Sci.* 85: 782-786. (c.f. Tuzun *et al.*, 1989).
- Kuc, J. (1982). Induced immunity to plant diseases. *Biosciences*. 32: 854-860.
- Kuc, J. (1990). Immunization for the control of plant disease in: Hornbg, D. (ed.). *Biological control of soil – borne pathogens*. CAB International wallingford 355-373.
- Kuc, J. (1995). Systemic induced resistance. *Aspects of Applied Biology*. 42, 235-242.
- Laun, N. (1998). Induced resistance: first experiences with Bion in vegetables. *Germuse. Munchen*. 34 (5): 294-297.

- Li Swahuju, Ma. D., Pang J., Huo Z., Li S., Ma D., Pang J. and Z. Huo (2000). Induced effect of salicylic acid on the activity of several enzymes and disease resistance of cucumber. *Acta Agriculturae Boreali Sinica*. 15 (2): 118-122.
- Li, Kloepper, J. W. and S. Tuzun (1995). Induction of systemic resistance in cucumber against Fusarium wilt by plant growth promoting *Rhizobacteria*. *Phytopathology*, 85 (6): 695-698.
- Liu, L., J. W. Kloepper and S. Tuzun (1995). Induction of systemic resistance in cucumber against Fusarium wilt by plant growth promoting *Rhizobacteria*. *Phytopathology*, 85 (6): 695-698.
- Luo G., J. T. Cui, Y. Gao and L. Zhang (1997). Relationship between sugar, lignin content and resistance to powdery mildew of cucumber. *Acta Phytopathologica Sinica* 27(1): 65-69.
- Mahgoob, A. E. A. and S. A. M. Zaghlool (2002). Effect of salicylic and Jasmonic acids on the response of tomato plants to root rot nematode *Meloidogyne incognita*, infection. *Annals of Agricultural Science (Cairo)*. 47 (3): 1107-1119.
- Malamy, J., J. P. Carr, D. Klessig, and I. Raskin (1990). Salicylic acid: a likely endogenous signal in the resistance response of Tobacco to virus infection. *Science*, 250, 1002.
- Malhotra, B., J. C. Onyilagha, B. A. Boham, G. H. N. Towers, D. James, J. B. Harbone and C. J. French. (1996). Inhibition of tomato ringspot virus by flavonoids. *Phytochemistry*. 43 (6): 1271-1276.
- Mariela, O. and H. Christina (2001). The complexity of pathogen defense in plants. *J. Plant Physiol*, 27 (1 – 2), 101-109.
- Masami M., S. Keita and S. Takeo and S. Yoshiteru. (2003). Powdery mildew resistance in cucumber. *JARQ*, 37 (1): 7-14. (c. f. <http://www.jircas.affrc.go.jp>).
- Matta A., G. I. Abattista and L. Ferraris (1988). Stimulation of 1, 3 glucanase and chitinase by stresses that induced resistance to Fusarium Wilt in Tomato. *Phytopath. Medit.* 27: 45-50.
- Mauch F. and L. A. Staehlin (1989). Functional implications of the subcellular localization of ethylene-induced and β -1,3 glucanase in bean leaves. *The Plant Cell*. 1, 447-457.
- Maxwell D. P. and D. F. Bateman (1967). Changes in the activity of some oxidases in extracts of *Rhizoctonia* infected bean hypocotyls in relation to lesion maturation. *Phytopathology*, 57: 132-136.
- Mazzanti De Castanon M. A, R. E. Alvarez, and M. G. Carbrera-De-Alvarez, (1987). Contribution to the knowledge of the etiology of powdery mildew of *Cucurbitaceae* grown in North-East Argentina. *Fitopaiologia*, 12 (1): 21-29.
- McNally, D. J., K. V. Wurms, C. Labbe, R. R. Belanger (2003). Synthesis of C. glycosyl flavonoid phytoalexins as a site-specific response to fungal penetration in cucumber. Ltd, Oxford, UK. 63 (6): 293-303.
- Mètraux, J. P. (2001). Systemic acquired resistance and salicylic acid: Current State of knowledge. *European J. of plant pathology* 107: 13-18.
- Mètraux, J. P. and T. Boller (1986). The local and systemic induction of chitinase in cucumber plants in response to viral, bacteria and fungal infections. *Physiol. Mol. Plant Pathology*, 28: 161-169.
- Mètraux, J. P., H. Singer, J. Ryals, E. Ward, M. Wyss-Benz, J. Gaudin, K. Rashdorf, E. Schmid, W. Blum and B. Inverardi (1990). Increase in salicylic acid at the onset of systemic acquired resistance in cucumber, *Science*, 250, 1004.

- Metwally, M. M. M. (2004). Resistance induction against disease of Faba bean crop. Ph. D. Thesis. Faculty of Agriculture, Suez Canal Univ. 159 pp.
- Meuwly, P., W. Molders, A. Buchala and J. P. Mètraux (1995). Local and systemic biosynthesis of salicylic acid in infected cucumber plants. *Plant Physiology*, 109 (3): 1107-1114.
- Meuwly, P., W. Molders, K. Summermatter, L. Sticher, J. P. Mètraux, M. Geibel, D. Treutter and W. Feucht (1994). Salicylic acid and chitinase in infected cucumber plants. *International Symposium on Natural Phenol in Plant Resistance*, Volume I, 13-17 Sep., 1993, Weih Enstephan, Germany. *Acta Horticulturae*, 381: 371-374.
- Michael, D., K. Walter, D. Bob and S. Theodor (2001). Induced resistance in plants by chemicals. *European Journal of Plant Pathology* 107: 19-28.
- Mills, P. R., E. J. Gussin and R. K. S. Wood, (1986). Induction of resistance in cucumber to *Colletotrichum lagenarium* by 6-benzylaminopurine. *Journal of Phytopathology-Phytopathologische Zeitschrift*. 117 (1): 11-17.
- Mirostera, O., J. Siegrist and H. Buchenauer (2002). Mechanisms of phosphate induced disease resistance in cucumber. *European Journal of Plant Pathology*, 108: 345-353.
- Misaghi, I. J. (1982). *Physiology and Biochemistry of plant-pathogen interactions*. Plenum press, New York and London. 287 pp.
- Mohamed, Y. F., M. Bardin, P. C. Nicot and M. Pitrat (1995). Causal agents of powdery mildew of cucurbits in Sudan. 79 (6): 634-636.
- Monreal, J. and E. T. Reese (1969). The chitinase of *Serratia marcescens*. *Canadian J. of Microbiology*, 15: 689-696.
- Moreno, R. Q. and S. Romero, (1991). Cleistothecal formation in Squash var castilla (*Cucumis moschata*) by *Sphaerotheca fuliginea* in Jesus Maria. Aguas Calientes. *Revista-Mexicana-de-Fitopatologia*, 9: 94-96. (c. f. International Internet, CAB).
- Mosa A. A. (1997). Effect of foliar application of phosphates on cucumber powdery mildew. *Annals of agricultural Science (Cairo)* 42: 1, 241-255.
- Moshe, R. and R. Reuven (2000). Prior inoculation with non-pathogenic fungi induces systemic resistance to powdery mildew on cucumber plants: *European Journal of Plant Pathology* 106: 633-638.
- Moshe, R., A. Vladimir and R. Reuven (1997). A foliar spray of micro nutrient solutions induces local and systemic protection against powdery mildew (*Sphaerotheca fuliginea*) in cucumber plants. *European Journal of Plant Pathology*, 103: 581-588.
- Mucharromah, E., and J. Kuc (1991). Oxalate and phosphates induced systemic resistance against diseases caused by fungi, bacteria and viruses in cucumber. *Crop Protection*. 10: 4, 265-270.
- Nabil, A. A. A. (1999). Effect of chemical and heat treatments of seeds on Squash infection by cucumber mosaic virus (CMV). *Assiut. Journal of Agricultural Sciencs*. 30 (4): 193-206.
- Naik, S. T. (1995). Role of sugars and phenols in improving resistance in *Hibiscus* spp. against *Colletotrichum abelmoschii*. *Indian Forester*, 121: 306- 308. (c.f. *Pl. Pathol.*, 1996-1997).
- Okuno, T., M. Nakayama, N. Okajima and I. Furusawa, (1991). Systemic resistance to Downy mildew and appearance of acid soluble proteins in cucumber leaves treated with biotic and abiotic inducers. *Annals of the Phytopathological Society of Japan*. 57: 203- 211.

- Ongena, M., F. Daayf, P. Jacques, P. Thonart, N. Benhamou, T. C. Paulitz and R.R. Belanger, (2000). Systemic induction of phytoalexins in cucumber in response to treatments with *Pseudomonas fluorescens*. Plant Pathology. 49: 4, 523-530.
- Orober, M., S. J. Sigrist, H. Buchenauer, H. Lyr, P. E. Russell, H.W. Dehne and H. D. Sisler (1998). Induction of systemic acquired resistance in cucumber by foliar phosphate application. Modern fungicides and antifungal compounds II. 12th international Reinhard Sbrunn Symposium, Friedrichroda, thuringia, Germany, 24th-29th may 1998. 339 -348.
- Pasini, C., F. D'Aquila, P. Curir, and Gullino, M. L. (1997 b). Effectiveness of antifungal compounds against rose powdery mildew (*Sphaerotheca pannosa* var. *rosae*) in glasshouses. Crop Protection, 16 (3): 251-256.
- Pasini, C., F. D'Aquila, P. Curir, M. L. Gullino and A. Carnero (1997a). Alternative strategies to control rose powdery mildew. (Albajes, R. (ed).) integrated control in protected crops Mediterranean climate. Proceedings of the Meeting at Tenerife, Canary Islands, 3-6 November 1997. Bulletin- OILB- SROP., (4): 84-87.
- Pe'er S. and Y. Cohen (1987). Sugar accumulation in tobacco plants systemically protected against blue mold (*Peronospora tabacina*). Phytoparasitica. 15 (4): 307-315.
- Peixoto C. B., G. J. Tanner, M. A. Galli (2000). Test of de um inductor de resistencia na cultura da batata (*Solanum Tuberosum* L.). Ecosystema, 25 (2): 215-216.
- Prachi S. R. T. and M. B. Singh (2002). Salicylic acid induced insensitivity to culture filtrate of *Fusarium oxysporum* f.sp *zingiberi* the cells of Zingiber officinal Roscoe. European Journal of Plant Pathology. 108: 31-39.
- Procter, J. T. A. (1981). Stomatal conductance changes in leaves of McIntosh apple trees and after fruit removal. Can. J. Bot, 59: 50-53.
- Radwan, Nabila, A. M. (2000). Implication of certain physical and chemical treatment to improve disease resistance of parley to powdery mildew. Ph.D. Thesis, Fac., Agric., Cairo Univ. 146 pp.
- Raskin, T. (1992). Role of salicylic acid in plants. Annu. Rev. Plant Physiology. 43: 439-463.
- Ratanam, S. V., P. N. Reddy, K. C. Rao, S. K. Raju and J. H. Kumar (2001). Effect of salicylic acid and Bion seed treatment in induction of systemic resistance against sunflower leaf blight, *Alternaria helianthi* Hansf. Tub. et Nish. Indian J. of Plant Protection. 29 (2): 79-81.
- Rauscher M., A. L. Adam, S. Wirtz, R. Guggenheim, K. Mendgen and H. B. Deisi (1999). PR-1 protein inhibits the differentiation of rust infection hyphae in leaves of acquired resistant broad bean. Plant- Journal. 19 (6): 625-633.
- Remus-Borel, W., J. G. Menzies and R. R. Belanger (2005). Silicon induced antifungal compounds in powdery mildew infected wheat. PMPP Physiological and Molecular Plant Pathology, 66 (3): 108-115.
- Reuveni M., V. Agapov and R. Reuveni (1995). Induced systemic protection to powdery mildew in cucumber by phosphate and potassium fertilizers: effects of inoculum concentration and potassium inoculation treatment. Canadian J. of Plant Pathology. 17 (3): 247-251.
- Reuveni M., V. Agapov and R. Reuveni (1993). Induction of systemic resistance to powdery mildew and growth increase in cucumber by phosphates. Biological Agriculture and Horticultural, 9: 305-315.

- Reuveni M., V. Agapov and R. Reuveni, (1994). Foliar sprays of phosphates inducers growth increase and systemic resistance to *Puccinia Sorghi* in maize. Plant Pathology. 43, 245-250.
- Reuveni, M., V. Agapov and R. Reuveni (1997). A foliar spray of micronutrient solutions induces local and systemic protection against powdery mildew (*S. fuliginea*) in cucumber plants. European J. of Plant Pathology, 103: 581-588.
- Rezende, J. A. M., J. K. Brown, A. L. Lourencao and I. P. Bedendo, (1997). Observation on the reduced incidence of powdery mildew on Zucchini squash silvered leaves in Brazil. Fitopathologica. Brasileira. 22: 4, 568.
- Ride, J. P. (1983). Cell wall and other structural barriers in defense. In Calloz J. A. (ed). Biochemical plants pathology. John Wiley and Sons. New Yourk.
- Ried J. D. and D. M. Ogryd-Ziak (1981). Chitinase over producing mutant of servatia marcesces. Apple. and Environ. Microbiol. 41: 664-669.
- Rikin, A., M. Waldman, A. E. Richmond and A. Dovrat (1975). Hormonal regulation of morphogenesis and cold resistance: modification by abscisic acid and by gibberellic acid in alfalfa (*Medicago sativa L.*) seedlings. J. of Expermental Botany, 26 (2): 175-183.
- Ross, A. F. (1961). Systemic acquired resistance induced by localized virus infections in plants. Virology, 13: 340-358.
- Ruberto, G., A. Renda, M. Piattelli, P. Rapisardo and A. Starrantino (1997). Essential oil of two new pigmented citrus hybrids, Citrus clementina and Citrus sinensis. J. Agric. Food Chem. American Chemical Society, 45: 467-471.
- Sailaja, P. R. and A. R. Podile (1998). A phytoalexin is modified to less fungitoxic substances by crown rot pathogen in groundnut (*Arachis hypogaea L.*) Indian J. of Eperimental Biology.36 (6): 631-634.
- Sakhabutdinova, A. R., D. R. Fatkhutdinova, M. V. Bezrukova and F. M Shakirova.(2003). Salicylic acid prevents the damaging action of stress factors on wheat plants. Bulg. J. Plant Physiology, 314-319.
- Sallam Minass E. A. (1997). Studies on the leaf rust of wheat in Egypt. Ph. D. Thesis, Fac. Agric., Zagazig Unvi. 119 pp.
- Sang, G. K. and Y. Y. Joo, (1992). Changes in protein patterns resulting from infection of rice leaves by *Xanthomonase oryzae*. Molec. Plant Microbe Interacts. 5 (4): 356-360.
- Schneider, S. and W. R Ullrich (1994). Differential induction of resistance and enhanced enzyme activities in cucumber and tobacco caused by treatment with various abiotic and biotic inducers. Physiological and Molecular Plant Pathology, 45: 291-304.
- Schneider, S. and W. R. Ullrich (1990). Metabolic changes associated with induced systemic resistance to fungal and bacterial plant disease, (Abstr) Rev. Plant Path. 69: 4783.
- Segrist, J., D. Glenewinkel, C. Koll, and M. Schmidtke (1997). Chemically induced resistance in green been against bacteria and fungal pathogens. J. Plant Dis. Prot 104, 599-610.
- Sharma,, G. K. and M.W. Khan (1991). Observations on occurrence and identity of powdery mildew of cucurbits in Tamil Nadu. Indian Phytopathology, 44 (1): 45-51.
- Shi, Zhongliang, Qiu, Song Ying., Ma, Aiping. XU, Gang Yuang. Wu, Jiping and Lu, LaHu. (2002). Studies on mechanisms of biochemical resistance to blossom midge (*Sitodiplosis mosellana Gelin*) in wheat. Journal of Triticeae Crop, 22 (4): 63-65.
- Sidhan, G. S, Indra Hooda and D. R. Parashar (1999). Sources of resistance to Cercospor leaf spot in mung bean and biochemical parameters for resistance. J. Mycol. and Plant Pathol, 29: 130-132. (c. f. Rev. Plant. Pathol. 1157).

- Silva, L. H. C. P, M. L. V. Resende, R. M. Souza, J. R. Campos and A. M. S. Castro (2003). Induction of resistance against *Xanthomonas vesicatoria* in tomato by acibenzolar-s-methyl. *Summa Phytopathologica*. 29 (2): 177- 181.
- Smith-Becker J., N. T. Been, and J. O. Becker (2003). Acibenzolar-S-methyl induces resistance to *Colletotrichum lagenarium* and cucumber mosaic virus in cantaloupe. *Grope Protection*. 22 (5): 769-774.
- Snell, F. D. and C. T. Snell (1953). Colrimetric methods of analysis including some turbidimetric and nephelometric methods. D. van Nostrand Company Inc. Toronto-New York London. Vol. 111, 606 pp.
- Soylu, E. M., S. Soylu, and O. Baysal, (2003). Induction of disease resistance and antioxidant enzymes by acibenzolar-s-methyl against bacterial canker (*Clavinacter michiganensis* subsp. *Michiganensis* in tomato. *Journal of plant pathology*. 85: 3, 175-181.
- Staub, J. E. and L. K. Crubaugh (2001). Cucumber inbred line USDA 6632E. Cucrbit Genetics Cooperative. No. 24: 6-7.
- Sticher, L., B. Mauch-Mani, and J. P. Mètraux (1997). Systemic acquired resistance. *Annu. Rev. Plant Pathology*, 35: 235-270.
- Stromber, A and S. Brishammar (1991). Induction of systemic resistance in potato (*Solanum tuberosum* L.) plants to late blight by local treatment with *Phytophthora infestans*, *Phytophthora cryptogea* or dipotassium phosphate. *Potato Reaserch*. 34: 219-225.
- Sukhwant, S., P. P. Singh., P. S. Bedi and S. Singh (1994). Effect of different chemicals on infective potential of Teliospores of *Neovossia indica*. *Indian J. Agric Sci.*, 64 (3): 199-201 (c.f. CAB Abstr. 1995).
- Sutton, B. C. S. and M. Shaw (1986). Changes in two ribonuclease isozymes during rust infection of flax cotyledons. *Plant Physiol.*, 69: 205-209.
- Talanova, V. V. and A. F. Titov (1994). Endogenous abscisic acid content in cucumber leaves under the influence of unfavourable temperatures and salinity. *J. of Experimental Botany*, 45 (7): 1031-1033.
- Thomas, W. and R. A. Ducher (1924). The calorimetric determination of carbohydrates in plants by picric acid reduction method I. the estimation of reducing sugars and sucrose. *J. Am. Soc.*, 46: 1662-1669.
- Toppan, A, D. Roby and M. T. Esquerre-Tugaye (1982). Cell surface in plant microorganism interactions. *Plant Physiology*, 70: 82-86.
- Townsend, G. R. and J. W. Heuberger (1943). Methods for estimating losses caused by disease in fungicide experiments. *Pl. Dis. Repter*. 27: 340-343.
- Tsay, J. G. and B. K. Tung (1992). Ecology and control strategy of powdery mildew of cucurbits. *Chih Wu Pao Hu Hsueh Hui K'An*. 34 (2): 139-148. (c. f. International Internert, CAB).
- Tuzen, S., M. N. Rao, U. Vogeli, C. L Schardi and J. Kuc (1989). Induced systemic resistance to blue mold: Early induction and accumulation of β -1, 3 glucanase, chitinase and other pathogenesis-related protein (b-proteins) in immunized tobacco. *Phytopathology*, 79: 979-983.
- Tuzun, S. (2001). The relationship between pathogen-induced resistance (ISR) and muligenic (horizontal resistance in plants. *European Journal of plant pathology*. 107 (1): 85-93.

- Tyuterev, S. L. and S. A. Tarlakovsii (1994). Dynamics of the accumulation of free amino acids in wheat plants under influence of a disease resistance inducer. Sel' Skokhozyastvennaya Biologiya. 3, 120-127. (c. f. International Internert, CAB).
- Uozumi, T. and H. Yoshii (1952). Some observations on the mildew fungus affecting cucurbitaceous plants. Ann Phytopath. Soc. Japan 16: 123-126.
- Van Loon L. C. (1997). Induced resistance in plants and the role of pathogenesis-related proteins. European journal of plant pathology. 103: 753-765.
- Van Peer, R., G. J. Niemann and B. Schippers (1991). Induced resistance and phytoalexin accumulation in biological control of *Fusarium* wilt of Carnation by *Pseudomonas* sp. strain WC5417 r. Phytopathology 81: 728-734.
- Velkov, N. and S. Masheva. (2002). Species and races composition of powdery mildew on cucurbits in Bulgaria. Cucurbit Genetics Cooperative. 25, 7-10. (c. f. International Internert, CAB).
- Vogt, M, and H. Buchenauer (1997). Enhancement of biological control by combination of antagonistic fluorescent pseudomonas strains and resistance inducers against damping off and powdery mildew in cucumber. Zeitschrift Fur Pflanzenkrankheiten and Pflanzenschutz, 104 (3): 272-280.
- Walter, D. R., A. F. Mitchell, J. Hampson and A. Mcpherson, (1993). The induction of systemic resistance in barley to powdery mildew infection using salicylic acid and various phenolic acids. Annals of Applied Biology, 122: 451-456.
- Yang Rong Jin, Peng XinXiang, Jiang ZiDe, Guo Zhen Fei, Li MingQi, Yang RJ, Peng XX, Jiang ZD, Guo ZF and Li MQ (2000). Studies on the mechanism of oxalate induced resistance to anthracnose in cucumber leaves. Acta phytopathologica Sinica. 30 (2): 153-157.
- Yarwood, C. E. (1978). History and Taxonomy of powdery mildew in spencer, 1978. The powdery mildew. 1-32 pp. Academic Press, London.
- Ye, X. S., N. Strobel and J. Kuc (1995). Induced systemic resistance (ISR): activation of natural defense mechanisms for plant disease control as part of integrated pest management (IPM). In: Reuveni R(ed) Novel Approaches to Integrated pest Management (pp 95-113) CRC Press, Boca Raton, Florida, USA
- Ye, X. S., Pan, S. Q. and J. Kuc (1990). Association of pathogenesis related proteins and activities of peroxidase, B- 1,3 glucanase and chitinase with systemic induced resistance to blue mould of tobacco but not systemic tobacco. Physiological and Molecular Plant Pathology, 36: 523-531.
- Yedidia, I., N. Benhamou, Y. Kapulnik and I. Chet, (2000). Induction and accumulation of PR proteins activity during early stages of root colonization by the mycoparasite *Trichoderma harzianum* strain T-203. Plant Physiology and Biochemistry, 38 (11): 863-873.
- Yun, Xing Fu, Li, Rong Xi, Ma, LiGo, Huo, Xiu Wen, Yun, XF, Li, Rx, Ma, LG and Huo, Xw. (1997). Induced resistance to downy mildew in cucumber by chemicals. Acta. Phytopathologica Sinica. 24 (2): 159-163.
- Zahng Zongshen, Peng Xiniang, Jiang Zide, Xu DaGao, Li MingQi, ZhangZs, Peng XX, Jiang Z, XU DG and Li MQ (1998). The systemic induction of peroxidase by oxalate in cucumber leaves. Acta Phytopathologica Sinica. 28 (2): 145-150.
- Zaracoritis, C. (1965). Attemp to identify powdery mildew fungi by conidial characters. Trann. Brit. Mycology. Soc., 48: 553-558.

Zielinska, M. and M. Michniewicz (2001). The effect of calcium on the production of ethylene and abscisic acid by fungus *Fusarium culmorum* and by infected wheat seedlings
Acta Physiologiae Plantarum, 23 (1): 79-85.