


A decorative border of black shamrocks curves along the left side of the page, starting from the top and ending near the bottom left.

REFERENCES

A decorative border of black shamrocks curves along the bottom right of the page, starting from the bottom center and extending towards the right edge.

REFERENCES

- *Abenavoli, T., Rubber, S.; Fisher, V.; Axclrod, H. and Zuckerman, K.*

(1981):

Exercise testing with myocardial scintigraphy in asymptomatic diabetic males.

Circulation 63:54.

- *Acharya, U.; Shekhar, C. and Anamd, S. (1991):*

Lack of pain during myocardial infarction in diabetics is autonomic dysfunction responsible?

Am.J. Cardiol. 68:793.

- *Aronow, W.S. and Cassidy, J. (1975):*

Five years follow up of double master's test, maximal treadmill stress test and resting and post exercise apex cardiogram in asymptomatic persons. Circulation 52:616-8.

- *Aronow, W.; Anthany, D. and Epstein, S. (1992):*

Prevalence of silent myocardial ischaemia detected by 24-hours ambulatory ECG, and its association with new coronary events at 40 month follow up in elderly diabetic and non diabetic with coronary artery disease.

Am.J. Cardiol. 69:555.

- Assey, M.E.; Walters, G.L. and Hendix, G.H. (1987):

Incidence of acute myocardial infarction in patients with exercise induced silent myocardial ischaemia.

Am. J. Cardiol, 59:457.

- Attali, J.R.; Sachs, R.N; Valensi, P.; Palsky, D.; Tellier, P.; Vullpillat, M.; Lanfranch, J. and Sebayon, J. (1988):

Asymptomatic diabetic cardiomyopathy a non invasive study.

Diabetes Res. Clin. Pract. 4 : 183-190.

- Biagini, A.; Mazzei, M.G. and Carpeggioni, C.(1982):

Vasospastic ischaemic mechanism of frequent asymptomatic transient ST segment changes during continuous E.C.G. monitoring in selected unstable angina patients.

Am. H.J. 101:13-20.

- Bonow, R.O., Bacharch, S.L. Green, M.V. (1987):

Prognostic implications of symptomatic versus asymptomatic (silent) myocardial ischaemia induced by exercise in mildly symptomatic and asymptomatic patients with anginographically documented coronary artery disease.

Am. J. Cardiol. 60 778.

- Bradley, R.F.; Morble, A.; Withe, P. and Kraull, L.P. (1971):

Cardiovascular disease in joslin's diabetes mellitus, philadelphia, lea and febiger. publishers P 417

- Breisblatt, W.M.; Weiland, F.L. and McLain, J.R. (1988):

Usefulness of ambulatory radionuclide monitoring of left ventricular function early after acute myocardial infarction for predicting residual myocardial ischaemia.

Am. J. Cardiol. 62: 1005-1010.

- Bruce, R.A. (1969):

Stress testing in screening for cardiovascular disease.

Bull N.Y. Acad. Med. 45:1288-1295.

- Coupbell M-J and Machin D (1993) :

Medical statics- A common sense approach 2nd edition.

John Wiley & sons.

Chichester-New york, Brisbane, Toronto-singapore.

- Cecchi, A.C. and Dovellini, F. (1983):

Silent myocardial ischaemia during ambulatory.

E.C.G. monitoring in patients with effort angina.

J. Am. Coll. Cardiol. 1:934.

- Chiariello, M.; Indolfi, C. and condorelli, M.(1982):

La cardiopatia ischaemia silentenel diabetica in patri P.L.,

Cardiopatia ischaemia silente. Firenze P. 283.

- Cohn, P.F. (1985):

Silent myocardial ischaemia classification. prevalence and prognosis.

Am.J. Med. 79:2.

- Cohn, P.F. (1986):

Silent myocardial ischaemia and infarction.

Marcel Dekker, Inc., New York P 95.

- Cohn, P.F. (1989):

Mechanisms to explain the absence of pain in diabetes mellitus is a factor? In Cohn P.F. (ed) silent myocardial ischaemia and infarction ed 2.

New York Marcel Dekker PP 87-51.

- Connor, E. and Wingard, D.L. (1983):

Sex differential in ischaemic heart disease mortality in diabetics.

A prospective population study.

Am. J. epidemiol. 118 : 489-496.

- Cumming, G.R.; Samm, J.; Barys, K.L. and Kich, L. (1975):

Electrocardiographic changes during exercise in asymptomatic Men 3 year follow up.

Can. Med. Assoc. J. 112 : 57-81.

- Daniel W.W. (1987):

Biostatistics: A Foundation for analysis in the health sciences - 4th Ed. John Wiley & Sons.

- *Deanfield, J.E.; Maseri, A. and Selwyn, A.P. (1983):*

Myocardial ischaemia during daily life in patients with stable angina
its relation to symptoms and heart rate changes.

Lancet 2: 753-788.

- *Dean Field, J.E.; Shea, M. and Ribiero, P. (1984):*

Transient ST segment depression as a marker of myocardial

ischaemia during daily life.

Am. J. Cardiol. 54:1195-1200.

- *DeFronzo, R. (1987):*

Effect of physical training on insulin action in obesity.

Diabetes 36:1379.

- *Devine, S.M.; Laedtke, A.J. and Zelis, R. (1981):*

Coronary artery disease in diabetic patients in:

Clinical cardiology and diabetes, Vol 11. New york,

Future Publishing Co; PP 1, 87.

- *Devlin, J.T. ; Hirshman, M.; Harton, E.D. and Harton, E.S. (1986):*

Enhanced peripheral and splenchnic insulin sensitivity in NIDDM
men single baut of exercise.

Diabetes 36: 434-9.

- *Dewood, M.A. and Rozanski, A. (1986):*

Long term prognosis of patients with and without silent ischaemia.

Circulation 74(2):50

- Diamond, G.A. and Forrester, J.S. (1979):

Analysis of probability as an aid in the clinical diagnosis of coronary artery disease.

N. Engl. J. Med. 300:1350-1358.

- Doyle, J.T. and Kinch, S.H. (1970):

The prognosis of an abnormal electrocardiographic stress test.

Circulation 41:545 - 53.

- Droste, C. and Roskamm, H. (1983):

Experimental pain measurement in patients with asymptomatic myocardial ischaemia.

J. Am. Coll. Cardiol, 1:940.

- Ellested, M.H. and Kuan, P. (1984):

Nalozone and asymptomatic ischaemia failure to induce angina during exercise testing.

Am. J. Cardiol, 54:982.

- Erikssen, J. and Thaulow, E. (1984):

Follow up of patients with asymptomatic myocardial ischaemia in Rutish houser W, Roskamm H. (eds):

Silent myocardial ischaemia springer-verlag, Berlin, 156.

- Erupean Atherosclerosis Society Study on optimal Level of Serum Cholesterol (1987):

Lipid Revial Vol. I. No. 2 P. 11.

- Ewing, D. (1990):

Assessment of cardiovascular effects in diabetic autonomic neuropathy.

Ann. Int. Med. 92:308.

-Ewing D, Borsey D, Bellavere F, Clarke B (1981):

Cardiac autonomic neuropathy in diabetes; comparison of measures

of R-R interval variation.

Diabetologia 21:18.

-Ewing D, Campbell I, Murray A, Neilson J, Clarke B (1978):

Immediate heart - rate response to standing simple test for autonomic neuropathy in diabetes.

Br. Med. J. 1:145.

- Ewing D, Martyn C, Young R, Clarke B (1985):

The value of cardiovascular autonomic function tests; 10 year experience in diabetes.

Diabetes care; 8:491.

- Faerman, L. (1977):

Autonomic neuropathy and painless myocardial infarction in diabetic patients: histological evidence of their relationship.

Diabetes 26:1147.

- *Falcone, C.; Dservi, S. and Poma, E. (1971):*

Clinical significance of exercise induced silent myocardial ischaemia in patients with coronary artery disease.

J. Am. Coll. Cardiol. 9:295-299.

- *Falcone, C.; Guasti, L. and Corsica, G. (1987):*

Correlation between beta endorphin levels and anginal pain in patients with coronary artery disease.

J. Am. Coll. Cardiol. 9:176 A.

- *Fleg, J.; Lewis, C.; Costa, J. and Edward, E. (1990):*

Prevalence and prognostic significance of exercise induced silent myocardial ischaemia detected by thallium scintigraphy and E.C.G. in asymptomatic volunteers.

Circulation 81:428.

- *Froelicher, Jr. V.F., Tomas, M.M.; Pillow, C. and Lancaster, M.C. (1979):*

Epidemiologic study of asymptomatic men screened by maximal treadmill testing for latent coronary artery disease.

Am. J. Cardiol. 34: 770-6.

- *Gottlieb, S.O. (1989):*

Analysis of ST segment data scanning and interpretation.

J. Myocardial Ischaemia 1:55-61.

- *Gottlieb, S. (1991):*

Asymptomatic ischaemia in angina.

Clinical implications Cardiology Clinics Vol. 9 No.1 P 49.

- *Gottlieb, S.O. Gerstenblith, G. and Achuff, A.C. (1988):*

Ischaemia ST segment changes by ambulatory Holter predict one year mortality in high risk post infarct patients.

JAMA 259:1030.

- *Gottlieb, S.O.; Weisfeldt, M. L. and Ouyong, P. (1986):*

Silent ischaemia a marker for early unfavorable outcomes in patients with unstable angina.

N. Engl. Med. 314:1216.

- *Hauser, A.M.; Gangad Haran, V. and Ramos, R.G. (1985):*

Sequence of mechanical, E.C. G. and clinical effects of repeated coronary artery occlusion in human beings: echocardiographic observations during coronary angioplasty.

J. Am. Coll, Cardiol. 5:193.

- *Kannel, W.B.; Dannenberg, A.L. and Abbott, R.D. (1985):*

Un recognized myocardial infarction and hypertension the framingham study.

Am. H.J. 109: 585.

- *Kannel W, McGee D (1979):*

Diabetes and cardiovascular disease. Framingham study.

JAMA; 291:2035.

- *Kostmer G-M clin -chem (1976):*

22. 625.

- Kowalchuk, G.J. and Nesto, R.W. (1989):

Silent myocardial ischaemia mechanisms and rationale for therapy.

Am. J. Med. 86 (Suppl IA) : 9-13.

- Kralowski, A.S.; Kasinski, E.J.; Warrom, J.H.; Leland, O.S.; Busick, E.J.; Asmal, A.C.; Rand, L.I.; Christlieb, A.R.; Bradley, R.F. and Kahn, C.R. (1987):

Magnitude and determinants of C.A.D. in juvenile onset IDDM.

Am. J. Cardiol. 59:750-5.

- Landt, K.W.; Campaigne, B.N.; James, F.W. and Sperling, M.A. (1985):

Effects of exercise training in insulin sensitivity in adolescents with type II diabetes.

Diabetic care 8:461-5.

- Langer A, Freeman M, Steiner G, Armstrong P (1991):

Detection of silent myocardial ischaemia in diabetes mellitus.

Am.J. Cardiol . 67:1073.

- Lemp, G.; Zwag, R. and Hughes, J. (1987):

Association between the severity of diabetes and coronary atherosclerosis.

Am. J. Cardiol. 60 : 1015.

- Levin A (1966):

A simple test of cardiac function based upon the heart rate changes induced by Valsalva manoeuvre.

Am. J. Cardiol. 18:90.

- Levy, R.D.; Shapira, L.M. and Wright, T.C. (1986):

The hemodynamic significance of asymptomatic ST segment depression assessed by ambulatory pulmonary artery pressure monitoring.

Br. H. J. 56:526 - 530.

- Machac, J.; Technetium-99 Isanatriate (1989):

A perfusion or a viability agent? (editorial)

J. Am. Coll. Cardiol. 14:1685-1688.

- Malliani, A. (1986):

The elusive link between transient myocardial ischaemia and pain.

Circulation 73:201.

- Matehant-B; Umachandtan. V; Stenveson-R; Kopel man - PG; Timmis-AD (1994):

Silent myocardial ischaemia; role of subclinical neuropathy in patients with and without diabetes.

J. Am. Coll. Cardiol 22(5): 1433-7.

- **Margalis, J.R.; Kanne, W.B.; Feinleib, M.; Dowber, T.R.; M.C. and Namarh, P.M. (1973):**

Clinical features of unrecognized myocardial infarction silent and asymptomatic.

Am. J. Cardiol. 32:1.

- **Mark, D.B.; Halkky, A. and Califf, R.M. (1989):**

Painless exercise ST-segment deviation on the treadmill long term prognosis.

J. Am. Coll. Cardiol. 14:885-892.

- **Masseri, A.; Chierchia, S. and Davies, G. (1985):**

Mechanisms of ischaemic cardiac pain and silent myocardial ischaemia.

Am. J. Med. 29:7.

- **Medaliie, J.H. and Goldbaur, U. (1976):**

Unrecognized myocardial infarction, five year incidence, mortality and risk factors.

Ann. Intern. Med. 84: 526-531.

- **Mody, F.V.; Nademanec, K. and Intracat, V. (1988):**

Severity of silent myocardial ischemic in ambulatory E.C.G. monitoring in patients with stable angina pectoris; Relation to prognostic determinants during exercise stress testing and coronary angiography.

J. Am. Coll. Cardiol. 12: 1169-1176.

- Nable, E.G.; Tocco, M.N. and Selwyn, A.P. (1987):

Characteristics and significance of ischaemia detected by ambulatory E.C.G. monitoring.

Circulation 75 (Suppl V); V-74-V-83.

- Nademanee, Intarachat, V. and Kasephson, M.A. (1987):

Prognostic significance of silent myocardial ischaemia in patients with unstable angina.

J. Am. Coll. Cardiol. 10:1.

- Naka, M.; Hiramatsu, K. and Aizawa, Z. (1992):

Silent myocardial ischaemia in patients with NIDDM as judged by treadmill exercise testing and angiography.

Am. Heart. J. 123:46.

- Nave, M.P.H.; Elizabeth Barret Connor and Deborah L. Weingard (1990):

Resting E.C.G. abnormalities suggestive of asymptomatic ischaemic heart disease associated with NIDDM in defined population.

Circulation 81: 889-906.

- Nesto, R.W. and Phillips, R.T. (1986):

Asymptomatic myocardial ischaemia in diabetic patients.

Am. J. Med. 80 (Suppl 4 C): 40.

- *Nesto, R.W. and Kowolchuk, G.J. (1987):*

The ischaemic cascade; Lemporal sequence of haemodynamic, electrocardiographic and symptomatic expression of ischaemia.

Am. J. Cardio, 57:23 C.

- *Nesto, R.; Phillips, R. and Hill, T. (1988):*

Angina and exertional myocardial ischaemia in diabetic and non diabetic patients assessment by thallium scintigraphy.

Ann. Intern. Med. 108:170.

- *Niakan, E. (1986):*

Silent myocardial infarction and diabetic cardiovascular autonomic neuropathy.

Arch. Intern. Med. 146:2229.

- *Nikkila, E. (1984):*

Prevention of progression of coronary atherosclerosis by treatment of hyperlipidemia. A seven year prospective angiographics study.

B.M.J. 289:220.

- *Opasich, C.; Cobelli, F, and Assandri, J. (1984):*

Incidence and prognostic significance of symptomatic and asymptomatic exercise induced ischaemia in patients with recent myocardial infarction.

Cardiology 71:284.

- ***O'sullivan, JJ. Conroy-RM, Macdonald K. McKema TJ, Mauret BJ. (1991):***

Silent ischaemia in diabetic men with autonomic neuropathy.

Br. Heart. J. (66) 4: 313-15.

- ***Pan, W.H.; Cedres, L.C.; Liuk, Dyer, A.; Scheenberg, J.R. Shekilla, R.B.; Shekille, R.B.; Stomler, R.; Smith D.; Colette, P. and Stomler, J. (1986):***

Relationship of clinical diabetes and asymptomatic hyperglycemia to risk of coronary heart disease mortality in men and women.

Am. J. Epidemiol. 123: 504-516.

- ***Parmley, W. (1985):***

Silent ischaemia. Tip of the ice berg.

Am. J. Med. 79 (Suppl 3 A):1.

- ***Patriarmian, J.O. and Bradley, R.F. (1965):***

Acute myocardial infarction in 258 cases of diabetes mellitus mortality and 5-year survival.

N. Engl. Med. 273:455.

- ***Pierce, G.N.; Beamish, R.E. and Dhalla, N.S. (1988):***

Dysfunction of the cardiovascular system in diabetes, in pierce G.N., Beamish, R.E.; Dhalla, N.S. (eds).

Heart dysfunction in diabetes.

Boca Raton, Fla, C.R.C. Press, P. 51.

- Pyarals, K. and Laskaso, N. (1983):

Macrovascular disease in diabetes mellitus, in man II, puorala K.,
Tauscher, A (eds): diabetes in epidemiological prospective.
Edinburgh, Churchill Livingstons, PP. 208-209.

- Pyorala, K.; Laakso, U. and Usitupa, M. (1987):

Diabetes and atherosclerosis: an epidemiological view Diabetes
Met. Rev. 3: 463-524.

- Ranjadyan, K.; Unachandron, V. and Mill, G. (1990):

Prolonged anginal perceptual threshold in diabetes effects on
exercise capacity and myocardial ischaemia.
J. Am. Coll. Cardiol. 16:120.

- Ratner, S.J. (1988):

Clinical utility of myocardial perfusion imaging with thallium-201.
Cardiovascular Rev. Rep. 4: 42-46.

- Reismann, S.; Landheim, M. and Stamlof, H. (1985):

Asymptomatic patients with exercise TL 201 hypoperfusion:
identification of a high risk subset.
Circulation 72:75.

**- Roscoman, R.H.; Freedman, M.; Jenkins, D.; Strous, R.; Wurmm, M.
and Kositche K.R. (1967):**

Clinically unrecognized myocardial infarction in the western
collaborative group study.
Am. J. Cardiol. 19: 776-782.

- *Scott, R.C. (1975):*

Diabetes and the heart.

Aim. R.J. 90:283.

- *Selwyn, A.P., Deanfield, J. and Shea, M. (1986):*

Different pathophysiology of painful and silent myocardial ischaemia.

Circulation 74:57.

- *Serryus, P.A.; Winjs, W. and V.D. Brand M. (1984):*

Left ventricular performance, regional blood flow, wall motion and lactate accumulation during transluminal angioplasty.

Circulation 70:75.

- *Sheps, D.S.; Adam, K.F. and Hinderliter, A. (1987):*

Endorphins are related to pain perception in coronary artery disease.

Am. J. Cardiol. 59:523.

- *Smith, J.; Buckels, L. and Marcus, F. (1983):*

Clinical characteristics and results of non invasive tests in 60 diabetics after myocardial infarction.

Am. J. Med. 75:217.

- *Staut, R.W.; Bierman, E.L. and Rose, R. (1975):*

Effect of insulin in the proliferation of cultured primate arterial smooth muscle cells.

Circ. Res. 36: 319-27.

- Taggart T, H. and Stout, R.W. (1980):

Control of DNA synthesis in cultured vascular endothelial and smooth muscle cells response to serum. Platelet deficient serum lipid free serum insulin and Aestrogens.
Atherosclerosis 37: 549-57.

- Theroux, P.; Waters, D.D and Halphen, C. (1979):

Prognostic value of exercise testing soon after myocardial infarction.
N. Engl. J. and Med. 301:341.

- Tzivoni, D.; Gavish, A. and Zin, D. (1988):

Prognostic significance of ischaemic episodes in patients with previous myocardial infarction.
Am. J. Cardiol. 62: 661-664.

- Van Rijn (1981):

Effect of naloxone a specific opioid antagonist exercise induced angina.
Circulation 64 Suppl. N: 149.

- Watkins, P.J. (1977):

Facial sweating after food, A new sign of diabetic autonomic neuropathy. Br. Med. J. Vol., 1 P. 583.

- Watkins, P.J., and muckay, J.D. (1982):

Cardiac denervation in diabetic autonomic neuropathy An. inter.
Med 92 (2pt2) :304-307.

- Ward, J. D: Diabetic Neuropathy (1989):

British Medical Bulletin 45 (1) : 111 - 126.

- Weidinger, F.; Rannerle, A. and Sochar, H. (1986):

Role of beta endorphins in silent myocardial ischaemia.

Am. J. Cardiol. 58:428.

- Weiner, D.A.; Ryan, R.; M.C. and Cobe, C. (1987):

Significance of silent myocardial ischaemia during exercises testing
in patients with coronary aetery disease.

Am. J. Cardiol. 59:724.

- Wein̄er, D.A.; Ryan, T.J. and Mc Cabe, C.H. (1988):

Comparison of coronary artery by pass surgery and medical therapy
in patients with exercise induced silent myocardial ischaemia a
report from the coronary artery surgery study (CASS) registry.

J. Am. Coll. Cardiol. 12: 595-599.

- Weiner, D.A.; Tomas, J.; Parsonne, L. and Bernand, F. (1991):

Significance of silent myocardial ischaemia during exercise testing
in patients with diabetes: A reported form the coronary artery
surgery study registry.

Am. J. Cardiol. 68:729.

- *Weitzman, S.; Wagner, G.S.; Heiss, G.; Haney, T.L. and Sloman, G. (1932):*

Myocardial infarction site and mortality in diabetes.

Diabetes Care 5:31.

- *Wohlgelehter, D.; Joffe, C.C. and Cabin, H.S. (1987):*

Silent ischaemia during coronary occlusion produced by balloon

inflation: Relation to regional myocardial dysfunction.

J. Am. Coll. Cardiol. 10:491.

- *Yaung R. and Chanl (1984):*

Gastrointestinal tract complication of diabetes mellitus.

Arch. Int. Med. 44. 1251-1255.

- *Young D.S. Pestaner. L.C. and Gibberman (1975):*

V. Clin. Chemi. 21 Na5.