

Reference

Asiri S, Hasham A, al Anazy F, Zakzouk S, Banjar A
Tympanosclerosis : rreview of literature and incidence among
patients with middle –ear infection. J Laryngol Otol 1999; 113:
1076-80.

Babonis TR, Weir MR, Kelly PC. Impedance tympanometry
and acoustic reflectometry at myringotomy. Pediatrics 1991; 87:
475-480.

**Bayazit YA, Ozer E, Kara C, Gokpinar S, Kan Likama M
and Mumbuc.** An analysis of the single-stage tympanoplasty
with over-under lary grafting in tympanosclerosis. Otol.
Neurotol. 2004; 25(3): 211-4.

Bell EF. History of vitamin E in infant nutrition. Am J Cli Nutr
1987; 46: 183.

Bieri JG, Corash L, Hubbard VS. Medical uses of vitamin E.
N Eng1 J Med 1983; 308(18): 1063.

Boxall, j., Proops, D., and Michaels, L.: The specific
locomotiveactivity of tympanic membrane and cholesteatoma
epithelium in tissue culture. J. Otolaryngol. 17(4): 104 -144,
1988.

Chow CK. Vitamin E. In: Chen LH, ed. Nutritional aspects of aging. Vol I. Boca Raton, FL: CRC Press, 1986.

C. Mattson, K. Magnuson, S. Hellstrom. Myringosclerosis caused by increased oxygen concentration in traumatized tympanic membranes. *Ann. Otol. Rhinol. Laryngol* 1995; 104: 625-632.

Cüneyd Uneri, Tekin Baglam, Mine Yazici. The effect of vitamin E treatment on the edevelopment of myringosclerosis after ventilation tube insertion. *nternational Journal of : diatric,- Otorhinolaryngology* 2006; 70: 1045-1048.

Dempster JH, Mackenzie K. Tympanometry in the detection of hearing impairments associated with otitis media with effusion. *Clin Otolarngol* 1991;16:157-159

D. N. Riley, S. Herberger, G. McBride and K. Law. Myringotomy and ventilation tube insertion: a ten year follow up. *J. Laryngol. Otol.* 1997; 111: 257-261.

Fanton JC and Ward PA: Role of oxegen-derived free radicals and metablites in leukocyte-dependent inflammatory reaction 1982.

Forseni Flodin M. Macrophages and possible osteoclast differentiation in the rat bullar bone during experimental acute otitis media, with reference to tympanosclerosis. *Otol Neurotol* 2001; 22: 771-5.

Forseni M, Bagger- Siorback D, Hultcrantz S: A study of inflammatory mediators in the human tympanosclerotic middle ear. *Arch Otolaryngol Head Neck Surg*, 2001;127:559-564.

Forseni M, Hultcrantz M. Possible inflammatory mediators if tympanosclerosis development. *Int J Pediatr Otorhinolaryngol* 2002; 63(2): 149-54.

Friedman EM, Sprecher RC, Simon S, Dunn JK. Quantitation and prevalence of tympanosclerosis in a pediatric otolaryngology clinic. *Int J Pediatr Otorhinolaryngol* 2001; 60: 205-11.

Gibb AG. "Tympanosclerosis" *Acta. Otorhinolaryngologica*, Belgica 1971; 25: 956.

Gorur K, Ozcan C, Polat A, Unal M, Tamer L, Cinel I. The anti-oxidant and anti-apoptotic activities of selenium in the prevention of myringosclerosis in rats. *J Laryngol Otol* 2002; 116: 426-9.

Govaerts P, Jacob W and Marquet J. Histological study of the thin replacement membrane of humane tympanic membrane perforations .Acta Otolaryngol. (stockh.) 105: 297-302, 1988.

Grievink EH, Peters SAF, van Bon WHJ, et al. The effect of early bilateral otitis media with effusion on language ability : Aprospective cohort study. J Speech Hearing Res 1993; 36: 1004-1012.

Gutteridge JM and Halliwell B: Free redicals and antioxidants in the year 2000.Ahistorical look to the future 2000. **Ann N Y Acad Sci**; 899: 136.

Hallfrisch J, Muller DC, Singh VN. Vitamin A and E intakes and plasma concentrations of retinol, beta- carotene and alphatucopherol in men and women of the Baltimore Longitudinal Study of Aging. Am J Clin Nutr 1994; 60:176.

Harrison H, Fixsen A, Vickers A. A randomized comparison of homeopathic and standard care for the treatment of glue ear in children. Complementary Ther Med 1999;7:132-135.

Hemlin C, Carenfelt C, Papatziamos G. Single dose of betamethasone in combined medical treatment of secretory otitis media. Ann Otol Rhinol Laryngol 1997; 106:359-363.

Johnston LC, Feldman HM, Paradise JL, Bernard BS, Colborn DK, Casselbrant ML and Janosky JE. Tympanic membrane abnormalities and hearing levels at the ages of 5 and 6 years in relation to persistent otitis media and tympanostomy tubes insertion in the first 3 years of life: a prospective study incorporating and randomized clinical trial. *Pediatrics* 2004; 114(1): 58-67.

Karlidag T, Ilhan N, Kaygusuz I, Keles E and Yalcin S. Comparison of free radicals and antioxidant enzymes in chronic otitis media with and without tympano-sclerosis. *Laryngoscope* 2004; 114(1): 85-9.

Koc A, Uneri C. Genetic predisposition for tympanosclerotic degeneration. *EUR Arch Otorhinolaryngol* 2002;259:180-3.

Koc A, Uneri C. Sex distribution in children with tympanosclerosis after insertion of a tympanostomy tube . *Eur Arch Otorhinolaryngol* 2001;258:16-9.

Koehler KM, Garry PJ. Nutrition and aging. *Clin Lab Med* 1993; 13: 433.

Kuczkowski J, Bakowska A & Mikaszewski B: Microstructure of tympanic membrane in chronic otitis media:

immunohistochemical evaluation. *Otolaryngol Pol.*; 58(2): 275-80, 2004.

M.H. Bhaya, P.A. Schachern, T. Morizono, M.M. Paperella, Pathogenesis of tympanosclerosis, *Otolaryngol. Head Neck Surg.*109 (1993): 413-420.

Machlin LJ. Vitamin E. In: Machlin LJ, ED. Handbook of vitamins. 2nd ed . New York: Marcel Dekker, 1991.

Makiishi-Shimobayashi C, Tsujimura T, Sugihara A, et al. Expression of osteopontin by exudates macrophages inflammatory tissues of the middle ear: a possible association with development of tympanosclerosis, *Hear Rs* 2001; 153: 100-7.

Mandel EM, Casselbrant ML, Rockette HE, et al. Efficacy of antimicrobial prophylaxis for recurrent middle ear effusion. *Pediatr Infect Dis* 1996;15:1074-1082.

Mannaerts GP and Pvan Vveldhoven PP: Role of proxisomes in mammalian metabolism. *Cell Biochem Funct*; 10:141, 1992.

Marchisio P, Principi N, Passali D, et al. Epidemiology and treatement of otitis media with effusion in children in their first

year of primary school. Acta Otolaryngol (Stockh) 1998; 118: 557-562.

Marie FF: Inflammatory mediators and immunocompetent cells in the middle ear with particular regards to otitis media and tympanosclerosis. Acta Otolaryngol; 120(5): 577-82, 2000.

Mattsson C, Berggren D, Hellstrom S. Myringotomized tympanic membranes cultured in vitro do not develop myringosclerosis. Acta Otolaryngol (Stockh) 2002;122:168-72.

Mattsson C, Johansson C, Hellstrom S. Myringosclerosis develops within 9h of myringotomy. ORL J Otorhinolaryngol Relat Spec 1999;61:31-6.

Mattsson C, Stierna P, Hellstrom S: Treatment of myringosclerosis after myringotomy, Am J Otol 2000;21:804-8.

Maw AR, Bawden R. The long term outcome of secretory otitis media in children and the effects of surgical treatment: a ten year study. Acta Otorhinolaryngol Belg 1994; 48: 317-324.

Maw R, Bawden R. Spontaneous resolution of severe chronic glue ear in children and the effect of adenoidectomy,

tonsillectomy, and insertion of ventilation tubes (grommets).
BMJ 1993; 306: 756-760.

Maw R, Wilks J, Harvey I, et al. Early surgery compared with watchful waiting for glue ear and effect on language development in preschool children : a randomized trial. Lancet 1999; 353: 360-363.

Mayes PA. Biologic oxidation. In: Murray R, Granner D, Mayes P and Rodwell V, eds, Harber's Biochemistry, 25 edition, California: Appelton Lang, 2000.

McCord JM and Fridovich I: Superoxide dismutase: and enzymatic function for erythrocuprein (hemocuprein). J Biol Chem; 224: 6049, 1969.

M. Tos, P. Bonding, G. Poulsen. Tympanosclerosis of the drum in secretory otitis after insertion of grommets. A prospective, comperative study, J. Laryngol. Otol. 97(1983): 489-496.

Nasr GF, Nassar M, El-Begermy M, Kassim SK and Khalil R: Review of literature. In : Retrospective study of antioxidant levels in chronic sinusitis patients, Thesis, Ain Shams University Publication, Cairo, Egypt (2002).

Nelson JS, Fischer VW. Vitamin E. In:Barker BM, Bender DA, eds. Vitamins in Medicine. 4th ed. Vol 1. London: William Heinemann, 1981.

O'Donoghue: Tympanic epithelium: An ultrastructural, experimental and clinical study .J.R . Soc. Med.77:758-760, 1984.

Orlin MN, Effgen SK, Handler SD. Effects of otitis media with effusion on gross motor ability in preschool-aged children: preliminary finding. Pediatrics 1997;99:334-337.

Ozcan C , Polat G, Gorur K, Talas DU, Unal M, Cinel I. The inhibitory effect of topical N-acetylcysteine application on myringosclerosis in perforated rat tympanic membrane. Int J Pediatr Otorhinolaryngol 2002;63:179-84.

Ozcan C, Polat G, Gorur K, Talas DU, Bagdatoglu O, Cinel I, The effect of local administration of N-acetylcysteine in perforated rat tympanic membrane; an experimental study in myringosclerosis, Pharmacol Res 2002;45:5-9.

Paradise JL, Bluestone CD, Colborn DK, et al. Adenoidectomy and adenotonsillectomy for recurrent acute otitis media . JAMA 1999; 282: 945-953

Paradise JL, Dollaghan CA, Campbell TF, et al. Language, speech sound production, and cognition in three-year-old children in relation to otitis media with effusion in their first three years of life. Pediatrics 2000; 105: 1119-1130.

Paradise JL, Feldman HM, Campbell TF, et al: Early versus delayed insertion of tympanostomy tubes for persistent otitis media on developmental outcomes at the age of three years in relation to prerandomization illness patterns and hearing levels, Pediatr Infect Dis J 2003;22:309-14.

Parke DV: Nutritional antioxidants and Disease prevention: mechanisms of action. In: Basu TK, Temple NJ and Garg ML, eds. Antioxidants in human Health, Chapter 1,1 edition, Wallingford, UK: CAB International publishing 1999.

Pellett FS, Cox LC, MacDonald CB. Use of acoustic reflectometry in the detection of middle ear effusion. J Am Acad Audiol 1997;8:181-187

Pesci and Pickett: Genetic organization and enzymatic activity of superoxide dismutase from the microaerophilic human pathogen; *Helicobacter pylori*. *Gene*; 143 (1): 1, 1994.

Pirodda A, Ferri GG, Bruzzi C, Marini M & Faggioli G: Possible relationship between tympanosclerosis and atherosclerosis. *Acta Otolaryngol.*; 124(5): 574-6, 2004.

P.J. Dawes, B.J. Bingham, R. Ryhs, M.V. Griffiths. Aspirating middle ear effusions when inserting ventilation tubes; does it influence post-operative otorrhoea, tube obstruction or the development of tympanosclerosis? *Clin. Otolaryngol.* 16 (1991): 457-461.

Preston K. Pneumatic Otoscopy: Review of the literature. *Issues in Comprehensive Pediatric Nursing* 1998; 21: 117-128

Rach GH, Zielhuis GA, van Baarle PW, et al. The effect of treatment with ventilating tubes on language development in preschool children with otitis media with effusion. *Clin Otolaryngol* 1991;16:128-132

Reidpath DD, Glasziou PP, Del Mar CB. Systematic review of autoinflation for treatment of glue ear in children. *BMJ* 1999; 318: 1177-1178.

Roberts JE, Burchinal MR, Zeisel SA. Otitis media in early childhood in relation to children's school-age language and academic skills, *Pediatrics* 2002;110:696-706.

Rosenfeld RM, Mandel EM, Bluestone CD. Systemic steroids for otitis media with effusion in children. *Arch Otolaryngol Head Neck Surg* 1991;117:984-989.

Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA: The effect of ventilation tubes on language development in infants with otitis media with effusion: a randomized trial, *Pediatrics* 2000;106:E42.

Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA. Generalizability of trial results based on randomized versus nonrandomized allocation of OME infants to ventilation tubes or watchful waiting. *J Clin Epidemiol* 2001; 54: 789-94.

Rovers MM, Straatman H, Ingels K, van der Wilt GJ, van den Broek P, Zielhuis GA: The effect of short- term ventilation tubes versus watchful waiting on hearing in young children with persistent otitis media with effusion: a randomized trial. *Ear Hear* 2001; 22: 191-9.

Russel JD & Giles JJ: Tympanosclerosis in the rat tympanic membrane: an experimental study. *Laryngoscope.*; 112(9): 1663-6, 2002.

S.S. Costa, M.M. Paparella, P.A. Schachern, T.H. Yoon, B.P. Kimberly, Temporal bone histopathology in chronically infected ears with intact and perforated tympanic membranes, *Laryngoscope* 102, (1992): 1229-1236.

Saunders, R., and Weider, D.: Tympanic membrane sensation. *Brain* 108:387-404, 1985.

Schilder AG: Assessment of complications of the condition and of the treatment of otitis media with effusion. *Int J Pediatr Otorhinolaryngol* 1999; 49 (suppl 1) S 247-S251.

Shriberg LD, Friel-Patti S, Flipsen P, et al. Otitis media, fluctuant hearing loss, and speech- language outcomes: a preliminary structural equation model. *J Speech Language Hear Res* 2000; 43: 100-120.

Sies H: Oxidative stress from basic research to clinical application. *Am J Med*; 91 (suppl 3c) :31s, 1991.

Simsons, M.R.: Acoustic impedance tests. Ear disease, deafness and dizziness, Goodhill, V. Laryngoscope, 87:112-114, 1979.

Southern PA and Powis G: Free radicals in medicine: chemical nature and biologic reactions. Mayo Clin proc: 63:381, 1998.

S. Polat, O. Ozturk, C. Uneri, M. Yuksel, G. Haklar, S. Bozkurt and S. Kullu. Determination of reactive oxygen species in myringotomized tympanic membranes: effect of vitamin E treatment. Laryngoscope 2004; 114: 720-725.

Spratley JE, Hellstrom SO, Mattsson CK, Pais-Clemente M, Topical ascorbic acid reduces myringosclerosis in perforated tympanic membranes, A study in the rat, Ann Otol Rhinol Laryngol 2001; 110: 585-91.

Statland B. Vitamins and minerals: passing fads or keys to health? Med Lab Observ 1993; 25(12): 21.

Stool SE, Berg AO, Berman S, et al. Otitis media with effusion in young children. Clinical practice guideline. Number 12. Rockville, Maryland: Agency for Health Services, US Department of Health and Human Services, 1994.

Takahashi H, Honjo I, Hasebe S, et al. The diagnostic and prognostic value of eardrum mobility in OME. *Eur Arch Otol Rhinol Laryngol* 1999; 256: 189-191.

Tolbert NE: Metabolic pathways in peroxisomes and glyoxysomes. *Ann Rev Biochem*; 50: 133, 1981.

Traber MG, Ingold KU, Burton GW, Kayden Hj. Absorption and transport of deuterium- substituted 2R, 4', 8'R-alpha-tocopherol in human lipoproteins. *Lipids* 1988; 23:791.

Tracy JM, Demain JH, Hoffman KM, et al. Intranasal beclomethasone as an adjunct to treatment of chronic middle ear effusion. *Ann Allergy Asthma Immunol* 1998;80:198-206.

Van Balen FAM, Aarts AM, De Melker RA. Tympanometry by general practitioners: reliable? *Int J Pediatr Otorhinolaryngol* 1999;48:117-123.

Varsano IB, Volovitz BM, Grossman JE. Effect of naproxen, a prostaglandin inhibitor, on acute otitis media and persistence of middle ear effusion in children. *Ann Otol Rhinol Laryngol* 1998; 89: 389-392.

Vernon-Feagans L, Emanuel DC, Blood I. the effect of otitis media and quality of day care on children's language development. J Appl Dev Psychol 1997;18:395-409.

Vernon-Feagans L, Manlove EE, Volling BL. Otitis media and the social behaviour of day-care-attending children. Child DEV 1996;67:1528-1539.

Wajnberg, J.: The true shape of the tympanic membrane .J. Laryngol .Otol.101:538-541, 1987

Wielinga EW, Kerr AG. Tympanosclerosis. Clin Otolaryngol Allied Sci 1993; 18(5): 341-9.

Wonchang I. "Tympanosclerosis". Acta. Otolaryngoloigca, Stockholm 1969; 68: 62.

Woodford HP and whitehead TP: Is measuring serum antioxidant capacity clinically usetull? Ann Clin Biochem; 35: 48, 1998.

Yariktas M, Doner F, Dogru H, Yasan H and Delibas N. The role of fre oxygen radicals on the development of otitis media with effusion. Int J Pediatr Otorhinolaryngol 2004; 68(7): 889-94.