

REFERENCES

1. Brandstrup B, Tonnesen H, Beir Holgersen R, Hjortso E, Ording H, Lindorff - Larsen K, Rasmussen MS, Lanng C, Wallin L, Iversen LH, B, Pott F; Danish Study Group on Perioperative Fluid Therapy. Effects of intravenous fluid restriction on postoperative complications: comparison of two perioperative fluid regimens. A randomized assessorblinded multicenter trial. *Ann Surg* 2003;238:641- 8.
2. Nisanevich V, Felsenstein I, Almog G, Weissman C, Einav S, Matot I. Effect of intraoperative fluid management on outcome after intraabdominal surgery. *Anesthesiology* 2005; 103:25-32.
3. *Hepatobiliary Pancreat Dis Int* 2008; 7:J67-372.
4. Sauaia A, Moore FA, Moore EE, Moser KS, Brennan R, Read T a accordance with the recommendation above, RA, et al. Epidemiology of trauma deaths: A reassessment. J consideration should be given to administration en *Trauma* 1995;38:185-93.
5. Acosta JA, Yang JC, Winchell RJ, Simons RK, Fortlage DA, indicated in the prehospital setting, crystalloid in a level 1 trauma centre. *J Am Coll Surg* 1998; 186:528-33.
6. Gillham MJ, Pan' MJ. Resuscitation for major trauma. *Curr Opin* targets could be established, that will maintain better *Anaesthesiol* 2002; 15:167-72.
7. Prough Ds, White RT: Acidosis associated with perioperative saline administration: dilution or delusion? *Anesthesiology* 2000;93:1 167-9.
8. Liskaser FJ, Bellomo R, Hayhoe M, et al: Role of pump prime in the etiology and pathogenesis of cardiopulmonary bypass-associated acidosis *Ariesthesiology* 2006;93: 1170-3.

9. Siker D. Paediatric fluids, electrolytes and nutrition. In: George A Gregory, editor. *Paediatric Anaesthesia*. Churchill Livingstone 4th edition, 2002: 85-116.
10. Joshi GP. Intraoperative fluid restriction improves outcome after major elective gastrointestinal surgery. *Anesth Analg*. 2006; 101:601-605.
11. Cook-Sather SD, Harris KA, Chiavacci R, Gallagher PR, Schreiner MS. A liberalized fasting guideline for formula fed infants does not increase average gastric fluid volume before elective surgery. *Anesth Analg* 2003; 96: 965-969.
12. Lobo DN, Macafee DAL, Allison SP. How perioperative fluid balance influences postoperative outcome. *Best Prac Res Clin Anaesthesiol*. 2006; 20:439-455.
13. Chertow GM, Burdick E, Honour M, Bonventre JV, Bates DW. Acute rXkidney injury, mortality, length of stay, and costs in hospitalized patients. *J Am Soc Nephrol*. 2005; 16:3365-3370.
14. Sear JW. Kidney dysfunction in the postoperative period. *Br J Anaesth*. 2005; 95:20-32.
15. Holte K, Kehlei H. Fluid therapy and surgical outcomes in elective 1' surgery: A need for reassessment in fast-track surgery. *J Am Coll Surg*. 2006; 202:971-989.
16. Bundgaard-Nieisen M, Holte K, Secher NH, Kehlet H. Monitoring of perioperative fluid administration by individualized goal-directed therapy. *A eta Anaesthesiol ScandL* 2007; 51:331 -340.
17. Holte K, Sharrock NE, Kehlet H. Pathophysiology and clinical implications of perioperative fluid excess. *Br J Anaesh* 2003; 89:622-32.

18. Nisanevich V, Felsenstein I, Almogy G, et al. Effect of intraoperative fluid management on outcome after intraabdominal surgery. *Anesthesiology* 2005; 103: 25-32.
19. Berne R M, levy M N. *Physiology*. 5th ed. Philadelphia: Mosby, 2005.
20. Halperin M l, goldstein M B. *Fluid electrolyte and acid base physiology*. 3rd ed. Philadelphia: W B saunders, 1999.
21. Ganong W F. *Review of medical physiology*. 17th ed. London: Prentice Hall, 1995.
22. Guyton AC, Hall JE. The body fluid compartments: extracellular and intracellular fluids; interstitial fluid and edema. In: *Textbook of Medical Physiology*. 10th ed. Philadelphia: WB Saunders Company, 2000:264–278.
23. Bundgaard M. The three dimensional organization of tight junctions in a capillary endothelium revealed by serial-section electron microscopy. *J Ultrastruct Res* 1984; 88:1–17.
24. Rippe B, Rosengren B-I, Venturoli D. The peritoneal microcirculation in peritoneal dialysis. *Microcirculation* 2001; 8:303–320.
25. Michel CC, Curry FE. Microvascular permeability. *Physiol Rev* 1999; 79:703–761.
26. Rippe B, Haraldsson B. Transport of macromolecules across microvascular walls: the two-pore theory. *Physiol Rev* 1994; 74:163–219.
27. Rippe B, Rosengren B-I, Venturoli D. The peritoneal

- microcirculation in peritoneal dialysis. *Microcirculation* 2001; 8:303–320.
28. Ringer S. Regarding the action of the hydrate of soda, hydrate of ammonia, and the hydrate of potash on the ventricle of the frog's heart. *J Physiol* 1880;3: 195–202.
29. Haljamae H. Use of fluids in trauma. *Int J Intens Care* 2000; 6:20–30.
30. Tonnesen AS. Crystalloids and colloids. In: Ronald MD, Miller D, eds. *Anesthesia*. 4th ed. New York, Edinburgh, London, Madrid, Melbourne, Milan, Tokyo: Churchill Living stone, 1994: 1595–1617.
31. Svensen C, et al. Stability of the interstitial matrix after crystalloid fluid loading studied by volume kinetic analysis. *Br J Anaesth* 1999; 82:496–502.
32. Wiig H, Rubin K, Reed RK. New and active role of the interstitium in control of interstitial fluid pressure: potential therapeutic consequences. *Acta Anaesth Scand* 2003; 47:111–121.
33. Drobin D, Hahn RG. Volume kinetics of Ringer's solution in hypovolemic volunteers. *Anesthesiol* 2000; 90:81–91.
34. Morgan TJ, Venkatesh B, Hall J. Crystalloid strong ion difference determines metabolic acid-base change during in vitro hemodilution. *Crit Care Med* 2002; 30:157–160.
35. Kellum JA. Saline-induced hyperchloremic metabolic acidosis. *Crit Care Med* 2005; 30:259–261.

36. Stephens R, Mythen M. Optimizing intraoperative fluid therapy. *Acta Anaesthesiol Scand* 2006; 16:385–392.
37. Wilkes NJ, et al. The effects of balanced versus saline based hetastarch and crystalloid solutions on acid-base and electrolyte status and gastric mucosal perfusion in elderly surgical patients. *Anesth Analg* 2005; 93:811–816.
38. Waters JH, et al. Normal saline versus lactated Ringer's solution for intraoperative fluid management in patients undergoing abdominal aortic aneurysm repair: an outcome study. *Anesth Analg* 2003; 93:817–822.
39. Reid F, et al. (Ab) normal saline and physiological Hartmann's solution: a randomized double-blind cross over study. *Clin Sci* 2006; 104:17–24.
40. Arieff AI. Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *New Engl J Med* 1986; 314:1529–1535.
41. Strandberg P, Hahn RG. Volume kinetics of glucose 2.5% given by intravenous infusion after hysterectomy. *Br J Anaesth* 2005; 94:30–38.
42. Haggstrom J, Hedlund M, Hahn RG. Subacute hyponatraemia after transurethral resection of the prostate. *Scand J Urol Nephrol* 2001; 35:250–251.
43. Vincent JL. Fluid management: the pharmacoeconomic dimension. *Crit Care* 2000; 4 (suppl2): S33–S35.
44. Lindow J, Wijdicks EF. Phenytoin toxicity associated with

- hypoalbuminemia in critically ill patients. *Chest* 1994; 105:602–604.
45. Mimoz O, Soreda S, Padoin C, Tod M, Petitjean O, Benhamou D. Ceftriaxone pharmacokinetics during iatrogenic hydroxyl ethyl starch-induced hypoalbuminemia: a model to explore the effects of decreased protein binding capacity on highly bound drugs. *Anesthesiology* 2003; 93:735–743.
46. Quinlan GJ, Margaron MP, Mumby S, Evans TW, Gutteridge MC. Administration of albumin to patients with sepsis syndrome: a possible beneficial role in plasma thiol repletion. *Clin Sci* 1998; 96:459–465.
47. Zoellner H, Hou JY, Lavery M, et al. Inhibition of microvascular endothelial apoptosis in tissue explants by serum albumin. *Microvascular Res* 1999; 57:162–173.
48. Von Hoegen I, Waller C. Safety of human albumin based on spontaneously reported serious events. *Crit Care Med* 2001; 29:994–996.
49. Treib J, Haass A, Pindur G. Coagulation disorders caused by hydroxyl ethyl starch. *Thromb Haemost* 1997; 78:974–983.
50. Strauss RG, Stump DC, Henriksen RA. Hydroxy ethyl starch accentuates von Willen brand's disease. *Transfusion* 1985; 25:235–237.
51. Mardel SN, Saunders FM, Allen H, et al. Reduced quality of clot formation with gelatin-based plasma substitutes. *Br J Anaesth* 1998; 80:204–207.

52. Shoemaker WC. Comparison of the relative effectiveness of whole blood transfusions and various types of fluid therapy in resuscitation. *Crit Care Med* 1976; 4:71–78.
53. Cohn JN, Luria MH, Daddario RC. Studies in clinical shock and hypotension. V. Hemodynamic effects of dextran. *Circulation* 1967; 35:316–326.
54. Kreimeier U, Frey L, Messmer K. Small-volume resuscitation. *Curr Opin Anaesthesiol* 1993; 6: 400–408.
55. Wolf MB. Plasma volume dynamics after hypertonic fluid infusing in nephrectomized dog. *Am J Physiol* 1971; 221:1392–1395.
56. Kramer GC, et al. Hyperosmotic-hyperoncotic solutions. *Bailliere's Clin Anaesthesiol* 1997; 11(1):143–161.
57. Kramer GC, Townsend M, Deyo DJ. Hypertonic Resuscitation Website, 1997.
58. Vassar MJ, Holcroft JW. Use of hypertonic-hyperoncotic fluids for resuscitation of trauma patients. *J Intensive Care Med* 1992; 7:189–198.
59. Svensen C, Hahn RG. Volume kinetics of Ringer solution, dextran 70, and hypertonic saline in male volunteers. *Anesthesiology* 1997; 87(2):204–212.
60. Drobin D, Hahn RG. Kinetics of isotonic and hypertonic plasma volume expanders. *Anesthesiology* 2002; 96:71–80.
61. Mazzoni MC, et al. Amiloride-sensitive Na^+ pathways in capillary endothelial cell swelling during hemorrhagic shock. *J*

- Appl Physiol 1992; 73(4):1467–1473.
62. Junger WG, et al. Hypertonic saline resuscitation: a tool to modulate immune function in trauma patients? Shock 1997; 8:235–241.
 63. Kolsen-Petersen JA. Immune effect of hypertonic saline: fact or fiction. Acta Anaesthesiol Scand 2004; 48:667–678.
 64. Kramer GC. Hypertonic resuscitation: physiological mechanisms and recommendations for trauma care. J Trauma-Injury Infection and Critical Care 2003; 54(5):589–599.
 65. Riddez L, et al. Lower dose of hypertonic saline dextran reduces the risk of lethal rebleeding in uncontrolled hemorrhage. Shock 2001:1–6.
 66. Sandham JD, et al. A randomized, controlled trial of the use of pulmonary-artery catheters in high risk surgical patients. N Eng J Med 2003; 348(1):5–13.
 67. Riddez L, et al. Lower dose of hypertonic saline dextran reduces the risk of lethal rebleeding in uncontrolled hemorrhage. Shock 2001:1–6.
 68. Alam H, et al. Hypertonic saline: intraosseous infusion causes myonecrosis in a dehydrated swine model of uncontrolled hemorrhagic shock. J Trauma 2002; 52:18–25.
 69. Sloan EP, Koenigsberg MD, Bickell WH, et al. The use of diaspirin cross-linked hemoglobin (DCLHb) solution in the hospital management of hemorrhagic shock [abstr78]. Acad Emerg Med 1995;2.

70. Proceedings of IBC's Conference on Blood Substitutes and Related Products. Cambridge :1997.
71. Dubick MA. Prolonged hypotensive resuscitation HBOCs, Yeahor Nay? Swine experiments. Advanced Technology Applications for Combat Casualty Care (ATACCC). St. Pete Beach, FL, 2003.
72. Handrigan M. Prolonged hypotensive resuscitation HBOCs, Yeahor Nay? Rat experiments. Advanced Technology Applications for Combat Casualty Care (ATACCC). St. Pete Beach, FL, 2003.
73. Winslow RM. Methods and compositions for optimization of oxygen transport by cell-free systems. US Patent 6, 054, 427, The Regents of the University of California, 2000.
74. Mc Carthy MR, Vandegriff KD, Winslow RM. The role of facilitated diffusion in oxygen transport by cell-free hemoglobins : implications for the design of hemoglobin based oxygen carriers. *Biophys Chem* 2001; 92:103–117.
75. Winslow RM. Current status of blood substitute research: towards a paradigm. *J Int Med* 2003; 253:508–517.
76. Vandegriff KD, Malavalli A, Wooldridge J, Lohman J, Winslow RM. MP4, a new non vasoactive PEG-Hb conjugate. *Transfusion* 2003; 43:509–516.
77. Jackson J, Bolte RG. Risks of intravenous administration of hypotonic fluids for pediatric patients in ED and prehospital settings: let's remove the handle from the pump. *Am JE merg*

- Med 2000; 18:269–270.
78. Carcillo JA, Davis AL, Zaritsky A. Role of early fluid resuscitation in pediatric septic shock. *JAMA* 1991; 266:1242–1245.
 79. Berry F. Practical aspects of fluid and electrolyte therapy. In :Berry F, ed . *Anesthetic Management of Difficult and Routine Pediatric Patients*. New York: Churchill Living stone, 1986: 107–135.
 80. Baines DB, Murrell D. Preoperative hypoglycaemia, propranolol and the Jervell and Lange–Nielsen syndrome. *Paediatr Anaesth* 1999; 9:156–158.
 81. Bush GH, Steward DJ. Severe hypoglycaemia associated with preoperative fasting and intraoperative propranolol. A case report and discussion. *Paediatr Anaesth* 1996; 6:415–417.
 82. Steward DJ, DaSilva CA, Flege IT. Elevated blood glucose levels may increase the danger of neurological deficit following profoundly hypothermic cardiac arrest [letter]. *Anesthesiology* 1988; 68:653.
 83. Bush GH, Steward DJ. Can persistent cerebral damage be caused by hyperglycaemia? *Paediatr Anaesth* 1995; 5:385–387.
 84. Steingrub JS, Mundt DJ. Blood glucose and neurologic outcome with global brain ischemia. *Crit Care Med* 1996; 24:802–806.
 85. Bush GH, Steward DJ. Can persistent cerebral damage be caused by hyperglycaemia ? *Paediatr Anaesth* 1995; 5:385–

- 387.
86. Wong W, Fok TF, Lee C, et al. Randomised controlled trial: comparison of colloid or crystalloid for partial exchange transfusion for treatment of neonatal polycythaemia. *Arch Dis Child Fetal Neonatal* 1997; 76:F115–F118.
 87. Deak in CD, Hicks IR. A Bor ABC: prehospital fluid management in major trauma. *J Accid Emerg Med* 1994; 11:154–157.
 88. Sander Jensen K, Marving J, Secher NH, et al. Does the decrease in heart rate prevent a detrimental decrease of the end-systolic volume during central hypovolemia in man? *Angiology* 1990; 41: 687–695.
 89. Price HL, Deutsch S, Marshall BE, Stephen GW, Behar MG, Neufeld GR. Hemodynamic and metabolic effects of hemorrhage in man, with particular reference to the splanchnic circulation. *Circ Res* 1966; 18:469–474.
 90. Jackson R, Reid JA, Thorburn J. Volume preloading is not essential to prevent spinal-induced hypotension at caesarean section. *Br J Anaesth* 1995; 75(3):262–265.
 91. Gajraj NM, Victory RA, Pace NA, Van Elstraete AC, Wallace DH. Comparison of an ephedrine infusion with crystalloid administration for prevention of hypotension during spinal anesthesia. *Anesth Analg* 1993; 76(5):1023–1026.
 92. Wang BW, Chiou YH, Chen WB, Peng TY, Leung HK. Intravenous pretreatment of hypertonic saline can prevent

- systemic hypotension induced by spinal anesthesia. *Acta Anaesthesiol Sin* 1997; 35(2):85–90.
93. French GW, White JB, Howell SJ, Popat M. Comparison of pentastarch and Hartmann's solution for volume preloading in spinal anaesthesia for elective caesarean section. *Br J Anaesth* 1999; 83(3):475–477.
94. Morgan PJ, Halpern SH, Tarshis J. The effects of an increase of central blood volume before spinal anesthesia for cesarean delivery: a qualitative systematic review. *Anesth Analg* 2001; 92(4):997–1005.
95. Rout CC, et al. Are evaluation of the role of crystalloid preload in the prevention of hypotension associated with spinal anesthesia for elective cesarean section. *Anesthesiology* 1993; 79(2):262–269.
96. Tsen LC, et al. Hemodynamic effects of simultaneous administration of intravenous ephedrine and spinal anesthesia for cesarean delivery. *J Clin Anesth* 2000; 12(5):378–382.
97. Morgan PJ, Halpern SH, Tarshis J. The effects of an increase of central blood volume before spinal anesthesia for cesarean delivery: a qualitative systematic review. *Anesth Analg* 2001; 92(4): 997–1005.
98. Wilson D, et al. Preoperative dextrose does not affect spinal induced hypotension in elective cesarean section. *Can J Anaesth* 1999; 46(11):1024–1029.
99. Vercauteren MP, et al. Hydroxyethylstarch compared with

- modified gelatin as a volume preload before spinal anesthesia for caesarean section. *Br J Anaesth* 1996; 76:731–733.
100. Miller RD, ed. *Anesthesia*. 4th ed. New York: Churchill Livingstone, 1994.
101. Scheingraber S, Rehm M, Sehmisch C, et al. Rapid saline infusion produces hyperchloremic acidosis in patients undergoing gynecologic surgery. *Anesthesiology* 1999; 90:1265–1270.
102. Wilkes NJ, Woolf R, Mutch M, et al. The effects of balanced versus saline-based hetastarch and crystalloid solutions on acid–base and electrolyte status and gastric mucosal perfusion in elderly surgical patients. *Anesth Analg* 2001; 93:811–816.
103. Gan TJ, Bennett-Guerrero, Phillips-Bute B, et al. Hextend, a physiologically balanced plasma expander for large volume use in major surgery: a randomized Phase III clinical trial. *Anesth Analg* 1999; 88:992–998.
104. Haljamae H, Dahlqvist M, Walentin F. Artificial colloids in clinical practice: pros and cons. *Bailliere's Clin Anaesthesiol* 1997; 11:49–79.
105. Lorenz W, Duda D, Dick W, et al., and the Trial Group /Mainz /Marburg. Incidence and clinical importance of perioperative histamine release: randomized study of volume loading and antihistamines after induction of anaesthesia. *Lancet* 1994; 343:933–940.
106. Waagstein L, Haljamae H, Ricksten SE, Sahlman L. Effects of

- hypertonic saline on myocardial function and metabolism in non ischemic and ischemic isolated working rat heart. *Crit Care Med* 1995; 23:1890–1897.
107. Haljamae H. Crystalloids versus colloids: the controversy. *NATA Textbook*. Paris: R & J Editions Medicale, 1999:27–36.
108. Laxenaire MC, Charpentier C, Feldman L. Anaphylactoid reactions to colloid plasma substitutes: incidence, risk factors, mechanisms. A French multicentre prospective study. *Ann Fr Anesth Reanim* 1994; 13:301–310.
109. Ring J, Messmer K. Incidence and severity of anaphylactoid reactions to colloid volume substitutes. *Lancet* 1977; 1:466–469.
110. Hedin H, Richter W. Pathomechanisms of dextran induced anaphylactoid / anaphylactic reactions in man. *Int Arch Allergy Appl Immun* 1982; 68:122–126.
111. Arfors K E, Buckley PB. Pharmacological characteristics of artificial colloids. *Bailliere's Clin Anaes thesiol* 1997; 11:15–47.