

REFERENCES

- Abeshouse, G.A.; Goldstein, R.B. and Abeshouse, B.S (1959):** Adrenal cysts: review of the literature and report of three cases. *J.Urol.*, 81:711-719.
- Achilles, E.; Padberg, C.B.; Holl, K.; Kloppel, G. and Schroder, S. (1991):** Immunocytochemistry of paragangliomas- Value of staining for S-100 protein and GFAP in diagnosis and prognosis. *Histopatholog .*, 18 : 453-458.
- Aguilo, G. ;Trump, D.L. ; Baylin, S.B. and Sallen, S.E. (1991) :** Watery diarrhea syndrome in an adult with ganglioneuroma-pheochromocytoma :identification of VIP, calcitonin,and catecholamines and assessment of their biologic activity. *Cancer*,40:1526-1532.
- Ando, S.; Suzuki,M.; Yamamoto,,N.; Lida,T. and Kimura, H. (2004) :** The prognostic value of both neuron-specific enolase (NSE) and Cyfra21-1 in small cell lung cancer. *Anticancer, Res.*,3b :1941-1946.
- Angus, B., Kiberu, S. and Purvis, J. (1988):** Cytokeratins in cervical dysplasia: a comparative study immunohistochemical staining using monoclonal antibodies NCL-5D3, CAM5.2, and PLL1.*J. Pathol.*, 155:71-75.
- Aoyama, C; Qualman, S.T.; Regan, M. and Shimada. (1990):** Histopathologic features of composite ganglioneuroblastoma. Immunohistochemical distinction of the stromal component is relater to prognosis. *Cancer*, 65: 255-64.

August, C. ; August, K. ;Schroeder, S. ; Bahen, H.A. and Hinz, R. (2004) :CGH and CD/MIB-1 immunohistochemistry are helpful to distinguish metastasized from nonmetastasized sporadic pheochromocytomas. *Mod. J. Pathol.*, 17: 25-256.

Barzon, L. ; Masi, G. ; Fincati, K. ; Pacenti, M. Fallo, F. and Palu,G. (2005) :Shift from conn's syndrome to cushing's syndrome in a recurrent adrenocortical carcinoma. *Eur. J. Endocrinol.*, 5 : 629-636.

Battifora, H. (1992): Assessment of antigen damage in immunohistochemistry. The vimentin internal control. *Am. J. clin. Pathol.*, 96: 669-671.

Benn, D.E.; Dwight, Richardson, A.L.; Delbrige, L.; Bambach, C.P., Stowasser, M.; Gordon; R.D.;;Marsh, D.J. and Robinson, B.G. (2000): Sporadic and familial pheochromocytomas are associated with loss of at least two discrete intervals on chromosome 1p. *Cancer res.*, 24: 7048-7051.

Bennett, B.D.; Mckenna, T.J.; Hough, A.J.; Dean, R. and page, D.L. (1980) : Adrenal myelolipoma associated with cushing's disease. *Am. J.*

Bertagna , C. and Orth ,D.N. (1981) : Clinical and laboratory findings and results of therapy in 58 patients with adrenocortical tumors admitted to a single medical center (1951-1987) . *Am. J.*, 71 : 855-875.

Biausque, E.; Li, H.Z. and Banks, P.M. (2003): Diagnostic applications of an immunoperoxidase method in hematopathology.*J. Histochem cytochem.*,27:1192 B: Shift from cann's syndrome to cushing's syndrome in a recurrent adrenocortical carcinoma.) *Eur.j.Endocrinol.*,153 :629-636.

- Bornstien, S.R.; Stratakis, C.A. and Milovic, N. (1999)** : Adrenocortical tumors : recent advances in basic concepts and clinical management. *Ann. Intern. Med.*, 9 :759-771.
- Bove K.E. and Mcadmas, A.J. (1981)** : Composite ganglioneuroblastoma. An assessment of the significance of histological maturation in neuroblastoma diagnosed beyond infancy. *Arch. Pathol. Lab. Med.*,105:325-330.
- Bretan, P.N. Jr. and Lorig, R. A.(1989)**: Adrenal imaging. Computed tomographic Scanning and magnetic resonance imaging. *Urol. Clin .North Am.*; 16:505-513.
- Brossard, J. ; Bernstein , M .L. and Lemieux , B. (1996)**: Neuroblastoma: an enigmatic disease. *Br. Med. Bull.*,4: 787-801.
- Bundy, R.; Green, A.A.; Furman, W.L.; and Stephens, C.A. (1982)**:Virtilizing tumours of the adrenal gland in childhood. Report of eight cases.*J.Pediatr. Surg.*,4:291-296.
- Burke , C.R.; Egan ,M. ; Raafat ,F. ; Crocker, J. and Williams ,D. (1989)** :Comparative study of the degree of differentiation of neuroblastoma and mean numbers of nucleolar organizer regions. *J. Clin. Pathol.*,41 :527-531.
- Caccamo, D.; Katsetos , C .D.; Herma, M . M .; Frankfurter, A.; Collins ,V.P. and Rubinstein, L. J. (1989)**: Immunohistochemistry of aspontaneous murine ovarian teratoma with neuroepithelial differentiation. Neuron – associated B – tubulin as a marker for primitive neuroepithelium. *Lab. Invest .*, 60: 390-398.

- Card, R.; Brodeur, G.M.; Azar, C. and Brother, M. (1992) :**
Neuroblastoma .Effect of genetic factors on the prognosis and
treatment., 70 :1685-1694.
- Carney, J.A. (1992):** A drenal gland. In Sternberg S.S., ed: Histology for
pathologists, New York, , Raven press.
- Cerovic, S.; Cizmic,M.; Milovic,N. ;Ajdinovic,B. and Brajuskovic,G.
(2002) :** Pheochromocytomas as adrenal gland incidentalomas Srp
.Arh.Celok. Lek.,2: 47-51.
- Cheema ,J. ;Archer, K.F. ; Mack, E. ;Carlson. I.H. and Ulu, J. (1981)
:**Virilizing adrenal ganglioneuroma. JAMA., 239 :2273-2274
- Cohen, K.; Gottesman ,L. and Brennan, M. (1986):** Adreno-cortical
carcinoma. Surgery, 100: 1170-1177
- Colombat . F. ;Eisenhofer, G. ;Goldstein ,D.S. ;Kopin, I.J. and Crout,
J.R. (2000) :**Pheochromocytoma :rediscovery as catecholamine –
metaabolizing tumor. Endocrinol. Pathol.,14 : 193-212
- Condom, E.; Villabona, C.M. and Gomez, J.M. (1985):** Adrenal
myelolipoma in a woman with Congenital 17 – hydroxylase
deficiency. Arch. Pathol. Lab. Med .,109: 1116-1117
- Conn , J.W. ;Knopf , R.F. and Nesbit , R.M. (1964) :**Clinical
characteristics of primary aldosteronism from an analysis of 145
cases . Am. J. Surg.,107 :159-172
- Cooper, D.; Schermer, A. and sun ,T.T. (1985):** Biology of the disease
classification of human epithelia and their neoplasms using
monoclonal antibodies to keratins: Strategies, applications
andlimitations.inverst.,52:243-248

- Correa. P. and Chen, V.W.(1995)** : Endocrine gland cancer. *Cancer*,75: 338-352.
- Cote, J. Cordon – Cardo, C.; Reuter, V.E., and Rosen, P.P. (1990)** : Immuno-pathology of adrenal and renal cortical tumors. Coordinated change in antigenexpression is associated with neoplastic conversion in the adrenal cortex. *Am. J. pathol.*,136: 1077-1084.
- Cuschier, A. ;Kloppel, G. and Farndon, J. (1995)** : In *Essential Surgical Practice* . 3 rd. ed. By Vol.3(British library cataloguing).
- David, S.; Goldstein , G. E.; John, A. and Karel , P. (2004)** : Diagnosis and localization of pheochromocytoma . *Am. J. Pathol.*,43 :907-910
- De Moerloose, B. (2005)** :The prognostic significance of P-glycoprotein in children with acute lymphoblastic leukemia and neuroblastoma. *Verh. K. Acad. Geneeskd Belg.*,1 :45-54.
- Dejong, A.; Van Vark, M. and Abus-Lutterc. (1987)**: Pleomorphic rhabdomyosarcoma in adults: immunohistochemistry as a tool for its diagnosis. *Hum. Pathol.*,18:298-303.
- Dekker, A. and Oehrle, J.S. (1971)**: Hyaline globules of the adrenal medulla of man. A produce of lipid peroxidation ? *Arch. Pathol.*, 91 : 353-364.
- Del Gaudio, A. and Solidaro, G. (1986)** : Myelolipoma of the adrenal gland: report of two cases and areview of the literature. *Surgery*, 99: 293-301
- Denk, H.; Krepler, R. and Artlieb, U .(1983)**: Proteins of intermediate filaments: An immunohistochemical and biochemical approach to the classification of soft tissue tumors. *Am. J. Pathol.*, 10 :193-208.

- Diman, V.; Huglo, D.; Nocaudie, M.; pattou ,F.;Proye ,C. and Wemeau, J.L. (1992):** [Chromogranin A assay and (13) I-MIBG scintigraphy for diagnosis and-follow-up of pheochromocytoma. *J. Nud.Med.*,42: 993-997.
- Domonski, H.A.(2005):** Fine needle aspiration of ganglioneuroma. *Diagn. Cytopathol.*, 39 : 363-366.
- Doss, K.H.; Dralle, H. and Bocker, W. (1998):** Immunohistochemical and electron microscopic analysis of adenomas of the thyroid gland. A comparative investigation of hot and cold nodules. *Virch .Arch. Pathl. Anat.*, 571 :374-385.
- Douglas , W. ; Franquemont , M.D. ; Stacey , E. ; Mills . E. S. and Lack , E. E. (1994) :**Immunohistochemical detection neuroblastomatous foci in composite adrenal pheochromocytoma-neuroblastoma .*Am. J. Clin. Pathol.*,102 :163-170
- Eder, U. Fishcer,-Colrie, R.; Kogner, P. and Leitner, B. (1998) :** Levels and molecular forms of chromogranins in human childhood neuroblastoma and ganglioneuromas.*Neuroscilett.*, 1 :17-20.
- Elias,J.M.; Margiotta,M. and Gabroc,D. (1989) :**Sensitivity and detection efficiency of the peroxidase anti peroxidase(PAP), avidin-biotin complex (ABC) and peroxidase-labeled avidin-biotin (LAB) methods. *Am. J. Pathol.*,92:62-71
- El-Mekresh ,M.M.; Abdel-Gawad ,M.; El-diasty,T.; El-Baz, M. and Ghoneim, M. A(1996):** Clinical, radiological and histological features of adrenal myelolipoma: review and experience with a further eight cases. *Official Journal of the Pan African Urological Surgeons' Association (PAUSA) .*, 2 :72-85 .

- El-Naggar , A.K. ; Evans , D.B. and Macky , B. (1991) :**Oncocytic adrenal cortical carcinoma . *Ultrasr.pathol.*,15 : 549-556
- Erlandson ,R.A. and Reuter, V.E (1991) :** Oncocytic adrenal cortical adenoma. *Ultrastruct .pathol.*, 15: 539-547
- Evans , H.L. ; D'Angio ,G.J. ; Sather , H.N. and Vassilopoulou-Sellin , R. (1996) :**Adrenal cortical neoplasms . A study of 56 cases . *Am. J. Clin. Pathol.*,1 :76-86
- Favia, G.; Lumachi, F.; Carraro, P. D'Amico, D.F. (1995):** Adrenocortical carcinoma. Our experience. *Minerva Endocrinol.*,1 :95-99
- Feng, C.; Li, H.Z.; Yan, W.G.; Luo, Y.F. and Gao, J.L. (2005) :** The expression and significance of chromogranin A and synaptophysin in adrenal gland tumors. *Zhonghua. Nei. Ke. Za. Zhi.*,8 :486-488
- Ferrari, L.; Seregni, E.; Baletta, E.; Aliberti, G.; pallotti, F.; DellaTorre, S.; Luksch, R.; Fogt, F.; Vargas, M.P.; Zhuang, Z. and Merino, M.J. (1998) :** Utilization of molecular genetics in the differentiation between adrenal cortical adenomas . *Am.J.Pathol.*, 5:245-252.
- Foster, D.G. (1966):** Adrenal cysts. *Arch. surg.*,92:131-143.
- Franquemont, D.W.; Mills, S.E and Lack, E.E.(1994):** Immunohistochemical detection of neuroblastomatous foci in composite adrenal pheochromocytom and neuroblastoma. *Am. J. clin. Pathol.*, 102: 163-170.

- Gabrilove, J.L. ; Sharma, D.C.; and Woftiz, H.H.(1986) :** Feminizing adrenocortical tumors in the male. A review of 52 cases including a case report *Medicine.*, 44: 37-79.
- Gaffey, M.J.; Traweek, S.T. and Mills, S.E.(1992):** Cytokeratin experssion in adrenal cortical neoplasia: an immunohistochemical and biochemical study with implications for the differential diagnosis of adrenocortical, hepatocellular, and renal cell carcinoma. *Hum .pathol.*,23:144-153.
- Gestblom ,C. ; Grynfeld, A. ; Ora ,I.; Prtoft, E.; Larsson, C. and Axelson,K. (1999) :** The basic helix-ioop-helix transcription factors and HAND, amarker gene for the developing human sympathetic nervous system, is expressed in both high and low stage neuroblastomas. *Lab. Invest.*, 1 :67-79.
- Ghandur, K. ;Gabbiani, G.; Kapanci, Y. and Barazzone, P. (1979):** Immunohistochemical identification of intermediate sized filament in human neoplastic cells: A diagnostic acid aid for the surgical pathology. *Am. J. Pathol.*, 104:206 -210.
- Gicquel, C.; Bertagna, X.; Gaston, V.; Coste, J.; Louvel, A.; Bertherat, J.; Chapuis, Y.; Duclos, J.M.; Ploun, P.F.; Luton, J.P. and Bouc, Y.L. (2001) :** Molecular markers and long-term recurrences in alarge cohort of patients with sporadic adrenocortical tumors. *Cancer Res.*,61 :6762-6767.
- Gille Bouloux ,P.M (1992):** Pheochromocytomas and related tumors. In: Grossman, A., ed. *Clinical endocrinology*. London: Blackwell scientific, 459-485.

- Gimm, O. ; Armanios, M.; Dziema, H.; Neumann, H.P. and Eng, C. (2000):** Somatic and occult germ-line mutations in SDHD , a mitochondrial complex II gene, in nonfamilial pheochromocytoma. *Cancer. Rec.*,24: 6822-6825.
- Giovanella, L. (2005) :** Serum chromogranin-A assay in differential diagnosis of incidentally discovered adrenal masses. *Anticancer Res.*, (3A):1547-1550.
- Gordon, H. and Sweets, H.H. (1936) :**A sampling method for the silver impregnation of reticulum . *Am. J. path .*,12:545-552.
- Grim, C.E., Weinberger; M.H. and Higgins, J.T(1977):.** Diagnosis of secondary forms of hypertension. *JAMA .*, 237:1331-1336.
- Guido Bellezza, P.; Giansanti,M.; Cavaliere, A. and Sidoni, A. (2004) :** Pigmented pheochromocytoma of the adrenal gland. *Arch. Pathol. Lab. Med.*,128 : 125-128.
- Haak, H.R.; Fleuren, G.J. Hermans , J.; Cobben, L. and Fleuren , G.L. (1995) :** Neuroendocrine differentiation of adrenocortical tumors . *Cancer*,75 :860-864.
- Hachitanda, Y.; Ishimoto, K.; Hata, J. and Shimada, H. (1994):** One hundred neuroblastomas detected through a mass screening system in japan. *Cancer*,74 :3223-3226.
- Handa, U.; Khullar, U. and Mhan, H.(2005):** Pigmented pheochromocytoma : report of acase with diagnosis by fine needle aspiration. *Acta. Cyto.*, 9: 421-423.
- Handler,J. (2003):** Case studies in hypertension. Adrenal carcinoma. *J.Clin. Hypertens(Greenwich)*, 5 :355-358.

- Harris , H.F. (1900) :** On the rapid conversion of haematoxylin into haematin in staining reaction . J.Appl. Microsc. Lab. Meth. , 3:77.
- Hasegawa, T.; Hirose,T. ; Ayala, A.G. ; Ito , S. ; Tomaru , U. ; Shimoda ,T. and Hirohashi , S. (2001):** Adult neuroblastoma of the retroperitoneum and abdomen : clinicopathological distinction from primitive neuroectodermal tumor .Am. J. Surg. Pathol., 7: 918-924.
- Hedeland, H., Ostberg, G. and Hokfelt, B. (1988) :** On the prevalence of adrenocortical adenomas in an autopsy material in relation to hypertension and diabetes. Acta. Med. Scand., 184:211-218.
- Henley, D.J.; Van Heerden, J.A.; Grant, C.S.; Carney, J.A. and Carpenter, P.C. (1983) :** Adrenal Cortical Carcinoma – a continuing Challenge. Surger.,94: 926-931.
- Hervas, Benito, I.; Ravis Sanchez,A.; Bello Arques, P. ; Canete ,A. ; Fernandez, J.M. ;Savra Quiles, A.; Gonzalez cabezas, P.; Ruiz Rodriguez ,J.C.,;Castell,V.; Perez Pastor, J.L. ; Monfort ,J.A. and Mateo Navarro, A.(2001) :** Value of 123I-MIBG scanning neuron specific enolase and serum ferritin in the diagnosis and follow-up of patients with neuroblastoma Rev.ESP. Med. Nud.,20 :369-376.
- Hinze, R.; Machens, A.; Schneider, U.; Dralle, H. and Rath, F.W. (2000):** Simultaneously occurring liver metastases of pheochromocytoma and medullary thyroid carcinoma-a diagnostic pitfall with clinical implications for patients with multiple endocrine neoplasia type 2a. Pathol. Res. Pract., 7 :477-481.

Hoang, M. P.; Ayala, A. G. and Albores-Saavedra, J. (2002): Oncocytic adrenocortical carcinoma : a morphologic , immunohisto-chemical and ultrastructural study of four cases . Mod. Pathol., 9:973-978.

Hogan, T.F.; Gilchrist, K,W.; Westring , D.W. and Citrin , D.L.(1980) : A clinical and pathological study of adrenocortical carcinoma: therapeutic implications. Cancer, 45: 2880-2883.

Horiuchi, A.; Muraji, T.; Tsugawa, C.; Satho, S.; Misu, H.; Mabuchi, O.; Hotchkiss, R.D. (1948) :A microchemical reaction resulting in the staining of polysaccharide structures in fixed tissue preparation . Arch. Biochem., 16 :131-141.

Hough, A.J. ; Hollified, J.W.; page , D.L. ; Hartmann ,W.H. (1979): Prognostic factors in adrenal cortical tumors. A mathematical analysis of clinical and morphological data. Am. J. clin. pathol . 72: 390-399.

Hsiao, C.C. ; Huang, C.C. ; Sheen, M. ;Tai , M.H., Chen , C.M. ; Huang, L.L. and Chuang, J.H.(2005): Differential expression of delta-Like gene and protein in neuroblastoma, ganglioneuroblastoma and ganglioneuroma. Mod.pathol.,18: 656-562.

Hsu, S.M.; Raine, L. and Fanger, H. (1981) : A comparative study of the peroxidase-antiperoxidase method and an avidin-biotin complex method for studying polypeptide hormones with radioimmunoassay antibodies. Am.J. Clin. Pathol.,75: 734-738.

Huynh, T.T.; Pacak,K.; Brouwers, F.M.; Worrell, R.A.; Cleary, S.and Eisenhofer, G. (2005): Different expression of catecholamine transports in pheochromocytomas from patients with Von Hippel Lindau syndrome and multiple endocrine neoplasia type 2 . Eur. J. Endocrinol., 4 :551-563.

Isobe, T. and Akuyama, M. (1981): Structural relation of two S-100 Protein in bovin brain : Subunit composition of S-100a protein . Eur. J. Biochem., 115:469-474.

Ito, Y.; Arai , K.; Ryushi ; yoshida, H.; Tomoda , C.; Uruno, T.;Kuma, K. and Miyauchi , A. (2005) : S100 expression is significantly linked to differentiation of thyroid carcinoma. Pathol. Res. Pract., 201:551-556.

Jahanson, N.; Vaalamo, M. and Grenman, R. (1985): Expression of collagenase-3 in squamous cell carcinomas of the head and neck. Am.J.pathol., 151:499-508.

Joshi , V.V. ; Silverman, J. F.; Altshula , G . and Cantor , A. B. (1993): Systematization of primary histopathologic and fine – needle aspiration cytologic features of neuroblastic tumors. A report from the pediatric Oncology group. Hum. pathol .; 24: 493-504.

Kalidindi, R.S. and Hattingh, L. (2005): Bilateral giant adrenal myeliolipomas. Abdom.Imaging. In press.

Kamari, Y.; Sharabi, Y.; Leiba, A.; peleg, E. and Apter, S. (2005) :Peripartum hypertension from pheochromocytoma : a rare and challenging entity. Am.J.Hypertens., 18 : 1306-1312.

Kamic, R.; Jeffrey, L.; Medeiros, L. and Lawrence, M. (1990) : Adrenal cortical neoplasms; Histologic diagnosis; prognostic indicators; Immunohistochemistry; DNA content analysis; Molecular genetics). Am .J.clin. .Pathol.,97:73-83.

Kanagawa,K. and Fujita,M. (2004) : Thoracic neuroblastoma : outcome of incomplete resection. Pediatr. Surg. Int. ; 9 :714-728.

- Kazantesva, R.L.; Ball ,R.Y.; ixon, A .K . and Apsimon, A.T.(2002) :**
Metastatic transitional cell carcinoma of the bladder causing Addison's disease. J. Urol.,37: 986- 988.
- Key, F.M.; Hashida, Askari, H.A.; Sieber, W.H. and Fetterman, G.H.(1967):** Virilizing tumors of the adrenal cortex.. Am. J. Dis. Child. 115:4455-4460.
- Kikuchi,K. and Saito,M. (1999) :**Ganglion-cell tumor of the filum terminae
- Kliwer, J.A. ; Nouguchi, T. and Takeno, S. (19):** Histological and histopathological methods : theory & practice, 3rd edition:pp 390-430.
- Koike, K.; Lihara , M.; Kanbe, M.; Omi ,Y.; Aiba, M. and Obara ,T. (2003) :** Adult-type ganglioneuroblastoma in the adrenal gland treated by a laparoscopic resection : report of a case .Surg. Today. 10:785-790.
- Kruggelw, T.; Shimosto, Y. ;Adachi, T.; Cantor , A.B. and Arai, K. (1985):** Immunohistochemical and ultrastructural analysis of medullary carcinoma of thyroid in relation to hormone production. Am. J. Pathol., 89:555-560.
- Kumar, R.M. and Kumar, L.A. (1993):** Estrogen receptor in Immunohistochemistry. J. Histoche.cytochem. 26:803-810.
- Lack, E.E. (1994) :** Pathology of adrenal and extra – adrenal paraganglia. Major problems in pathology, vol 29. Philadephia: W.B. Saunders.

Lack, E.E. (1997):Tumors of the adrenal gland and extra-adrenal paraganglia .In : Atlas of Tumor Pathology. Washington ,D.C. ,Armed forces Institute of pathology .

Lack, E.E. ; Page, D.L. and Weiss, L.M.(1990) : Recommendations for Reporting of tumors of the adrenal cortex and medulla. Hum. pathol . 30: 887-890.

Lack, E.E.; Kozakwich, H.P. and Oerlel, J.E. (1997) : Embryology, developmental anatomy, and selected aspects of non-neoplastic pathology In: Lack, E.E.; ed. Pathology of the adrenal glands.Lack, E.E. edition. New York: Churchill Livingstone. 1-74.

Lack, E.E.; Travis, W.D. and Oerlel, J.E. (1999) :Adrenal cortical meoplasms. In lack ,E.E .(ed.) pathology of the adrenal glands. Churchill Livingstone, New York, P.115-171.

Lawson, C.; Fisher, C. and Gatter, K. (1987): An immunohistochemical study of differentiation in malignant fibrous histoma. Histopathology, 303-701-74.

Lenders, J.W. ; Pacak, K. ; Walther, M.M. ;Linehan, W.M. ;Mannelli, M.; Friberg, P. and Goldstein, D.S. (2002) :Biochemical diagnosis of Pheochromocytoma :which test is best ? JAMA. 287 :1427-1434.

Lenders, J. W.; Eisenhofer, G.; Mannelli, M. and Pacak, K.(2005):Pheochromocytoma. Lancet.86: 665-675.

Lewinsky, B. S.; Grigor, K.M.; Symington, T. and Neville , A.M. (1974) : The clinical and pathologic features of non-hormonal adrenocortical tumors . Report of 20 new cases and review of the literature . Cancer . 33 : 778-790.

- Li, M. and Weing, B.M. (2000):** Adrenal oncocytic pheochromocytoma. Am. J. surg. Pathol.,11: 1552-1557.
- Libe, R. and Bertherat , J. (2005):** Molecular genetics of adrenocortical tumors, from familial to sporadic disease. Eur.J.Endocrinol.;153 477-487.
- Linnoila, R.I.; Keiser, H.R. and Steinberg, S.M. (1990):** Histopathology of benign versus malignant sympathoadrenal paragangliomas. Clinicopathologic study of 120 cases including unusual histologic features. Hum. pathol ., 21: 1168-1180.
- Liu, H.G.; Zhang, S.Z. and He, C.Y. (2003):** Study on pathological features and diagnosis, differential diagnosis of olfactory neuroblastoma . Zhonghua Bing. Li. Xue. Za. Zhi., 5:432-436.
- Lloyd, R.V.; Sisson, J.C.; Shapiro, B. and Verhofstad, A.A.(1986) :** Immunohistochemical localization of epinephrine, norepinephrine, catecholamine–synthesizing enzymes, and chromogranin in neuroendocrine cells and tumors. Am. J. pathol ., 125: 45-54.
- Logie, A.; Boudou, P.; Boccon-Gibod, L.; Baudin, E. Vassal, G. Schlumberg, M.; Le Bouc, Y. and Gicquel, C. (2000):** Establishment and characterization of a human adrenal cortical carcinoma xenograft Model. Endocrinology, 9: 3165-3171.
- Luton, J.P.; Cerdes, S. ; Billaud, L.; Thomas, G.; Guihaume, B.; Bertagna, X. ; Laudat , M. H.; Louvel , A. ; Chapuis, Y.; Blondeau, P. ; Bonnin, A. and Bricaire, H. (1990) :** Clinical features of adrenocortical carcinoma, prognostic factors and the effect of mitotane therapy, N.Engl.J.MEd., 322:1195-1202.

- Manger, W.M. and Gifford, R.W. (2002)** : Pheochromocytoma. J. Clin. Hypertens. (Greenwich) .,4: 62-72
- Matsushima, A.Y.; Stabolcs, M.J.; Moulton, T.; O'Toole, K.M. and Knowles, D.M. (1994)**: Immunohistochemical evaluation of neuroblastic tumors correlates with clinical prognosis [Abstract]. Mod. Pathol.,7: 54 A.
- Mazurkiewicz, A.A.; Toezko, Z.; Karas, K. and NagusZewski, T.; (2005)** : {Huge adrenal pseudocyst }. Wiad .Lek ., 58: 255-257.
- Medeiros, L.J. and Weiss ,L.M. (1992)** : New development in the pathologic diagnosis of adrenal cortical neoplasms. Am .J. clin .pathol.,97: 73-83.
- Medeiros, L.J.; Katsas, G.G. and Balogh, K. (1985)** : Brown fat and adrenal pheochromocytoma. Association of coincidence ? Hum. pathol., 16: 970-972.
- Mendlesohn , B.L.; Frater, W. and Mitchell , B.S. (1978)**: The use protedytic enzymes to improve immunoglobulin staining by PAP technique. J .Histochem.; 11: 345-351
- Miettinen, M. (1993)**: Keratin immunohistochemical: update of applications and pitfalls. Pathol., Annual; 28:Part1. Appleton and Lang, Norwak and connecticut.
- Miettinen, M.; Letho,V.P. and Virtanen, I. (1984)**: Antibodies to intermediate filament proteins used in the diagnosis and classification of human tumours. Ultrastruct. Pathol.,7:83-89.

- Miettinen, M.; Letho, V.P. and Virtanen, J. (1985) :** Immunofluorescence microscopic evaluation of the intermediate filament expression of the adrenal cortex and medulla and their tumors. *Am. J. Pathol.*, 118: 360-366.
- Miller, R.W. ; Young, J.L. Jr. and Novakovic, B. (1995) :** Childhood cancer. *Cancer*, 75: 395-405.
- Misugi, K. ; Aoki, I. Kikyo, S. ; Sasaki, Y.; Tsunoda, A. and Nakajima, T. (1985) :** Immunohistochemical study of neuroblastoma and related tumors with anti S-100 protein antibody. *Pediatr Pathol.*, 3: 217-226.
- Molenaar, W.M. ; Baker, D.L.; Pleasure, D.; Lee, V.M. and Trojanowski, J.Q. (1990) :** The neuroendocrine and neural profiles of neuroblastomas, ganglioneuroblastomas, and ganglioneuromas. *Am. J. Pathol.*, 136: 375-382.
- Moreno, A.M.; Castilla-Guerra, L.; Martinez-Torres, M.C.; Fernandez, E and Galera-Davidson, H. (1999) :** Expression of neuropeptides and other neuroendocrine markers in human pheochromocytomas. *Neuropeptides*, 2 :159-163.
- Motta-Ramirez, G.A.; Remer, E.M.; Herts, B.R.; Gill, I.S. and Hamrahian, A.H. (2005):** Comparison of CT findings in symptomatic and incidentally discovered pheochromocytomas. *AJR. Am. J. Roentgenol.*, 3: 684-688.
- Nadji, M. and Morales, A.R. (1983):** Immunoperoxidase techniques: A practical approach to tumor diagnosis. ch.. 1, PP1-10. Ascp Chicago.

Nakano, M. (1988) : Adrenal cortical carcinoma. A clinicopathological and immunohistochemical study of 91 autopsy cases. Acta. Pathol., 38: 163-80.

Nelms, K .; Diner, E.K.; Lack, E.E.; Patel, S.V.; Ghasemian, S.R. and Verghese, M. (2004): Retroperitoneal ganglioneuroma encasing the celiac and superior mesenteric arteries. Scientific world journal.18 : 974-977.

Neville, A.M. Symington,T. and O'Here, M.J. (1985) : In the human adrenal cortex. Patholgy and Biology. An integrated Approach. New York, Spinger-verlage.

O'hare, M.J.; Monaghan, P. and Neville, A.M. (1979) : The pathology of adrenocortical neoplasia: A correlated structural and functional approach to the diagnosis of malignant disease. Hum .pathol. ;10: 137-154.

O'Leary, T.J.; Liotta, L.A. and Gill, J.R. (1982): Pigmented adrenal nodules in Cushing 's syndrome . Arch. Pathol. Lab. Med.,106 :257-262.

Odagiri,E. (2005) : neuron specific enolase. Nippon, Rinsho.,8 :717-719.

Oppedal, B.R.; Brandtzage, P. and Kemshead, J.T. (1987) : Immuno-histochemical differentiation of neuroblastomas from other small round cell neoplasms of childhood using a panel of mono – and polyclonal antibodies. Histopathology, 11:363-374.

Pacak, K. (2005): Pheochromocytoma. Lancet, (9486): 665-675.

- Pace ,V. ; Perents , E. and Germann , P. G. (2002) :** Pheochromocytomas and ganglioneuromas in the aging rat : morphological and immunohistochemical characterization . *Toxicol. Pathol.*,4 :492-500
- Pagani, A.; Forni , M.; Tonini, G.P.; Papotti, M. and Bussolati, G. (1992) :** Expression of members of the chromogranin family in primary neuroblastomas. *Diagn. Mol. Pathol.*,1: 16-24.
- Page, D.L.; Delellis, R.A. and Hough ,AJ. Jr. (1986):** Tumors of the adrenal. *Atlas of tumor pathology, 2nd series, Fascicle 23.* Washington, D.C.: Armed forces Institute of pathology.
- Papavasiliou, C.; Gouliamos, A.; Deligiori, E.; Vlahos,L. and Cambouris, T. (1990):**Masses of myeloidipose tissue : Radiological and clinical considerations. *Int.J. Radiat. Oncol. Biol. Phys.*,19:985-990.
- Paterson, D.R.; Bill, A.H., Jr., and Kirland, I.S. (1969) :** Neuroblastoma trends in time.*J. Pediatr. Surg.*; 4:244-251.*Pathol.*; 9 :1119-1128
Pathol. ; 115 :458-468.
- Pavolov, I.F. (2004) :** Effects of antibodies against S-100 antigen in Ultralow doses (Proprotien-100) on acquisition of avoidance response in rats. *Bull. Exp. Biol. Med.*;138:556-558.
- Pedersen, W.A.; Becker, L.E. and Yeger , H. (1993) :** Expression and distribution of peripherin protein in human neuroblastoma cell lines. *Int. J .cancer*, 6:145-152.

Pelagi, D.M.; Heatfield, B.M.; Anthony, R.L. and Trump, B.E. (1984): Immunohistochemistry of cytoskeleton of human prostatic epithelium. Evidence of disturbed organization in neoplasia. Am.J. pathol., 126:384-390.

Perents , E. ; Nakagawa , Y. ; and Ross , G. W. (1987) : Expression of epithelial membrane antigen in perineurial cells and their derivatives :an immunohistochemiccal study with multiple markers. Acta. Neuropathol.;

Perez, L.; Surin, M.R.; Collins, B.; Kogan, D.; Toukova, I.L. and Shurin , G.V. (2005): Comparative analysis of CD1a , S-100, CD83, and CD11c human dendritic cells in normal, Premalignant, and malignant tissues. Histol. Histopathol., 20: 1165-1172.

Peuchmaur, M. (2004) : Peripheral neuroblastic tumors : anatomo pathological classification. Ann. Pathol. ; 6 :556-567.

Plouin, P.F.; Gimenez-Roqueplo, A.P.; La batide Alanore, A.; Salenave, S. and Duclos, J.M. (2000): Recent progress in the diagnosis, prognostic evaluation and treatment of pheochromocytomas. Rev. Med. Intern.,12: 1075-1085.

Racine, S.X. and Berthet, J.P. (2005) : Diagnosis and management of a pheochromocytoma discovered with laparoscopic CO(2) insufflation. Ann. Fr., 8:451-456.

Raikhlin ,N. T. ; Baronin , A. A. ; Smirnova , E. A. and Petrov , S. V. (2002) : Immunohistochemical identification of hormonally-inactive tumors of adrenal gland . Vopr. Onkol.,6 :684-690

Saad, E. and Sokair S. (1995): Hanbook-tumor markers. First ed., printed in khattab press. kaser El-Nile St., Cairo, Egypt.

Saeger, W. (2000) : Histopathological classification of adrenal tumors. Eur. J. Clin. Invest.; 3 : 58-62.

Saeger, W. (2003) : { Adrenocortical tumors }. Pathol.,4 :272-279.

Saeger, W. ; Fassnacht , M. ;Chita ,R. ;Prager , G. ; Nies , C. Lorens . K. ; Simon , D. and Reincke, M. (2003) : High diagnostic accuracy of adrenal core biopsy :results of the German and Austrian adrenal network multicenter trial in 220 consecutive patients. Hum. Pathol.,34: 180-186.

Salmenkivi, K.; Haglundc Arola, J. and Heikkila, P. (2001) : Increased experssion of Ineascin in pheochromocytomas correlates with malignancy. Am. J. surg .pathol25: 1419-1423.

Sasano, H.; Nose, M. and Sasano,N. (1989) : Lectin histochemistry in adrenocortical hyperplasia and neoplasms with emphasis on carcinoma. Arch. Pathol. Lab. Med .;113: 68-72.

Sasano, N and Sasano, H. (1991) : The adrenal cortex. In: Kovacs k, Asa SL, eds. Functional edocrine pathology. London: Blackwell Scientific., 546-584.

Sasano, H.; White, P.C. New, M.L. and Sasano, N. (1988) :Immunohistochemical localization of cytochrome. P-450c21 in human adrenal cortex and its relation to endocrine function. Hum. Path., 19:181-187.

Sasano, H.; Suzuki ,T.; Sasano ,T., Kameya, T.; Sasano, N. and Nagula, H.(1991) : Adrenal cortical oncocytoma. A true nonfunctioning adrenocortical tumors. Am .J .surg. pathol., 15: 949-956.

Schmid, K.W.; Schroder,S.; Kirchmair, R.; Tots, M.; Bocker, W. and Fischer-Colbrie, R. (1994) : Immunohistochemical demonstration of chromogranin A, chromogranin B, and secretgranin II in extra-adrenal paragangliomas. Mod. Pathol., 3 :347-353.

Schroder,S.; Komminoth, P.; Padberg, B. and Heitz, P.U. (1995) : Morphological typing evaluation of tumor dignity and prognosis and etiologic classification of adrenomedullary and adrenocortical neoplasia. Patholge., 5 :307-314.

Schteingart,D.E.;Doherty,G.M.;Gauger,P.G.;Giordano,I.J.;Hammer, G.D.;KorobKin,M. and Worden ,F.P.(2005):Management of patients with adrenal cancer : recommendations of an international consensues conference. Endocr.Retat.Cancer.;12: 667-680.

Scott, H.W., Jr.; Oates, J.A.; Nies, A.S.; Burko, H.; Page, D.L. and Rhamy, R.K (1976) : Pheochromocytoma: Present diagnosis and management. Ann. Surg.;183:587-598.

Shah,S. and Ravindranath,Y. (1998) :Neuroblastoma. Indian. J. Pediatr. 5 :691-705

Sheeler, L.R. (1989) : Cushing's syndrome, Urol. Clin North Am. 16: 477-482.

Shimada, H.; Chatten ,J. and Newton , W.A. Jr. (1984) :
Histopathologic prognostic factors in neublastic tumors: definition of
suptypes of ganglioneuroblastoma and age linted classification of
neuroblastomas. JNCI.;73: 405-416.

**Shimada, H.; Nakagawa, A.;peters, J.;Wang, H. Luken, J.N.;Siegel,
S.E. and Seeger, R.C. (2004) :** TrkA expression in peripheral
neuroblastic tumors : prognostic significance and biological
relevance. Cancer. 101(8): 1873-1881.

Shioda, Y . and Kumagai, M.A. (2005) : Tumor markers of childhood
cancer. Gan.. To Kagaku Ryoho.; 32: 699-704.

Silva, E.G.; Mackay ,B. and Samaan, N.A. (1982) : Adrenocortical
carcinomas: An ultrostructural study of 22 cases. Ultrastruct pathol.;
3: 1-7.

Song,S.Y.; Parks,E.R.; Kim,S.R. and Suh,Y.I. (2004) : Oncocytic
adrenocortical carcinomas : A pathological and
immunohistochemical study of four cases in comparison with
conventional adrenocortical carcinomas. Pathol. Int. ; 8 :603-610.

**Stridsberg, M . ; Oberg K, Li .Q.; Engstrom, U. and Lundqvist,G.
(1995):** Measurments of chromogranin A, Chromogranin B (Secreto-
granin I), chromogranin Cl (Secretogranin II) and pancreastatin in
plasma and urine from patients with carcinoid tumours and endocrine
pancreatic tumours. J. Endocrinol . ;144: 49-59.

- Takashima, .; Kuramistu,Y.; Yokoyama, Y.; Lizuka, N.; Okita, K.;Oka, M. and Nakamura, k. (2005):** Overexpression of alpha enalase in hepatitis C virus-related hepatocellular carcinoma : association with tumor progression as determined by proteomic analysis. *Proteomics.S .*, (6):1686-1692.
- Tang, C.K. and Gray ,G.F. (1975) :** Adrenocortical neuplasms. Prognosis and morphology. *Urology.*, 5: 691-695.
- Tatic, S.; Havelka, M.; Paunovic, I., Botic, V., Diklic, A.; Brasanac, D. and Jankovic, R. (2002):** Pheochromocytoma-pathohistologic and immunohistochemical aspects. *Srp. Arch. Celok. Lek.*, 2: 7-13.
- Thomas, P.; Battifora, H.; Manderino ,G.L. and Patrick, J.(1987) :** A monoclonal antibody against neurone – specific enolase. Immunohisto-chemical comparison with a polyclonal antiserum. *Am. J. clin. Pathol.* ,88: 146-152.
- Tischler , A.S. ; Mobtaker , H. and Mann , K. (2000) :** Anti-lymphocyte antibody leu-7 (HNK-1) recognizes a constituent of neuroendocrine granule matrix . *J. Histochem. Cytochem.* ;34 :1213-1216
- Tissier, F.; Cavard, C.; Groussin ,L.; Perlemoin ,K.; Fumey, G.; Hognere,A.M.; ReneCorail, F.; Jullian, E.; Gicquel, C. and Beragana, X.(2005) :** Mutations of beta-catenin in adenocortical tumors : Activation of the Wnt signaling pathway is a frequent event in both benign and malignant adrenocortical tumor .*cancer Res.*;(17):7622-7627.
- Tong, A.L.; Zeng, Z.P., Yang, D.; Li, H.Z. and Li, M. (2005) :** Clinical analysis of 25 patients with bilateral pheochromocytomas. *Zhonghua,Nei. Ke. Za. Zhi.* ,10 : 751-754.

- Tornoczky, T.; Kalman, E.; Kajtar, P.G.; Pearson,A.D.; Board, J. and Shimada, H. (2004):** Large cell neuroblastoma : a distinct phenotype of neuroblastoma with aggressive clinical behaviour. *Cancer*, 2:390-397.
- Torrise,A.; Carillio,G.; Libra,M. and Zafam,S. (2003) :** Solitary ganglioneuroma of the ileo-cocol valve. *Pathologica.* ; 4 :192-195.
- Triche, T.J. ; Ross, W.E. and Chan, H.S. (1987) :** Glycogen-containing neuroblastoma with clinical and histopathologic features of Ewing's sarcoma . *Cancer*,41 :1425-1432.
- Tsai, S.T.; Jin, Y.T.; Tsai, W.C.; Wang,S.T.; Lin,Y.C.; Chang,M.T. and Wu,L.W. (2005) :** S-100, a potential marker for early recurrence in early-stage oral cancer. *Oral, Oncol.*,4:349-357.
- Unger, P.D.; Cohen, J.M.; Thung, S.N.; Gordon, R. and Dikman , S.H. (1991):**Lipid degeneration in a pheochromocytoma histologically mimicking an adrenal cortical tumor .*Arch. Pathol. Lab. Med.* , 114:892-894.
- Van Heerden, J.A. ; Sheps, S.G. ; Hamberger, B. ; Sheedy, P.F.;Poston, J.G. and Re Mine, W.H (1982) :** Pheochromocytoma: Current status and changing trends. *Surgery.*, 91:367-375.
- Van Slooten, H . Schaberg, A . Smeenk, D . and Moolenaar, A .J.(1985) :** Morphologic characteristics of benign malignant adrenocortical tumors. *Cancer*, 55: 766-73.
- Viola, K.V. and Sosa, J.A. (2005):** Current advances in the diagnosis and treatment of pancreatic endocrine tumors. *Cuit.Opin.Oncol.*, 17: 24-27.

- Vogelbaum, M.A.; Masaryk, T.; Mazzone, P.; Mekhail, T.; Fazio, V.; Kanner, A. and Janigro, D. (2005)** : S-100 beta as a predictor of brain metastases : brain versus cerebrovascular damage. *Cancer*,4 :817-824.
- Wallace, C.A.; Hallman, J.R. and Sangueta, O.P. (2003)** :Primary cutaneous ganglioneuroma : a report of two cases and literature review. *Am. J. Dermatopathol.*,239 :239-242.
- Weiss, L.M. (1984)** : Comparative histologic study of 43 metastasizing and nonmetastasizing adrenocortical tumors. *Am. J. surg.pathol.*, 8:163-169.
- Weiss, L.M. ; Medeiros, L.J. and Vickery, A.L. (1989)** : Pathologic feature of prognostic significance in adrenal cortical carcinoma. *Am. J. surg. pathol .*, 13: 202-206.
- Wieneke, J.A.; Thompson, L.D. and Heffess, C.S. (2003)** : Adrenal cortical neoplasms in the pediatric population : A clinicopathologic and immunophenotypic analysis of 83 patients. *Am. J. Surg. Pathol.*,7 : 867-881.
- Wilson, B.S.; Lloyed, M.D.; Mila Blaives, M.D. and Barry, S. (1984)** :Detection of chromogranin in neuroendocrine cells with a monoclonal antibody. *Am. J. Pathol.*, 5:367-371.
- Wirnsberger, G.H.; Becker, H.; Ziervogel, K. and Höfler, H. (1992):** Diagnostic immunohistochemistry of neuroblastic tumors. *Am .J. sur. pathol.*, 16: 49-57.
- Wooten, M.D. and King, D.K. (1993):** Adrenal cortical carcinoma. Epidemiology and treatment with mitotane and areview of literature. *Cancer*, 72 : 3145-3155.

Young, J.L. ; Ries ,L.G. ; Silverberg, E.; Horm, J.W. and Miller , R.W. (1986) :Cancer incidence , survival ,and mortality for children younger than age 15 years. *Cancer*, 58 :598-602.

Young, J.L. and Miller , R.W. (1975) :Incidence of malignant tumors in U.S. children. *J. pediatr.*,86: 254-258.

Yu, C.H.; Chen, H.H.; Liu, C.M. ; Jeng, Y.M.; Wang, Y.P.; Liu, B.Y.; Sun,A. and Chiang, C.P. (2005): HMB-45 may be a more sensitive marker than S-100 or Melan-A for immunohistochemical diagnosis of primary oral and nasalmucosal melanomas. *J. Oral.Pathol. Med.*, 9: 540-545.

Zerbini, C.; Kozakewich, H.P.; Weinberg, D.S.; Mudt, D.J.; Edwards,. and Lack, E.E. (1992) : Adreno-cortical neoplasm in childhood and adolescence: analysis of prognostic factors including DNA content. *Endocr. Pathol.*,3:116-128.