

Sikora, R. A. and Hoffmann, H.S. (1993):

Biological control of plant parasitic nematodes with plant-health- promoting rhizobacteria. Proceedings of Beltsville symposium XVII, Agriculture Research Service, U.S. Department of Agriculture, Beltsville, Maryland, May 2-6 No. 166-172.

Small, E., Jurzysta, M. and Nozzolillo, C. (1990):

The Evolution of hemolytic saponin content in wild and cultivated alfalfa (*Medicago sativa* Fabaceae) Economic Botany 44: 226-235.

Spizizene, J. (1958):

Transformation of biochemically deficient strains of *Bacillus subtilus* by deoxyribonucleate.
Pathology and Microbiology 44: 1072-1078.

Stahl, E. (1969):

"Thin Layer Chromatography" Edited by Egon Stahl, Springer- Verlag Berlin. Heidelberg. New York.

Stirling, G. R. (1984):

Biological control of *Meloidogyne javanica* with *Bacillus penetrans*.
Phytopathology 74: 55-60.

Stirling, G. R (1991):

Biological control of plant parasitic nematodes. CAB International; Walling ford, OX an OX 108 DE., UK. 282 pp;
39.pp

- Sukul, N.C., Das, P.K. and De, G.C. (1974):
Nematicidal action of some edible crops.
Nematologica 20: 181-191.
- Talmadge, K. and Gilbert, W. (1982).
Cellular location affects protein stability in *Escherichia coli*.
Proc. Natl. Acad. Sci. U.S.A. 79: 1830-1833.
- Tchounwou, P. B., England, Jr. A. J. and Malek, E. A. (1991):
Toxicity evaluation of ammonium sulfate and urea to 3
developmental stages of fresh water snails.
Arch. Environ. Contam. Toxicol., 21: 359-364.
- Thompson, J. P., Domingo, E. and Wirmen, R.C. (1980):
Pseudolysogeny of *Azotobacter* phages.
Virology 102: 267-285.
- Tibelius, K. H., Robson, R.L. and Yates, M. G. (1987):
Cloning and characterization of hydrogenase from
Azotobacter chroococcum.
Mol. Gen. Genet. 206: 285-290.
- Tortolero, M., Santero, E., and Casadesus J. (1983):
Plasmid transfer mobilization of nif markers in *Azotobacter vinelandii*.
Microbios lett. 22: 31-35.
- Tsukamoto, C., Kikuchi, A., Kudou, S., Harada, K., Kitamura, K. and
Okubo, K. (1992):
Group A acetyl saponin- deficient mutant from the wild soy bean.
Phytochemistry 31: 4139-4142.

- Tsukamoto, C., Kikuchi, A. Harada, K., Kitamura, K. and Okubo, IX.
(1993):
Genetic and Chemical poly morphisms of saponins in soybean seed.
phytochemistry. 34: 1351-1356.
- Uchida, Y., Takahashi, T., Sato, N. and Japan, (1975):
The characteristics of the antibacterial activity of garlic.
Abtibiotic., 28 (4)., 638-642 (C. F. Chem. Abster., 1976 (84):
111547n).
- Uhlenbrock, J. H. and Bijole, J.D. (1958):
Investigations on nematocides, Isolation and structure of a
nematicidal principale occuring in Tagetes roots.
Helminth. 27: 179-
- Varshney, I.P., Beg. M.F.A., Sankaram, A.V. B. (1985):
Saponins and sapogenins from *Quillaria saponaria* .
Fito Terapia. 56: 254-256.
- Wackernagel, W. (1973):
Genetci transformation in *E. coli*.; the inhibitory role of the rec
BC. DN ase.
Biochem. Biophys. Res commun. 51: 306-311.
- World Health organization (WHO) (1992):
Special program for research and training in tropical diseases.
Page 29-36.

Wyss, O.N. and Nimeck, M. W. (1962):

Interspecific transduction in *Azotobacter*.

Fed . Proc. 21: 456.

Yang, D. C. and Choi, K.T. (1991):

Patterns of soluble protein, reducing sugar and ginsenosides in transformed calli of ginseng (*Panax ginsenge* C. A. Meyer[*P. pseudoginseng*]).

Korean Journal of Ginseng Science 15: 124-130.

Yeates, G.W. (1981):

Nematode population in relation to soil environmental factors: a review.

Pedobiologia 22: 213.

Zaki, M. J. and Maqbool, M.A. (1991):

Combined efficacy of *Pasteuria penetrans* and other biocontrol agents on the control of root-knot nematode on okra.

Pak. J. Nematol. 9: 49-52.