

References

- Abdel-Hameid, N.A.H. (2007):** Physiological and histopathological alterations induced by phenol exposure in *Oreochromis aureus* juveniles. Turk. J. Fish. Aquat. Sci., 7: 131-138.
- Abd-El-Reheem, A.M.A. (2003):** Effect of feeding fish oil and vegetable oils on the serum levels of some inorganic ions; cholesterol and some hormones in white rats (*Rattus rattus*). J. Egypt. Ger. Soc. Zool., 40A:55-70.
- Abd-El-Reheem, A.M.A. (2000):** Comparative effects of vegetable oils and fish oil on some physiological parameters in the female rat (*Rattus rattus*). Al-Azhar Bull. Sci., 11(2):197-208.
- Adamopoulos, P.N.; Papamichael, C.M.; Zampelas, A. and Moulopoulos, S.D. (1996):** Cholesterol and unsaturated fat diets influence lipid and glucose concentrations in rats. Biochem. Physiol., 113(B): 659-663.
- Akahane, K.; Furuhashi, K. and Onodera, T. (1986):** Simultaneous occurrence of hypercholesterolemia and hemolytic anemia in rats fed cholesterol diet. Life Sci., 39(6):497-505.

Alberts, B.; Bray, D.; Lewis, J.; Raff, M.; Roberts, K. and Watson, J. D. (1994): Molecular biology of the cell. 3rd Ed., Garland publishing Co., Philadelphia, USA. Pp. 482.

Angeles, Z.M.; Ana, B.; Henri, G.; Paul, H. and Martinez, A.J. (1999): Alterations in carbohydrate and lipid metabolism induced by a diet rich in coconut oil and cholesterol in a rat model. J. Am. Coll. Nut., 18(1): 36-42.

Anton, F.H.S.; Marc, J.T.M.M. and Panl, M.J.S. (1989): Efficacy and tolerability of simvastatin (MK-733). Am. J. Med., 87: 39- 43.

Aprahamian, T.; Bonegio, R.; Rizzo, J.; Perlman, H.; Lefer, D.J.; Rifkin, I.R. and Walsh, K. (2006): Simvastatin treatment ameliorates autoimmune disease associated with accelerated atherosclerosis in a murine lupus model. J. Immunol., 177: 3028–3034.

Bayorh, M.A.; Ganafa, A.A.; Eatman, D.; Walton, M. and Feuerstein, G.Z. (2005): Simvastatin and losartan enhance nitric oxide and reduce oxidative stress in salt-induced hypertension. Am. J. Hypertens., 18(11):1496-1502.

Bellosta, S.; Bernini, F. and Ferri N. (1998): Direct vascular effects of HMG-CoA reductase inhibitors. Atherosclerosis, 137: 101–109.

Benjamin, M.M. (1970): ‘Outline of veterinary pathology’. (2nd. Ed.) The Iowa state University press, Ames, Iowa, USA. pp. 85-86.

Bernini, G.P.; Argenio, G.F.; Gasperi, M.; Vivaldi, M.S.; Franchi, F. and Salvetti, A. (1994): Effects of long-term simvastatin treatment on testicular and adrenal steroidogenesis in hypercholesterolemic patients. J. Endocrinol. Invest., 17(4) :227-33.

Böhm, M.; Herrmann, W.; Wassmann, S.; Laufs, U. and Nickenig, G. (2004): Does statin therapy influence steroid hormone synthesis. Z. Kardiol., 93(1): 43-48.

Britton, K.E.; Quinn, V.; Brown, B.L. and Ekins, R.P. (1975): A strategy for thyroid function tests. Br. Med. J., iii:350-352.

Brown, W.E.L. and Hill, A.V. (1923): The oxygen-dissociation curve of blood and its thermodynamical basis. Pro. Roy. Soc. London, 94:297-334.

Buchwald, H.; Menchaca, J.H.; Michalek, N.V.; Rohde, D.T.; Donald, B.H. and O’Dea, J.T. (2000a): Plasma cholesterol: An influencing factor

in red blood cell oxygen release and cellular oxygen availability. *J. Am. Coll. Surg.*, 191(5): 490-497.

Buchwald, H.; O'Dea, J.T.; Menchaca, J.H.; Michalek, N.V. and Rohde, D.T. (2000b): Effect of plasma cholesterol on red blood cell oxygen transport. *Clin. Exp. Pharmacol. Physiol.*, 27:951-955.

Çalışkan, S.; Çalışkan, M.; Kuralay, F. and Önvural, B. (2000): Effect of simvastatin therapy on blood and tissue ATP levels and erythrocyte membrane lipid composition. *Res. Exp. Med.*, 199:189-194.

Cazana, d-L. F.J.; Puyol, R. M.; Perez-Caballero, J.; Jimenez, J.A. and Duarte, M.A. (1990): Effects of dietary hyperlipidemia-hypercholesterolemia on rat erythrocytes. *Int. J. Vitam. Nutr. Res.*, 60(4): 392-397.

Chen, J.R.; Chiou, S.F.; Suetsuna, K.; Yang, H.Y.; and Yang, S.C. (2003): Lipid metabolism in hypercholesterolemic rats affected by feeding cholesterol-free diets containing different amounts of non-dialyzed soybean protein fraction. *J. Nutr.*, 19: 676–680.

Cherniack, R.M. and Cherniack, L. (1983): Respiration in health and disease (3rd. Ed), W.B. Saunders Co., Philadelphia, pp. 65-143.

- Choi, J.W. and Pai, S.H. (2004):** Influences of hypercholesterolemia on red cell indices and erythrocyte sedimentation rate in elderly Persons. Clin. Chim. Acta, 341:117-121.
- Chowienczyk, P.J.; Watts, G.F.; Cockcroft, J.R. and Ritter, J.M. (1992):** Impaired endothelium-dependent vasodilatation of forearm resistance vessels in hypercholesterolemia. Lancet, 340:1430–2.
- Ciesla, B. (2007):** Hematology in practice, F. A. Davis Company, Philadelphia, PA 19103, pp.17.
- Cindy, L.C.; Peter, H.J.; Wadi, N.S.; Eleamor, D.L.; Miguel, A.Q.; Shelley, W.S. and James, B.Y. (1988):** Rhabdomyolysis and renal injury with lovastatin use report of two cases in cardiac transplant recipients. JAMA, 260(2): 239-241.
- Cooksey, J. and Reilly, P. (1977):** Effect of hypercholesterolemia on the hemoglobin-oxygen dissociation curve of the rabbit. J. Surg. Res., 22: 23-27.
- Cooper, A.R.; Leslie, H.M.; David, K. and Detweiler, K.D. (1980):** Red cell cholesterol enrichment and spur cell anemia in dogs fed a cholesterol-enriched atherogenic diet. J. Lipid Res., 21:1082-1089.

Czakó, L.; Szabolcs, A.; Vajda, Á.; Csáti, S.; Venglovecz, V.; Rakonczay J., Z.; Hegyi, P.; Tiszlavicz, L.; Csont, T.; Pósa, A.; Berkó, A.; Varga, C.; Ilona, S.V.; Boros, I. and Lonovics, J. (2007): Hyperlipidemia induced by a cholesterol-rich diet aggravates necrotizing pancreatitis in rats. *Euro. J. Pharmacol.*, 572: 74–81.

Dabbagh, A.J.; Mannion, T.; Lynch, S.M. and Frei, B. (1994): The effect of iron overload on rat plasma and liver oxidant status in vivo. *Biochem. J.*, 300: 799-803.

Dabbagh, A.J.; Shwaery, G.T.; Keaney, J.F.; Jr. and Frei, B. (1997): Effect of iron overload and iron deficiency on atherosclerosis in the hypercholesterolemic rabbit. *Arterioscler. Thromb. Vasc. Biol.*, 17: 2638-3645.

Daghini, E.; Chade, A.R.; Krier, J.D.; Versari, D.; Lerman, A. and Lerman, L.O. (2006): Acute inhibition of the endogenous xanthine oxidase improves renal hemodynamics in hypercholesterolemic pigs. *Am. J. Physiol. Regul. Integr. Comp. Physiol.*, 290: 609–615.

Dawson, W.R. and Poulson, T.L. (1962): Oxygen capacity of lizard blood. *Arch. Environ. Health*, 1:92-98.

de Pinieux, G.; Chariot, P.; Ammi-Saïd, M.; Louarn, F.; Lejonc, J. L.; Astier, A.; Jacotot, B. and Gherardi, R. (1996): Lipid-lowering drugs and mitochondrial function: effects of HMG-CoA reductase inhibitors on serum ubiquinone and blood lactate/pyruvate ratio. *Br. J. Clin. Pharmacol.*, 42:333-337.

Drabkin, D.L. (1964): Spectrophotometric studies: XIV. The crystallographic and optical properties of the hemoglobin of man in comparison with those of other species. *J. Biol. Chem.*, 164:703-723.

Droubay, P.E. and Puppione, D.L. (1980): Dietary fat-induced postprandial lipemia: effect on arterial oxygen saturation and plasma lactate, triglyceride, and cholesterol levels in subjects with angina pectoris. *Am. J. Clin. Nutr.*, 33:1199-1207.

Dumas, B.T. and Biggs, H.G. (1972): In standard methods of clinical chemistry (Vol. 7). Academic press, New York, pp. 175.

Dumas, M.; Muller, S.; Gouin, F.; Baros, F.; Viriot, M. and Stoltz, J. (1997): Membrane fluidity and oxygen diffusion in cholesterol-enriched erythrocyte membrane. *Arch. Biochem. Biophys.*, 341(1):34-39.

Edwards, M.J. and Martin, R. (1966): Mixing technique for the oxygen hemoglobin equilibrium and Bohr effect. *J. Appl. Physiol.*, 21(6): 1898-1902.

Eissa, S.M.; Ziada, G.A.; Marie, M.S.; El-Shafey, A.A.M. and Hasheesh, W.S. (1988a): Regulation of acid-base status of blood of two rodents in relation to the adaptation to desert habitat. *Proc. Zool. Soc. A.R.E.*, 4th. Sci. Conf., 16:313-324.

Eissa, S.M.; Ziada, G.A.; Mohamed, H.A.; El-Shafey, A.A.M. and Hasheesh, W.S. (1988b): Factors affecting oxygen loading of blood of two Egyptian rodents inhabiting different habits. A: Temperature and pH. *Proc. Zool. Soc. A.R.E.*, 4th. Sci. Conf., 16:301-311.

El-Fiky, M.A. (1999): Ameliorative effects of the royal jelly and ascorbic acid on the hypercholesterolemia-induced complications in male rats. *Egypt. J. Zool.*, 32:1-26.

El-Gendy, A.; Tohamy, A. and El-Zawahry, B. (2000): Cholesterol-methionine combination increases the risk of hyperlipidemia and serum glucose in male albino rats. *J. Union Arab Biol., Cairo*, 14(A): 299-310.

- El-Shafey, A. A. M. (1998):** Effect of ammonia on respiratory functions of blood of tilapia zilli. *Comp. Bioch. Physiol.*, 121(A):305-313.
- El-Shafey, A.A.M. (1994):** Effect of praziquantel on the blood gases and oxygen equilibrium curve of the rats. *J. Egypt. Ger. Soc. Zool.*, 13(A):141-163.
- El-Shafey, A.A.M. (1990):** The effect of reserpine on the respiratory function of blood of the Nile grass rat (*Arvicanthis niloticus*). *Proc. Zool. Soc. A.R.E.*, 21:345-360.
- El-Shafey, A.A.M. and Seleim, M.M.E. (2002):** Effect of carbosulfan on blood gases and acid-base status of blood of albino rat. *Bull. Fac. Sci. Zagazig Univ.*, 24(1):291-302.
- Endo, A.; Tsujita, Y.; Kuroda, M. and Tanzawa, K. (1977):** Inhibition of cholesterol synthesis in vitro and in vivo by ML-236B, competitive inhibitors of 3-hydroxy-3-methylglutaryl-coenzyme A reductase. *Eur. J. Biochem.*, 77(1):33-36.
- Evan, A.S. (1989):** Management of hypercholesterolemia: Approach to diet and drug therapy. *Am. J. Med.*, 87 (4A): 20-27.

Farver, T.B.; Haskins, S.C. and Patz, J.D. (1986): Cardiopulmonary effects of acopromazine and the subsequent administration of ketamine in the dog. *Am. J. Vet. Res.*, 47(3):631-635.

Feng, Y.; Zhu, Y.; Chen, X.; Sha, J.; Fan, L. and Chen, Q. (2005): Effects of Diet-Induced Hypercholesterolemia on Testosterone-Regulated Protein Expression in Mice Liver. *J. Nanosci. Nanotechnol.*, 5(8): 1273-1276.

Finley, P.R.; Schiffman, R.B.; Williams, R.J. and Lichti, D.A. (1978): Cholesterol in high-density lipoprotein: use of Mg^{2+} /dextran sulfate in its enzymic measurement. *Clin. Chem.*, 24: 931-933.

Folin, O. and Wu, H. (1919): A system of blood analysis. *J. Biol. Chem.*, 38: 81-110.

Forsythe, W.A.; Miller, E.R.; Hill, G.M.; Romsos, D.R. and Simpson, R.C. (1980): Effect of dietary protein and fat sources on plasma cholesterol parameters, LCAT activity and amino acid levels on tissue lipid content. *J. Nutr.*, 110(12):2467-2479.

Fossati, P.; Prencipe, L. and Berti, G. (1980): Use of 3,5-dichloro-2-hydroxybenzenesulfonic acid/4-aminophenazone chromogenic system in

direct enzymic assay of uric acid in serum and urine. Clin. Chem., 26: 227-231.

Fredrickson, D.S.; Levy, R.I. and Less, R.S. (1967): New Engl. J. Med., 176:34. Cited after Stanbio laboratory, Boerne, Texas, USA.

Friedewald, W.T.; Levy, R.I. and Fredrickson, D.S. (1972): Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. Clin. Chem., 18: 499-502.

García-Fuentes, E.; Gil-Villarino, A.; Flor-Zafra, M. and García-Peregrín, E. (2002): Dipyridamole prevents the coconut oil-induced hypercholesterolemia: A study on lipid plasma and lipoprotein composition. Int. J. Biochem. & Cell Biol., 34:269–278.

Gey, K.F. (1993): Prospects for the prevention of free radical disease, regarding cancer and atherosclerosis disease. Bri. Med. Bull., 49(3):679-699.

Goldstein, J. L. and Brown, M. S. (1987): Regulation of low-density lipoprotein receptors: implications for pathogenesis and therapy of hypercholesterolemia and atherosclerosis. Circulation, 76(3): 504-507.

- Gornal, A.G.; Bardawil, C.J. and David, M.M. (1949):** Determination of serum proteins by means of the Biuret reaction. *J. Biol. Chem.*, 177: 751-766.
- Gray, L.H. and Steadman, J.M. (1964):** Determination of the oxyhemoglobin dissociation curves for mouse and rat blood. *J. Physiol.*, 175:161-171.
- Gross, L.P.; Margaret, L.R.; Dale, V.B. and Marian A.P. (1991):** Platelet hypersensitivity in cholesterol-fed rabbits: enhancement of thromboxane A₂- dependent and thrombin-induced, thromboxane A₂- independent platelet responses. *Atherosclerosis*, 88:77-86.
- Grundy, S.M. (1987):** Dietary therapy for different forms of hyperlipoproteinemia. *Circulation*, 76(3):523-528.
- Hass, M.J.; Reinacher, D.; Li, J.P.; Wong, N.C. and Moorandian, A.D. (2001):** regulation of apo A1 gene expression with acidosis: requirement for transcriptional repressor. *J. Mol. Endocrinol.*, 27:43-57.
- Havel, R.J. (1986):** Functional activities of hepatic lipoprotein receptors. *Ann. Rev. Physiol.*, 48: 119-134.

Heibashy, M.I.A. (2000): Hypolipidemic effect of taurine and L-carnitine on rats fed a high cholesterol diet. *J. Union Arab Biol., Cairo*, 14(A): 11-23.

Heller, F.R.; Desager, J.P. and Harvengt, C. (1981): Plasma lipid concentrations and lecithine; cholesterol acyl transferase activity in normolipidemic subjects given fenofibrate and colestipol. *Metabolism*, 30(1):67-71.

HIŞMİOĞULLARI, A.A.; Bozdayi, A.M. and Rahman, K. (2007): Biliary lipid secretion. *Turk. J. Gastroenterol.*, 18 (2): 65-70.

Homma, Y. (2004): Predictors of atherosclerosis. *J. Atheroscler. Thromb.*, 11(5):265-270.

Hoyos, M.; Guerrero, J.M.; Perez-Cano, R.; Olivan, J.; Fabiani, F.; Garcia-Pergañeda, A. and Osuna, C. (2000): Serum cholesterol and lipid peroxidation are decreased by melatonin in diet-induced hypercholesterolemic rats. *J. Pineal Res.*, 28:150–155.

Imaeda, A.; Kaneko, T.; Aoki, T.; Kondo, Y.; Nakamura, N.; Nagase, H. and Yoshikawa, T. (2002): Antioxidative effects of fluvastatin and its metabolites against DNA damage in streptozotocin-treated mice. *Food Chem. Toxicol.*, 40:1415-1422.

Jayasooriya, A.P.; Masanobu, S.; Chizuko, Y.; Mikio, K.; Kyosuuke, Y. and Nobuhiro, F. (2000): Effects of *Momordica charantia* powder on serum glucose levels and various lipid parameters in rats fed with cholesterol-free and cholesterol-enriched diets. J. Ethnopharmacol., 27:331-336.

Jonathan, A. and Tobert, M.B. (1986): Therapeutic response to lovastatin (Mevinolin) in non familial hypercholesterolemia. JAMA, 256(20): 2829-2834.

Kamata, K.; Sugiura, M.; Kojima, S. and Kasuya, Y. (1996): Restoration of endothelium-dependent relaxation in both hypercholesterolemia and diabetes by chronic taurine. Eur. J. Pharm., 303: 47-53.

Keiichi, N.; Masaru, K.; Yoshito, S.; Hiroaki, Y.; Hiroto, N.; Jiro, F.; Itsuro, H. and Mitsuhiro, O. (1998): Myopathy induced by HMG-Co-A reductase inhibitors in rabbits: A pathological, electrophysiological, and biochemical study. Toxicol. Appl. Pharmacol., 152: 99-106.

Kerckhoffs, D.A.; Brouns, F.; Hornstra, G. and Mensink, R.P. (2002): Effects on the human serum lipoprotein profile of beta-glucan, soy protein and isoflavones, plant sterols and stanols, garlic and tocotrienols. J. Nutr., 132(9):2494-2505.

- Koter, M.; franiak, I.; Strychalska, K.; Broncel, M. and Chojnowska-Jezierska, J. (2004):** Damage to the structure of erythrocyte plasma membranes in patients with type-2 hypercholesterolemia. *Int. J. Biochem. & Cell Biol.*, 36:205–215.
- Koyama, T.; Araiso, T. (1988):** Diffusion pathway of oxygen in ox lung. *Adv. Exp. Med. Biol.*, 222:63-68.
- Kunitama, M.; Komatsu, N. and Miura, Y. (1996):** Effect of the inhibitors of 3-hydroxy-3-methylglutaryl coenzyme-A reductase, pravastatin and simvastatin, on colony formation by hematopoietic progenitor cells. *Rinsho Ketsueki.*, 37(11): 1314-1317.
- LaRosa, J.C.; Hunninghake, D.; Bush, D.; Criqui, M.H.; Getz, G.S.; Gotto, A.M.; Jr, G.S.M.; Rakita, L.; Robertson, R.M. and Weisfeldt, M.L. (1990):** The cholesterol facts. A summary of the evidence relating dietary fats, serum cholesterol, and coronary heart disease. A joint statement by the American Heart Association and the National Heart, Lung, and Blood Institute. The Task Force on Cholesterol Issues, American Heart Association. *Circulation*, 81: 1721-1733.
- Lasser, N.L.; Roheim, P.S.; Eldestein, D. and Eder, H.A. (1973):** Serum lipoproteins of normal and cholesterol-fed rats. *J. Lipid Res.*, 14: 1-8.

- Leaf, D.A.; Connor, W.E.; Illingworth, D.R.; Bacon, S.P.; Sexton, G. (1989):** The hypolipidemic effects of Gemfibrozil in type V hyperlipidemia. *JAMA*, 262(22): 3154-3160.
- Lee, H.S.; Ahn, H.C. and Ku, S.K. (2006):** Hypolipemic effect of water extracts of *Picrorrhiza rhizoma* in PX-407 induced hyperlipemic ICR mouse model with hepatoprotective effects: A prevention study. *J. Ethnopharmacol.*, 105(3): 380-386.
- Liao, J.K. (2002):** Beyond lipid lowering: the role of statins in vascular protection. *Int. J. Cardiol.*, 86:5-18.
- Luneva, O.G.; Brazhe, N.A.; Maksimova, N.V.; Rodnenkov, O.V.; Parshina, E.Y.; Bryzgalova, N.Y.; Maksimov, G.V.; Rubin, A.B.; Orlov, S.N. and Chazov, E.I. (2007):** Ion transport, membrane fluidity and haemoglobin conformation in erythrocyte from patients with cardiovascular diseases: Role of augmented plasma cholesterol. *Pathophysiology*, 14: 41–46.
- Mahfouz, M.M. and Kummerow, F.A. (2000):** Cholesterol-rich diets have different effects on lipid peroxidation, cholesterol oxides, and antioxidant enzymes in rats and rabbits. *J. Nutr. Biochem.*, 11:293–302.

Mahley, R.W. and Holcombe, K.S. (1977): Alterations of the plasma lipoproteins and apoproteins following cholesterol feeding in the rat. *J. Lipid Res.*, 18: 314-324.

Marcia, B.A. and Carlos, A.M.dL. (2003): Effects of chronic high fat diets on renal function and cortical structure in rats. *Exp. Toxic. Pathol.*, 55: 187–195.

Martin, M.J.; Hulley, S.B.; Browner, W.S.; Kuller, L.H. and Wentworth, D. (1986): Serum cholesterol, blood pressure, and mortality: implications from a cohort of 361,662 men. *Lancet*, 2:933–936.

Martinet, W.; Knaapen, M.W.; De Meyer, G.R.; Herman, A.G. and Kockx, M.M. (2001): Oxidative DNA damage and repair in experimental atherosclerosis are reversed by dietary lipid lowering. *Circulation Res.*, 88 (7): 733–739.

Matos, S.L.; de Paula, H.; Pedrosa, M.L.; dos Santos, R.C.; de Oliveira, E. L.; Júnior, D.A.C. and Silva, M.E. (2005): Dietary models for inducing hypercholesterolemia in rats. *Braz. Arch. Biol Technol.*, 48(2):203-209.

Menchaca, J.H.; Michalek, N.V.; Rohde, D.T.; Hirsch, A.T.; Tuna, N. and Buchwald, H. (2004): Improvement of blood oxygen diffusion capacity

and angina symptoms by cholesterol lowering with simvastatin. J. Appl. Res., 4(3):410-418.

Menchaca, J.H.; Michalek, N.V.; Rohde, D.T.; O'Dea, J.T. and Buchwald, H. (1998): Decreased blood oxygen diffusion in hypercholesterolemia. Surgery, 124: 692-698.

Milionis, H.J.; Kakafika, A.I.; Tsouli, S.G.; Athyros, V.G.; Bairaktari, E.T.; Seferiadis, K.I. and Elisaf, M.S. (2004): Effects of statin treatment on uric acid homeostasis in patients with primary hyperlipidemia. Am. Heart J., 148: 635-640.

Mol, M.A.E.; Desmet, R.C.; Terpestra, A.H.M. and West, C.E. (1982): Effect of dietary protein and cholesterol on cholesterol concentration and lipoprotein pattern in the serum chickens. J. Nutr., 112:1029-1037.

Murray, R.K.; Granner, D.K.; Mayes, P.A. and Rodwell, V.W. (2003): Harper's illustrated biochemistry (26th. Ed.), McGraw-Hill Companies, USA. Chap. 26 Pp. 219.

Nakaya, N.; Homma, Y.; Tamachi, H.; Shigematsu, H.; Hata, Y. and Goto, Y. (1987): The effect of CS-514 on serum lipids and apolipoproteins in hypercholesterolemic subjects. JAMA, 257(22): 3088-3093.

Nanami, H.; Oda, H. and Yokogoshi, H. (1996): Antihypercholesterolemic action of taurine on streptozotocin-diabetic rats or rats fed a high cholesterol diet. *Taurine*, 2: 561-568.

Nestle, P.J.; Reardon, M. and Billington, T. (1979): In vivo transfer of cholesteryl esters from high density lipoproteins to very low density lipoproteins in man. *Biochem. Biophys. Acta*, 57: 403-407.

Nicoll, D.; McPhee, S.J.; Pignone, M.; Chou, T.M. and Detmer, W.M. (2001): Pocket guide to diagnostic tests. Third edition, McGraw-Hill Companies, USA. pp. 115-116.

Nikolić, M.; Stanić, D.; Antonijević, N. and Niketić, V. (2004): Cholesterol bound to hemoglobin in normal human erythrocytes: a new form of cholesterol in circulation. *Clin. Biochem.*, 37:22– 26.

Olson, E.B.; JR and Dempsey, J.A. (1978): Rat as a model for human like ventilator adaptation to chronic hypoxia. *J. Appl. Physiol.*, 44(5):763-769.

Ozansoy, G.; Güven, C.; Ceylan, A.; Can, B.; Aktan, F.; Öz, E. and Gönü, B. (2005): Effects of simvastatin treatment on oxidant/antioxidant state

and ultrastructure of streptozotocin-diabetic rat lung. *Cell Biochem. Funct.*, 23: 421–426.

Paget, G.E. and Barnes, J.M. (1964): Evaluation of drug activities: pharmacometrics. 1st. Ed., Laurence and Bacharach publishing Co., Academic press, New York. Pp. 133-166.

Pahan, K. (2006): Lipid-lowering drugs. *Cell. Mol. Life Sci.*, **63(10)**: 1165–1178.

Pernadones, C.A.; Illera, V.A.; Peckham, D.; Stunz, L.L. and Ashman, R.F. (1993): Regulation of apoptosis in vitro in mature Murine spleen T cells. *J. Immunol.*, 151:3521-3529.

Pirat, A.; Zeyneloglu, P.; Aldemir, D.; Yücel, M.; Özen, Ö.; Candan, S. and Arslan, G. (2006): Pretreatment with simvastatin reduces lung injury related to intestinal ischemia-reperfusion in rats. *Anesth. Analg.*, 102:225–32.

Pittman, R.C. and Steinberg, D. (1984): Sites and mechanisms of uptake and degradation of high density and low density lipoproteins. *J. Lipid Res.*, 25: 1577-1585.

- Pond, W.G.; Yen, J.T.; Mersmann, H.J. and Haschek, W.M. (1986):** Comparative effects of dietary protein and cholesterol-fat content on genetically lean and obese pigs. *J. Nutr.*, 116: 1116-1124.
- Pulicini, T.; Elchebly, M.; Dusserre, E.; Berthezene, F. and Ponsin, G. (1995):** Cholesterol ester transfer and high-density lipoprotein conversion in normolipidemic, hypercholesterolemic, and hypertriglyceridemic non-insulin- dependent diabetics. *Biochem. Mol. Med.*, 55: 54-60.
- Rabini, R.A.; Polenta, M.; Staffolani, R.; Tocchini, M.; Signore, R.; Testa, I. and Mazzanti, L. (1993):** Effect of hydroxymethylglutaryl-CoA reductase inhibitors on the functional properties of erythrocyte membranes. *Exp. Mol. Pathol.*, 59:51-57.
- Reitman, S. and Frankel, S. (1957):** A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.*, 4:105-119.
- Richmond, W. (1973):** Preparation and properties of a cholesterol oxidase from *Nocardia sp.* and its application to the enzymatic assay of total cholesterol in serum. *Clin. Chem.*, 19: 1350-1356.

Ritzel, U.; Leonhardt, U.; Näther, M.; Schäfer, G.; Armstrong, V.W. and Ramadori, G. (2002): Simvastatin in primary biliary cirrhosis: effects on serum lipids and distinct disease markers. *J. Hepatol.*, 36(4):454-458.

Russell, D.W. (1992): Cholesterol biosynthesis and metabolism. *Cardiovasc. Drugs Ther.*, 6(2):103-110.

Salem, M.M. and Zaahkouk, S.A. (2000): Effect of high fat and protein diet on some physiological parameters related to lipid and protein metabolism in albino rats. *J. Egypt. Ger. Soc. Zool.*, 33A:279-290.

Scheen, A.J. (1998): How I study hypertriglyceridemia. *Rev. Med. Liège.*, 53: 103-105.

Shalaby, M.A.; El Zorba, H.Y. and Kamel, G.M. (2004): Effect of α -tocopherol and simvastatin on male fertility in hypercholesterolemic rats. *Pharmacol. Res.*, 50:137-142.

Shappell, S.D.; Murray, J.A.; Nasser, M.G.; Wills, R.E.; Torrance, J.D. and Lenfant, C.J.M. (1970): Acute change in hemoglobin affinity for oxygen during angina pectoris. *N. Engl. J. Med.*, 282: 1219-1224.

Shin, M.J.; Cho, E.Y.; Jang, Y.; Lee, J.H.; Shim, W.H.; Cho, S.Y.; Rim, S.J.; Kang, S.M.; Ha, J.W.; Ko, Y.G.; Kim, S.S.; Park, H.Y. and Chung, N. (2005): A beneficial effect of simvastatin on DNA damage in 242T allele of the NADPH oxidase p22phox in hypercholesterolemic patients. Clin. Chim. Acta, 360:46–51.

Siemieniuk, E. and Skrzydlewska, E. (2005): Coenzyme Q10: its biosynthesis and biological significance in animal organisms and in humans. Postepy. Hig. Med. Dosw., 59:150-159.

Sinet, M.; Muffat, J.M.; Henzel, D.; Renault, G. and Pocidal, J.J. (1984): Performance of hypothermic isolated rat heart at various levels of blood acid-base status. J. Appl. Physiol., 56(6):1526-1532.

Smith, P.F.; Grossman, S.J.; Gerson, R.J.; Gordon; L.R.; Deluca, J.G.; Majka, J.A.; Wang, R.W.; Germershausen, J.I. and MacDonald, J.S. (1991): Studies on the mechanism of simvastatin-induced thyroid hypertrophy and follicular cell adenoma in the rat. Toxicol. Pathol., 19(3):197-205.

Snedecor, G.W. (1971): Statistical methods, 14th. Ed. The Iowa state collage press, Ames, Iowa, USA.

- Steinbach, J.H.; Blackshear, P.L.J.; Varco, R.L. and Buchwald, H. (1974):** High blood cholesterol reduces in vitro blood oxygen delivery. *J. Surg. Res.*, 16:134-139.
- Stepuro, T.L. and Zinchuk, V.V. (2006):** Nitric oxide effect on the hemoglobin-oxygen affinity. *J. Physiol. Pharmacol.*, 57(1): 29-38.
- Straneva, J. E. and Briddell, R. A. (2000):** Cholesterol affects megakaryocytopoiesis. *Abst. Exp. Hematol.*, 28: 31–131.
- Tabacco, A.; Meiattini, F.; Moda, E. and Tarli, P. (1979):** Simplified enzymic/colorimetric serum urea nitrogen determination. *Clin. Chem.*, 25: 336-337.
- Tarnawski, R. And Tomasik, A. (1994):** Changes of the activity of urea cycle enzymes extracted from rat liver after surgical reduction of stomach capacity and after feeding a high-fat diet. *Arch. Tierenakr.*, 46(2): 165-171.
- Thompson, G.R.; Judith, F.; Jenkinson, M. and Iris, T. (1986):** Efficacy of mevinolin as adjuvant therapy for refractory familial hypercholesterolemia. *Q. J. Med.*, 60(232):803-811.

- Tikkanen, M.J. and Nikkila, E.A. (1987):** Current pharmacological treatment of elevated serum cholesterol. *Circulation*, 76 (3): 529-533.
- Tiwari, A. (2006):** An overview of statin-associated proteinuria. *Drug Discovery Today*, 11(9): 458-464.
- Tobert, M.B. (1987):** New developments in lipid-lowering therapy: the role of inhibitors of hydroxylmethylglutaryl Coenzyme A Reductase. *Circulation*, 76(3): 534-538.
- Tovar, A.M.F.; Pecly, I.M.D.; Rangel, E.P.; Melo-Filho, N.M.; Mourão, P.A.S. and Jr, M.L. (2007):** Experimentally induced metabolic acidosis in rabbits modulates the interaction of aortic glycosaminoglycan with plasma low-density lipoprotein—An interesting observation about the association of acidosis and atherosclerosis. *Atherosclerosis*, 192:33-39.
- Trinder, P. (1969):** Determination of blood glucose using 4-aminophenazone. *J. Clin. Path.*, 22(2):158-161.
- Turbino-Ribeiro, S.M.L.; Silva, M.E.; Chianca, D.A.; Jr.; de Paula, H.; Cardoso, L.M. and Colombari, E. (2003):** Iron overload in hypercholesterolemic rats affects iron homeostasis and serum lipids but not blood pressure. *J. Nutr.*, 133: 15-20.

- Valenti, S.; Cuttica, C.M.; Fazzuoli, L.; Giordano, G. and Giusti, M. (2001):** Biphasic effect of nitric oxide on testosterone and cyclic GMP production by purified rat leydig cells cultured in vitro. *Int. J. Androl.*, 22(5): 336-341.
- Vaziri, N.D. and Liang, K. (2004):** Effects of HMG-CoA reductase inhibition on hepatic expression of keycholesterol-regulatory enzymes and receptors in nephrotic syndrome. *Am. J. Nephrol.*, 24: 606–613.
- Vega, G.L. and Grundy, S.M. (1987):** Treatment of primary moderate hypercholesterolemia with lovastatin (mevinolin) and colestipol. *JAMA*, 257(1): 33-38.
- Velho, J.A.; Okanobo, H.; Degasperi, G.R.; Matsumoto, M.Y.; Alberici, L.C.; Cosso, R.G.; Oliveira, H.C.F. and Vercesi, A.E. (2006):** Statins induce calcium-dependent mitochondrial permeability transition. *Toxicol.*, 219: 124–132.
- Wilson, M.D. and Rudel, L.L. (1994):** Review of cholesterol absorption with emphasis on dietary and biliary cholesterol. *J. Lipid Res.*, 35:943-955.

Yokogoshi, H.; Mochizuki, H.; Nanami, K.; Hida, Y.; Miyachi, F. and Oda, H. (1999): Dietary Taurine Enhances Cholesterol Degradation and Reduces Serum and Liver Cholesterol Concentrations in Rats Fed a High-Cholesterol Diet. *J. Nutr.* 129: 1705-1712.