

## SUMMARY

The present work carried out to investigate the ability of *S. platensis* to produce antimicrobial substance, by the following objectives.

- I. Determination of the best solvent for bioactive metabolites extraction
- II. Studying some growth conditions (temperature, pH, light intensity, light duration and aeration) affecting *S. platensis* growth and its antimicrobial activities.
- III. Production, extraction, purification and identification of the antimicrobial substance produced by *S. platensis*.

**The following results were obtained:**

- 1- The biomass production using Aiba & Ogawa medium was gradually increased from the 1<sup>st</sup> day to the 9<sup>th</sup> day with maximum biomass production of 111.70 mg drt wt. /100 ml occurred.
- 2- Standard sources of nitrogen (2.5 g/l NaNO<sub>3</sub>) and phosphate (0.5 g/l K<sub>2</sub>HPO<sub>4</sub>) exhibited the maximum biomass production.
- 3- Ethanol was the best water miscible solvent to extract the antimicrobially active compound from the whole culture (cells with exometabolites) as well as ethyl acetate and diethyl ether (water immiscible).
- 4- The maximum biomass production (132.27 mg dry wt./100 ml) and antimicrobial activities of *S. platenis* whole culture against *Staphylococcus aureus* recorded the maximum inhibition zone of 35.33 mm while culture filtrate recorded

inhibition zone of 35.67 mm against *Candida albicans*. The antibiotic extract from cells grown at 35 °C exhibited the largest inhibition zone against *Pseudomonas aeruginosa* (30.0 mm).

- 5- pH 9.0 showed the maximum biomass production of 92.13 mg dry wt./100 ml and the maximum inhibition zone produced from *S. platensis* whole culture (was 48. mm against *Candida albicans* and 39.67 mm ,29.67 mm against *Candida albicans* by culture filtrate and cells extract respectively.
- 6- Biomass production of *S. platensis* exhibited the maximum value at light intensity of 2.0 and 2.25 klux without significant difference.
- 7- Antimicrobial activity by whole culture, culture filtrate and cells extraction showed the maximum inhibition zone against *Candida albicans*, 34.67, 40.0 and 29.67 mm, respectively.
- 8- The aerated medium showed the maximum mass production ( $94.73 \pm 0.1154$  mg dry wt./ 100 ml). the antimicrobial activities produced from aerated *S. platensis* medium recorded the highest inhibition zones resulted from whole culture, filtrate and extracted cells.
- 9- Antibiotic produced by *S. platensis* was extracted from the culture medium after 9 days of incubation at 30 °C, pH 9.0 and 2.5 Klux using diethyl ether (water immiscible solvent). It was obvious from the bio-autography that, antibiotic is one compound.
- 10- The physical properties of the antibiotic shown that it was greenish yellow in colour without characteristic odour. Its

melting point ~ 40 °C, soluble in chloroform, methanol, diethyl ether while sparingly soluble in acetone.