

CONTENTS

Title	Page
List of Tables	vi
List of Figures	ix
INTRODUCTION	1
REVIEW OF LITERATURE	4
1. Root rot disease of french bean and soybean.	4
2. Occurrence of <i>Fusarium solani</i> in french bean and soybean	6
3. Biology and ecology of <i>F. solani</i> f.sp. <i>phaseoli</i>	7
4. Resistance against fungal pathogens (especially <i>F. solani</i> f.sp. <i>phaseoli</i> and <i>C. lindemuthianum</i>)	9
4.1. Resistance in french bean (<i>Phaseolus vulgaris</i> L.) and soybean (<i>Glycine max</i> L.).	10
4.1.1 Role of phytoalexins in resistance	14
4.1.1.1. Phytoalexins and resistance in french bean	14
4.1.1.2. Phytoalexin production and disease resistance in soybean	16
4.1.1.3. Mechanism(s) of phytoalexin production	18
4.1.1.4. Accumulation of phytoalexins production	19
4.1.1.5. Toxicity of phytoalexins	22
4.1.1.6. Metabolism of phytoalexins	23
5. Induction of resistance in french bean and soybean (prime mechanisms for induction of resistance)	25
5.1. Role of biotic elicitors	25
5.1.1. Prior inoculation with viable organisms	25
5.1.2. Prior treatment with microbial extracts	26
5.2. Role of abiotic elicitors (temperature)	26

5.2.1. Induction of resistance by heat-transfer	26
5.2.2. Effect of temperature on suppression of resistance or induction of susceptibility.	28
5.3. Mechanisms of induced resistance	29
5.3.1. Role of hypersensitive response (HR)	29
5.3.2. Relative roles of glyceollin, lignin and the hypersensitive response (HR).	31
5.3.3. Role of pathogenesis-related (PR) proteins.	33
5.3.3.1. Systemic-induced resistance (SIR) associated with PR proteins.	36
5.3.3.2. Role of peroxidases	38
6. Role of RAPD-PCR in estimating the resistance of different plant cultivars.	40
MATERIALS AND METHODS	42
Experimental procedures	42
I. Plant material	42
I.1. Source of seeds	42
I.2. Growth of plants.	42
II. Fungal material	43
II.1. Source of fungal cultures used in this study	43
II.2. Media	44
(1) Potato Dextrose Agar Medium (PDA)	44
(2) <i>Colletotrichum</i> sp. Growth Medium	44
(3) Czapek Dox Medium	44
II.3. Cultivation technique	45
II.4. Maintenance of cultures	45
II.5. Sub-culturing	45

III. Method of inoculation	45
III.1. Inoculation preparation	45
III.2. Inoculation techniques	46
III.2.1. Growing plant inoculation technique	46
III.2.2. Hypocotyl inoculation technique	46
III.2.3. Bean pods inoculation technique	47
III.2.4. Challenging technique	47
IV. Anatomical examination using light microscopy	48
IV.1. Preparation of sections	49
IV.2. Histochemical staining of sections	49
V. Phytoalexins estimation techniques	49
V.1. Extraction method for phytoalexin phaseollin from french bean	49
V.2. Extraction method for phytoalexin glyceollin from soybean	50
V.3. Chromatographic techniques	50
V.3.1. TLC (Thin Layer Chromatography) technique.	50
V.3.2. HPLC (High Performance Liquid Chromatography) technique	51
VI. Protein detection (SDS-PAGE of proteins electrophoresis)	52
VI.1. Extraction of protein	52
VI.2. Quantification of extracted proteins	52
VI.3. Reagents of the SDS-PAGE	54
VI.4. Sample preparation	55
VI.5. Gel casting	55
VI.6. Gel electrophoresis	56
VI.7. Molecular weight estimation of protein subunits	57
VII. SDS-PAGE for peroxidase iso-enzymes	57

VIII. Random amplified polymorphic DNA (RAPD) analysis	58
VIII.1. Genomic DNA isolation, purification and quantification	58
Reagents	58
VIII.2. Genomic DNA extraction for RAPD analysis	61
VIII.3. Quantification and qualification of DNA	62
VIII.4. RAPD-PCR amplification conditions	62
VIII.5. Agarose gel electrophoresis	64
EXPERIMENTAL RESULTS	66
I. Pathogenicity test	67
II. Induced resistance against the target pathogens <i>F. solani</i> f.sp. <i>phaseoli</i> , <i>F. solani</i> f.sp. <i>glycine</i> and <i>C. lindemuthianum</i>	84
II.1. Infection of french bean and soybean intact hypocotyls by <i>F. solani</i> f.sp. <i>phaseoli</i> (the causal pathogen of root and hypocotyls rot of french bean and soybean).	87
II.2. Infection of french bean (<i>Phaseolus vulgaris</i> L.) and soybean (<i>Glycine max</i> L.) intact hypocotyls by <i>F. solani</i> f.sp. <i>glycine</i> and <i>C. lindemuthianum</i> (the causal pathogen of bean anthracnose), separately.	128
III. Phytoalexins production	132
IV. Total soluble protein analysis (SDS-PAGE of proteins electrophoresis)	139
IV.1. French bean (<i>Phaseolus vulgaris</i> L.) cultivars	139
IV.2. Soybean (<i>Glycine max</i> L.) cultivars.	140
IV.3. The original and reisolated isolates of the target pathogen.	141

V. Peroxidase induction in french bean (<i>Phaseolus vulgaris</i> L.) and soybean (<i>Glycine max</i> L.) in response to microbial challenge.	156
V.1. French bean (<i>Phaseolus vulgaris</i> L.) cultivars	156
V.2. Soybean (<i>Glycine max</i> L.) cultivars.	158
VI. RAPD and genetic stability	168
VI.1. French bean (<i>Phaseolus vulgaris</i> L.) cultivars	168
VI.2. Soybean (<i>Glycine max</i> L.) cultivars	176
VI.3. <i>Fusarium solani</i> f.sp. <i>phaseoli</i>	184
DISCUSSION	193
SUMMARY AND CONCLUSION	210
REFERENCES	214
ARABIC SUMMARY	1