

Contents

Title	Page
Introduction	1
Literature Review	4
1.Causal pathogen characteristics.	4
2.Detection and identification of <i>R. solanacearum</i> .	8
3.The effect of host-root-derived chemical signals between plants and microbes	15
4.Collection of plant root exudates.	22
5.Root exudates composition.	23
6.Difference between root exudates of potato varieties.	26
7.Chemotactic movement of <i>R.solanacearum</i> towards host root exudates.	27
8.Effect of different soil amendments (abiotic inducer) on pathogenicity of <i>R.solanacearum</i>	28
8.1.Effect of different concentrations of Salicylic acid on the pathogenicity of <i>R. solanacearum</i> .	28
8.2.Compounds associated with disease resistance in β -aminobutyric acid (β ABA)-treated plants.	30
8.3.Effect of soil amended with Calcium oxide and Urea on the population of <i>R. solanacearum</i> .	33
9.Effect of non host plants on the population of <i>R. solanacearum</i> in soil.	35
Material And Methods	
1.Materials.	37
2.Isolation And Identification of bacteria (<i>R. solanacearum</i>).	37
3.Isolation process	38
4.Identification method	39
5.Sensitivity of the different potato cultivars to the infection severity by <i>R. solanacearum</i> bacterium.	44
5.1.Root Exudates collection	45
5.2.Analysis of root exudates composition by GC mass apparatus.	47
6.Studying the chemotactic movement of bacteria towards potato root exudates	47
7.Effect of soil amendments (Abiotic factors) on the population of <i>R. Solanacearum</i> .	48
8.Effect of non host plants root exudates (biotic elicitor) on causative agent of potato brown rot.	54
9.Studying the antimicrobial properties of both root exudates of (Sweet basil & Chinese chive) on <i>R. solanacearum</i> bacterium on basal medium.	56
Results	
1.Isolation and identification of <i>R. solanacearum</i> isolates collected from different habitats.	58
2.Identification of <i>R. solanacearum</i> isolates collected from different localities using classical	62

methodology (biochemical test).	
3.Sensitivity of different potato cultivars to the infection by <i>R. solanacearum</i>	67
4.Count of <i>R. solanacearum</i> bacteria ($\times 10^3$)/ 100 μ l attracted by different potato cultivar root exudates on semi selective media of South Africa (SMSA)	68
5.Effect of different concentrations of salicylic acid on the pathogenicity of <i>R. solanacearum</i> for tomato plants after 10 days from soil infestation and count of bacteria on SMSA media for two bacterial load.	70
6.Effect of different concentrations of salicylic acid on the pathogenicity of <i>R. solanacearum</i> for potato plants After 14 days from soil infestation	72
7.Count of <i>R. solanacearum</i> in potato cultivars roots after salicylic acid concentrations treatment on SMSA media.	73
8.Effect of different concentrations of β -amino butyric acid on <i>R. solanacearum</i> pathogenicity in tomato plants after 5 and 10 days from soil infestation and count of bacteria in tomato roots on SMSA media.	74
9.Effect of different concentrations of β -ABA on <i>R. solanacearum</i> pathogenicity in potato plants after 7 and 14 days from soil infestation.	76
10.Effect of soil amended with(calcium oxide with urea) on <i>R. solanacearum</i> pathogenicity on tomato and potato plants after different periods from soil infestation.	77
11.Count of <i>R. solanacearum</i> bacterium in soil amended with (calcium oxide with urea) cultivated with potato plants after time course on SMSA media	79
12.Effect of both Chinese chive and sweet basil plants on population of <i>R. solanacearum</i> by counting bacteria in infested soil with time course.	80
13.Effect of Chinese chive and sweet basil root exudates on cultured <i>R. solanacearum</i> on basal media.	80
15.Chemical analysis of root exudates of different potato cultivars in different types of soil by GC mass detector.	82
16.Chemical analysis of root exudates of Spunta cultivar after different soil amendments by GC mass.	84
Discussion	91
Summary	103
References	106
Appendix	128
Arabic summary	

