

References

- Aley, P.; French, E. and Nydegger, U. (1994).** Methods of greenhouse inoculation with *Pseudomonas solanacearum* to select resistant potato clones. Fitopatologia 29:20 (Abstract).
- Anonymous, (1994).** Bacterial wilt: the disease and its causative agent *Pseudomonas solanacearum* (eds. Hayward, A. and Hartman, G.). 259 p. CAB International, Wallingford (UK).
- Bacilio-Jimenez (2003).** Chemical characterization of root exudates from rice (*Oryza sativa*) and their effects on the chemotactic response of endophytic bacteria. Plant Soil 249, 271–277.
- Bais, H.(2001).** Root-specific metabolism: the biology and biochemistry of underground organs. In Vitro Cell. Dev. Biol. Plant 37: 730–741.
- Bais, H.; Walker, T.; Schweizer, H. and Vivanco, J. (2002).** Root specific elicitation and antimicrobial activity of rosmarinic acid in hairy root cultures of sweet basil (*Ocimum basilicum L.*). Plant Physiol Biochem 40: 983-995.
- Bais, H.; Park, S. and Weir, T.; (2004).** How plants communicate using the underground information superhighway. Trends in Plant Science, 9 : 26-32.
- Bellstedt, D. and Van der, M. (1989).** The development of ELISA kits for the detection of *Ps. solanacearum* bacterial wilt in potatoes. In: Potato Research Symposium (Aart appelnavorsing simposium), 64-69, South Africa.
- Buddenhagen, I.; Sequeira, L. and Kelman, A. (1962).** Designation of races of *Pseudomonas solanacearum*. Phytopathology 52:726.
- Buddenhagen, I. and Kelman, A. (1964).** Biological and physiological aspects of bacterial wilt caused by *Pseudomonas solanacearum* . Annu. Rev. Phytopathol. 2: 203-230.

- Buee, M. (2000).** The pre-symbiotic growth of arbuscular mycorrhizal fungi is induced by a branching factor partially purified from plant root exudates. *Mol. Plant–Microbe Interact.* 6: 693–698.
- Chamsai, J.; Buchenauer, H.; Orober, M. and Siegrist, J. (1998).** DL-3 aminobutyric acid induces localized cell death, salicylic acid, PR proteins and resistance in tomato and tobacco. International Workshop on Pathogenesis-Related Proteins in Plants Signaling Pathways and Biological Activities, 5th. Aussois, France.
- Chet, I. and Mitchell, R. (1976).** Ecological aspects of microbial chemotactic behaviour. *Annu. Rev. Microbiol.* 30: 221-239.
- Cohen, Y. (1994a).** Local and systemic control of *Phytophthora infestans* in tomato plants by DL-3-amino-n-butanoic acids. *Phytopathology* 84:55-59.
- Cohen, Y. (1994b).** 3-Aminobutyric acid induces systemic resistance against *Peronospora tabacina*. *Physiol. Mol. Plant Pathol.* 44:273-288.
- Cohen, Y.; Niderman, T.; Mosinger, E. and Fluhr, R. (1994).** β -aminobutyric acid induces the accumulation of pathogenesis-related proteins in tomato (*Lycopersicon esculentum* Mill.) and resistance to late blight infections caused by *Phytophthora infestans*. *Plant Physiol.* 104:59-66.
- Cohen, Y. (1996).** Induced resistance against fungal diseases by aminobutyric acids. in: *Modern Fungicides and Antifungal Compounds*. H. Lyr, P. E. Russel, and H. D. Sisler, eds. Intercept, Andover, 461-466.
- Cohen, Y. (2000).** Methods for protecting plants from fungal infection. U.S. Patent 6: 51-75.
- Cohen, (2001).** Synergistic mixtures of an amino acid. Israel Patent 123722.
- Coleno, A.; Hingand, L. and Rat, B. (1976).** Some aspects of the serology of *Ps. solanacearum* E.F. Smith and Application of serology for studying

bacterial wilt. In: Sequeira L. and Kelman A. (eds.), Proceeding of the First International Planning Conference of the Ecology and Control of Bacterial Wilt Disease Caused by *Ps. solanacearum*, 110-119.

Collinge, D. and Slusarenko, A. (1987). Plant gene expression in response to pathogens; Plant Mol. Biol. 9 :389 – 410.

Collins, C. and Patricia, M. (1984). Identification methods. In Microbiological Methods. 5th Ed., Butterworth, Borough Green, Sevenoaks, Kent, England, 56-113.

Cowan, S. and Steel, R. (1974). Manual for the Identification of Medical Bacteria. Second edition, Cambridge Univ., Press. Cambridge, PP. 238.

Crowell, D.; John, M.; Russell, D. and Amasino, R . (1992) Characterization of a stress-induced developmentally regulated gene family from soybean; Plant Mol. Biol. 18 :459–466.

Dakora, F. (2003). Defining new roles for plant and rhizobial molecules in sole and mixed plant cultures involving symbiotic legumes. New Phytologist. 158:39-49.

Degenhardt, J.; Gershenzon, J.; Baldwin, I. (2003). Attracting friends to feast on foes: engineering terpene emission to make crop plants more attractive to herbivore enemies. Current Opinion in Biotechnology, 14 : 169-176.

Delaney, T.; Uknes, S.; Vernooji, B.; Friedrich, L.; Weymann, K.; Negrotto, D.; Gaffney, T.; Gut-Rella, M.; Kessmann, H.; Ward, E. and Ryals, J. (1994). A central role of salicylic acid in plant disease resistance; Science 266 :1247–1250.

Deweert, S. (2002). Flagella-driven chemotaxis towards exudates components is an important trait for tomato root colonization by *Pseudomonas fluorescens*. Mol. Plant–Microbe Interact. 15: 1173–1180.

- Deweger, L.; Vander, V.; Wijfjes, A.; Bakker, P.; Schippers, B. and Lugternberg, B. (1987).** Flagella of a plant growth-stimulating *Pseudomonas fluorescence* strain are required for colonisation of potato roots. J. Bacteriol. 169: 2769-2773.
- Diagnostic protocols for regulated pests, (2004).** *Ralstonia solanacearum* European and Mediterranean Plant Protection Organization 2004 OEPP/EPPO, *Bulletin OEPP/EPPO Bulletin* 34, 173 –178.
- Dicke, M. and Hilker, M. (2003).** Induced plant defences: from molecular biology to evolutionary ecology. Basic and Applied Ecology, 4 : 3-14.
- Dixon, R. (2001)** Natural products and plant disease resistance. Nature 411:843–847.
- Dowson, W. (1957).** Plant disease due to bacteria. Second edition. Cambridge at the University Press, PP. 232.
- Duckett, C.; Oparka, K.; Prior, D.; Dolan, L.; and Roberts, K. (1994).** Dye-coupling in the root epidermis of *Arabidopsis* is progressively reduced during development. 120:3247–3255.
- Eden, G.; Supriadi, N. and Hunt, P. (1987).** Serological relationship between the xylem limited bacterium causing sumatra disease of cloves in Indonesia and *Ps. solanacearum*. In: Civerolo E., Collmer A., Davis R. and Gillaspie A. (eds.), Plant Pathogenic Bacteria. Proceedings of the Sixth International Conference on Plant Pathogenic Bacteria, Maryland, 2-7 June 1985: 357-363.
- El-Didamony, G.; Ismail, A. ; Sarhan, M. and Abdel-Aziz, S. (2003).** Identification and pathogenicity of bacterial caused wilt of potatoes in some types of Egyptian soil. Egyptian Journal of Microbiology. 38: 89-103.
- El-Mishad, A. (2002).** Manual of Practical Microbiology. Commercial press, Kalyoub, Egypt.

- Elphinstone, J. and Aley, P. (1993).** Integrated control of bacterial wilt of potato in the warm tropics of Peru. Pages 276-283 in: Bacterial Wilt. ACIAR Proc. 45. G. L. Hartman and A. C. Hayward, eds. Australian Centre for International Agricultural Research, Canberra, Australia.
- Elphinstone, J. (1996).** Survival and possibilities for extinction of *Ps. solanacearum* (Smith) in cool climates. Potato Research, 39: 403-410.
- Elphinstone, J.; Hennessey, J.; Wilson, J. and Stead, D. (1996).** Sensitivity of different methods for the detection of *Pseudomonas solanacearum* in potato tuber extracts. *Bulletin OEPP/EPPO Bulletin* 26: 663–678.
- Engelbrecht, M. (1994).** Modification of a semi-selective medium for the isolation and quantification of *Pseudomonas solanacearum*. ACIAR Bacterial Wilt Newsletter, 10: 3-5.
- Estabrook, E. and Yoder, J. (1998).** Plant-plant communications: rhizosphere signaling between parasitic angiosperms and their hosts. *Plant Physiol* 116: 1-7.
- European and Mediterranean Plant Protection Organization. *Ralstonia solanacearum*, (2004).** Diagnostic protocols for regulated pests. *Bulletin OEPP/EPPO Bulletin* 34:173 –178.
- Fahy, P. and Persley, G. (1983).** Plant bacterial disease. A Diagnostic Guide. Academic Press, New York.
- Farag, N.; Stead, D. and Janse, J. (1999).** Short Communication. *Ralstonia (Pseudomonas) solanacearum* race 3, biovar 2, Detected in Surface (irrigation) Water in Egypt. Short Communication Potato Brown Rot Project (PBRP), Dokki, Cairo, Egypt. *Journal of Phytopathology*. 147: 478- 485.
- Flores, H.; Vivanco, J. and Loyola-Vargas, V. (1999).** "Radicle" biochemistry: the biology of root-specific metabolism. *Trends Plant Sci* 4: 220-226.

- Fujii, Y. (1989).** Prospective views of biological control by allelochemicals. Jpn. J. Soil Sci. Plant Nutr. 60:240-245.
- Gaffney, T.; Friedrich, L.; Vernooij, B.; Negrotto, D.; Nye, G.; Uknes, S.; Ward, E.; Kessmann, H. and Ryals, J. (1993).** Requirement of salicylic acid for the induction of systemic acquired resistance; Science 261: 754–756.
- Gilpatrick J. (1969).** Role of ammonia in the control of avocado rootrot with alfalfa meal soil amendment. Phytopathology 59:973-978.
- Griffin, G.; Hale, M. and Shay, F. (1976).** Nature and quantity of sloughed organic matter produced by roots of axenic peanut plants. Soil Biol Biochem 8: 29-32
- Grodzinsky, A.(1992).** Allelopathic effects of *cruciferous* plants in crop rotation. Allelopathy: Basic and Applied Aspects. Chapman and Hall, London. 77-86
- Gunawardena, U. and Hawes, M. (2002).** Tissue-specific localization of root infection by fungal pathogens: role of root border cells. Mol. Plant–Microbe Interact. 15:1128–1136.
- Hara, H. and Ono, K. (1983).** Ecological studies on the bacterial wilt of tobacco caused by *Pseudomonas solanacearum* E.F. Smith. Bull. Okayama Tob. Exp. Stn. 42: 127-138.
- Harnack, W. and Schwarz, J. (1966).** Fungicidal compositions. UK Patent 1: 408-507.
- Harrison, D. and Freeman, H. (1961).** Bacterial wilt of potatoes. II. Serological relationship of two strains of *Ps. solanacearum* and a culture of *Corynebacterium sepdonicum*. Australian Journal of Agricultural Research, 12: 872-877.
- Harsh Pal, B.; Sang-Wook, P.; Tiffany, L.; Weir1, R. ; Callaway, M. and Jorge, M.(2004).** How plants communicate using the underground information superhighway. Trends in Plant Science. 9:26-32.

- Hawes, M.; Gunawardena, U.; Miyasaka, S. and Zhao, X. (2000).** The role of root border cells in plant defense. *Trends Plant Sci* 5: 128-133 .
- Hayward, A. (1964).** Characteristics of *Pseudomonas solanacearum*. *J. Appl. Bacteriol.* 27:265-277.
- Hayward, A., (1991).** Biology and Epidemiology of a bacterial wilt caused by *Pseudomonas solanacearum*. *Annual Review of Phytopathology*, 29: 65-87.
- Hayward, A.(2000).** *Ralstonia solanacearum*. In: Encyclopedia of Microbiology, Lederberg J. (ed.), Academic Press. 4: 32-42.
- He, L.; Sequeira, L. and Kelman, A. (1983).** Characteristics of *Pseudomonas solanacearum* from China. *Plant Dis.*, 67: 1357-1361.
- Heather, A. (2005).** *Ralstonia solanacearum*: Requirement for Soilborne Plant Pathogens, Springer 2005: PP 728.
- Hirsch, A., Bauer W., Bird D., (2003).** Molecular signals and receptors: controlling rhizosphere interactions between plants and other organisms. *Ecology*, 84 : 858-868.
- Hoaglands, D. And Arnon, D. (1938)** *Circ.-Calif. Agric. Exp. Sta.* 347.
- Huang, H. , and Huang, J. (1993).** Prospects for control of soilborne plant pathogens by soil amendment. *Curr. Topics Bot. Res.* 1:223-235.
- Huang, J .;Chang, F. and Wang, C.(1997).** Characterization of a lily tapetal transcript that shares similarity with a class of intracellular pathogenesis-related (IPR) proteins; *Plant Mol. Biol.* 34 :681–686.
- Hwang, B. ; Sunwoo, J. ; Kim, Y. and Kim, B. (1997).** Accumulation of beta-1,3- glucanase and chitinase isoforms, and salicylic acid in the DL-beta-amino-n-butyric acid-induced resistance response of pepper stems to *Phytophthora capsici*. *Physiol. Mol. Plant Pathol.* 51:305-322.
- Jacobs, M. and Gerstein, M. (1960).** Hand-book of Microbiology. Van D. Nostrand Co., Inc., New York, USA, PP. 139-202.

- Jakab, G.; Cottier, V.; Toquin, V.; Rigoli, G.; Zimmerli, L.; Metraux, J. and Mauch- Mani, B. (2001).** β -Aminobutyric acid-induced resistance in plants. *Eur. J. Plant Pathol.* 107:29-37.
- Janse, J. (1988).** A detection method for *Pseudomonas solanacearum* in symptomless potato tubers and some data on its sensitivity and specificity. *EPPO Bulletin*, 18: 343-351.
- Janse, J. (1996).** Potato brown rot in Western Europe – history, present occurrence and some remarks on possible origin, epidemiology and control strategies. *Bulletin of OEPP*, 26: 679-695.
- Janse, J.; Van den, B.; Ephinstone, J.; Simpkins, S.; Tjou-Tam, S. and Van Vaerenbergh, J. (2004).** Introduction to Europe of *Ralstonia solanacearum* biovar 2, race 3 in *Pelargonium zonale* cuttings. *Journal of Plant Pathology* 87: 147–155.
- Jaunet, T. and Wang, J. (1999).** Variation in genotype and aggressiveness of *Ralstonia solanacearum* race 1 isolated from tomato in Taiwan. *Phytopathology*, 89: 320-327.
- Jeun, Y. (2000).** Immunolocalization of PR protein P14 in leaves of tomato plants exhibiting systemic acquired resistance against *Phytophthora infestans* induced by pretreatment with 3-aminobutyric acid and reinoculation with tobacco necrosis virus. *Z. Pflanzenkrankh. flanzenschutz* 107:352-367.
- Jeun, Y. and Buchenauer, H. (2001).** Infection structures and localization of the pathogenesis- related protein AP24 in leaves of tomato plants exhibiting systemic acquired resistance against *Phytophthora infestans* after pre-treatment with 3-aminobutyric acid or tobacco necrosis virus. *J. Phytopathol.* 149:141-153.
- Jeun, Y. ; Siegrist, J. and Buchenauer, H. (2000).** Biochemical and cytological studies on mechanisms of systemically induced resistance to *Phytophthora infestans* in tomato plants. *J. Phytopathol.* 148:129-140.

- Jian, Y. and Caitilyn, A. (2006).** Chemotaxis Is Required for Virulence and Competitive Fitness of the Bacterial Wilt Pathogen *Ralstonia solanacearum*. Journal of Bacteriology American Society for Microbiology. 188: 3697–3708.
- Jing, Q. (1999).** Allelopathic suppression of *pseudomonas solanacearum* infection of tomato (*Lycopersicon Esculentum*) in a tomato-chinese chive (*Allium tuberosum*) intercropping system. journal of chemical ecology, 25.
- Jones, D. (2003).** Associative nitrogen fixation and root exudation – what is theoretically possible in the rhizosphere? Symbiosis 35:19–38
- Kalburtji, L. and mosjidis, J. (1993).** Effects of sericea lespedeza root exudates on some perennial grasses. J. Range Manage. 46:312-315 .
- Kang S. and Mills A.. (2004). Soil bacterial community structure changes following disturbance of the overlying plant community. Soil Science. 169 :55-65.
- Kang, Y.; Mao, G. He-ly, V.; Mehan, K. and Mcdonald, D. (1994).** Factors involved in virulence and pathogenicity of *Ps. solanacearum* and their role in pathogenesis. Groundnut Bacterial Wilt in Asia. Proceeding of the Third Working Group Meeting 4-5 July 1994. Oil Crops Research Institute, Wuhan, China, 53-62; 23 ref.
- Karganilla, A. and Buddenhagen, I. (1972).** Development of selective medium for *Pseudomonas solanacearum*. Phytopathology, 62: 373-376.
- Kelman, A. (1954).** The relationship of pathogenicity in *Pseudomonas solanacearum* to colony appearance on a tetrazolium medium. Phytopathology, 44: 693-695.
- Kelman, A.; Hartman, G.; and Hayward, A. (1964).** Introduction. In: Hayward A. and Hartman G. (eds.), Bacterial Wilt: The disease and its causative agent *Pseudomonas solanacearum*. CAB International, Wallingford, UK. 1-7.

- Kempe, J. and Sequeira, L. (1983).** Biological control of bacterial wilt of potatoes attempts to induce resistance by treating tubers with bacterial. *Plant dis.*, 67:499.
- Kessmann, J.; Staub Hofmann, T.; Maetzke, T.; Harzog, J.; Ward, E.; Uknes, S. and Rayls, J. (1994).** Induction of Systemic Acquired resistance in plants by chemicals . *Annual review of plant pathology* 32:439-459.
- King, E.; Ward, M. and Raney, D. (1954).** Two simple media for the demonstration of pyocyanin and fluorescin. *J. Lab. Clin. Med.*, 44: 301-307.
- Klement, Z.; Rudolph, K. and Sands, D. (1990).** *Methods in Phytobacteriology*, Academic Kiado, Budapest, PP. 568.
- Kneer, R. (1999).** Characterization of the elicitor-induced biosynthesis and secretion of genistein from the roots of *Lupinus luteus* L. *J. Exp. Bot.* 50: 1553–1559.
- Kojima, E. and Buddenhagen, I. (1969).** Antigenic relationships of strains of *Ps. solanacearum*. *Phytopathology*, 59: 1035-1036 (abstract).
- Krieg, N. and Holt, J. (1984).** *Pseudomonas solanacearum*. *Bergey's Manual of Systematic Bacteriology*, vol. 1, Williams and Wilkins, Baltimore, London.
- Kuniyasu, K. (1989).** Suppression of soil-borne diseases by soil amendment with cruciferous crops. *Agric. Hortic.* 64:955-959.
- Lelliott, R. and Stead, D. (1987).** *Methods for diagnosis of bacterial diseases of plants. Methods in Plant Pathology*, vol. 2, Ed. T.F. Preece. Blackwell Scientific Publication 216 PP.
- Limpens, E. and Bisseling, T., (2003).** Signaling in symbiosis. *Current Opinion in Plant Biology*, 6 : 343-350.

- Lopez, M. (1993).** Mecanismos de resistencia inducidos por compuestos quimicos frente a la podredumbre blanda en tuberculos de patata. Ph.D. thesis, Universidad Complutense de Madrid , spain , 215pp.
- Louws, F.; Fulbright, D.; Stephens, C. and De Bruijn, F. (1994).** Specific genomic fingerprints of phytopathogenic *Xanthomonas* and *Pseudomonas* pathovars and strains generated with repetitive sequences and PCR. Applied and Environmental Microbiology, 60 : 2286 -2295.
- MacLennan, D.; Kuc, J.; and Williams, E. (1963).** Chemotherapy of the apple scab disease with butyric acid derivatives. Phytopathology 53:1261-1266.
- Mahmoud, S.; Abdl-Hafez, A.; El-Sawy, M.; Bishay, F. and Farag, N. (1978)** Bacterial wilt disease of potato: Root exudates in relation to *Pseudomonas solanacearum* growth. Agricultural Research Review. 56:138-148.
- Makino, T.; Ono, T.; Muso, E. and Honda, G. (1998).** Inhibitory effect of *Perilla frutescens* and its phenolic constituents on cultured murine mesangial cell proliferation. Planta Med; 64: 541–545.
- Marschner, H. (1991).** Root-induced changes in the availability of micronutrients in the rhizosphere. In: Waisel Y., Eshel A., Kakafi U. (eds.), Plant roots, 503.
- Marfa, M. ; Marfa, J. ; Javier, Z. ; Jesus, L. and Sanchez and Ramona ,B. (2001).** effect of acetylsalicylic acid on soft rot produced by *Erwinia Carotovora* subsp . *Carotovora* in potato tubers under green house conditions . potato research ,44: 197-206.
- Mark, A. ;Czarnot, A.; Rex, N.; Franck, E. ;Dayan, C.; Nimbal, I. and Leslie, A. (2001).** mode of action, localization of production, chemical nature, and activity of sorgoleone: a potent psii inhibitor in *sorghum* spp. root exudates symposium weed technology.. 15 :813–825

- Marineau, C.; Matton, D. and Brisson, N. (1987).** Differential accumulation of potato tuber mRNAs during the hypersensitive response induced by arachidonic acid elicitor; *Plant Mol. Biol.* 9: 335–342.
- Matton, D. and Brisson, N. (1989).** Cloning, expression and sequence conservation of pathogenesis related gene transcripts of potato; *Mol. Plant-Microbe Interact.* 2 :325–331.
- Matton, D.; Bell, B. and Brisson, N. (1990).** Nucleotide sequence of a pathogenesis-related gene of potato; *Plant Mol. Biol.* 14 :863–865.
- Matton, D.; Prescott, G.; Bertrandd, C.; Camirand, A and Brisson, N. (1993).** Identification of cis-acting elements involved in the regulation of the Pathogenesis-related gene STH-2 in potato; *Plant Mol. Biol.* 22: 279–291.
- McCarter, S. (1991).** Bacterial wilt in compendium of tomato diseases. Jones, I.B.; R.E. Stall and T.A. Zitter (eds), American Phytopathological Society, Minneosate, USA, 28-30.
- Mccully, E. (1999).** Roots in soil: unearthing the complexities of roots and their rhizospheres. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 50, 695–718.
- Mertaux, J.; Ryals, J.; Wards, E.; Wyss-Benz, M.;Gaudin, J.; Rashdorf ,K.; Schmid, E. ; Blum, W. and Inverardi, B. (1990).** Increase in salicylic acid at the onest of systemic acquiured resistance in cucumber, *Science* 250: 1004-1006 .
- Michel, V. and Mew, T. (1998).** Effect of a soil amendment on the survival of *Ralstonia solanacearum* in different soils. *Phytopathology* 88: 300-305.
- Mithofer, A. (2002).** Suppression of plant defence in *rhizobia*–legume symbiosis. *Trends Plant Sci.* 7: 440–444.
- Murry, M.; Zang, D.; Schneider, M. and De Bruijin, F. (1995).** Use of repetitive sequences and the polymerase chain reaction (Rep-PCR) to fingerprinting the genomes of *Frankia* isolates. *Symbiosis*, 19: 223-240.

- Nakashima, J. and Nydegger, U. (1986).** Detection de *Ps. solanacearum* mediante las tecnicas serologicas de latexy ELISA. Fitopatologia, 21: 46-51.
- Narasimhan, K. (2003).** Enhancement of plant–microbe interactions using a rhizosphere metabolomics-driven approach and its application in the removal of polychlorinated biphenyls. Plant Physiol. 32:146–153.
- Nardi, S. (2000).** Soil organic matter mobilization by root exudates. Chemosphere 5: 653–658.
- Nesmith, W. and Jenkins, S. (1979).** A selective medium for the isolation and quantification of *Pseudomonas solanacearum* from soil. Phytopathology, 69: 182-185.
- Neumann, G. and Martinoia, E. (2002).** Cluster roots – an underground adaptation for survival in extreme environments. Trends Plant Sci. 7:162–167.
- Nielsen, L. and Todd, F. (1945).** Preliminary evaluation of some soil disinfestants for controlling Southern bacterial wilt of potatoes. Am. Potato J. 22:197-202.
- Nimbal, C.; Pedersen, J. ; Yerkes, C. ; Weston, L. and Weller, S. (1996).** Phytotoxicity and distribution of sorgoleone in grain sorghum germplasm. J. Agric. Food Chem. 44:1343–1347.
- Okabe, N. (1969).** Population changes of *Pseudomonas solanacearum* and soil microorganisms in artificially infested natural field soils. Bull. Fac. Agr. Shizuoka Univ., 19: 1-29.
- Oort, A. and Van Andel, O. (1960).** Aspects in chemotherapy. Mededel. Opz. Gent. 25:961-992.
- Ovadia, A. (2001).** Mode of action of 3-aminobutyric acid in systemic acquired resistance. Ph.D. thesis. Bar-Ilan University, Ramat-Gan, Israel.

- Palva, T.; Hurting, M. ; Saindrenan, P. and Palva, E. (1994).** salicylic acid induced resistance to *Erwinia Carotovora subsp. Carotovora* in tobacco molecular plant microbe interaction 7:356-363.
- Papavizas, G. and Davey, C. (1963).** Effect of amino compounds and related substances lacking sulfur on *Aphanomyces* root rot of peas. Phytopathology 53:116-122.
- Papavizas, G. (1966).** Suppression of *Aphanomyces* root rot of peas by cruciferous soil amendment. Phytopathology 56:1071-1075.
- Pastrik, K; Elphinstone, J. and Pukall, R. (2002).** Sequence Analysis and Detection of *Ralstonia solanacearum* by Multiplex PCR Amplification of 16S–23S Ribosomal Intergenic Spacer Region with Internal Positive Control. European Journal of Plant Pathology, 108: 831-842.
- Patrick, Z. (1986).** Allelopathic mechanisms and their exploitation for biological control. Can. J. Plant Pathol. 8: 225-228.
- Pegg, K. and Moffett, M. (1971).** Host range of the ginger strain of *Pseudomonas solanacearum* in Queensland. Aust. J. Exp. Agric. Anim. Husb. 11: 696-698.
- Peters, N. (1986).** A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. Science 233: 977–980.
- Phillips, D. (2000).** Biosynthesis and release of rhizobial nodulation gene inducers by legumes. In Prokaryotic Nitrogen Fixation: A Model System for the Analysis of a Biological Process (Triplett, E.W., ed.) pp. 349–364, Horizon Scientific.
- Phillips, D. ; Ferris, H.; Cook, D. and Strong, D., (2003).** Molecular control points in rhizosphere food webs. Ecology. 84: 816-826.
- Picard, C.; Ponsonnet, C.; Paget, E.; Nesme, X and Simonet, P. (1992).** Detection and enumeration of bacteria in soil by direct DNA extraction and

polymerase chain reaction. *Applied and Environmental Microbiology* 58: 2717-2722.

Poussier, S.; Cheron, J.; Couteau, A. and Luisetti, J. (2002). Evaluation of procedures for reliable PCR detection of *Ralstonia solanacearum* in common natural substrates. *Journal of Microbiological Methods* 51: 349-359.

Power, R.(1983). Relationship between the soil environment and tomato resistance to bacterial wilt (*Pseudomonas solanacearum*): 4. Control Methods. *Surinaamse Landbouw* 31:39-47.

Pradhanang, P.; Elphinstone, J. and Fox, R.(2000). Identification of crop and weed hosts of *Ralstonia solanacearum* biovar 2 in the hills of Nepal. *Plant Pathology* 49:403–413.

Prior, P. and Bérémis, M. (1990). Induction de la résistance au flétrissement bactérien dû à *Pseudomonas solanacearum* E F Smith chez un cultivar de tomate réputé sensible. *Agronomie* 10:391-401.

Putnam, A.(1986). Allelopathy: Can it be managed to benefit horticulture? *HortScience* 21:411-413.

Radoslava, M. and Harro, j. (2006). the effect of host-root-derived chemical signals on the germination of parasitic plants. marcel dicke and willem takken (eds.), *chemical ecology: from gene to ecosystem*, 39-54 springer. printed in the netherlands.

Raskin, I. (1992). Role of salicylic acid in plants . *Annual review of plant pathology and plant molecular Biology* 43:439-463.

Rasmann, S.; Kollner, T. and Degenhardt J., (2005). Recruitment of entomopathogenic nematodes by insect-damaged maize roots. *Nature*, 434: 732-737.

- Rasmussen, J.; Hammerschmidt, R; and Zook, M. (1991).** Systemic induction of Salicylic acid accumulation in cucumber after inoculation with *Pseudomonas syringae* pv . *Syringae* . plant pathology 97:1342-1347.
- Raviv, A. (1994).** The mode of action of β - aminobutyric acid in inducing resistance in tomato plants against late blight. Ph.D. thesis. Bar-Ilan University, Ramat-Gan, Israel.
- Rice, E. (1995).** Biological Control of Weeds and Plant Diseases: Advances in Applied Allelopathy. University of Oklahoma Press, Oklahoma.
- Robinson, A. (1993).** Serological detection of *Ps. solanacearum* by ELISA. In: Hartman G. and Hayward A., (eds.), Bacterial Wilt. Proceedings of an International Symposium, Kaohsiung, Taiwan, ROC, 28-30 October, 1992. ACIAR Proceedings 45: 54-61.
- Rohde, (1960).** Acetylcholinesterase in plant-parasitic nematodes and an anticholinesterase from asparagus. *Helminthol. Soc.* 27:121-123.
- Rovira, A. (1973).** Zones of exudation along plant roots and spatial distribution of microorganisms in the rhizosphere . *pestic . sci* 4 : 361 – 366.
- Rush, C. and Lyda, S. (1982).** Effects of anhydrous ammonia on mycelium and sclerotia of *Phymatotrichum omnivorum*. *Phytopathology* 72:1085-1089.
- Ryals, J.; Neuenschwander, U. ; Willits, G. ; Molina, A. ; Steiner, H. and Hunt ,M. (1996).** Systemic acquired resistance; *Plant Cell* 8: 1809.
- Ryan, P. and Delhaize, E. (2001).** Function and mechanism of organic anion exudation from plant roots. *Annu Rev Plant Physiol Mol Biol* 52: 527-560.
- Scarpati, M. and Oriente, G. (1958).** Isolation and constitution of rosmarinic acid from *Rosmarinus officinalis*. *Ricerca Sci*; 28: 2329–2333
- Schaad, N. (1980).** Laboratory guide for identification of plant pathogenic bacteria. The American Phytopathological Society, Printed in USA.

- Schaad, N; Cheong, S.; Tamaki, S.; Hatziloukas, E. and Panopoulos, N. (1995).** A combined biological and enzymatic amplification (Bio-PCR) technique to detect *Ps. syringae* pv. *phaseolicola* in bean seed extracts. *Phytopathology*, 85: 243-284.
- Scher, F.; Kloepper, J. and Singleton, C., (1985).** Chemotaxis of fluorescent *Pseudomonas* spp. To soybean seed exudates in vitro and in soil. *Can. J. Microbiol.* 31: 570-583.
- Sequeira, L. (1958).** Bacterial wilt of bananas: Dissemination of the pathogen. *Vol. 88, No. 4, 1998 305* gen and control of the disease. *Phytopathology* 48:64-69.
- Shehata and Nevein, A. (2001).** Studies on Biological Control of Potato Brown Rot Disease (M.Sc. Thesis). Department of Botany (Plant Pathology), Fac. of Science, Zagazig Univ. (Benha branch), PP. 69 - 158.
- Shailasree, S.; Sarosh, B.; Vasanthi, N. and Shetty H. (2001).** Seed treatment with β -aminobutyric acid protects *Pennisetum glaucum* systemically from *Sclerospora graminicola*. *Pest Manag. Sci.* 57:721-728.
- Siegrist, J.; Orober, M. and Buchenauer, H. (2000).** Beta-aminobutyric acid-mediated enhancement of resistance in tobacco to tobacco mosaic virus depends on the accumulation of salicylic acid. *Physiol. Mol. Plant Pathol.* 56:95-106.
- Smiley, R.; Cook, R. and Papendick, R. (1970).** Anhydrous ammonia as a soil fungicide against *Fusarium* and fungicidal activity in the ammonia retention zone. *Phytopathology* 60:1227-1232.
- Smith, T. (1944).** Control of bacterial wilt (*Bacterium solanacearum*) of tobacco as influenced by crop rotation and chemical treatment of the soil. *Circ. 692, U.S. Dep. Agric., Washington, DC.*

- Smith, J.; Offord, L.; Holderness, M. and Saddler, G. (1995).** Generic diversity of *Burkholderia solanacearum* race 3 in Kenya. App. Environ. Microbiol., 61: 4263-4268.
- Sree Vidya, C.; Manoharan, M. and Lakshmi Sita, G. (1999).** Cloning and characterization of salicylic acid-induced, intracellular pathogenesis-related gene from tomato (*Lycopersicon esculentum*). 24: 287-293.
- Steven, B. (2003).** Spudlines. Seed Issue. 41: 1-9.
- Sticher, L.; Mauch-Mani, B. and Mertaux, P. (1997).** Systemic Acquired Resistance. annual Review of plant pathology 35:235-270.
- Strider, D.; Jones, R. and Haygood, R. (1981).** Southern bacterial wilt of geranium caused by *Pseudomonas solanacearum*. *Plant Disease* 65: 52–53.
- Sun, S. and Huang, J. (1985).** Formulated soil amendment for controlling Fusarium wilt and other soilborne diseases. *Plant Dis.* 69:917-920.
- Tang, C.; Wat, C. and Towers, G. (1987).** Thiophanes and benzofurans in the undisturbed rhizosphere of *Tagetes patula* L. *Plant Soil* 98:93-97.
- Tans-Kersten, J.; Guan, Y. and Allen, C. (1998).** *Ralstonia solanacearum* pectin methylesterase is required for growth on methylated pectin but not for bacterial wilt. *Applied and Environmental Microbiology*, 65: 4918-4923.
- Tarah, S. (2004).** Interactions between Soil Microbial Communities and Plant Roots: A Minireview. *Soil and Crop Sciences*, Colorado State University.1-16.
- Trieu, A. (1997)** Gene expression in mycorrhizal roots of *Medicago truncatula*. In *Radical Biology: Advances and Perspectives on the Function of Plant Roots* (Flores, H.E. et al., eds), American Society of Plant Physiologists .498–500.
- Toshiaki, M.; Takahiko, O.; Eri, M.; Haruyoshi, Y.; Gisho, H. and Shigetake, S. (2000).** Inhibitory effects of rosmarinic acid on the

proliferation of cultured murine mesangial cells. Nephrol Dial Transplant 15: 1140-1145.

Tsai, Y. and Olson, B. (1991). Rapid method for direct extraction of DNA from soil and sediments. Appl. Environ. Microbiol., 57: 1070-1074.

Tsao, P. and Oster, J. (1981). Relation of ammonia and nitrous acid to suppression of *Phytophthora* in soils amended with nitrogenous organic substances. Phytopathology 71:53-59.

Vaughan, E. (1944). Bacterial wilt of tomato caused by *Pseudomonas solanacearum*. Phytopathology, 34: 443-458.

Van Andel, O. (1958). Investigations on plant chemotherapy II. Influence of amino acids on the relation plant-pathogen. Tijdschr. Planteziekten 64:307-327.

Van Etten, H. (1994). Two classes of plant antibiotics: phytoalexins versus phytoanticipins. Plant Cell 6: 1191–1192.

Van Kan, J. ; Cozijnsen, T.; Danhash, N. and De Wit, P. (1995). Induction of tomato stress protein mRNA by ethephon, 2,6-dichloroisonicotinic acid and salicylate; Plant Mol. Biol. 27 :1205–1213.

Van West, P. (2002). Oomycete plant pathogens use electric fields to target roots. Mol. Plant–Microbe Interact. 15: 790–798.

Vasse, J.; Frey, P. and Trigalet, A. (1995). Microscopic studies of intercellular infection and protoxylem invasion of tomato roots by *Pseudomonas solanacearum*. The American Phytopathological Society. 8: 241-251.

Vermeer, J. and McCully, M. (1982). The rhizosheath of Zea – new insight into the structure and development. Planta 156:45–61.

Vernooij, B.; Friedrich, L.; Morse, A.; Reist, R.; Kolditz-Jawhar, R.; Ward, E.; Uknes, S.; Kessmann, H. and Ryals, J. (1994). Salicylic acid is

not the translocated signal responsible for inducing systemic acquired resistance but is required signal transduction . plant cell 6:959-965.

Vincent, V. and Mew, T. (1998). Effect of a Soil Amendment on the Survival of *Ralstonia solanacearum* in Different Soils 88:300-305.

Walker, J. (1950). Plant pathology MC. Grow. Hill book Co. Inc. New York, Toronto, London (3rd ed., 1969).

Walker, J. (1971). Populations of *Pratylenchus penetrans* relative to decomposing nitrogenous soil amendments. J. Nematol. 3:43-49.

Walker, T. (2003). Metabolic profiling of root exudates of *Arabidopsis thaliana*. J. Agric. Food Chem. 51:2548–2554.

Walker, T.; Pal Bais, H.; Grotewold, E. and Vivanco, J.(2003). Root Exudation and Rhizosphere Biology. Plant Physiology. 132: 44-51.

Warner, S.; Scott, R and Draper, J. (1992). Characterization of a wound-induced transcript from the monocot asparagus that shares similarity with a class of intracellular pathogenesis-related (PR) proteins; Plant Mol. Biol. 19 :555–561.

Wenneker, M.; Verdel, M.; Van Beuningen, A.; Derks, J. and Janse, J. (1999). *Ralstonia (Pseudomonas) solanacearum* race 3 (biovar 2) in surface water and natural weed hosts: first report on stinging nettle (*Urtica dioica*). *European Journal of Plant Pathology* 105:307–315.

White, R. (1979). Acetylsalicylic acid (aspirin) Inducing resistance to tobacco mosaic virus in tobacco . Virology 99:410-412.

Wilson, I. (1997). Inhibition and facilitation of nucleic acid amplification. Applied and Environmental Microbiology 63: 3741-3751.

Wo, H.(2002). Flavonoids: signal molecules in plant development. Adv. Exp. Med. Biol. 505:51–60.

Wu, H. (2000). Allelochemicals in wheat (*Triticum aestivum* L.): variation of phenolic acids in root tissues. J. Agric. Food Chem. 48:5321–5325.

- Yabuuchi, E.; Kosako, Y.; Oyaizu, H.; Yano, I.; Hotta, H.; Hashimoto Y., Ezaki, T. and Arakawa, M. (1992).** Proposal of *Burkholderia* gen. nov. and transfer of seven species of the genus, *Pseudomonas* homology group. II to the new genus, with the type species *Burkholderia capacia* (Palleroni and Holmes 1981) Comb. nov. Microbial. Immunol., 36: 1251-1275.
- Yabuuchi, E.; Kosako, Y.; Yano, I.; Hotta, H. and Nishiuchi, Y. (1995).** Transfer of two *Burkholderia* and an *Alcaligenes* species to *Ralstonia* General Nov: Proposal of *Ralstonia pikettii* (Ralston, Palleroni, and Doudoroff 1973) comb. Nov, *Ralstonia solanacearum* (Smith, 1896) comb. Nov and *Ralstonia eutropha* (Davis, 1969) comb. Nov. *Microbiology and Immunology*, 39: 897-904.
- Yamamoto, M; Torikai, S. and Oeda, K (1997).** A major root protein of carrots with high homology to intracellular pathogenesis-related (PR) proteins and pollen allergens; *Plant Cell Physiol.* 9 :1080–1086.
- Yigal, R. (2002).** β -Aminobutyric Acid-Induced Resistance Against Plant Pathogens. *Plant Disease.* 86 : 448-457.
- Yoder, J. (1999).** Parasitic plant responses to host plant signals: a model for subterranean plant–plant interactions. *Curr. Opin. Plant Biol.* 2: 65–70.
- Yu, J. and Matsui, Y. (1994).** Phytotoxic substances in the root exudates of *Cucumis sativus* L. *J. Chem. Ecol.* 20: 21-13.
- Zheng, X. and Sinclair, J. (1996).** Chemotactic response of *Bacillus megaterium* strain B153-2-2 to soybean root and seed exudates. *Physiol. Mol. Plant Pathol.* 48: 21–35.
- Zimmerli, L.; Jakab, C.; Metraux, J. and Mauch-Mani, B. (2000).** Potentiation of pathogen- specific defense mechanisms in *Arabidopsis* by β -aminobutyric acid. *Proc. Nat. Acad. Sci. USA* 97:12920-12925.

Zimmerli, L.; Metraux, J. and Mauch- Mani, B. (2001). β -aminobutyric acid-induced protection of *Arabidopsis* against the necrotrophic fungus *Botrytis cinerea*. Plant Physiol. 126:517-523.