

REFERENCES

REFERENCES

- Adachi I, Kimura S, Abe K, (1984):
Production of multiple hormones in medullary carcinoma of the thyroid;
In : Seventh International congress of Endocrinology,
Amsterdam Oxford/Princeton : Excerpta Medica;
284-290.
- Addison R, Heward P, Kamal A, (1972):
Immunoenzymetric assay for quantitative measurement
of ferritin in serum.
Ann Otol Rhinol laryngol; 95 : 70-72.
- Al-Sarraf M, Shu T, Lai I, (1981):
Tumour markers in patients with epidermoid cancer of
head and neck and the effect of therapy.
Head Neck Surg; 3 : 246-251.
- Anderson NE, Cunningham M, Posner JB, (1984):
Autoimmune pathogenesis of paraneoplastic neurological
syndromes.
Clin Neurobiol; 91: 1168-1173.
- Amos P, Urist W, Speer K, (1968).
Cellular and humoral immune responses to human
urinary bladder carcinomas.
Int J Cancer; 5 : 310- 313.
- Arakawa F, Takakura F, Okamoto N, (1987):
Monoclonal antibodies against the carbohydrate
moiety of carcinoembryonic antigen and closely
related antigens.
Cancer Res; 47:66- 71.
- Barton FE, Cottal WI, Walker B, (1987):
The principle of chemosurgery supramyohyoid neck
dissections. Am J surg; 146: 439-440.
- Bast R, Klug T, St John E, et al. (1983):
Radioimmunoassay using a monoclonal antibody to
monitor the course of epithelial ovarian cancer.
N Engl J Med; 509:883-7.

- Baylin S, Mendelson G (1982):
Medullary thyroid carcinoma: a model for the study of
tumour progression and cell heterogeneity.
In: Owens A, Coffey D, Baylin S (eds) Tumour cell
heterogeneity-Origin and Implications. Academic
press, New York, P9.
- Beauduin M, Baldewyns P. (1981):
Comparative evaluation of the clinical relevance of
CEA and ferritin testing. in : Unlenbruck G, Wintzer
G. eds: CEA und andere Tumormarker.
Limburg: Tumor diagnostik verlag, 293- 8.
- Begent RHJ, (1985):
Recent advances in tumour imaging. Use of
radiolabelled antitumour antibodies.
Biochemica et Biophysica Acta; 780: 151 -166.
- Bhata Vdekkar M, Vora H, Shah G, (1987):
Significance of ferritin as a marker in head and neck
malignancies.
Tumori; 73: 59 - 63.
- Bhatia P, Singh M, Chakravorty M, (1971):
Enzymatic study in diagnosis of head and neck cancer.
Indian J Otolaryngol; 31: 76-79.
- Bhatia P, Singh h, Fous M, (1979):
A preliminary study of blood group substances A and B
in oral epithelium exhibiting atypia.
J Oral Pathol; 13 : 155- 165.
- Black G, Cassoni A, Gimelette D, et al. (1981):
serum thyroglobin in thyroid cancer.
Br Med J; 3 : 443 - 445.
- Blumgart L, (1971):
Tumours of the gut
Am J surg; 12 : 922 - 925.
- Bocca E, Pignataro O, Oldini C, (1983):
Supraglottic laryngectomy, 30 years experience.
Ann Otol Rhinol laryngol: 92: 14-18.

- Boesen A, Hokland P, Jensen O, (1984):
Acid phosphatase and acid estrase activity in
neoplastic and non-neoplastic lymphoid cells A semi
quantitative evaluation related to immunological
marker in 122 cases.
Scand J haematol; 32 : 313 - 322.
- Boorman K, Dodd B, Klein P, (1943):
Substances A, B, M, N and Rh : Their occurrence in
tissues and body fluids.
J pathol; 55 : 329 - 339.
- Botach H, Glatz J, Wenzel W, et al (1983):
Long term follow - up using serial serum
thyroglobulin determinations in patients with
differentiated thyroid carcinoma.
Cancer; 52 : 1850 - 1859.
- Boyd R, (1966)
Active immunotherapy for acute lymphoblastic
leukemia.
Lancet: 1 : 697-704.
- Bunn P A, (1988):
Paraneoplastic syndromes in Cecil textbook of
medicine by: Wyngaarden JB, Smith LH (eds), Published
by W.B. Saunders Co, : 1097.
- Burgdorf C, Mukai K, Posai J, (1981):
Immunohistochemical identification of factor VIII -
related antigen in endothelial cells of cutaneous
lesions of allergic vascular nature,
Am J clin Pathol; 75-171.
- Burge L, Goodwin G, Baylin S et al (1981):
Endocrine-related biochemistry in the spectrum of
human lung carcinoma.
J Clin Endocrinol Metab; 53: 422 -427.
- Burge L, Purnell M, Harris C, et al, (1986).
Immunocytochemical evaluation of human oesophageal
neoplasms and preneoplastic lesion for B chronic
gonadotropin and placental lactogen.
Cancer Res, 46: 2936-2943.

Burkitt D, (1966):

The african lymphoma - etiology and diagnosis.
J Roy coll surg, Edin; II; 170- 175.

Busnardo B, Gierilli E, Zarot P, et al, (1983):

Different diagnostic significance of carcinoembryonic antigen and calcitonin as tumour markers of medullary thyroid carcinoma.
Acta Endocrinol; 102 : 57 - 63.

Compbell JB, Crocker J, and shenoi PM <. (1988):

S 100 protein localisation in minor salivary gland tumours: an aid to diagnosis.
I J laryngol Otol; 102 : 905 - 908.

Caldarelli D, Javid H, lee I, (1988):

Neuroendocrine markers in paraganliomas of the head and neck.
Ann otol Rhinol laryngol; 94 : 555- 559

Caselitz J, Jaup T, Seifert G, (1981):

Lactoferrin and lysozyme in carcinomas of the parotid gland.
Virchows Arch (Pathol Anat): 394: 61 - 73.

Caselitz J, Becker J, Seifert G, (1983):

Coexpression of keratin and vimentin filaments in adenoid cystic carcinomas of salivary glands.
Virchows Arch (Pathol Anat): 403: 337- 344.

Cellews M, (1956):

Antibodies elicited by cancer in patients.
cancer; 8 : 409 - 415.

Cerrato M, Poggio E, Buossolati G, (1988):

Monoclonal antibodies against epithelial antigens in laryngeal carcinoma.
J laryngol Otol; 102: 713-715

Chaganti R, German I, Reto R, (1985):

Genetics in clinical onchology
NewYork, Oxford University Press; 305 - 315.

- Chen Q, Fan F, liu Z, (1986):
Biochemical markers of tumour diathesis and early cancer.
Cancer detect prevent; 9: 145 - 149.
- Clamettes S, Twomey R, Runke P, (1977):
In Vitro synthesis of functional ovarian sarcoma virus transforming-gene product.
Cancer; 26: 588 - 594.
- Cohenhim R, (1975):
Sequential evaluation of general immune comptence in cancer patients: correlation with clinical course.
Cancer; 35: 660 - 667.
- Collens V, (1956):
Simple and malignant tumour: an aid to diagnosis.
Am J Roentgen; 76: 988 - 992.
- Cortesina G, Part A, CerratoM, et al (1988):
Monoclonal antibodies against epithelial antigens in laryngeal carcinomas: An immunocy to chemical and clinicopathological investigation.
J laryngol Otol; 102 : 709 - 712.
- Cox D, Pierquin B, Chassagn R, (1988):
Towards consistent coccal control of certain malignant tumours
Radiology; 99: 661 - 667.
- Cragg S, Wagstaff M, Worwood M, (1980):
Sialic acid and the microheterogeneity of human serum ferritin.
Clin Sci; 58: 259 - 262.
- Crocker J, Jenkis R, Shah V, (1985):
Immunohistochemical demonstration of S- 100 protein in salivary gland neoplasms.
J pathol; 146: 115 - 121 .
- Damle S, Talvedkar R, Rause B, (1971):
The study of glycolytic enzymes in relation to cancer II.
Ind J cancer. 8 : 21 -- 29.

- Devidson I kamal A , lasser A, (1972):
Early immunologic diagnosis and prognosis of
carcinoma.
Am J clin pathol; 57: 715-730 .
- Dearnley D, Stafford D, (1988):
Treatment of " inoperable" neck nodes using surgical
clearance and postoperative interstitial irradiation.
Brit J surg; 75 : 62 - 64.
- Deconzo M, Howard P, Irish E, (1975):
Antigenic deletion and prognosis of patients with
stage A transitional cell bladder carcinoma.
J Urol; 114 : 874 - 787.
- Deland F, (1968):
Diagnosis of early squamous cell carcinoma in head
and neck.
J Radiol; 91 : 1195 - 1197.
- Dredsale JW, (1983):
Ferritin as a tumour marker.
Clin J Immunoassay, 6: 234-240.
- Eagan M, Crocke J, Nar P, (1988):
Localisation of salivary amylase and epithelial
membrane antigen in salivary gland tumours by means
of immunoperoxidase and immunogold-silver techniques.
J laryngol Otol; 102 : 242 -247.
- Ellis L, langlossM, Heffner K (1987):
Spindle cell carcinoma of the Aerodigestive Tract -
An immunohistochemical analysis of 21 cases.
Am J surg pathol; 11 (5) : 335 - 342.
- Fairly S, (1971):
Cutaneous hypersensitivity reaction to autologous
extracts of malignant melanoma cells.
Lancet; 1 : 116 -120.

Feichter G, Abel U, Haag D. (1987):

S- phase fractions and DNA - ploidy of oropharyngeal squamous epithelium carcinomas compared with histologic grade, stage, response to chemotherapy and survival.

Acta Otolaryngol (stockh); 104 : 377-384.

Femenser R, Gay I, weisel M, (1982)

Malignant transformation in inverted papilloma.

Ann Otol Rhinol laryngol; 94: 39-43.

Forhman LA, Szabo M, Belerowitz M, et al (1981):

Partial purification and characterisation of a peptide with growth hormone-releasing activity from extra-pituitary tumours in patients with acromegally. J clin Invest; 65 : 43 - 48 .

Frable MA. (1982):

Fine - needle aspiration biopsy revisited.

Laryngoscope; 92 : 1414-1418.

Fraumini J F. (1988):

Epidemiology of cancer, in Cesil textbook of medicine. by : Wyngarden JB, Smith L H (eds), Published by W.B. Saunders Co. : 1189.

Gadwin JD. (1957).

Carcinoid tumours:an analysis of 2837 cases. cancer; 36: 560- 569 .

Galliani P, Mariligit G, Hersh E. (1976):

Onchogenic DNA viruses: replication, tumour gene expression, and role in human cancer.

Cancer Res; 35 : 58-62.

Gatenby R, Mulhern CB, Strwatz J. (1988):

CT- Guided percutaneous biopsies of head and neck masses.

Radiology; 146 : 717 - 719.

Ghatei Mohammed, David R, Nicoll G, et al (1985):

Gastrin - releasing peptid -like immunoreactivity in medullary thyroid carcinoma.

Am J clin pathol; 84: 581 - 586.

- Giller M, Todaro P, Holmus S, (1978):
The role of immunotherapy in management of patients with malignant disease.
J surg; 3 : 3309 -312.
- Goel G, Kohli S, Lal H (1986):
Serum phosphohexose isomerase levels in patients with head and neck cancer.
J laryngol Otol; 100 : 581 - 585.
- Goyal A, shah G, Karelia H, (1987).
Significance of ferritin as a marker in head and neck malignancies.
Tumori; 73 : 59 -63.
- Grander D, Rothman M, Utiger F, (1979):
Serum thyroglobulin in normal subjects and patients with hyperthyroidism due to graves disease.
Clin Endocrinol; 11 : 585- 589.
- Greace A, vaughn K, Hainsworth D, (1988):
Advanced poorly differentiated carcinoma of unknown primary site: Recognition of a treatable syndrome.
Arch Intern Med; 104: 547- 555.
- Grey M (ed), (1985):
Cancer risks and prevention.
New York, Oxford University Press: 401 - 408.
- Griffith M, Beck C, (1963):
The value of serum phosphohexose isomerase as an index of metastatic breast carcinoma activity.
Cancer, 16: 1032-1041.
- Hodfield T R, Lange S, Wold R, (1970):
Serum induced lymphoid cell-mediated cytotoxicity against human transitional cell carcinomas of urinary tract.
J Natl Cancer Inst; 40 : 231- 237.
- Hamads S, Ishakawa N, Wada T, (1976):
Association of medullary carcinoma of the thyroid with carcinoembryonic antigen.
B, J Cancer; 34:111 -115.

- Hamada S. Hamads S, (1977):
Localisation of carcinoembryonic antigen in medullary
thyroid carcinoma by immunofluorescent techniques.
B.J cancer; 36:572-576.
- Hann H, levy M, Evans A, (1980):
Serum ferritin as aguide to therapy in neuroblastoma.
Cancer Res; 40: 1411-1413.
- Harbans L, Singh B, Wig U, et al (1988):
Serum immunoglobulin E levels in patients with head
and neck cancer.
J laryngol Otol; 102-434.
- Harwood A R, (1982):
Cancer of the larynx - the Toronto experience.
J Otolaryngol II, Supplement II; 1 - 221.
- Hattori M, Fukase M, yoshimi H et al (1978):
Ectopic production of human chorionic gonadotropin in
malignant tumours.
Cancer; 42: 2338-2333.
- Hazard T, Drysdale W, Alpert E, (1977):
Ferritinemia in cancer.
Nature; 265: 755-756.
- Herberman R. B. (1983):
Uses and Limitation of tumour markers. In:
Fishman witt, ed: Onchodevelopmental markers.
New York: Academic press: 409-418.
- Hiratuska H, Imamura M, Ishii Y, (1984):
Immunohistologic detection of lymphocyte subpopulations
infiltrating in human oral cancer with special
refrence to its clinical significance.
Cancer; 53: 2456-2466.
- Hirsch B, Jonson J. T, Rabin B.S, (1988):
Immunostimulation of patients with head and neck
cancer.
Arch Otolaryngol; 107 : 658-663.

Hueper R. (1954):
Structural representations of tumour-host
relationship in gastric carcinoma,
Surg Gynecol Obstet; 102: 599-605.

Hyamas, J, Langloss M, Dennis K, (1987):
Spindle-cell carcinoma of the aerodigestive tract.
Am J surg pathol; (5): 339-349.

Hyderman S, Bekesi R, Mink M, (1979):
Immunologic response in patients with disseminate
cancer.
Ann Surg; 168: 347- 357.

Irie R, Irie K, Morton L, (1976):
A membrane antigen common to human cancer and fetal
brain tissues.
Cancer Res; 36: 3510-3315.

Isaacson P, Jones D, B., Judd MA, (1979):
Alpha - 1 antitrypsin in human macrophages.
lancet; 8149: 964-5.

Ishikawa N, Hamada S. (1976):
Association of medullary carcinoma of the thyroid
with CEA.
Br J Cancer; 34 :11 -115.

Ishii K, Green H, (1973):
Lethality and adenosine for cultured mammalian cells
by interference with pyrimidine biosynthesis.
J Cell Sci: 13 : 429-439.

Jaffe E. A. (1977):
Endothelial cells and the biology of factor VIII.
N Engl J Med; 296: 377-383.

Toplin G, Jagatheesan A, (1962):
Serum glycolytic enzymes and acid phosphatase in
mammary carcinomatosis.
Brit Med J; 7 : 827-831.

- Khan J. Ibanez M. Luna A. (1986):
Report of one hundred tumours of the minor salivary glands.
Am J Surg; 112., 493-497.
- Kameya T, Besshot T. Tsumuraya M, et al. (1983):
Peptide hormone production by adenocarcinoma of the lung; its morphologic basis and histogenetic considerations.
Virchows Arch; 400: 245-257.
- Karja J, Syrjanen K, Nuutinen J. (1987):
Immuno - competent cell reaction in prognostic evaluation of squamous cell carcinoma of the lip.
Acta Otolaryngol (Stockh); 103: 640-643.
- Katopodis N. Hirshaut Y, Geller N. (1982):
Lipid associated sialic acid test for the detection of human cancer.
Cancer Res; 42: 5270-5.
- Kersrud F, Brandtzage P, Juap T. (1982):
Characterisation of epithelial elements in human major salivary glands by functional markers.
Histochemist; 66: 285 - 292.
- Khan T, Schwere E, Hanser Z. (1983):
Molecular cloning and characterisation of the DNA of a new human Papilloma Virus from a laryngeal carcinoma.
Internat J Cancer; 37: 61-65.
- Kindblom LG, Jacobsen B, Jacobsen M, (1982):
Immunohistochemical investigations of tumours of supposed fibroblastic -histocytic origin.
Hum pathol; 13 : 834-40.
- Kodama M, Fletchers S. Kermaen A J (1980):
The incidence of distant metastases among patients dying with head neck cancers.
J Surg; 30 : 827-833.

- Kohler G. Benz R. (1962).
Immunological enhancement of tomografts in mice:
a review.
Cancer REs; 18: 992 -996
- Knowar S, oldini R, kasim M. (1986)
the place of radiotherapy in the treatment of head
and neck cancer patients .
j laryngol Otol ; 84 : 867- 896 .
- Koroki M, Koga y, MastouKa y. (1983)
Antigenic reactivites of purified preparations of
cancinoembryonic antigen and normal antigens using
four different radioimmunoassay systems for CEA .
j Immunol Methods ; 60 : 221 _ 233 .
- Kuwano H, hashimoto H. (1985)
Atypical fibroxanthoma distingusishable from spindle
cell carcinoma in sarcoma-like skin lesions : a
clinicopatholgic and immuohistochemical study of
21 cases .
Cancer ; 55 : 172 _ 80 .
- Lal H, Madan G, kohli S, et al , (1987)
Serum enzymes in head neck cancer ll . aliestrase .
j laryngol Otol ; 101 : 819 _ 822 .
- Laszlo j. (1988)
Introduction to oncology , in : Cisil textbook of
medicine by wyngaarden j , Smith LH. (eds) Published
by W . B .
Saunders Co. : 1082.
- Lippman M, wells S, Baylin B et al (1982)
The prognostic signifigance of cellular heterogeneity
in medullry thyroid carcinoma .
j Clin Endocrinol Metab ; 54 : 233 _ 239 .
- Luger T A , Linkish W , knobler R ,(1983)
Serial determination of ferritin levels in patients
with malignant melanoma .
Oncology ; 40 : 263 _ 267 .

- Macario M, poggio E, prat M, (1988)
An immunohistochemical and clinicopathological investigation of laryngeal carcinomas .
j Laryngol , Otol ; 102 : 709 _ 712 .
- Mafee M F , (1982)
Dynamic C T and its application to Oto laryngology: head and neck surgery.
j Oto laryngol ; 11 : 307 _ 318 .
- Maier H, Adler D, Born A, (1987)
S_ phase fractions and D N A _ ploidy of head and neck squamous cell carcinoma .
Acta Otolaryngol (Stokh); 104 : 385 _ 387 .
- Maity R , Burma P , (1973)
Metabolic changes in breast carcinoma 11. phosphohexose isomerase and lactic dehydrogenase level in serum .
Ind j cancer ; 10 : 356 _ 329 .
- Mash j P, jeager P, loosli R, (1974)
Detection of recurrence of large bowel carcinoma by radio immunoassay of circulating carcinoembryonic antigen.
Lancet ; 2 : 535 _ 539 .
- Martino G , jannocci F, jacono G . (1982)
Ferritin and calcium in colo-rectal and lung cancer correlation with other types of neoplasms and non _neoplastic diseases .
Tumor diagnostic Ther ; 3 : 74 _78 .
- Matsunaga A, kuroi M, Higushi H, (1987)
Antigenic heterogeneity of carcinoembryonic antigen in the circulation .
cancer Res ; 47 : 56 _ 61 .
- Maxim G , veltri B, watne A, (1980)
serum ferritin as a tumour marker for patients with carcinoma of the head and neck (Abstr).
proc Am Assoc Cancer Res ; 21 : 233 _ 237 .

- Maxim E , Robert W veltri B, (1986)
serum ferritin as a tumour marker in patients with
squamous cell carcinoma of the head neck .
Cancer ; 57 : 305 _ 311 .
- Maxim E. Veltri W, (1986)
serum ferritin as a tumour in patients with
Squamous cell carcinoma of the head and neck .
cancer ; 57 : 305 _ 311 .
- Mia O j, Russel D , liu z , (1984)
polymaine in the urine of patients with malignant
tumours.
j Oncol ; 6 : 269 _ 273 .
- Miradith A, Randall C, Shaw j , (1987) .
Towards consistent local control of certain malignant
tumours.
j Laryngol Otol ; 101 : 1046 _1054 .
- Moll R, Franke WW, Schiller DL, (1982) .
The catalog of human cytokeratins: patterns of expre_
ssion in normal epithelia, tumours and cultured cells.
cell 31 ; 11 : 24 .
- Morton D, Samuel A, wells j, (1981) .
Immunobiology and immunotherapy of neoplastic
diseases.
Cancer Detect prevent ; 4 : 577 _ 597 .
- Muller C. (1969):
Prognosis of early metastases in bladder carcinomas.
Brit J Surg; 56: 625- 630.
- Murphy G. Flynn T. Rich G, (1984):
Involucrin expression in normal and neoplastic human
skin : A marker for keratinocyte differentiation.
J invest Dermatol; 82 : 453-457.
- Nakia M, Mori M, (1986):
Immunohistochemical distribution of monoclonal
antibodies against Keratin in papillomas and
carcinomas from oral and nasopharyngeal regions.
Oral surg oral med oral pathol; 62: 292-302.

- Nakazata A, Eveson J, Cawson R. (1982):
Tumours of the minor (oropharyngeal) salivary glands:
a dermographic study of 336 cases
J Oral Pathol; 14: 500-509.
- Nichols E, Kannagi R, Fuks A. (1985):
Carbohydrate determinants associated with
carcinoembryonic antigen.
J Immunol; 135 : 1911-1913.
- Nishahara H, Akedo H, Halter S. (1970):
Multienzyme patterns of serum adenosine deaminase by
agar gel electrophoresis: an evaluation of the
diagnostic value in lung cancer.
Clinica Chimica Acta; 30: 251-258.
- Nurula A, Morgan L, Padfield J, (1988):
NED survival in head and neck cancer: Post-mortem
correlations.
J laryngol Otol; 102: 194-197.
- Nuutinen J, Karja J, Kohama G I, (1987):
Local and distant metastases in patients with
surgically treated squamous cell carcinoma of the
lip.
Clin Otolaryngol; 6 : 415-419.
- O'Dorisio M, Falko M, Cataland S, et al (1982):
Gastrin responses to a test meal in patients with
familial medullary thyroid carcinomas.
J clin Endocrinol Metab; 54: 798-802.
- Okumara K, Toda T. (1971):
Regulation of homocytotropic antibody formation in
the rat:
Effect of Thymectomy and splenectomy.
J Immunol; 106 : 1091-1025.
- Pacini F, Gianni B, Taddei D. (1985):
Humoral markers for thyroid carcinoma.
Cancer detect prevent; 8 17-22.

ParattM, Spinnato G, Comoglio P, (1985):
Monoclonal antibodies against the human epidermoid carcinoma A 431.
Cancer Detect Prevent; 8 169-179.

Piovella F, Nalli G, Malamani G, et al (1987):
The ultrastructural localisation of factor VIII-
antigen in human platelets, megakaryocytes and endo-
thelial cells utilising a ferritin labelled antibody.
B, J haematol; 39: 209-213.

Pinoso L, Pilizzo R, Zorat P, (1972):
Different diagnostic significance of carcinoembryonic
antigen.
Acta Endocrinol (Suppl 252) 102: 57 - 59.

Pluygers P, Beauduin P, Baldwins E. (1986):
Tumour markekr s for cancer detection 1,
Cancer Detect Prevent 9; 495-504

Plumegart p ,P Sparks R, Spears K, (1971)
Relationship between neutrailisation of Epstein
Bart virus and antibodies to cell membrane antigens
induced by the virus .
j Natl Cancer Inst ; 45 : 989 - 993 .

poste G , Tzeng V, Doll j et al (1982)
Evalution of tumour cell heterogeneity during
progressive growth of individual lung metastases.
Proc Natl Acad Sci USA; 79: 6574-6578.

Prat M, Spinnuto R, Comoglio M, (1985):
Monoclonal antibodies against the human epidermoid
carcinoma A 431.
Cancer Detect prevent; 8 : 169-179.

Proctor M, Wallace E, (1988):
Head and neck oncology service -one year's work.
J laryngol Otol; 102: 428-431.

- Raczynka S. (1966):
Chemical immunosuppression and human cancer.
Cancer. 34: 1474-1479.
- Rashidi Ali, Varndell M, Facer P, et al, (1983):
Immunocytochemical localisation of Ketacalcin.
J Clin Endocrinol Metab: 57: 680-682.
- Raju G, Kakar P, Das K, et al (1988):
Role of fine needle aspiration biopsy in head and neck tumours.
J Laryngol Otol; 102 : 248-251.
- Regazzi M, Luna M, Lowery A. (1981):
Minor Salivary gland tumours: an aid to diagnosis
Am J Surg; 12: 516-522.
- Richer R, Begad R, Raine L, (1984):
Recent advances in tumour imaging. use of radiolabelled antitumour antibodies.
J Histochemist cytochemist; 29: 577 -580.
- Robertson GL, Bert T, Foster B, (1986):
Water metabolism - in Brenner B M, Rector F (eds):
the Kidney. 3 rd ed. W. B. Saunders Co.
Philadilphia: 385- 432.
- Robson RS, (1963):
Immune RNA therapy for renal cell carcinoma: survival and immunological monitoring.
Ann Surg; 184: 49 - 55
- Rogier H, Clamettes C, Laplanche A. et al. (1983):
The values of calcitonin and carcinoembryonic antigen in the treatment and management of non-familial medullary Thyroid carcinoma.
Cancer; 51; 855-862.
- Rosen S , Maggia F, Hershman M, et al (1975):
Placental proteins and their subunits as tumour markers
Ann, Intern Med; 82 : 71-83.

- Ruchiti C, Heinz A, Greber D, (1984):
Factor VIII-related antigen in malignant hemangioendothelioma of the thyroid: additional evidence for the endothelial origin of this tumour.
Am J Clin Pathol; 82: 474-480.
- Saad M F, Fritche A, Samaan N, (1984):
Diagnostic and prognostic values of carcinoembryonic antigen in medullary carcinoma of the thyroid.
J Clin Endocrinol Metab; 58: 889-894.
- Said W, Nash G, Sasson A F. (1983):
Involucrin in lung tumours: A specific marker for squamous differentiation.
Lab invest; 49: 563-568.
- Said W, Sasson A F, Shintaku P, (1984):
Involucrin in squamous and basal cell carcinomas of the skin: An immunohistochemical study.
J Invest Dermatol; 82: 449-452.
- Samaan N, Friesen H, Pearson H. (1973):
Serum placental lactogen levels during pregnancy and in trophoblastic disease.
J Clin Endocrinol; 26: 1303-1308.
- Schwartz M, Walsh S, West M, (1962):
Serum enzymes in disease x: glycolytic and oxidative enzymes and transaminases in patients with carcinoma of head, neck and oesophagus.
Cancer; 15 :922 - 930.
- Seifert G, Caselitz J. (1985):
Intermediate sized filament proteins (prekeratin, vimentin, desmin) in the normal parotid gland and parotid gland tumours.
Virchows Arch (Pathol Anat): 393:273-276.
- Shaw H J, (1988):
Surgical Salvage for squamous cancer involving the supraglottic larynx.
J laryngol Otol; 102: 704-708.

- Sidell N, Irie R, Morton D. (1979):
Oncofetal antigen 1: A target immune cytotoxic human
cancer.
Br J cancer; 40: 950-953.
- Sidell N, Morton D. (1979):
Immune cytotoxicity of human malignant melanoma by
antibody to oncofetal antigen 1. Complement
dependent cytotoxicity.
Cancer Immunol Immunother; 7: 151.
- Sikri K, Hamid A, Polak M, et al. (1985):
Medullary carcinoma of the thyroid: an
immunocytochemical and histochemical study of 25
cases using eight separate markers.
Cancer; 56 : 22481-2491.
- Silverman A, Alexander C, Chretien B. (1976):
CEA levels in head and neck cancer.
Cancer; 37: 2204-2241.
- Sloare M, Ormardo P. (1981):
Geographic pathology of thyroid disease.
Pathol Res Pract; 171 : 285-292.
- Spratt J. (1965):
Garcinoma of the uterus; an aid to early diagnosis.
Surg Clin N Amer; 45: 1103- 1113.
- Stell D, McCormik N. (1985):
Cancer of the head and neck: Are we doing any better?
Lancet; 2: 11127-1131.
- Stepman P, Robertson M, Christ D. (1979):
The vascular endothelium -pathobiologic significance,
a review.
- Sufrin G, Tritsch G, Krawznki J. (1978):
Adenosine deaminase activity in patients with
carcinoma of the bladder.
J Urol; 119: 343-346.

- Sun TT, Green H, (1978):
Keratin filaments of cultured human epidermal cells.
Formation of intermolecular disulfide bonds during
terminal differentiation.
J Biochem; 253: 2053- 2060
- Syrjanen K J. (1975):
Morphologic manifestations of tumour -host
relationships in association with breast, astric and
colorectal cavcinoma.
M.D. Thesis, Universty of Helsinki; 1- 86.
- Syrjanen K J, Kajara J, Lund D. (1987):
Epidermoid carcinoma of the lip. Histologic grading
in the clinical evaluation.
Acta Otolaryngol (Stockh); 103: 640-644.
- Tan S. Chen Q, Fan F, (1986):
Biochemical markers of tumour diathesis and early
cancer
Cancer detect prevent; 9: 145-149.
- Tarrot P., Shaw H J, (1987):
Partial laryngectomy for recurrent cancer after
irridiation.
Clin Otolaryngol; 3: 49-62.
- Taxy JB, Hidreg D F, Battifera H, (1983):
Nasopharyneal carcinoma: antikeratin immunohistochem-
istry and electron microscopy.
Am J clin pathol; 83: 320 - 325.
- Terry R, Shaw H J, Grey C, (1986):
Expression of low molecular weight cytokeratins in
the neoplastic vocal cord.
J laryngol Otol; 100: 1279-1282.
- Terry R M, Grey C, Bird C. C, (1986 b):
Aberrent expression of low molecular weight
cytokeratins in primary and secondary squamous cell
carcinoma of the head and neck.
J Laryngol Otol; 100: 1283 -1287.

- Tsutsumi Y, Osamura R, Watanabe K, et al. (1983):
Simultaneous immunohistochemical localisation of
gastrin releasing peptide and calcitonin in human
bronchial endocrine-type cells.
Virchows Arch; 400 : 163-171.
- Vaid P, Sharma D, Shastri K. (1974).
A clinicobiochemical study of phosphohexose
isomerase, alisterase, and sulfhydryl group in 25
normal and 35 Cancer suspects.
Ind J cancer; 11 : 448 - 456.
- Veltri R, Mengoli F, Maxim P. (1977):
Isolation and identification of human lung tumour
associated antigens.
Cancer Res; 37 : 1313 - 1322.
- Vilkano R, Brennan S, Brock P. (1983):
Radioimmunoassay for a monoclonal antibody -
defined tumour marker C A 19-9.
Clin chem. 29: 549-2.
- Voller A, Bedwell D, Brattlet A. (1976):
Enzyme immunoassays in diagnostic medicine.
Bull W H O, 53 : 55-56.
- Volm M, Snoka J, Wayss D. (1985):
D N A distribution in non-small-cell-lung carcinomas
its relationship to clinical behaviour.
Cytometry; 6 : 348 356.
- Wagener C, Breuer H. (1978):
Comparative studies on the radioimmunological
determination of carcinoembryonic antigen in tumour
tissue.
J Clin chem clin Biochem; 16: 323-327.
- Waldman T, Bull M, Jost M. (1974):
Serum immunoglobulin E levels in patients with
neoplastic disease.
J Immunol; 113: 379-386.

Walf J, Melvin K, Alzaadi A, A, et al, (1979):
C-cell hyperplasia preceding medullary thyroid carcinoma.
J Clin Pathol; 19: 114-117.

Walterl F, Israel M.S. (1974):
Introduction to oncology, Text book of general pathology, Edited by Churchill Livingstone Edinburgh and London .
Published by Longham Group Limited.

Watanabe T, Cox B, Hudson R. (1983):
Elevated serum secretory immunoglobulin A levels in patients with head and neck carcinoma.
Otolaryngol Head Neck Surg; 91: 136- 142.

Watne L. (1984):
Treatment alternatives: head and neck cancers
Cancer; 54: 2673-2678.

Wells S A, Meleweiz F, Ketcham A, (1978):
Delayed cutaneous hypersensitivity reactions to extracts of carcinomatous cells of the cervix uteri.
Surg Gynacol Obstet; 136: 717-721.

Weilant L, Pearson G R, Neil H B, (1981):
Elevation of the antibody responses to E.B.V. associated antigens as diagnostic markers in nasopharyngeal carcinoma in American populations.
Head Neck Surg; 3 : 258-260.

White M; (1960) :
Lymphocyte response in patients with head and neck cancers
Aarchi Otolaryngol 102 : 596 - 600

Wolf P L (1979):
Tumour associated markers: the importance of identification in clinical medicine.
New York: Masson.

Wig U, Sani A, Gupta V, (1983):
E-rosette forming cells in squamous cell carcinoma of
the larynx and laryngopharynx.
J Laryngol Otol; 97: 527-530.

Wilson B. S, (1960):
Detection of chromogranin in neuroendocrine cells
with a monoclonal antibody.
Am J path; 120: 186-190.

Yamamoto E, Miakawa A, Kohama G I, (1984):
Mode of invasion and lymph node metastases in
squamous cell carcinoma of the oral cavity.
Head Neck Surg; 6 : 938-947.

Yarde D E, Husa R, Garancis C, (1979):
Immunocytochemical localisation of human
choriogonadotropin in human malignant trophoblast,
Model for human choriogonadotropin secretion.
Lab invest; 40 391-398.