

REFERENCES

Adachi, K.; Chole, R. and Yee, J. (1991):

Indomethacin inhibition of middle ear bone resorption.

Arch otolaryngol head and Neck Surg; 117 : 267 -9.

Albert, D.H. and Snyder, F. (1983):

Biosynthesis of 1- alkyl - 2 - acetyl - sn - glycerol - 3 - phosphocholine (platelet activating factor).

J. Biol. Chem.; 258 : 97 - 102.

Albiin, N. (1984):

The anatomy of the eustachian tube.

Acta Otolaryngol (Stockh) Suppl. 414 : 34-37.

Albiin, N.; Hellstrom, S.; Salen, B. and Stenfors, L.E. (1983):

The stapedial artery in the rat a - microscopic study under normal conditions and in otitis media with effusion (OME).

Acta Anat.; 115 : 134 - 140.

Albiin, N.; Hellostron, S.; Salen, B.; Stenfors, L.E. and Soderberg, O. (1983):

The anatomy of the eustachian tube in rat : Amacro and microscopical study.

Anat. Rec., 207 : 513 - 521.

Albiin, N.; Hellstrom, S.; Salén B., Stenfors, L.E. and Wirell, S. (1985):

The vascular supply of the rat tympanic membrane.

Anat. Rec., 212 : 17 - 22.

Albiin, N.; Hellstrom, S.; Stenfors, L.E. and Crene, A. (1986):

Middle ear mucosa in rats and human.

Annals of Otology, Rhimology and laryngology.

Suppl. 125 - 95 : 2-15.

Anson, B.J. and Donoldson, J.A. (1981) :

Surgical anatomy of the tomporal bone. 3rd edn.

Philadelphia : W.B. Saunders.

Arnold, W. and Vosteen, K.H. (1975):

Die Reaktion der Millelohrschleimhaut ber tubenverschless.

Acta otolaryngol (stockh), Suppl. 330, 48.

Arnoux, B.; Durand, J.; Rigaud, M.; Vargaftig, B.B. and Benveniste, J. (1981):

Release of platelet activating factor (PAF-acether) and arachidonic acid metabolites from alveolar macrophages.

Agents Actions, 11 : 555-556.

Aursudkij, B.; Roger, D.F.; Evans, T.W.; Alton, EFWF, Chung, K.F.; Barnes, P.J. (1987):

Reduced tracheal mucus velocity in guinea pig in Vivo by platelet - activating factor.

Am. Rev. Respir. Dis.; 135 : 160.

Barbaro, J.F. and Zweifler, N.J. (1966):

Antigen-induced histamine release from platelets and rabbits producing homologous PCA antibody.

Proc. Soc. Exp. Biol. Med.; 122 : 1245-1247.

Barnes, P.; Chung, K. and Page, C. (1988):

Platelet - activating factor as a mediator of allergic disease.

J. Allergy Clin. Immunol., 81 : 919 - 34.

Barnes, P.J.; K.F. Chung and C.P. Page (1988):

Inflammatory mediators and asthma.

Pharmacol. Rev., 40 : 49.

Beery, Q.C. (1979):

Longitudinal assessment of eustachian tube function in children.

Laryngoscope, 89 : 1446.

Beery, Q.C.; Bluestone, C.D. and Cantekin, E.I. (1975):

Otologic history, audiometry and tympanometry as a case finding procedure for school screening.

Laryngoscope, 85 : 1976.

Benveniste, J.; Chignard, M.; Le Couedic, J.P. and Vargaftig, B.B. (1982):

Biosynthesis of platelet-activating factor (PAF-acether) II. Involvement of phospholipase A2 in the formation of PAF-acether and lyso-PAF-acether from rabbit platelets.

Throm. Res.; 25 : 375-385.

Benveniste, J.; Le Couedic J.P.; Polenosky, J.; Tence, M. (1977):

Structural analysis of purified platelet activating factor by lipase.

Nature, 269 : 170-171.

Bluestone, C.D. and Contekin, E.I. (1981) :

Panel on experiences with testing eustachian tube function.

Ann. Otol. Rhinol. laryngol, 90 : 552 -561.

Bluestone, C.D. and Doyle, W.J. (1985) :

Eustachian tube function.

Annals, Otol. Rhinol. laryngol. Suppl., (94) : 1-60.

Bluestone, C.D. and Klein (1983):

OME, Atelectasis and Eustachian dysfunction.

Ped. Otolaryngology, Philadelphia, W.B. Saund. Company, pp : 356 -512.

Camusi, G.; Agletta, M.; Malavasi, F.; Tetta, C.; Placebello, W.; Sanavia, F. and Bussolino, F. (1983):

The release of platelet activating factor from human endothelial cells in culture.

J. Immunol.; 131 : 2397 - 2403.

Camusi, G.; Tetta, C.; Coda, R.; Benveniste, J. (1981):

Release of platelet-activating factor in human pathology ; evidence for the occurrence of basophil degranulation and PAF release in systemic lupus erythematosus(SLE).

Lab. Invest. 44 : 241 - 251.

Cantekin, E.I. (1980):

Gas absorption in the middle ear.

Ann. Otol. Rhinol. Laryngol., 89 : 71.

Cauwenberge, P. and Bernstein, J. (1987):

Inflammatory mediators in middle ear disease.

In : Bernstein J., Orga P., eds, Immunology of the ear.

New York, NY : Raven Press : 338 -9.

Columbo, M.; Casolaro, V.; Warner, J.; Mac Glashan, D.;

Kagey Sobotka, A. and Lichtenstein, L. (1990):

The mechanism of mediator release from human basophils induced by platelet activating factor.

J.Immunol.; 145 : 3855-3861.

Cuss, F.M.; Dixon, C.M.S. and Barnes, P.J. (1986):

Effect of inhaled platelet activating factor on pulmonary function and bronchial responsiveness in man.

Lancet, 2 : 189-192.

Davis, W.B.; G.A. Fells, X.H. Wun, J.E. Fradek, A. Venet and R.G. Crystal (1984):

Eosinophil -mediated injury to lung paranchymal cells and inerstitial matrix a possible role of eosinophils in chronic inflammatory disorders of the lower respiratory tract.

J. Clin. Invest., 74 : 269.

De-Maria, T.B.; Briggs, B.R. and Lim, D.J. (1981):

Experimental OME following middle ear inoculation
of non-viable hemophilus influenzae.

Laryngoscope : 91 : 279 - 280.

***Dulfano, M.J.; Luc, C.K.; Beckas, M. and Wooten, B.
(1986):***

Ciliary inhibitory effect of asthma patients sputum.

Clinic. Sc., 63 : 393 - 396.

Elner, A. (1977) :

Quantitative studies of gas absorption from the
normal middle ear.

Acta Otolaryngol (Stockh); 83 : 25.

***Evans, T.W.; Chung, K.; Rogers, D.F. and Barnes, P.J.
(1987):***

Effect of platelet activating factor on airway vascular
permeability : possible mechanisms.

Appl. Physiol., 63 : 479 - 484.

Fawcett, D.N. (1994):

Bloom and Faweett, A textbook of Histology,
Chapman and Hall.
New York, London.

Frigas E. and G.J. Gleich (1986):

The eosinophil and the pathogenesis of asthma.
J. Allergy Clin. Immunol., 77 : 527.

Furukawa, M. Horii, T. and Oda, M.(1992):

PAF in Human Respiratory Disease.
J. Lipid Mediat., 5 : 191-192.

Furukawa, M.; Kubo, N. and Yamashita, T. (1995):

Biochemical evidence of platelet activating factor
(PAF) in human middle ear effusions.
Laryngoscope, 105 : 188 - 91.

Gacek, R.R. (1983):

Cupulolithiasis and posterior ampullary nerve
transection.
Annals of Otology, Rhinology and laryngology,
Suppl., 112 : 25 - 30.

Ganbo, T. and Hisamatsu, K. (1990):

Mucosal dysfunction and damage induced by platelet activating factor (PAF).

Acta Otolaryngol. (Stockh), 10 : 427-436.

Ganbo, T.; Hisamatsu, K.; Kikoushima, K.; Nakajima, M.;

Phar, H.I.; Murakami, Y. (1996):

Effect of platelet activating factor on mucociliary clearance of eustachian tube.

Ann. Otol. Rhinol. Laryngol. 105 : 140 - 145.

Gateau, O.; Arnoux, B.; Deriaz, H.; Viars, P. Benvenise, J.

(1984):

Acute effects of intrabracheal administration of PAF-acether (platelet activating factor) in humans.

Am. Rev. Respir. Dis., 133 : 129.

Giebink, G.S.; Juhn, S.K. and Weber, M.L. (1982):

The bacteriology and cytology of chronic otitis media with effusion.

Pediatr. Infect. Dis., 1 : 98-103.

Godfroid, J.J.; Heymans, F.; Michel, E.; Redeuith, C.;

Steiner, E. and Benveniste, J. (1980):

Platelet-activating factor (PAF-acether): total synthesis of 1-O-Octadecyl 2-O-acetyl sn-glycero-3-phosphorylcholine. FEBS Lett; 116 : 161 -4.

Gundersen, T. and Tonning, F.M. (1976):

Ventillating tubes in the middle ear.

Arch. otolaryngol., 102 : 198.

***Halonen, M.; Palmer, J.D.; Lohman, I.C.; McManus, L.M.
and Pinchard, R.N. (1980):***

Respiratory and circulatory alterations induced by
acetyl glyceryl ether phosphory choline, a mediator of
IgE anaphylaxis in the rabbit.

Am. Rev. Respir. Dis., 122 : 915-924.

***Hartung, H.P.; Parnham, M.J.; Winkleman, J.;
Engleberger, W. and Hadding, U. (1983):***

Platelet activating factor (PAF) induces the oxidiative
burst in macrophages.

Int. J. Immunopharmacol., 5 : 115 - 121.

Hebel, R.; Stromberg, M.W. (1976):

Anatomy of the laboratory rat.

PP. 55-57 and 145 - 149 edited by the Williams and

Wilkins Company, Baltimore.

Henocq, E.; Vargaftig, B.B. (1986):

Accumulation of eosinophils in response to
intracutaneous PAF- acether and allergen in man.

Lancet, 1 : 1378 - 1379.

Hentzer, E. (1969) :

Ultrastructure of the human tympanic membrane.

Acta. otolaryngologica, 63 : 376-390.

Hentzer, E. (1970) :

Ultrastructure of the normal mucosa in the human middle ear, mastoid cavities and Eustachian tube.

Ann Otol Rhinol Laryngol, 19 : 1143-1157.

Hentzer, E. (1972):

Ultrastructure of the middle ear mucosa in secretory otitis media (1) serous effusion.

Acta. Otolaryng, 73 : 394 - 401.

Holmquist, J. and Renvall, V.C. (1974):

Eustachian tube function in secretory otitis media

Arch. Otolaryngol. 99 : 59 - 61.

Honjo, I. (1985):

Clearance function of the Eustachian tube.

Ann. Otol. Rhinol. Laryngol., 94 (suppl), 120:29 - 30.

Honjo, I.; Ushiro, K.; Okazaki, N. and Kumazawa, T. (1981):

Evaluation of Eustachian tube function by contrast roentgenography.

Arch. Otolaryngol., 107 : 350 - 352.

Humphrey, J.H. and Jacques, R. (1954):

The histamine and serotonin contents of the platelets
and polymorphnuclear leucocytes of various species.

J. Physiol., 124 305 -10.

Humphrey, D.M.; McManus, L.M.; Satanchi, K.; Hanahan,

D.J. and Pinckard, R.N. (1982):

Vasoactive properties of acetylglycerol ether
phosphorylcholine and analogues.

Lab. Invest., 46 : 422 - 427.

Hussl, B. and Lim, D.J. (1974):

Experimental middle ear effusion on
immunoflourescent study.

Ann. Otol. Rhinol Laryngol., 83 : 332 - 342.

Hwang, S.B.; Lam, M.H. and Shen, T.Y. (1985):

Specific binding sites for platelet activating factor in
human lung tissue.

Biochem. Biophys. Res. Commun., 128 : 972 - 979.

Ingelstedt, S.; Ivarsson, A. and Jonson, B. (1967):

Quantitative determination of tubal ventilation during
changes in ambient pressure as during ascent and
descent in aviation,

Acta Otolaryngol (Stockh) [suppl] 228 : 31.

Jouvin-Marche, E.; Cerrina, J.; Coeffler, E.; Durous, P. and Benveniste, J. (1983):

Effect of the Ca^{++} antagonist nifedepine on the release of platelet activating factor (PAF-acether), slow reacting substances(SRS), and beta-glucoronidase from human neutrophils.

Eur. J. Pharmacol., 89 : 19-26.

Juhn, S.K. (1982):

Studies on middle ear effusion.

Laryngoscope, 92 : 287 - 291.

Jung, J. (1988):

Prostaglandins, leukotrienes, and other arachidonic acid metabolites in the pathogenesis of otitis media.

Laryngoscope, 98 : 980 - 993.

Jung, T.T.K.(1988):

Prostaglandines, leukotrienes and other arachidonic acid metabolites in pathogenesis of otitis media.

Laryngoscope., 98 : 980 - 93.

Jung, T.T.K.; Park, Y.M. and Schlund, D. (1990):

Effect of prostaglandin, leukotriene, and arachidonic acid on experimental otitis media with effusion in chinchillas.

Ann. Otol, Rhinol laryngol., 99 (suppl, 148) : 28 - 32.

Jung, T. and Rhee, C. (1991):

Otolaryngologic approach to the diagnosis and management of otitis media.

Otolaryngology Clinic North Am., 24 : 931 - 45.

Jung, T.T.K.; Shea, D.; Giebink, G.S. and Juhn, S.K. (1984):

Comparative histopathology of animal models of otitis media in chinchillas; a temporal bone study.

In : Lim DJ, Blue stone CD, Klei, JO, Nelson, JD, ed., Recent advances in otitis media with effusion.

Philadelphia, Pa : BC Decker : 91-6.

Kokko, E. (1974):

Chronic secretory otitis media in children.

Acta. otolaryngol (stockh), suppl., 327.

Lambrecht, G. and Parnham, M.J. (1986):

Kad surenone distinguishes between different platelet activating factor receptor Subtypes on macrophages and polymorphonuclear leucocytes.

Br. J. Pharmacol., 87 : 287 - 299.

Lamp, C.B. (1973):

Chronic secretory otitis media. Etiologic factors and pathologic mechanisms.

Laryngoscope, 83 : 276 - 291.

Larsen, P.L. and Tos, M. (1991) :

Polyp formation after experimental tubal obstruction
in Rat. Acta Otolaryngol (Stoka). 111 :926 - 933.

**Lee, T.C.; Lenihan, D.J.; Malone, B.; Roddy, L.L. and
Wasserman, S.I. (1984):**

Increased biosynthesis of platelet activating factor in
activated human eosinophils.
J. Biol. Chem., 259 : 5526 - 5530.

**Lellouch - Tubiana, A.; Lefort, J.; Pirotzky, E.; Vargaftig,
B.B. and Pfister, A. (1985):**

Ultrastructural evidence for extravascular platelet
receivement in the lung upon intravenous injection of
platelet activating factor (Paf-acether) to guinea pig.
Br. J. Exp. Pathol., 66 : 345 - 355.

**Levi, R.; Burke, J.A.; Guo, Z.G.; Hattori, Y.; Hoppens, C.M.;
McManus, L.M.; Hanahan, D.J. and Pinckard,
R.N. (1984):**

Acetylc Glycyl ether phosphoryl choline (AGEPC) a
putative mediator of cardiac anaphylaxis in guinea
pig.
Circ. Res., 54 : 117 - 124.

Lim, D.J. (1984):

Functional morphology of tubotympanum.

Acta. Otolaryngol (stockh). Suppl. 414 : 13-18.

Lim, D.J. and De-Maria, T.D. (1982):

Panel discussion. : pathogenesis of otitis media,

Bacteriology and immunology.

Laryngoscope 92 : 278 - 286.

Lim, D.J. and Shimada, T. (1971):

Secretory activity of normal middle ear epithelium.

Ann. otol., 80 : 319 - 329.

Liu, Y.S. (1976):

Microorganisms in chronic otitis media with effusion.

Ann. Otol. Rhinol. Laryngol., 85, "Suppl 25" : 249.

Lotner, G.Z.; Lynch, J.M.; Betz, S.J. and Henson, P.M. (1980):

Human neutrophil derived platelet activity factor.

J. Immunol., 124 : 676 - 684.

Lynch, J.M. and Henson, P.M. (1986):

The intracellular retention of newly synthesized platelet activating factor.

J. Immunol., 137 : 2653 - 2661.

McManus, L. (1986):

Pathobiology of platelet - activating factors.

Pathol. Immunopathol. Res., 5 : 104-17.

McManus, L.M.; Hanahan, D.J. and Pinckard, R.N. (1981):

Human platelet stimulation by acetylcetyl glyceryl ether phosphoryl choline.

J. Clin. Invest., 67 : 903 - 906.

McNicoll, W.D. and Scanlen, S.G. (1979):

Submucous resection : The treatment of choice in the Nose -Ear distress syndrome.

The Journal of Laryngology and Otology, 93 : 357 - 367.

Mencia-Huerta, J.M.; Lewis, R.A.; Razin, E. and Austen, K.F. (1983):

Antigen - initiated release of platelet - activating factor (PAF - acether) from mouse bone marrow - derived mast cells sensitized with IgE.

J. Immunol., 131 : 2959 - 2964.

Minami, T.; Kubo, N. and Tomoda, K. (1992):

Effect of various inflammatory mediators on eustachian tube patency.

Acta otolaryngol (stockh)., 112 : 680 - 685.

Morley, J.; Page, C.P. and Paul, W. (1983):

Inflammatory actions of PAF acether in guinea pig skin.

Br. J. Pharmacol., 80 : 503 - 509.

Naobumi, N.; Gieebink, G.S.; Daniel, Z.; Harada, T. and Juhn, S.K. (1991):

Early biochemical events in pneumococcal otitis media : Arachidonic acid metabolites in middle ear fluid.

Ann. Otol Rhinol. Laryngol, 160 : 385 - 388.

Nomura, Y. (1984):

Otological significance of the round window.

Advances in Oto-Rhinolaryngology.

No. 33. edited by C.R. Pfaltz Basel : Karger, PP. I - 184.

Ostfeld, E.J. and Silberberg, A. (1991):

Gas composition and pressure in the middle ear : a model for the physiological steady state.

Laryngoscope., 101 : 297.

Paparella, M.M.; Hiraide, F. and Juhn, S.K. (1970):

Cellular events involved in middle ear fluid production.

Ann. Otol. Rhinol. Layrngol, 79 : 766 - 779.

Paparella, M.M. and Koutroupas, S. (1982):

Exploratory tympanostomy revisited.

Laryngoscope, 95 : 531.

Parenter, L. and Flower, R.J. (1985):

Hydrocortisone and "macrocatin" inhibit the zymosan induced release of lyso-PAF from rat peritoneal leukocytes.

Life Sci., 36 : 1225 - 1231.

Pipkorn, U.; Karlsson, G.; Bake, B. (1984):

Effect of platelet activating factor on human nasal mucosa.

Allergy, 39 : 141 - 145.

Rhee, M.D. and Chung, K.U. (1993):

Otitis media induced by platelet activating factor.

Ann. Otol. Rhinol. Laryngol., 102.

Riu, R. (1966):

La physiologie de la trompe d'Eustache.

Paris, Libraire Anette.

Robinson, D. (1987):

Lipid mediators of inflammation.

Rheum Dis. Clin. North. Am., 13 : 385 - 405.

**Rogers, D.F.; Aursudkij, B.; Evans, T.W.; Belvisi, M.G.;
Chung, K.F. and Barnes, P.J. (1987):**

Platelet activating factor increases protein exudation
but not mucus secretion in guinea pig trachea in
Vivo.

Am. Rev. Respir. Dis., 135 : 160.

Sade, J. (1966):

Middle ear mucosa.

Arch otolaryng, 84 : 137 - 146.

Sade, J. (1971):

Mucociliary flow in the middle ear.

Ann. Otol. Rhinol. Laryngol., 80 : 336 - 341.

**Shaw, J.O.; Pinckard, R.N.; Ferrigni, K.S.; McManus, L.M.
and Hanahan, D.J. (1981):**

Activation of human neutrophils with 1-0-hexadecyl /
octadecyl-2-acetyl -sn-3-phosphorylcholine (platelet
activating factor).

J. Immunol, 127 : 1250 - 1255.

Silverstein, H.; Miller, G.F. and Linderman, R.C. (1966):

Eustachian tube dysfunction as a cause for chronic
secretory otitis media in children.

Laryngoscope, 76 : 259 - 273.

Smith, G.D.L. (1984):

Etiologic aspects and pathogenesis of persistent middle ear effusion.

Ann. Otol, Rhinol, laryngol, 5 : 286 - 290.

Stenfors, L.E.; Hellstorm, S. and Albiin, N. (1985):

Eustachian tube function. Middle ear clearance.

Ann. Otol. Rhinol, Laryngol, Suppl., 120 (94) :
30 - 31.

Stenfors and Raisanen (1991):

Immunoglobulin - coated bacteria in effusion from secretory and suppurative otitis media.

Ann. J. Oto. Laryngology, 12 : 161 - 164.

Takahashi, H.; Fujita, A. and Honjo, I. (1987) :

Site of eustachian tube dysfunction in patients with OME.

Ann. J. of Otolaryngol., 8 : 361 - 362.

Tence, M.; Polonsky, J.; LeCouedic, J.P. and Benveniste, J. (1980):

Pelease, purification, and characterization of platelet activating factor (PAF).

Biochimie, 62 : 251 - 259.

Tos, M. (1981):

Experimental tubal obstruction. Changes in the middle ear mucosa elucidated by quantitative histology.

Acta. otolaryngol (Stockh); 92 : 51.

Tos, M. (1985):

Histologic anatomy of the Eustachian tube and middle ear.

Ann. Otol. Rhinol. Laryngol., 94 : 9-11.

Valone, F.H.; Coles, E.; Reinhold, V.R. and Goetzl, E.J. (1982):

Specific binding of phospholipid platelet activating factor by human platelets.

J. Immunol., 129 : 1637 - 1641.

Valone, F.H. and Goetzl, E.J. (1983):

Specific binding by human polymorphonuclear leucocytes of the immunological mediator.

1-O-hexadecyl/octadecyl-2-acetyl-sn-glycero-3-phosphoryl choline.

Immunology, 48 : 141 - 149.

Van Cauwenberge, P.B. and Bernstein, J.M. (1987):

Inflammatory mediators in the middle ear diseases. In
: Immunology of the ear.

J. M. Bernstein and P.L. Ogra (Eds.). Raven Press,
New York, PP. 331 - 343.

Van Der Beek, J.M.H. and Kuijpers, W. (1984):

The mucosperiosteum of the middle ear in
experimentally induced sterile otitis media.

Acta. Otolaryngol. (Stockh) : Suppl. 414 : 71 - 79.

***Vargaftig, B.B.; Chignard, M.; Benveniste, J.; Leforte, J.
and Wal, F. (1981):***

Background and present status of research on platelet
activating factor (PAF-acether).

Ann. NY acad. Sci., 370 : 119 - 137.

***Vargaftig, B.B.; Lefat, J.; Chignard, M. and Benveniste, J.
(1980) :***

Platelet activating factor induces a platelet dependent
bronchoconstriction unrelated to the formation of
prostaglandin derivations.

Eur. J. Pharmacol., 65 : 185 - 192.

Veltri, R.W. and Sprinkle, P.W. (1973):

Serous otitis media. Immunoglobulin and lysozyme
levels in middle ear fluid and serum.

Ann. Otol. Rhinol. Laryngol., 82 : 297 - 301.

Wardlaw, A.J.; Mogbel, R.; Cromwell, O. and Kay, A.B. (1986):

Platelet activating factor : a potent chemokinetic factor for eosinophils.

J. Clin. Invest., 78 : 1701 - 1706.

Wirtz, H.; Lang, M.; Sannwald, U. and Hahn, H. (1986):

Mechanism of platelet activating factor (PAF) induced secretion of mucus from tracheal submucosal glands in ferret.

Fed. Proc., 45 : 418.

Yamashita, T.; Okazaki, N. and Kurmazaw, T. (1980):

Relation between nasal and middle ear allergy, experimental study.

Am. Otol- Rhinol, Laryngol., 29 (68) : 147 - 152.

Yasaka, T.; Boxer, L.A. and Bachner, R.L. (1982):

Monocyte aggregation and superoxide anion response to formyl methionyl - leucyl phenylalanine FMLP and platelet activating factors (PAF).

J. Immunol., 128 : 1939 - 1944.

Yellon, R.F.; Marucha, P.T. and Craven, R. (1991):

Characterization of cytokines present in middle ear effusions.

Laryngoscope, 101 : 165 - 169.
