

REFERENCES

Anderson, S.D. (1980):

"Some ECMR properties in relation to other signals from the auditory periphery". *Hear. Res.*, 2: 273-296.

Anderson, S.D. and Kemp, D.T. (1979):

"The evoked cochlear mechanical responses in laboratory Primate. *Arch. Otorhinolaryngol.*, 224: 47-54.

ANSI (American National Standard Institute), (1969):

"Specifications for audiometers S 3.6 "

New York: American National Standard Institute, Inc .

Arts, H.A.; Norton, S.J. and Rubel, F.W. (1990) :

"Influence of perilymphatic tetrodotoxin and calcium on hair Cell function ". *ARO Abstracts*.

Ashmore, J.F. and Brownell, W.E. (1986):

"Movement induced by electrical stimulation in outer hair cells isolated from the guinea pig cochlea" *.J.physiol* , 377: 41-47 .

Astrup, P., Kjeldsen, K. and Wanstrup, J. (1967):

"Enhancing influence of carbon monoxide on the development of atheromatosis in cholesterol fed rabbit " *.J.Atheroscler Res.*, 7, 343.

Auerbach, O. , Hammond, E.C. ,Kirman , D . and Garfinkel , L .

(1967):

"Histological changes in bronchial tubes of cigarette smoking dogs" , Cancer, N.Y ., 20 , 2055 .

Baldwin, M. and Watkin, p. (1992) :

"The clinical application of otoacoustic emissions in paediatric audiology." J. laryngol . otol ., 106 : 301- 306 .

Bennet , J. R . (1976) :

"Cigarette smoking and chest pain" . Brit . Med . J. 1: 97 .

Bentley,H.R. , and Burgan ,J.G. (1961) :

Cited by E. L. Wynder & Hoffmann (1967) Tobacco Manufactures , Standing Committee Research Papers 4,2nd ed.

Bogen , E.(1929) :

"The composition of cigarettes and cigarette smoke" .
JAMA , 93 : 110 .

Bonfils, P., Avan, P. and francois, M. (1990) :

"Clinical significance of otoacoustic emission": A perspective.
Ear and Hearing, 11(2): 155-158.

Bonfils P.; Bertrand Y. and Uziel, A. (1988a) :

"Evoked otoacoustic emissions: Normative data and presbycusis". *Audiology*, 27: 27-35.

Bonfils, P., Uziel, A. and Narcy, P. (1989) :

"The properties of spontaneous and evoked emission in neonates and children" .A preliminary report.
Arch. Otorhinolaryngol., 246: 249-251.

Bonfils, P., Uziel, A and Pugol R. (1988b) :

"Evoked otoacoustic Emissions form adults and infants":
Clinical application. *Acta. Otolaryngol.*, 105: 445-449.

Bray , P., and Kemp , D.T. (1987) :

"Advanced echo technique suitable for infant screening".
Br . J. Audio1., 12: 191- 204.

Brown .A.M , Kemp D.T (1984) :

"Suppressibility of 2 F1 –F2 stimulated acoustic emissions in Gabril & Man" .*Hear , Res* , 13 : 29 –37 .

Brown ,A .M ,Mc Dowell ,B. and Forge , A .(1989) :

"Acoustic distortion products can be used to monitor the effects of chronic gentamycin treatment" . *Hear .Res .* , 42 :143-156 .

Brown ,M .C , Nutall ,A.L of Masta , R. I (1984) :

”Intracellular recording from cochlear inner hair cells, effects of stimulation of the crossed olivocochlear efferents”.

Science,222,69-72.

Brownell, W. E. (1984) :

”Microscopic observation of the cochlear hair cell motility” .

Scanning Electron Micros.,111 : 1401-1406 .

Brownell, W.E.(1990) :

”Outer hair cell electromotility and otoacoustic emissions” .Ear and hearing ,11 (2) : 82 - 92.

Brownell , W.E and Kachar ,B. (1986) :

”Outer hair cell motility” . A possible electrokinetic mechanism

. In : Allen, J .B., Hall, J.L., Hubbard , A.E., Neely ,S.T.

and Tubis, A . (eds) . Perineural Auditory Mechanisms.

New York. springer-verlag . pp 369-376 .

Bundgaard T, Wildt J, and Elbrond O , (1994) :

”Oral squamous cell carcinoma in non users of tobacco and alchol”.

Clin ,Otolaryngol , 19 : 320.

Campbell , J .M . (1957) :

“Polycyclic hydrocarbons in cigar smoke” . B r. J. cancer ,11 , 192

Campbell .M.J and Machin D (1993) :

”Medical statistics” .A common sense approach . John Wiling
& Sons(Eds.) 2nd Ed . New York Chichester Brisbane ,Tronto
& Singapore.

Cane , A.M.,O ‘ Donoghu G. and Lutman , M.F. (1992) :

“The feasibility of using otoacoustic emissions to monitor cochlear
function during acoustic neuroma surgery” .Scand . Audiol . , 21:
173-176 .

Carrol ,P. & Ireland ,P.(1935) :

“Association of toxic deafness with toxic ambylopia due tobacco
and alcohol ”. Archives of otolaryngolo., 21,459

Carter ,W.L . , (1975) :

”Fixation of tobacco smoke aerosols for size distribution
studies” . J. of colloid and interface science ,53 :134 .

Cogna (2001) :

“ cigarette smoking as a risk factor of periodontal diseases ” .

A review mailto : info @ dental - smile .com .

Collet , L . (1993) :

”Use of otoacoustic emissions to explore the medial olivocochlear system in humans” . Br . J. Audiol ., 27 : 155-159.

Daniel W.W (!987) :

”Biostatistics ,A foundation for analysis in the health sciences ”
4th Ed John Wiley & Sons . New York Chichester Brisbane
,Tronto & Singapore .

Dijk Van ,P . and Wit , H.P . (1987) :

”The occurrence of click evoked otoacoustic emissions” (Kemp
echoes) in normal hearing ears .
Scand . Audiol . , 16 :62 – 64 .

Doll , R.& Hill ,A.(1964) :

”Mortality in relation to smoking ” : ten years observations of
British doctors .Brit. Med .J.1,1399 .

Domino , E.F. (1969) :

”Tobacco cigarette smoking and patellar reflex depression ” :
Clin . Pharmac . Ther . , 10 ,72 .

El-Sharabassy and Zaghloul (2001) :

"Paper presented at American Academy of Audiology" .San Diego,USA .

Evans , B.N. (1988):

"Asymmetries in outer hair cell electromechanical responses" .
Abstracts of the Midwinter Meeting of the association for
Research Otorhinolaryngology ,11: 29.

Evans , E.F. , Wilson , J.P. and Borewe , T.A. (1981) :

"Animal models of tinnitus" . In : Tinnitus CIBA Foundation
Symposium . Edited by : D . Evered and G. Lawrenson .
Pitman London . pp: 108-138.

Fabiani , M . (1993) :

"Evoked otoacoustic emissions in the study of the adult
sensorineural hearing loss ". Br . J. Audiol . , 27 : 131-137 .

Fabiani , M . , Baretto , F . and Gobbi , E . (1990) :

"Acute and chronic acoustic trauma in young adults studied with
evoked otoacoustic emissions" . I n : Montorsi M . , Zennaro , F.
(eds) . Specialized surgery . Bologna :Monduzzi Editore ,pp :24.

Fraumeni , J . F . (1975) :

”Persons at high risk of cancer” ,an approach to cancer ,
aetiology and 1 st .Ed Academic Press , New York .

Galal , A ., Youssef ,A . , and Salem ,E.S.(1973) :

”Nicotine levels in relation to pulmonary manifestations of Goza
and cigarette smoking” .Egypt .J. Chest . Dis & Tub 15 (2) : 141-
156 .

Glorig ,A.(1965) :

”Cigarette smoking and tinnitus” .Jama . ,192,269 .

Goldsmith ,J.R . and Landaw , S.A (1968) :

”Carbon monoxide and human health”. Science , 162 ,1352 .

Gold ,T. (1948) :

”Hearing 11: The physical basis of the action of the cochlea.

Proc. R . Soc. London Biol. Sci ., 135: 492-498.

Harris , F.P . and Probst ,R . (1992) :

”Transient evoked otoacoustic emissions in patients with

Meniere ' s disease” .Acta . Otolaryngol . (stockh) . , 112 : 36-44 .

Herxheimer , A. , Griffiths , R.L . , Hamilton , B, and Wakefield , M .
(1967) :

"Circulating effect of nicotine aerosol inhalation and cigarette
smoking in man" . Lancet ii : 754 .

Incze , J . , Vaughan , G . W , . P . and Stony , M.S (1982) :

"Premalignant changes in normal appearing epithelium in
patients with squamous cell carcinoma of upper aerodigestive
tract" . Am .J . S urg . , 144 : 401 .

Jarboe , C.H , . and Rosene , C.J (1961) :

Quoted by Shephard (1982): " Volatile products of pyrolysis of
nicotine" . J .Chem , Soc .19 :2455 –2458 .

Jarvik , M.E . (1979) :

"Tolerance to the effect of tobacco" .Natt : Ins . During abuse Res.
Monoger :Ser . 231 : 150 .

Jerger ,Jay . (1976) :

"Clinical experience with impedance audiometry" .
Arch. Otolaryngol . , 92,311-324 .

Johnsen N.J., Bagi , P . and Elberling , C . (1983) :

”Evoked acoustic emissions from the human ears” . 111 :

Finding in neonates ,Scand . audiol . , 12 : 17-24 .

Johnsen N.J. , Bagi , P., Parbo , J . Elberling ,C. (1988) :

” Evoked acoustic emissions from the human ears” . IV :

Final results in 100 neonates . Scand . Audiol . , 17 : 27-34 .

Johnsen , N.J ., and Elberling, C. (1982) :

” Evoked acoustic emissions from the human ears ”. 1 :

Equipment and response parameters .Scand . Audiol . , 11 , 3-12 .

Johnson N.J . , Parbo , J . and Elberling , C . (1989) :

”Evoked otoacoustic emission s from the human ears” . V .

Developmental changes . Scand . Audiol . , 18 : 59-62 .

Kamal ,M , Tawfik , S ,Shalaby A . Isaac, O . (2001) :

”The development of a hearing conservation program for bus drivers in Cairo transport authority ” unpublished doctoral thesis .

Kamal, N ;Hazah,N and Eldanasuri ,I(1996) :

”Otoacoustic emission measurementsin Meniere 's disease”.

Ain Shsms medical J. october

Karen J. Crick shanks , Ronald Klans , Barbara , Klim , Terry ., l. and Ted , S. (1998) :

"Cigarette smoking and hearing loss " . J . A m Med Ass ' n 1715
– 1719 .

Kemp , D.T. (1978) :

"Stimulated otoacoustic emissions from within the human
auditory system " . J. Acoustic .Soc .Am., 64: 1386-1391.

Kemp, D.T. (1980) :

"Towards a model for origin of cochlear echoes".
Hear . Res.,2: 533-548.

Kemp, D .T.(1981) :

"Physiologically active cochlear micromechanics-one
source of Tinnitus". In: Tinnitus CIBA Foundation
Symposiun . Edited by : D.Evered and G. Lawrenson.
Pitman , London . PP: 54-81 .

Kemp, D.T.(1982) :

"Cochlear echoes : Implications for noise induced hearing
loss". In :New perspectives on noise induced hearing loss
. Edited by :R.P. Hamernik , D. Henderson and R. Salvi .
Raven , New york . PP: 189-207 .

Kemp, D. T. (1986) :

"Otoacoustic emissions , traveling waves and cochlear mechanics" . Hear . Res . , 22: 95 –194 .

Kemp , D.T. , Bray , P. , Alexander , L. and Brown , A.M. (1986) :

"Acoustic emissions cochleography" –practical aspects .

Scand . Audiol . Supp . , 25 : 71 –82 .

Kemp , D.T. , Bray , P . and Ryans , M . (1990) :

"A guide to effective use of otoacoustic emissions" .

Ear and Hearing , 11 (2) : 93-105 .

Kemp , D.T. and Brown A.M. (1984) :

"Ear canal acoustic and round window electrical correlates of $2f_1$ – f_2 distortion generated in the cochlea" .Hear .Res . , 13 : 39-46 .

Kemp ,D.T.and Chum , R.A. (1980) :

"Properties of the generator of stimulated acoustic emissions".

Hear .Res . , 2 : 213 - 232 .

Kim, D.O. (1982) :

"Efferent neural control of cochlear mechanisms ?

Olivocochlear bundle stimulation effects cochlear

biomechanisms non linearity" .Hear . Res . , 6 : 171- 182 .

Kirikaie, I . , Nomura. Y. Shitara, T .Kobayashi ,T. (1962) :

”Sudden deafness due to burgers disease” . Archives of
otolaryngology ,75,502 .

Koide , Y . (1958) :

”Introductory studies on chemical physiology of labyrinth” .

Acta . Med . Biol . , 6: 1 .

Kumar , R . and Lader , M . (1981) :

”Nicotine and smoking” :Curr . Dev. Psychopharmacol ,6 : 127 .

La Charriere , D . (1875) :

”Are you smoking more , what about hearing loss?

In S.L Shapiro MD (1964) Eye Ear Throat Monthly 96-100 .

Lader , M . (1978) :

”Nicotine and smoking behaviour”, Br. J .Clin . Pharmacol . 5:289 .

Lareo AC, et al ,(1992) :

”History of previous nasal diseases and sinonasal cancers”, a case –
control study .Laryngoscope 102 : 439.

Laurance , D.R . (1980) :Clinical pharmacology” 5 th ed . Churchill – Living stone , London . , p . 465.

Linnaeus , C (2001) : Species platerium ” . In Google search , www.
”Smoking and hearing loss” .

Long , C.R . and Tubis, A , (1988) :

”Modification of spontaneous and evoked otoacoustic
Emissions and associated microstructure by aspirin
consumption”. J. Acoustic . Soc . , Am . , 84 : 1343 –1353 .

Lousbury – Martin ,B.L., Martin ,G .K . and Coats , A .C (1991a):

”Physiology of the auditory and vestibular system” . in
Ballenger J .J (ED) Diseases of the nose , ear , throat , head &
neck surgery , Lea and Febiger , Philadelphia . CH ,46 : 948-1005 .

Mac Carthy , R . L . and Shklar , G .(1980) :

”Diseases of oral mucosa ”. 2 nd Ed .
Lea and febiger ,Philadelphia , p : 16 , 178 and 182 .

Marston , L . , Sterrett ,M .& Mcl Ennan ,R. (1960) :

”Effect of cigarette smoking on tympanic membrane admittance
characteristics” .Ear & Hearing , 1,267.

Martin , G.K ., and Lousbury – Martin , B.L . (1993) :

”Clinical testing of distortion product otoacoustic emissions”.

Ear and Hearing , 11 (1) : 11-21.

Martin , G.K., nd Lousbury - Martin ,B.L. Probst R.and

Coats.A.C. (1987) :

”Acoustic distortion products in rabbit ear canal 11”.

Sites of origin revealed by suppression contours and pure tone exposure . Hear .Res ., 28 : 101-108.

Martin, G.K. ,Probst , R . and Lousbury –Martin ,B.L.(1990)

”Otoacoustic emission in human ears” : Normative

findings . Ear and Hearing , 11(2) : 106 –120.

Maxon , A.B., . White , K.R. and Vohr , B.R. (1993) :

”Using transient evoked otoacoustic emissions for neonatal hearing screening” . Br . J . Audiol . , 27 : 149 – 153 .

Mc Cartney , S , Peter , A and Barbara , W : (1994) :

”Central auditory processing in school age children prenatally exposed to cigarette smoke” .Neurotoxicol :3 , 269-276 .

Mc Fadden, D. and Plattsmier, H.S (1984):

"Aspirin abolishes spontaneous otoacoustic emissions".

J. Acoustic Soc .Am., 76:443-448.

Mc Nally , W.D. (1932) :

"The tar in cigarette smoke and its possible effects" .

Am . J. Cancer , 16 : 1502 .

Metha , F.S , (1981) :

"Early detection of oral cancer and precancerous lesion"

2nd Ed : Basic Dental Research Unit , Tata Institute Bombay , P .90

Metha , F.S , Sohair , B.E . and Deftary, D .K . (1972) :

"A correlative histological study of carcinoma and epithelial

atypia of palate among Indian reverse smoker" .

Brit . J . Cancer , 26 : 230 .

Mumpower , R.C. , Lewis , J .S , and Touey , G.P. (1962) :

Quoted by Shephard (1982): " Determination of carbon monoxide in cigarette smoke by gas chromatography" . Tobacco Sci .6:142-145 .

Naguli (2001) : "Environ Health project ". Aug 109 suppl , 4 : 578 –

588 .

National Health Survey (1967) :

"Cigarette smoking and health characteristics in the U.S". July 1964-June ,1965, Washington , National center for health statistics .10 ,No.34.

National committee of mother and childhood , (April, 2001) : " Report. of National conference" ,Cairo,Egypt.

Neely , S.T. and kim,D.O (1983) :

"An active cochlear model showing sharp tuning and high Sensitivity". Hear . Res . , 9 : 123 –130

Newsome , J .R . , and Keith , C .H . (1965) .

"Variation of the gas phase composition within a burning cigarette" .Tobacco Sci . 65-69. Quoted by Shephard (1982) .

Nizietic ,B . & Alterj ,A.(1952) :

"Quoted from hearing loss in adults : relation to age ,sex ,exposure to loud noise and cigarette smoking ".arch .Environ .Health ,29 , Au .1974 .

Nolan , M.and Tucker , I .G . (1981) :

"Functional hearing loss in children" .

J . Br . Assoc . Teach . Deaf , 5 : 2 – 10 .

Norton , S.J.and Neely , S.T. (1987) :

”Tone –burst evoked otoacoustic emissions from normal hearing subjects”. J . Acoust. Soc . Am . 81 : 1868- 1972 .

Norton , S , J. , Mott, J.B. and Champlin ,C.A. (1989) :

”Behavioral spontaneous otoacoustic emissions following intense acoustic stimulation” . Hear . Res. , 38 : 243-258 .

Otto ,G. (1923) :

” Nicotine Poisoning of the inner Ear” ,33 Laryngoscope
(4) 262-266 April .

Paymaster , J . C . (1956) :

”Cancer of buccal mucosa ”, A clinical study of 650 cases in Indian patients . Cancer, 9 : 431 .

Pindborg , J .J . (1981) :

”Oral cancer and precancer” .Reprinted 1st ed . , John .Wright
and Sons Ltd .Bristol . pp . 12 and 136 .

Pindborg , J .J . Barmes , D . and Road Peterson , B . (1968) :

”Epidemiology and histology of oral leukoplakia and leukoderma Cancer” . , 22 : 379 .

Pindborg , J. J . and Sirsal , S. M . (1966) :

”Oral submucous fibrosis” .Oral Surgery , 22 : 746 .

Plinkert , P.k. (1992) :

”Early detection of inner ear hearing loss by evoked otoacoustic emissions in high risk infants and after administration of ototoxic drugs” .Abstracts of first National Conference on ECOG , OAEs and intraoperative monitoring . Wuzburg .sept . 20-24.

Probst , R . , Casts , A.C. ,Martin , G.K. and Lousbury

-Martin , B.L. (1986) :

”Spontaneous , click and tone –burst evoked otoacoustic emissions from normal ears” . Hear , Res . , 12 : 261 –275 .

Probst , R . , Lousbury – Martin , B.L . and Martin , G.K. (1987) :

”Otoacoustic emissions in ears with hearing loss” .

Am . J . Otolaryngol . , 8 : 73 –81 .

Probst, R., Lousbury –Martin ,B.L and Martin G.K ., (1991):

”A review of otoacoustic emission ”.

J. Accoust ,Soc .Am ., 80 (5).

Ramenten ,N. , Rakienten , M .L . , Faldman , D. and Doykin , M.J.
(1952) :

”Mamalian ciliated respiratory system” , A.M.A .

Arch .of otolaryngol . American medical association Chicago
10 , ILL ., 56 : 494 .

Reshef , I . Attias , J . and Furst , M . (1992) :

”Evoked otoacoustic emissions in testing noise induced
sensorineural and functional hearing loss ”.

Abstracts of first European Conference on Audiology .
Cambridge , 12-24 .

Robinson,J.C ., and Forbes ,W .F. (1975) .

”The role of carbon monoxide in cigarette smoking”

1.carbon monoxide yield from cigarettes .

Arch . Environ . Health 30 : 425 –434 .

Robin ,L .(1977) :

”Sensorineual deafness” . Scott-Brown, diseases of the ear , nose
,throat .Ballantyne , J & Groves .J . Archives of otolaryngology
,82,236 .

Roth , M (1946) :

"The use of tobacco" . I n John . H .(ed)the use of tobacco . Its physical, intracellular, moral effects on the human system
(1967) p . 102 .

Salem ,E.S . , Abdel Hakim ,M . , and Hanafi ,A. (1973) :

"Goza versus cigarette smoking among patients with Pulmonary disease" .Egypt .J . Chest Dis &Tub . 16 (2) : 121-140 .

Salem ,E.S . , and Abdel latif ,E.A .A .(1974).

"Physical and physiologic studies on the etiogenesis of chronic airways obstruction in Goza smokers".
Egypt .J . chest Dis . & Tuub . 17 (1) : 1-10 .

Salomon ,G. Groth , J . and Anthonisen ,B . (1993) :

"Preliminary results and consideration in hearing screening of newborns based on otoacoustic emissions" .
Br . J . Audiol . , 27 : 139-141 .

Samwell , Z . (1973) :

"Correlation of smoking history with hearing loss" . J . Am Med Ass'n (8) Feb . 920 .

Santos-Sacch, J. and Dilger, J.P. (1988) :

"Whole cell currents and mechanical response of isolated outer hair cells". Hear .Res ., 35 : 143-150 .

Schloth , E . (1982) :

"Mechanical and otoacoustical influences on spontaneous otoacoustic emissions". Hear . Res . , 11 : 285-293 .

Shapiro,S . (1964) :

"Are you smoking more but hearing loss". Ear ,nose & throat .Mon. 43.96

Shenoi , p .(1979) :

"Vascular lesions of the inner ear". Scott – brown, Disease of the ear , nose & throat .Ballantyne ,J. &Groves ,(ed) , (pub .) ,London , vol ,2.

Shephard , R . J . (1982):

"The risks of passive smoking". Croom Helm – London
Camberra p p 17-21 .

Siegel, J. H. and Kim, D. O (1982) :

"Cochlear biomechanics vulnerability to acoustic trauma and other alterations, as seen in neural responses and ear canal sound pressure". In : new perspective on noise induced hearing loss : R.P. Hamernik, D. Haderson and R. Salvi. Raven (Eds.), New York . pp : 137 –151 .

Siegelaub, A., Friedman, G., Adour, K. & Seltzer, C. (1974) :

"Hearing loss in adults": relation to age, sex, exposure to loud Noise and cigarette smoking. Arch. Environ. Health. 29, 107 .

Solan, C. (1988) :

"Auditory processing, speech reception, and language comprehension". What is the connection? Symposium on central auditory processing, August, 18-19 .

Soliman, S. (1976) :

"speech discrimination audiometry using Arabic phonotactically balanced words". Ain Shams Med. J. 27, 27-30

Soliman, S. (1985) :

"Development of SSW test".

Paper presented at the 8 th Ain Shams Annual congress

Soliman ,S, Tawfik ,S. & Dawoud , M .(1989) :

"The effects of smoking on hearing" . Ain Shams Medical
Journal 246-256

Steinberg , R.A. and Jeffery , R.N . (1956):

Quoted by Shephard (1982): "Effect of micronutrient
deficiencies on nicotine formation by tobacco in water culture" .
Plant physiol .31; 377-382 .

Stevens , J. C . (1988) :

"Click evoked otoacoustic emissions in normal and hearing
impaired adults" . Br . J . Audiol . , 22: 45 –49 .

Stevens , J. C .Hutchinson , J . , and Connell , J . (1990) :

"Click evoked otoacoustic emissions in neonatal screening" .
Ear and Hearing , 11 (2) : 128-133 .

Stevens , J. C., Webb , H.D. , Smith , M.E. Bulfin , J . T

and Ruddy .H. (1987) :

"A comparison of otoacoustic emissions and brainstem electric
response audiometry in normal newborn and babies admitted
to a special care baby unit". Clin phys ,physiol, 8: 95-104.

Tachmes, L . , Fevnandez , R.J. and Sackner ,M.A. (1978) :

”Haemodynamic effect of smoking cigarettes of high and low nicotine content” . Chest 74 : 243 .

Tanaka , Y . Suzoki , M . and Inoue , T . (1990) :

”Evoked otoacoustic emissions in sensorineural hearing impairment” :its clinical applications .

Ear and Hearing , 11 (2) : 134-143 .

Traynelis , S.T. and Dingledines, R .(1989) :

”Role of extracellular spaces in hyperosmotic suppression of potassium induced electrographic seizure” .

J . Neurophysiol . , 61 : 927 – 938 .

Tso , T.C.and Gari , G.B. (1972) :

”Effect of tobacco characteristics on cigarette smoke composition”

In 1 .Shmeitz (ed) ,The chemistry of tobacco and tobacco smoke – Plenum Press .New York p.p 53-55 .

U.S.Public health Service (1968) :

”Supplement to public health service”. Publication N0.1969,P.32 .

U.S. Surgeon General (1972) .

"The health consequences of smoking".U.S.department of Health
Education and Welfar,Arlington,va. Quoted by shephard (1982) .

Uziel , A . 1990) :

"Les oto-emissions acoustiques" . Applications cliniques.

Ann . otolaryngol . (Paris) . , 107:48-50 .

Van Caneghem, D .(1954) :

"Quoted from hearing loss in adults :relation to age , sex ,exposure
to loud noise and cigarette smoking".Arch.Environ,Health.27,Aug
.1974 .

Wald ,.N .J (1976) :

"Smoking tables in carbon monoxide" .Brit .Med .J. , 1: 434 .

Weiss , W (1970) :

"How smoking affects hearing" .Medical Times J , 84 -89 .

Weston ,T ; (1964) :

"Presbycusis" : A clinical study , J . laryngol otol. 273 - 286 .

Weybrew ,J. A ., Wotz , W.G ., and Long ,R.C.(1972) :

”Projected changes in the composition of bright (flue-cured)
to tobacco”. in I . Schmeltz (ed.). The chemistry of tobacco and
tobacco smoke , plenum press , New – York pp 38-39 .

Wier.C.C., Pasanen , E.G and Mc Fadden, F.(1988) :

”Partial dissociation of spontaneous otoacoustic emissions and
distortion product during aspirin use in humans”.

J.Acoust. Soc. Am ., 84:230-237.

Wilson , J . (1980 a):

”Evidence of cochlear origin of acoustic emissions threshold , fine
structure and tonal tinnitus” . Hear .Res . , 223-253 .

Wilson , J.P. (1984) :

”Otoacoustic emissions and hearing mechanisms” .

Rev. Laryngol ., 105 : 179- 191.

Wilson , J. P. and Evans . E. F. (1983) :

”Effects of frusemide , flexidil , noise and tone over stimulation on
evoked otoacoustic emissions in cat” .Proc.Int.physiol.Sci., 15,100 .

Wit , H . P . and Ritsma , R . J . (1980) :

”Evoked acoustical responses from the human ear”: Some
experimental results .Hear. Res . , 2: 253-261.

World Health Organization WHO, (1975):

”Smoking and its effects on health”, WHO technical report
, Series N0, 568 .WHO, Geneva, 1975.

Wynder ,E.L. , Hoffman ,D. (1967) :

Quoted by Shephard (1982) : ” Tobacco and tobacco smoke”,
Academic press, New York.

Wynder E.L. Stillman SD: (1977) :

”Comparative epidemiology of tobacco related cancers”.
Cancer research .37: 4608.

Zelman ,S . (1973) :

”correlation of smoking history with hearing loss”. J .American
Medical Association ,223,920 .

Zwicker ,E .(1983) :

”Delayed evoked otoacoustic emissions and their suppression
by Gaussian shaped pressure impulses ”.Hear .Res .,11 : 59-71 .

Zwicker , E. and Manley , G . (1981) :

” Acoustical responses and suppression period patterns in guinea pigs” . Hear . Res . , 4 : 43-52 .

Zwislocki , J. (1962) :

”Analysis of the middle ear function” , I : Input impedance .

J. Acoust . Soc . Am . , 34 : 1514 – 1523 .