

References

- **Abbo A., Navon-Venezia S., Hammer-Muntz O., and Krichali T (2005):**
Multidrug –resistant *Acinetobacter baumannii*. CDC, Emerg Infect Dis; 11(1).
- **Abel F.L (1989):**
Myocardial function in sepsis and endotoxin shock. AmJ physiol; 257; R1265-R1281.
- **Abo-Alam M.S., Badrawy M.M., El-Rashidy N.G., and Morsy M.A (1998):**
Bacteriologic study on neonatal septicemia M.Sc.Thesis.
- **Adam B., Baillie G.S., and Douglas L.J (2002):**
Mixed species biofilms of *Candida albicans* and *Staphylococcus epidermidis*. J.Med.Microbiol; (51):344-9.
- **Adams-Chapman I., and Stoll B.J (2006):**
Neonatal infection and long-term neurodevelopmental outcome in the preterm infant. Curr Opin Infect Dis; 19(3):290-7.
- **Afsharpaiman S.H., and Mamishi S (2008):**
Diagnosis of bacteremia in febrile patients: PCR versus other routine methods. Tehran university medical journal; 66(3):186-175.
- **Al-Zwaini E.J.K (2002):**
Neonatal septicaemia in the neonatal care unit, Al-Anbar governorate, Iraq. Eastern Mediterranean Health Journal; 8, No 4 and 5.

- **Amann R. I., Ludwig W., and Schleifer K.H (1995):**

Phylogenetic identification and in situ detection of individual microbial cells without cultivation. Microbiol. Rev. 59:143–169.

- **Andrade, Soraya S., Bispo, Paulo J.M., Gales., and Ana G (2008):**
Advances in the microbiological diagnosis of sepsis. Shock. 30 (7) Suppl; 1:141 -46.

- **Anoop P., Anjay M.A., Vincent J., and Girija B (2003):**

Fulminant Citrobacter meningitis with multiple periventricular abscesses in a three-month –old infant. Braz J Infect Dis; 7(6) Salvador.

- **Ascuitto R.J., Geber M.A., and Cates K.L (1985):**

Buffy coat smears of blood drawn through central venous catheters as an aid to rapid diagnosis of systemic fungal infections. J Pediatr; 106:445.

- **Ashok K. Yadav., Wilson C.G., Prasad P.L., and Menon P.K (2005):**

Polymerase Chain Reaction in Rapid Diagnosis of Neonatal Sepsis. J Indian Pediatric; 42:681-84.

- **Asinidi A.A., Archibong E.L., and Mannan N.B (2002):**

Mother-infant colonization and neonatal sepsis in prelabor rupture of membranes. J Saudi Med; 23(10):1270-4.

- **Aurangzeb B., and Hameed A (2003):**

Neonatal sepsis in hospital –born babies: bacterial isolates and antibiotic susceptibility patterns. J Coll Physicians Surg Pak; 13(11):629 32.

- **Back N.A., Linemann C.C.Jr., and Pfaller M.A (1993):**

Recurrent epidemics caused by a single strain of erythromycin - resistant staphylococcus aureus. The importance of molecular epidemiology: JAMA; 270:1329.

- **Back N.A., Linnemann C.C., Staneck J.L., and Kotugal U.R (1996):**

Control of methicillin-resistant *Staphylococcus aureus* in a neonatal intensive care unit: use of intensive microbiologic surveillance and mripirocin. *Infect Control Hosp Epidemiol*; 17:227-31.

- **Badr K.F (1997):**

Glomerulonephritis: Roles for lipooxygenase pathways in pathophysiology and therapy. *J Curr Opin Nephrol Hyperten*; 6:111-118.

- **Badrawi N., Naguib H.F., El-Kholy A.A., and Fahmy S.S (2005):**

Epidemiological study of early –onset sepsis. M.D. Thesis.

- **Badrawi N., Wasef M.A., Ihsan I.A., and Gergis M.N (2001):**

Detection of different microbiological species responsible for early –onset sepsis .M.Sc .Thesis.

- **Baird D (1996):**

Staphylococcus: cluster forming Gram-positive cocci. In: Collee J.G., Marmion, B.P., Fraser A.G., and Simmons A., eds. *Mackie and Mac carthey practical medical microbiology*, pp 245-274. Churchill livingstone, Newyork.

- **Baltimore R.S (2004):**

Pseudomonas, *Burkholderia*, and *Stenotrophomonas* .In *Nelson Textbook of pediatrics*; 17th ed: 933-7. Edited by Behrman R.E., Kliegman R.M., and Jenson H.B. By W.B. Saunders.

- **Bang A.T., Bang R.A., Bactule S.B., Reddy H.M., and Deshmukh M.D (1999):**

Effect of home based neonatal care and management of sepsis on neonatal mortality: field trial in rural India. *Lancet*; 354:1955-61.

- **Barassi A., Pallotti F., and Melzi d’Eril G (2004):**

Biological Variation of procalcitonin in healthy individuals .Clinical Chemistry; 50:1878.

- **Bennet R., Eriksson M., and Zetterstrom R (1981):**

Increasing incidence of neonatal septicemia causative organism and predisposing risk factors Act Ped.Scand; 70-207.

- **Bennet R., Eriksson M., and Zetterstrom R (1987):**

Etiology of neonatal septicemia in relation to prior antibiotic treatment. Acta Ped.Scand, 76-673.

- **Benson D.A., Karsch-Mizrachi I., Lipman D.J., Ostell J., and Wheeler D.L (2007):**

GenBank. Nucleic Acids Res., 35(Database issue), 21–25.

- **Bick R.L (1989):**

Disseminated intravascular coagulation .Semin Thromb Haemosta ;15:1-98.

- **Bode-Thomas F., Ikeh E.I., Pam S.D., and Ejeliogu E.U (2004) :**

Current aetiology of neonatal sepsis in Jos University Teaching Hospital. Niger J Med ;13(2):130-5.

- **Boguski M.S., Lowe T.M., and Tolstoshev C.M (1993) :**

dbEST—database for ‘expressed sequence tags’. Nature Genet., 4, 332–333.

- **Bone R.C (1991):**

The pathogenesis of sepsis. Ann Intern Med;(b):115:457-469.

- **Bosshard P.P., Abels S .Zbinden ., Bo“ttger R.E.C., and Altwegg M (2003):**

Ribosomal DNA Sequencing for Identification of Aerobic Gram-Positive Rods in the Clinical Laboratory (an 18-Month Evaluation).J.Clinical Microbiol 41:4134-4140.

- **Boyle R.J., Chandler B.D., and Stonestree B.S (1978) :**

Early identification of sepsis in infants with respiratory distress. Pediatrics ;62 :744.

- **Boyer K.M., and Gotoff S.P (1986):**

Prevention of early-onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis. N Engl J Med; 314:1665-1669.

- **Braslavsky I., Hebert B., Kartalov E., and Quake S.R (2003):**

[Sequence information can be obtained from single DNA molecules.](#) Proc. Natl. Acad. Sci. U.S.A. **100** (7): 3960–4.

- **Brezis M., and Rosen S (1995):**

Hypoxia of the renal medulla.Its implication for disease .N Eng J Med; 332:647-655.

- **Brown D.R., Kutler D., Rai B., Chan T., and Cohen M (1995):**

Perinatal /neonatal clinical presentation bacterial concentration and blood volume required for a positive blood culture .J.Perinatal.15:157-159.

- **Brozanski B.S., Jones J.G., Krohn M.J., and Jordan J.A (2006):**

Use of Polymerase Chain reaction as a diagnostic tool for neonatal sepsis can result in a decrease in use of antibiotics and total neonatal intensive care unit length of stay.Journal of Perinatology, 26:688-692.

- **Buttery J.P (2002):**

Blood cultures in newborns and children: Optimizing an every day test
Arch Dis Child Fetal Neonatal ;87:F25-8.

- **Carlos C.C., Ringertz S., and Rylander M (1991):**

Nosocomial staphylococcus epidermidis septicaemia among very low birth weight neonates in an intensive care unit .J Hosp Infect; 19:201.

- **Carrigan S.D., Scott G., and Tabrizian M (2004) :**

Toward Resolving the Challenges of Sepsis Diagnosis. Clinical Chemistry; 50:1301-14.

- **Chang Y.L., Tseng C.P., and Tsay P.K (2004):**

Procalcitonin as a marker of bacterial infection in the emergency department: an observational study .Crit Care; 8(1):P12-20.

- **Cheesbrough M (2000):**

Biochemical tests to identify bacteria.In: Cheesbrough M,ed.District laboratory practice in Tropical countries, part1,pp:63-70.Cambridge university press,uk.

- **Cheisa C., Panero A., and Osborn J (2004):**

Diagnosis of neonatal sepsis: A clinical and laboratory challenge.Clinical Chemistry; 50:279-87.

- **Church G.M (2006) :**

Genome for all .Sci .Am.294(1):46-54.

- **Cimino G.D., Metchette K.C., Tessman J.W., Hearst J.E., and Isaacs S. T (1991):**

Post-PCR sterilization: a method to control carryover contamination for the polymerase chain reaction. Nucleic Acids Res. 19:99-107.

- **Cole F.S (1991):**

Bacterial infection of the newborn: In: Schaffer and Avery's disease of the newborn. Editors, HW, Ballard RA and Avery ME, 6th ed. WB Saunders company, Philadelphia; p350.

- **Cole F.S (1998):**

Bacterial infection of the newborn. In: Tauesch HW and Ballard RA, editors. Avery's Diseases of new born 7th ed. WB Saunders; 490-508.

- **Collee J.G., Duguid J.P., Fraser A.G., Marmion B. P., and Simons A (1996):**

Test for identification of bacteria. In: Collee J G, Fraser A G, Marmion B P, and Simon S A, eds. Mackie and Mac cartney practical medical microbiology., pp:314-325. Churchill Livingstone, New York.

- **Colle J.G., and Marrow (1996):**

Specimen collection, culture containers and media. In: Collee JG, Fraser A.G, Marmion B.P, Simmons A (eds.): Mackie and Mac cartney practical Medical microbiology. New York: Churchill livingstone, pp:95-111.

- **Da Silva O., Ohlsson A., and Kenyon C (1995):**

Accuracy of leukocyte indices and C-reactive protein for diagnosis of neonatal sepsis: a critical review. Pediatr Infect Dis J; 14:362 -6.

- **DeFilippes F. M (1991):**

Decontaminating the polymerase chain reaction. BioTechniques 10:26-30.

- **Dennis A. Benson., Ilene Karsch-Mizrachi., David J. Lipman., James Ostell ., and David L. Wheeler* (2008):**

GenBank. Nucleic Acids Research, 36:25-30.

- **Edberg S. C (1985):**

Principles of nucleic acid hybridization and comparison with monoclonal antibody technology for the diagnosis of infectious diseases. Yale J. Biol. Med. **58**:425–442.

- **Edwards J.R., Ruparel H., and Ju J (2005):**

Mass-spectrometry DNA sequencing. Mutation Research **573** (1-2): 3–12.

- **Edwards M.S., and Baker C.J (2004) :**

Sepsis in the newborn. In Krugman's Infectious Disease of Children, 11th ed :545-61. Edited by Gershon A.A., Hotez P.J., Katz S.L. By Mosby.

- **Eisenstein B.I (1990):**

New molecular techniques for microbial epidemiology and the diagnosis of infectious diseases J.I.D/April, 161:595-602.

- **El-badrawy A., El-Sebaie D., Khairat S., and Shible S.F (2003):**

A study of microbiological pattern of neonatal sepsis .M.Sc.Thesis.

- **El-Maraghy S., Iskander I., Abdel.Ghani H., and Bazzi R.A (1998):**

Value of Scd14 levels as a diagnostic and prognostic tool in neonatal sepsis
, M.Sc.Thesis.

- **Erlich H.A., David. Gelf., and Sninsky J.J (1991):**

Recent advances in the polymerase chain reaction. Science.252:1643-1651.

- **Eskandar R.M., Badrawi N.H., El-Attar S., and El-Falaky N.M (2003):**

Epidemiological study of neonatal sepsis in a neonatal intensive care unit
M.D.Thesis.

- **Faden H.S (1976):**

Early diagnosis of neonatal bacteremia by buffy-coat examination. J Pediatr; (6): 1032-4.

- **Flenday V., Woodgate P., Street P (2004):**

Intramuscular penicillin for the prevention of early onset group B streptococcal infection in newborn infants. Cochrane Database System Rev; (3):CD 003667.

- **Foca M., Jakob K., and Whittier S (2000):**

Endemic *Pseudomonas aeruginosa* Infection in a Neonatal Intensive Care unit. The New England Journal of Medicine; 343(10):695-700.

- **Fong Y., Marano M.A., and Moldawer L.L (1990):**

The acute splanchnic and peripheral tissue metabolic response to endotoxin in humans .J Clinical Invest; 85:1896-1904.

- **Franz A.R., Steinbach G., and Kron M (1999):**

Reduction of unnecessary antibiotic therapy in newborn infants using interleukin -8 and C-reactive protein as markers of bacterial infections. Pediatrics; 104:447-53.

- **Frederick S.Nolte., Janis M.Williams., Robert C.Jearis., Josephine A.Morello., Cindy D.Leitch., Scott Matushek., Lynne D.Schwabe., Francis Dorigan., and Frank E.Kocka (1993):**

Multicenter Clinical Evaluation of a Continuous Monitoring Blood Culture System Using Fluorescent-Sensor Technology (BACTEC 9240) Journal of Clinical Microbiology; 31:552-557.

- **Freedman J., Goldmann D.A., and Smith N.E (1990) :**

Association of intravenous lipid emulsion and coagulase –negative staphylococcal bacteremia in neonatal intensive care units. N Engl J Med; 323:301-8.

- **Freij B.J (2004):**

Management of neonatal herpes simplex virus infections. Indian J Pediatr; 71(10):921-6.

- **Freij B.J., and McCracken G.H (1999):**

Acute infections. In the textbook of neonatology: Pathophysiology and Management of the newborn, 8th ed : 1189-230 .Edited by Avery G.B., Fletcher M.A.,and MacDonald M.G.By Lippincott Williams and Wilkins.

- **Frobes B.A., Sahm D.F., and Weissfeld A.S (1998):**

Gram Negative bacilli and coccobacilli, MacConkey positive, oxidase negative.In: Frobes B.A,Sham D.F,and Weissfeld A.S(eds):Bailey and Scotts Diagnostic microbiology. Missouri: Mosby. Inc,pp501-527.

- **Gerdes G (2004):**

Diagnosis and management of bacterial infections in the neonates. Ped. Clin.N.Am; 51:939-59.

- **Gilbert W., and Maxam A (1973):**

[The nucleotide sequence of the lac operator](#). Proc. Natl. Acad. Sci. U.S.A. 70 (12): 3581–4.

- **Go C.H (2004):**

Escherichia Coli Infections Emedicine Retrieved from <http://www.emedicine.com/med/topic734.htm>.

- **Goldman D.A., Leclair J., and Maccone A (1975):**

Bacterial colonization of neonates admitted to an intensive care environment.J Pediatr; 9:288.

- **Gordon R.C., Barrett F.F., and Clark D.J (1972):**

Influence of several antibiotics, singly and in combination, on the growth of *Listeria monocytogenes*. *J pediatr*, 80:667.

- **Gottorf P.S (1992):**

Infections of the new born. In: Nelson EW, Behrman ER, Kliegman MR and Vaughan CV, editors. *Nelson Textbook of pediatrics*, 14th ed. Philadelphia, WB Saunders, 502.

- **Gottorf P.S (1996):**

Neonatal Sepsis and meningitis. In: *Textbook of pediatrics*, 528-536.

- **Gottorf P .S (2000):**

Infections of the newborn infants. In *Nelson Textbook of pediatrics*, 16th edition :538-52 Edited by Behrman R.E ., Kliegman R.M., and Jenson H.B. By W.B .Saunders.

- **Graca A., Silverio C., and Ferrerira J.P (2004):**

Congenital or neonatal cytomegalovirus infection. *Acta Med Port*; 17(4):335-40.

- **Greisen k., Loeffelholz M., Purohit A., and Leong D (1994):**

PCR primers and probes for the 16S r RNA gene of most species of pathogenic bacteria, including bacteria found in cerebrospinal fluid. *J.Clin. Microbiol.* 32: 335-351.

- **Griffin M.P., and Moorman J.R (2001):**

Towards the early diagnosis of neonatal sepsis and sepsis like illness using novel heart rate analysis. *Pediatrics*; 107:97-104.

- **Griffin M.P., O'Shea T.M., and Bissonette E.A (2003):**

Abnormal heart rate characteristics preceding neonatal sepsis and sepsis like illness. *Pediatr. Res*; 53:920-26.

- **Guerina N.G (1992):**

Bacterial and fungal infection In: Cloherty JP and Stark AR (edit), manual of neonatal care. 3rd Ed. Little and Brown Company, Boston; p146.

Guerina N.G (1998):

Infections: Bacterial and fungal Infections. In Manual of Neonatal Care 4th ed : 239-327. Edited by Cloherty J.P.and StarkA.R.By Little, Brown and Co.

- **Guibourdenche J., Bedu A., and Petzold L (2002):**

Biochemical markers of neonatal sepsis : value of procalcitonin in the emergency setting. Ann Clin Biochem; 39:130-5.

- **Guyer R. L., and Koshland D. E (1989):**

The molecule of the year. Science 246:1543-1546.

- **Hall N (2007):**

Advanced sequencing technologies and their wider impact in microbiology. J. Exp. Biol. **210** (Pt 9): 1518–25.

- **Hanna G.J., Johnson V.A., and Kuritzkes D.R (2000):**

[Comparison of sequencing by hybridization and cycle sequencing for genotyping of human immunodeficiency virus type 1 reverse transcriptase](#). J. Clin. Microbiol. **38** (7): 2715–21.

- **Hansen T and corbet A (1998):**

Neonatal pneumonias.In: Tauesch HW and Ballard RA, editors. Avertys Diseases of the Newborn 7th ed .WB Saunders; 648-658.

- **Haque K.N., Khan M.A., and Kerry S (2004):**

Pattern of culture-proven neonatal sepsis in a district general hospital in the united kingdom.Infect control Hosp Epidemiol; 25(9):759-64.

- **Hashim M.E., El.Abid N., Abdel-Ghani H., and Fahmy S.S (2000):**

The value of estimation of the complement level in neonatal sepsis.
M.Sc.Thesis.

- **Hatzidaki E., and Gourgiotis D (2005):**

Interleukin-6 in preterm premature rupture of membranes as an indicator of neonatal outcome. Acta Obstet Gynecol Scand; 84:632-8.

- **Hickey S.M., and McCracken J.R (1997):**

Interleukin-6 in preterm premature rupture of membranes as an indicator of neonatal outcome. Acta Obstet Gynecol Scand; 84:632-8.

- **Hollenberg S.M., Cunnion R.E., and Parillo J.E (1991):**

The effect of tumor necrosis factor on vascular smooth muscle.Chest; 100:1133-1137.

- **Hosnpud J.D (1993):**

The effects of interleukin-1 on myocardial function and metabolism.Clin Immunol Immunopathol; 68:175-180.

- **Huang E.T.S., Chu Y.W., and Leung C.M (2001) :**

Epidemiology and Infection Control Implications of Acinetobacter spp. In Hong Kong. J Clin Microbiol, 39(1): 228-34.

- **Huang Y.S., Wang S.M., Liu C.C., and Yang Y.J (2002) :**

Invasive Escherichia coli infection in infancy: clinical manifestation, outcome, and antimicrobial susceptibility. J Microbiol Immunol Infect; 35:103-8.

- **Hudson L (1995):**

New therapies for ARDS, Chest, 108suppl.1:79S-91S.

- **Hyde T.B., Hilger T.M., Reingold A., Farley M.M., O'Brien K.L., and Schuchat A (2002):**

Active Bacterial Core Surveillance (ABCs) of the Emerging Infections Program Network. Trends in incidence and antimicrobial resistance of early-onset sepsis: population-based surveillance in San Francisco and Atlanta. *Pediatrics*.; 110: 690-695.

- **Innis M. A (1989):**

PCR protocols: a guide to methods and applications. Academic Press, Inc., San: Diego, Calif.

- **Innis M.A., Gelfand D.H., Sninsky J.J., and Whit T.J (1990):**

PCR Protocols :Aguide to methods and applications .Academic Press, SanDiego, Calif.

- **Isscs D (2003):**

A ten years multi centre study of coagulase negative staphylococcal infection in Australasian neonatal units. *Arch Dis Child Fetal Neonatal* ed; 88 :P89-93.

- **John Wiley., and Sons Ltd (2005):**

Essential Microbiology Textbook. Great Britain by Antony Rowe, Ltd, Chippenham, Wiltshire; P79:89.

- **Jon Jonasson., Margaretha Olofsson., and Hans- JugMonstein (2002):**

Classification, Identification and subtyping of bacteria based on pyro - sequencing and signature matching of 16S rRNA fragments.*APMIS* 110:263-272.

- **Jones E.M .,mcCulloch S.Y., and Reeves D.S (1994):**

A 10 year survey of the epidemiology and clinical aspects of listeriosis in a provincial English City. *J Infect*; 29:91.

- **Jordan J.A (1996):**

Identification of human parvovirus B19 infection in idiopathic non immune hydrops fetalis. Am.J.Obstet.Gynecol. 174:37-42.

- **Jordan .Jeanne. A., Butchko .Allyson. R., and Durso Mary .Benth (2005):**

Use of Pyrosequencing of 16S r RNA Fragments to Differentiate between Bacteria Responsible for Neonatal Sepsis. Journal of Molecular Diagnostic, 7:105-110.

- **Jordan J. A., Laughner Jones., and Durso M. B (2009):**
Utility of Pyrosequencing in Identifying Bacteria Directly from Positive Blood Culture Bottles. J. Clin Microb; 47: p 368–372.

- **Jordan J.A., and Durso M.B (2000):**

Comparison of 16S r RNA Gene PCR and BACTEC 9240 for Detection of Neonatal Bacteremia. Journal of Clinical Microbiology, 38:2574-2578.

- **Julou-Schaeffer G., Gray G.A., Fleming I., Schott C., Parratt J.R., and Stoclet J.C (1990):**

Loss of vascular responsiveness induced by endotoxin involves L-arginine pathway. AM J Physiol; 259:H1038-H1043.

- **Kaloyanides G.J (1994):**

Antibiotic –released nephrotoxicity. Nephrol Dial Transplant; 9:130-134.

- **Kanus W.A., Sun X., Hakin R.B., and Wanger D.R (1994):**

Evaluation of definition for adult respiratory distress syndrome. AmJ Respir Crit Care Med; 150:311-317.

- **Karlowicz M.G., Buescher E.S., and Surka A.E (2000):**

Fulminant late onset sepsis in a neonatal intensive care unit, 1988-1997, and the impact of avoiding empiric vancomycin therapy. Pediatrics 106:1387-1390.

- **Karthikeyan G., and Premkumar K.N (2001):**

Neonatal Sepsis: Staphylococcus aureus as the predominant pathogen.
Indian J Pediatr; 68(8):715-7.

- **Kaufman D., and Fairchild K.D (2004) :**

Clinical Microbiology of Bacterial and Fungal Sepsis in Very –Low –
Birth –Weight Infants. Clin Microbiol Rev; 17(3):638-80.

- **Kawai J., Shinagawa A., Shibata K., Yoshino M., Itoh M., Ishii Y.,
Arakawa T., Hara A., and Fukunishi Y (2001):**

Functional annotation of a full-length mouse cDNA collection. Nature,
409, 685–690.

- **Kawczynski P.P., and Piotrowski A (2004):**

Procalcitonin and C-reactive protein as a marker of neonatal
sepsis .Ginekol Pol; 75(6):439-44.

- **Kellogg J.A., Ferrentino F.L., and Goodstein M.H (1997):**

Frequency of low level bacteremia in infants from birth to two months of
age. Pediatr Infect Dis J; 16(4):381-5.

- **Kelly C.J., and Neilson E.G (1996):**

Tubulointerstitial disease. In: Brenner EM, editor. The kidney, 5th ed
philadelphia:Saunders, pp1655-1679.

- **Kenon C., Overturf G., and Bessman S (1996):**

Granulocyte colony –stimulating factor as a marker for bacterial infection
in neonates.J Pediatr; 128:765-9.

- **Keown B.J (1994):**

An acetylated (Nucleas free) bovine serum albumin in a PCR buffer
inhibits amplification.Biotechniques.17 (2):246-248.

- **Khudyakov Y.E., Gaur I., Singh J., Patel P., and Fields H.A (1994):**

Primer specific solid phase detection of PCR products. Nucleic Acid Res. 22(7):1320-1321.

- **Kilani R.A., and Basamad M (2000):**

Pattern of proven bacterial sepsis in a neonatal intensive care unit in Ritadh-Saudi Arabia: a 2-years analysis. J Med Liban; 48(2):77-83.

- **Kimberlin D.W (2002):**

Meningitis in the neonate. Curr Treat Options Neurol; 4(3):239-48.

- **Kimura K., Suzuki S., Wachimo J., and Kurokawa H (2008):**

First molecular characterization of group B streptococci with reduced penicillin susceptibility Antimicrobial Agents and chemotherapy. August.p.2890-2897. vol.52.No.8.

- **Kite P., Millar M.R., and Gorham P (1988):**

Comparison of five tests used in diagnosis of neonatal bacteremia. Arch Dis Child; 63:639.

- **Klein J.O (2001):**

Bacterial sepsis and meningitis. In Infectious Diseases of the fetus and newborn infant 5th Ed: 943-98. Edited by Remington J .S and Klein J.O. By W.B. Saunders.

- **Koehler W (2002) :**

Web page change and persistence - a four-year longitudinal study. J. Am. Soc. Inform. Sci. Technol., 53, 162–171.

- **Koneman E.W., Allen S. D., Janda W .M., Schreckenberger R. C., and Winn W.C (1997):**

CHARTS. In: Koneman E W, Allens S D, Janda W M, Shreckenberger R C and Winn W C, eds.pp:1295-1395.Lippincot, Philadelphia.

- **Kumar A., Thota V., Dee L., Olsen J., Uretz E., and Parillo J.E (1996):**

Tumour necrosis factor-and interleukin-1 β are responsible for in vitro myocardial cell depression induced by human septic shock serum. JExp Med., 183: 949-958.

- **Kwok S., and Higuchi R (1989):**

Avoiding false positives with PCR. Nature. 339:237-238.

- **Ladhani S., Konana O.S., Mwarumba S., and English M.C (2004):**

Bacteremia due to Staphylococcus aureus .Arch Dis Child; 89: 568-71.

- **Laforgia N., Coppola B., Carbone R., Grassi A., Mautone A., and Iolascon A (1997):**

Rapid detection of neonatal sepsis using polymerase chain reaction. Acta Pediatr 86:1097-1099.

- **Laurent T., Markert M., and Von Flidner V (1994):**

CD11b/CD18 expression, adherence, and chemotaxis of granulocyte in adult respiratory distress syndrome. Am J Respir Crit Care Med; 149: 1534-1538.

- **Lee H.K., Lee W.G., and Cho S.R (2000):**

Evaluation of different blood culture media in neonatal sepsis. J Microbiol Immunol Infect; 33:165-8.

- **Levy I., Rubin L.G., Vasishtha S., Tucci V., and Sood S.K (1998) :**

Emergence of candida parapsilosis as the predominant species causing candidemia in children.Clin Infect Dis; 26:1086-8.

- **Levy R.D., Hindley T., Burman R., Halder S.A., and Blease J (1995):**

Unusual cause of pseudomonal infection. Br Med J; 310:258.

- **Li J.X., Oliver J.R., La C.Y., and Phillips J.B (1995):**

Delayed thromboxane tumour necrosis factor- α but not leukotriene inhibition, attenuates prolonged pulmonary hypertension in endotoxaemia. Am J Med Sci (b); 310:103-110.

- **Linda E.A., Elie C.M., and Edman J.C (2000):**

Medical mycology. In medical microbiology. Edited Brook, G.F., Butel, J.S., Ornston, L.N. Twentieth edition. Copy – right by Appleton & Lange chapter 54 page 55 – 556.

- **Longo M. C., Berninger M. S., and Hartley J. L (1990):**

Use of uracil DNA glycosylase to control carry-over contamination in polymerase chain reaction. Gene 93:125-128.

- **Lucey D., Dolan M. J., Moss C. W., Garcia M., Hollis D., Wegner G.S., Morgan G., Almeida R., Leong D., and Greisen K. S (1992):**

Relapsing illness due to *Rochalimaea henselae* in immunocompetent hosts: implication for therapy and new epidemiological associations. Clin. Infect. Dis. 14:683–688.

- **Lynn D.Schwabe., Richard B.Thomson.JR,KK.Flint., and Frank P.Koontz (1995):**

Evaluation of BACTEC 9240 Blood Culture System by Using High-Volume aerobic Resin Media. Journal Of Clinical Microbiology, 33:2451-2453.

- **Lynn R.Burton., Gwendolyn R.Wilson ., and Paul G.Engelkirk (2001):**

Microbiology for the health science Textbook 7th ed, Awoller Kluwer Company, Philadelphia, 2001; p 342.

- **Madaminov M.S., Mamatkulov I.Kh., and Akhmedova M.D (2003):**
Plasmid profile of common neonatal sepsis causative agents. Like Sprava; (5-6): 46-8.
- **Madoff L.C., Paoletti L.C., and Tai J.Y (1994):**
Maternal immunization of mice with group B streptococcal type III polysaccharide beta C protein conjugate elicits protective antibody to multiple serotypes J Clin Invest; 94:286.
- **Maggadia R., Melvin P., and Weinstein M (2001):**
Laboratory diagnosis of bacteremia and fungemia. Infect Dis Clin N Amer; 15 (4).
- **Mahmood A., Karamat K.A., and Butt T (2002) :**
Neonatal sepsis: high antibiotic resistance of the bacterial pathogens in a neonatal intensive care unit in Karachi. J Pak Med Assoc; 52(8):348-50.
- **Main E.K., and Slagle T (2000):**
Prevention of early-onset invasive neonatal group B streptococcal disease in a private hospital setting: the superiority of culture-based protocols. Am J Obstet Gynecol.; 182:1344-1354.
- **Makhoul R.I., Smolkin T., Sujov P., Kassis I., Tamir A., Shalginov R., and Sprecher H (2005):**
PCR –Based diagnosis of neonatal staphylococcal bacteremias. Clinical microbiological J.P. 4823-4825.
- **Mammen E.F (1998):**
The haematological manifestations of sepsis. J Antimicrob Chemother; 431 supp.A: 17-24.
- **Manucha V., Rusia U., Sikka M., Faridia M., and Madan N (2005):**

Haematological parameters and C – reactive protein in the detection of neonatal sepsis .J.Pediatr. Child Health; 38:459- 464.

- **Margulies M., Egholm M., Altman W.E (2005):**

["Genome sequencing in microfabricated high-density picolitre reactors"](#).
Nature 437 (7057): 376–80.

- **Marks J.D., Maks C.B., and Luee J.M (1990):**

Plasma tumour necrosis factor in patients with septic shock: mortality rate, incidence of adult respiratory distress syndrome, and effect of methylprednisolone administration. Am Rev Respir Dis; 141: 94-91.

- **Mart – Leblond A., Grirnaud L., and David S (2004):**

Evaluation of a rapid Immunochromatographie assay for identification of Candida albicans and Candida dubliniensis. J.Clin.Microbiol.42 (11):4956 – 4960.

- **Matuschak G.M (1997):**

Liver dysfunction in sepsis .In:Fein AM,Abraham EM,Balk RA,Bernard GR, Bone RC, Dantzer DR and Fink MP,editors .Sepsis and Multiorgan failure .Baltimore:William and Wilkins,p.168-188.

- **Mc Cord J.M., Gao B., Leff J., and Flores S.C (1994):**

Neutrophil-generated free radicals: possible mechanisms of injury in adult respiratory distress syndrome. Environ Health Perspect; 10:57-60.

- **Mc Kenn T.M (1990) :**

Prolonged exposure of rat aorta to low levels of endotoxin in vitro results in impaired contractility .J Clin Invest;86:160-168.

- **McDonald L.C., Walker M., Carson L., and Arduino M (1998) :**

Outbreak of Acinetobacter spp.blood stream infections in a nursery association with contaminated aerosols and air conditioners .Pediatr Infect Dis J; 17(8):716-22.

- **Mizock B.A., Sabelli H.C., and Dubin A (1990):**

Septic encephalopathy: Evidence for altered phenylalanine metabolism and comparison with hepatic encephalopathy. Arch Intern Med; 150:443-449.

- **Mohsen L., Hagraas A.M., Khalaf-Allah O., and Aziz N.S (2001):**

Interleukin-1 receptor antagonist in neonatal sepsis .M.S.c.Thesis.

- **Monstein H.J., Kihlström E., and Tiveljung A (1996):**

Detection and identification of bacteria using in- house broad range 16S rDNA PCR amplification and genus-specific DNA hybridization probes, located within variable regions of 16S r RNA genes. APMIS; 104:451–8.

- **Morgan .Almeida R., Leong D., and Greisen K. S (1992):**

Relapsing illness due to Rochalimaea henselae in immunocompetent hosts: implication for therapy and new epidemiological associations. Clin. Infect. Dis. 14:683–688.

- **Morira M.E.L (2005):**

Controversies about the management of invasive fungal infections in very low birth weight infants. J Pediatr (Rio J); 81(1 Suppl):S52-8.

- **Morita J.Y.O., Brien K.L., and Schuchat A (1999):**

Prevention of perinatal group B streptococcal infections. Pediatr Infect Dis J; 18:279-180.

- **Mullis K. B (1990):**

The unusual origin of the polymerase chain reaction. Sci. Am. 240:56-65.

- **Mullis K.B and Faloona F.A (1987):**

Specific synthesis of DNA in vitro via a polymerase catalysed chain reaction. In methods in enzymology (R.wv.ed) Academic Press, Sandiego.Calif.,Vol ,155 , pp.335-350.

- **Mullis K.B., Faloona F.A., Scharf., and Saiki S (1986):**

Specific enzymatic amplification of DNA in vitro: Polymerase chain reaction .Cold Spring Har bar symp.Quant.Biol.51:263-273.

- **Myers J.P., and Linnemann C.C (1982) :**

Bacteremia due to methicillin –resistant staphylococcus aureus. J Infect Dis, 145:532.

- **Ng P.C (2004):**

Diagnostic markers of infection in neonates. Arch Dis Child Fetal Neonatal Ed; 89:F229.

- **Ng P.C., Lee C.H., and Fok T.F (2000):**

Central nervous system candidiasis in preterm infants: limited value of biochemical markers for diagnosis. J Pediatr Child Health; 36:509 – 10.

- **Niveleau A., Drouet .Reynoud C., Fores F., Bruno C., and Pain C (1994):**

Polymerase chain reaction products containing 5-methyldeoxy cytidine; microplate immuno-quantitation method. Nucleic Acid Res. 22(24):5508-5509.

- **Noya F.J.D., and Baker C.K (1992):**

Prevention of group β streptococcal infection. Pediatr Infect; 6-41.

- **Nupponen I., Andersson S., and Jarvenpaa A.L (2001):**

Neutrophil CD11b expression and circulating interleukin-8 as diagnostic markers for early –onset neonatal sepsis .Pediatrics; 108:e12-7.

- **Olsen G. J., and Woese C. R (1993):**

Ribosomal RNA: a key to phylogeny. FASEB J. 7:113–123.

- **Osterhout G.J., Shull V.H., and Dick J.D (1991):**

Identification of clinical isolates of gram negative non fermentative bacteria by an automated cellular fatty acid identification system. J.Clin. Microbiol. 29: 1822 -1830.

- **Parsons G (1988):**

Development of DNA probe-based commercial assays. J. Clin. Immunoassay 11:152-158.

- **Pastor C.M., Billiar T.R., and Payen D.M (1995):**

Liver injury during sepsis, J Crit Care; 183-197.

- **Persing D. H (1991):**

Polymerase chain reactions: trenches to benches, 1991. J. Clin. Microbiol. 29:1281-1285.

- **Petros A., Lamb G., and Leane A (1994):**

Effects of nitric oxide synthase inhibitor in humans with septic shock. Cardiovasc Res; 28:34-39.

- **Philip A.G.S., and Hewitt J.R (1980):**

Early diagnosis of neonatal sepsis. Ped; 65:1036-1041.

- **Phillips E.A., Tapsall J.W., and Smith D.D (1980):**

[Rapid tube CAMP test for identification of Streptococcus agalactiae \(Lancefield group B\).](#) J.Clin. Microbiol. 12 (2): 135–7.

- **Pourcyrous M., Bada H.S., and Korones S.B (1993):**

Significance of serial C-reactive protein responses in neonatal infection and other disorders. *Pediatrics*; 92:431-5.

- **Powers D.L., and Mandell G.L (1974):**

Intraleucocytic bacteria in endocarditis patients. *JAMA*; 227-312.

- **Prins J.M., Buller H.R., and Kuijper E.J (1993):**

Once versus thrice daily gentamycin in patients with serious infections. *Lancet* 341:335-339.

- **Ralph D., Stuart L., and Morven S (1987):**

Infection in the fetus and neonatal and perinatal medicine. By the C.V Mosby CO.St.Louis(4th ed); p765-794.

- **Rebecca Clayton., Granger. Sutton., Paul S.Hinkle., Carol Bult., and Cherisields (1995):**

Intraspecific Variation in Small Subunit r RNA sequences in GenBank. Why single Sequences May Not Adequately Represent Prokaryotic Taxa. *International Journal of Systematic Bacteriology*, 45:595-599.

- **Relman D.A (1993):**

The identification of uncultured microbial pathogens. *J. Infect. Dis.* **168**:1–8.

- **Relman D. A., Loutit J. S., Schmidt T. M., Falkow S., and Tompkins L. S (1990):**

The agent of bacillary angiomatosis. An approach to the identification of uncultured pathogens. *N. Engl. J. Med.* 323:1573–1580.

- **Rennie J.M., and Robertson N.R (2002) :**

Infection. In *A Manual of Neonatal Intensive Care* 4th ed : 223-67. By Arnold, a member of the Hodder Headline Group.

- **Richard T., Boutin P., Arnould C., and Jetin A (1995):**

Efficient production of internal standard DNA for quantitative PCR using an automated sequencer. *Biotechniques*:18(2):212-213.

- **Rodwell R.L., Leslie A.L., and Tudehope D.L (1989):**

Evaluation of direct and buffy coat films of peripheral blood for the early detection of bacteremia. *Aust Pediatr J*; 25:83.

- **Ruth A., Lea R.F., Shaunte C.J., Bobby L.B., Trang T.T., and Jones V (2007):**

DNA pyrosequencing –Based bacterial pathogen identification in a pediatric hospital setting, *J.Clin.Microbiol.*, 45(9):2985-2992.

- **Ruvalo C., and Bauer C.R (1982) :**

Intrauterine acquired *Pseudomonas* infection in the neonate. *Clin Pediatr*; 21:664-7.

- **Saez-Lioren X., and McCracken G.H (1993):**

Sepsis syndrome and septic shock in pediatrics: Current concepts of terminology, pathophysiology and management. *J pediatr*, 123-497.

- **Saez-Lioren X., and McCracken G.H (1998):**

Perinatal bacterial disease. In textbook of Pediatric Infectious disease. 4th ed. p897. Edited by Feigin R.D. AND By W.B.Saunders.

- **Saiki R.K., Chang C.A., and Levenson C.H (1988):**

Diagnosis of sickle cell anemia and β -thalassemia with enzymatically amplified DNA and non radioactive allele –specific oligonucleotide probes.

- **Saiki R.K., Scharf S.S., Faloona F., Mullis K.B., Horne H.G., Erlich H.A and, Arnheim (1985):**

Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anaemia. Science, 230:1350-1354.

- **Saiki R.K., Walsh D.S., and Erlich H.A (1989):**

Genetic analysis of amplified DNA with immobilized sequence –specific oligonucleotide probes. Proc .Natl.Acad .Sci.U.S.A.86:6230-6234.

- **Sanger F (1980):**

[Determination of nucleotide sequences in DNA](#). Nobel lecture, 8 December 1980.

- **Sanger F., Nicklen S., and Coulson A.R (1977):**

[DNA sequencing with chain-terminating inhibitors](#). Proc. Natl. Acad. Sci. U.S.A. **74** (12): 5463–7.

- **Sarkar G., and Sommers S (1990):**

Shedding light on PCR contamination.Nature. (London):343:27.

- **Sarkar G., and Sommer S (1991):**

Parameters affecting susceptibility of PCR contamination to UV inactivation. Bio-Techniques 10:590-594.

- **Schmidt T. M., and Relman D. A (1994):**

Phylogenetic identification of uncultured pathogens using ribosomal RNA sequences. Methods Enzymol. **235**:205–222.

- **Schrag S., Gorwitz R., Fultz-Butts K., and Schuchat A (2002):**

Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. MMWR Recommendations Rep; 51(RR-11):1-22.

- **Selby D.M., Gautier G., and Luban N.L (1990):**

Overwhelming neonatal septicemia diagnosed upon examination of peripheral blood smears .Clin Pediatr; 29:706.

- **Seoud I., Ghobashi M., Iskander I., and Omar M (2001) :**

Nosocomial infection in Kasr El –Aini neonatal intensive care unit as a university hospital .M Sc.Thesis .

- **Shange S., Chen Z., and Yu X (2001):**

Detection of bacterial DNA by PCR and reverse hybridization in the 16S r RNA gene with particular reference to neonatal septicemia. Acta paediatr, 90:179-83.

- **Shaw C.K.I., Shaw P.I., and Thapalia A (2007):**

Neonatal sepsis bacterial isolates and antibiotic susceptibility patterns at a NICU in a tertiary care hospital in western Nepal: A retrospective analysis. Kathmandu University Medical Journal; 5:153-160.

- **Shendure J., Porreca G.J., and Reppas N.B (2005):**

Accurate multiplex polony sequencing of an evolved bacterial genome. Science **309** (5741): 1728–32.

- **Shenoy S., Antony G., and Shenoy U.V (2000) :**

Value of superficial cultures in diagnosing neonatal sepsis .Indian J Pediatr; 7(5):337-8.

- **Shinefield H.R., and Geme J.W (2001) :**

Staphylococcal Infections.In Infectious Diseases of the fetus and Newborn Infant ed: 1217-47.Edited by Remington J.S and Klein .J.O.By W.B.Saunders.

- **Smith L.M., Sanders J.Z., and Kaiser R.J (1986):**

Fluorescence detection in automated DNA sequence analysis. Nature **321** (6071): 674–9.

- **Smith M.W., Holmsen A.L., Wei Y.H., Peterson M., and Evans G.A (1994):**

Genomic sequence sampling: a strategy for high resolution sequence-based physical mapping of complex genomes. *Nat. Genet.*, 7, 40–47.

- **Solorzano-Sontos F., Miranda-Navales M.G., and Leanos-Miranda B (1999):**

A blood microculture system for the diagnosis of bacteremia in pediatric patients. *Scand J Infect Dis*; 30:481-3.

- **Stoker N.G (1990):**

The polymerase chain reaction and infectious disease: hopes and realities. *Trans.Roy.Soc . Trop.Med.Hyg*; 84:755.

- **Stoll B.J (2004):**

Infections of neonatal infant .In *Nelson Textbook of Pediatrics*, 17th ed: 623-40. Edited by Behrman R.E., Kliegman R.M., AND Jenson H.B .By W.B.Saunders.

- **Stoll B.J., Hansen N., and Fanaroff A.A (2002):**

Late-onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network. *Pediatrics*; 110(2 Pt 1):285-291.

- **Stoll B.J., Holman R.C., Schuchat A (1998):**

Decline in sepsis-associated neonatal and infant deaths in the United States, 1979 through 1994. *Pediatrics*.;102:e18.

- **Storm W (1981):**

Early detection of bacteremia by peripheral smears in critically ill newborns. *Acta Pediatr Scand*; 70:415.

- **Susan L. Lukacs., Kenneth C. Schoendorf., and Anne Schuchat (2004):**

Trends in Sepsis-Related Neonatal Mortality in the United States. J Pediatr Infect Dis 23(7):599-603, 2004.

- **Tan T.O., Mason J.M., and Shulman R.J (1993):**

Molecular epidemiology of Coagulase-negative Staphylococcus blood isolates from neonates with persistent bacteremia and children with central venous catheter infection. J Infect Dis; 169:1393.

- **Tan T.O., Musser J.M., and Shulman R.J (1994):**

Molecular epidemiology of Coagulase-negative Staphylococcus blood isolates from neonates with persistent bacteremia and children with central venous catheter infections .J Infect Dis; 169:1393.

- **Tang YI-Wei., Nicole M .Ellis., Marlene K.Hopkins., Douglas H.Smith., Deborah E.Dodge., and David H.Persing (1998) :**

Comparison of Phenotypic and Genotypic Techniques for Identification of unusual aerobic pathogenic gram negative bacilli. J Clin Microbiology. P.3674-3679.

- **Tarkin I.S., Henry T.J., and Fey P.I (2003):**

PCR rapidly detects methicillin-resistant staphylococci periprosthetic infection. J.Clin Orthop; 89-94.

- **Tirodker U.H., Nataro J.P., and Smith S (2003):**

Detection of fungemia by polymerase chain reaction in critically ill neonates and children. J. Perinatol; 23(2):117-22.

- **Tomashefski J.F (1990):**

Pulmonary pathology of the adult respiratory distress syndrome .J. Clin Chest Med; 1:593-619.

- **Townsend S.M., Pollack H.A., and Gonzalez-Gomez I (2003) :**

Citrobacter koseri Brain Abscess in the Neonatal Rat: Survival and Replication within Human and Rat Macrophages. J. Infection and Immunity; 71(10):5871-80.

- **Ueda K., Seki T., Kuda T., Yoshida T., and Kataoka M (1999):**
Two distinct mechanisms cause heterogeneity of 16S r RNA .J Bacteriol ;181 :78-81.
- **Valerie Cilia., Benedit Lafay., and Richard Christen (1996):**
Sequence Hetrogeneities Among 16S Ribosomal RNA Sequences, and Their Effect on Phylogenetic Analyses at the Species Level. J.Molecular Biology Evol.13:451-461.
- **Vigushin D., Pepys M., and Hawkins P (1993):**
Metabolic and scintigraphic studies of radioiodinated human C-reactive protein in health and disease .J Clin Invest; 91:1351-7.
- **Villanueva Uy.M.E., Briones C.R., and H.G.Uy (2003):**
Application of polymerase chain reaction in late onset neonatal sepsis. Pediatr. Res.53:313A.
- **Villari P., Sarnataro C., and Iacuzio L (2000):**
Molecular epidemiology of Staphylococcus epidermidis in a neonatal intensive care unit over a three –year period. J Clin Microbiol; 38:1740-6.
- **Vincent J.I (1998):**
Cardiovascular alteration in sepsis .J Antimicrob Chemother; 41 suppl A: 9-15.
- **Visintine A.M., Oleske J.M., and Nahmias A.J (1977):**
Listeria monocytogenes infection in infants and children. AM J Dis Child, 131:393.

- **Visser V.E., and Hall R.T (1980):**
Lumbar puncture in the evaluation of neonatal sepsis. J Pediatrics ; 96: 1063-1067.

- **Volante E., Moretti S., Pisani E., and Bevilacqua G (2004):**
Early diagnosis of bacterial infection in the neonate.J Matern Fetal Neonatal Med; 16 suppl2:13-6.

- **Wang A.M., Doyle M.V., and Mark D.F (1989):**
Qualitative evaluation of PCR products using a synthetic internal standard. Proc.Natl.Acad.Sci.U.S.A.86 : 9717-9721.

- **Washington J.A., and Ilstrup D.M (1986):**
Blood cultures: issues and controversies .Rev Infect Dis; 8:792.

- **Watt S., Lanotte P., and Mereghetti L (2003):**
Escherichia coli Strains from pregnant women and neonates: Intraspecies Genetic Distribution and prevalence of virulence factors .J Clin Microbiol; 41(5): 1929-35.

- **Weirich E., Rabin R.L., and Maldonado Y (1998):**
Neutrophil CD11b expression as a diagnostic marker for early –onset neonatal infection.J Pediatr; 132:445-51.

- **Wheeler D.L., Barrett T., Benson D.A., Bryant S.H., Canese K., Chetvernin V., Church D.M., DiCuccio M., and Edgar,R (2008):**
Database resources of the National Center for Biotechnology Information. Nucleic Acids Res., This issue (Database issue).

- **White T.J., Madej R., and Presingt D.H (1992):**

The polymerase chain reaction: clinical applications. *Advances in clinical Chemistry*. 29:161-169.

- **Wilson K.H., Blitchington R.B., and Greene R. C (1990):**

Amplification bacterial 16S ribosomal DNA with polymerase chain reaction. *J. Clin.Microbiol.* **28**:1942–1946. (Erratum, *J. Clin. Microbiol.* 29:666, 1991).

- **Woese C. R (1987):**

Bacterial evolution. *Microbiol. Rev.* **51**:221–271.

- **Woo P. C. Y., Teng J. L. L., Wu J. K. L., Leung F. P. S., Tse H., Fung A. M. Y., Lau S. K. P., and Yuen K.y (2009):**

Guidelines for interpretation of 16S rRNA gene sequence-based results for identification of medically important aerobic Gram-positive bacteria. *J Med Microbiol* 58: 1030-1036.

- **Yagupski P., and Nottle F.S (1990):**

Quantitative aspects of septicemia.*Clin.Microbol .Rev.*3:269-279.

- **Zangwill K.M., Schuchat A., and Wenger J.D (1992):**

Group B streptococcal disease in the united states 1990; Report from a multistate active surveillance system.*MMWR Morb Mortal Wkly Rep*; 41:25.

- **Zhang P., Xie M., and Spitzer J.A (1994):**

Hepatic neutrophil sequestration in early sepsis; Enhanced expression of adhesion molecules and phagocytic activity. *Shok*; 2:133-140.

- **Ziadi A.K., Huskins W.C., Thaver D., Bhutta Z.A., Abbas Z., and Goldman (2005):**

Acquired neonatal infections in developing countries. *Lancet*; 365:1175 - 1180.