

## References

- A report of the surgeon general. Physical activity and health.. US DHHS 1996.
- Adam, E, Melnick, JL, Probstfield, JL, et al. High level of cytomegalovirus antibody in patients requiring vascular surgery for atherosclerosis. *Lancet* 1987; 2:291.
- Adams, MR, Jessup, W, Hailstones, et al. L-Arginine reduces human monocyte adhesion to vascular endothelium and endothelial expression of cell adhesion molecules. *Circulation* 1997; 95:662.
- Adler, SP, Hur, JK, Wang, JB, et al. Prior infection with cytomegalovirus is not a major risk factor for angiographically demonstrated coronary artery atherosclerosis. *J Infect Dis* 1998; 177:209.
- Agmon, Y, Khandheria, BK, Meissner, I, et al. Independent association of high blood pressure and aortic atherosclerosis: A population-based study. *Circulation* 2000; 102:2087.
- Al Suwaidi, J, Higano, ST, Holmes, DR Jr, et al. Obesity is independently associated with coronary endothelial dysfunction in patients with normal or mildly diseased coronary arteries. *J Am Coll Cardiol* 2001; 37:1523.
- Ambrose, JA, Tannenbaum, MA, Alexopoulos, D, et al. Angiographic progression of coronary artery disease and the development of myocardial infarction. *J Am Coll Cardiol* 1988; 12:56.
- Anderson, JL, Carlquist, JF, Muhlestein, JB, et al. Evaluation of C-reactive protein, an inflammatory marker, and infectious serology as risk factors for coronary artery disease and myocardial infarction. *J Am Coll Cardiol* 1998; 32:35.
- Anderson, TJ, Meredith, IT, Charbonneau, F, et al. Endothelium-dependent coronary vasomotion relates to the susceptibility of LDL to oxidation in humans. *Circulation* 1996; 93:1647.
- Assmann, G, Shulte, H. Relation of high density lipoprotein cholesterol and triglycerides to instances of atherosclerotic coronary artery disease (The PROCAM Experience). *Am J Cardiol* 1992; 70:733.
- Balletshofer, BM, Rittig, K, Enderle, MD, et al. Endothelial dysfunction is detectable in young normotensive first-degree relatives of subjects

- with type 2 diabetes in association with insulin resistance. *Circulation* 2000; 101:1780.
- Barker, DJ, Osmond, C, Simmonds, SJ, Wield, GA. The relation of small head circumference and thinness at birth to death from cardiovascular disease in adult life. *BMJ* 1993; 306:422.
  - Barrett-Connor E, Slone S, Greendale G *et al.* The Post-menopausal Estrogen/Progestin Intervention Study: primary outcomes in adherent women. *Maturitas* 1997; 27: 261–74.
  - Barrett-Connor, EL, Bush, TL. Estrogen and coronary heart disease in women. *JAMA* 1991; 265:1861.
  - Barrett-Connor, Heart Disease in Women: Ancel Keys Lecture *Circulation* Vol. 95, No. 1 January 7, 1999:252–264.
  - Benetos, A, Zureik, M, Morcet, J, et al. A decrease in diastolic blood pressure combined with an increase in systolic blood pressure is associated with a higher cardiovascular mortality in men. *J Am Coll Cardiol* 2000; 35:673.
  - Benjamin, EJ, Levy, D. Why is left ventricular hypertrophy so predictive of morbidity and mortality?. *Am J Med Sci* 1999; 317:168.
  - Berger, JS, Roncaglioni, MC, Avanzini, F, et al. Aspirin for the primary prevention of cardiovascular events in women and men: a sex-specific meta-analysis of randomized controlled trials. *JAMA* 2006; 295:306.
  - Berliner, JA, Navab, M, Fogelman, AM, et al. Atherosclerosis: Basic mechanisms: oxidation, inflammation and genetics. *Circulation* 1995; 91:2488.
  - Besthesida et al.,: AHA/ACC Task Force Report on Prevention of Coronary Artery Disease. *J Am Coll Cardiol* 2000; 59:132.
  - Bhagat, K, Moss, R, Colier, J, Vallance, P. Endothelial "stunning" following a brief exposure to endotoxin: a mechanism to link infection and infarction?. *Cardiovasc Res* 1996; 32:822.
  - Blair SN. Changes in physical fitness and all cause mortality: a prospective study of healthy and unhealthy men. *JAMA* 1995; 73: 1093–8.

- Blair, SN, Kohl, HW III, Paffenbarger, RS Jr, et al. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 1989; 262:2395.
- Blake, GJ, Dada, N, Fox, JC, et al. A prospective evaluation of lipoprotein-associated phospholipase A(2) levels and the risk of future cardiovascular events in women. *J Am Coll Cardiol* 2001; 38:1302.
- Blankenberg, S, Rupprecht, HJ, Bickel, C, et al. Circulating cell adhesion molecules and death in patients with coronary artery disease. *Circulation* 2001; 104:1336.
- Blankenberg, S, Rupprecht, HJ, Bickel, C, et al. Cytomegalovirus infection with interleukin-6 response predicts cardiac mortality in patients with coronary artery disease. *Circulation* 2001; 103:2915.
- Blankenhorn DH, Hodis HN. George Lyman Duff Memorial Lecture. Arterial imaging and atherosclerosis reversal. *Arterioscler Thromb* 1994;14(2):177–92.
- Blum, A, Giladi, M, Weinberg, M, et al. High anti-cytomegalovirus (CMV) IgG antibody titer is associated with coronary artery disease and may predict post-coronary balloon angioplasty restenosis. *Am J Cardiol* 1998; 81:866.
- Boffetta, P, Garfinkel, L. Alcohol drinking and mortality among men enrolled in an American Cancer Society Prospective Study. *Epidemiology* 1990; 1:342.
- Bosma H, Marmot MG, Hemingway H, Nicholson A, Brunner EJ, Stanfeld S. Low job control and risk of coronary heart disease in the Whitehall II (prospective cohort) study. *Br Med J* 1997; 314: 558–65.
- Boushey, CJ, Beresford, SAS, Omenn, GS, et al. A quantitative assessment of plasma homocysteine as a risk factor for vascular disease: Probable benefits of increasing folic acid intakes. *JAMA* 1995; 274:1049.
- Bowman, TS, Sesso, HD, Ma, J, et al. Cholesterol and the risk of ischemic stroke. *Stroke* 2003; 34:2930.
- Brand, FN, McGee, DL, Kannel, WB, et al. Hyperuricemia as a risk factor of coronary heart disease: The Framingham Study. *Am J Epidemiol* 1985; 121:11.

- Brizzi, MF, Formato, L, Dentelli, P, et al. Interleukin-3 stimulates migration and proliferation of vascular smooth muscle cells: a potential role in atherogenesis. *Circulation* 2001; 103:549.
- Brown KA, Rowen M. Extent of jeopardized viable myocardium determined by myocardial perfusion imaging best predicts perioperative cardiac events in patients undergoing noncardiac surgery. *J Am Coll Cardiol*. 1993;21:325-330.
- Brown, DW, Giles, WH, Croft, JB. Left ventricular hypertrophy as a predictor of coronary heart disease mortality and the effect of hypertension. *Am Heart J* 2000; 140:848.
- Bui, MN, Sack, MN, Moutsatsos, G, et al. Auto-antibody titers to oxidized LDL in patients with coronary atherosclerosis. *Am Heart J* 1996; 131:663.
- Burian, K, Kis, Z, Virok, D, et al. Independent and joint effects of antibodies to human heat-shock protein 60 and Chlamydia pneumoniae infection in the development of coronary atherosclerosis. *Circulation* 2001; 103:1503.
- Burke, AP, Kolodgie, FD, Farb, A, et al. Healed plaque ruptures and sudden coronary death: evidence that subclinical rupture has a role in plaque progression. *Circulation* 2001; 103:934.
- Bush, TL, Cowan, LD, Barrett-Connor, EL, et al. Estrogen use and all-cause mortality. *JAMA* 1983; 249:903.
- Calle, EE, Thun, MJ, Petrelli, JM, et al. Body-mass index and mortality in a prospective cohort of U.S. adults [see comments]. *N Engl J Med* 1999; 341:1097.
- Camargo, CA, Stampfer, MJ, Glynn, RJ, et al. Prospective study of moderate alcohol consumption and risk of peripheral arterial disease in US male physicians. *Circulation* 1997; 95:577.
- Capron, L. Chlamydia in coronary plaques – hidden culprit or harmless hobo. *Nat Med* 1996; 2:856.
- Chesney AM. Social support and heart disease in women. Implications for intervention. In: Orth-Gomer K, Chesney AM, Wenger NK, eds. *Women, Stress and Heart Disease*. Mahwah, New Jersey: Lawrence Erlbaum Associates, Publishers, 1998: 165–84.

- Chobanian, AV, Bakris, GL, Black, HR, Cushman, WC. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report. *JAMA* 2003; 289:2560.
- Clarkson, P, Celermajer, DS, Donald, AE, et al. Impaired vascular reactivity in insulin-dependent diabetes mellitus is related to disease duration and low density lipoprotein cholesterol levels. *J Am Coll Cardiol* 1996; 28:573.
- Clarkson, P, Celermajer, DS, Powe, AJ, et al. Endothelium-dependent dilatation is impaired in young healthy subjects with a family history of premature coronary disease. *Circulation* 1997; 96:3378.
- Collaborative overview of randomised trials of antiplatelet therapy--I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. Antiplatelet Trialists' Collaboration. *BMJ* 1994; 308:81.
- Collins, R, Peto, R, MacMahon, S, et al. Blood pressure, stroke, and coronary heart disease. part 2, short-term reductions in blood pressure: overview of randomized drug trials in their epidemiological context. *Lancet* 1990; 335:827.
- Cooper, R, Cutler, J, Desvigne-Nickens, P, et al. Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: Findings of the National Conference on Cardiovascular Disease Prevention. *Circulation* 2000; 102:3137.
- Criqui, MH, Heiss, G, Cohn, R, et al. Plasma triglyceride level and mortality from coronary heart disease. *N Engl J Med* 1993; 328:1220.
- Culleton, BF, Larson, MG, Kannel, WB, Levy, D. Serum uric acid and risk for cardiovascular disease and death: The Framingham Heart Study. *Ann Intern Med* 1999; 131:7.
- Cybulsky, MI, Iiyama, K, Li, H, et al. A major role for VCAM-1, but not ICAM-1, in early atherosclerosis. *J Clin Invest* 2001; 107:1255.
- Danesh, J, Peto, R. Risk factors for coronary heart disease and infection with *Helicobacter pylori*: Meta-analysis of 18 studies. *BMJ* 1998; 316:1130.

- Danesh, J, Whincup, P, Walker, M, et al. Fibrin d-dimer and coronary heart disease: prospective study and meta-analysis. *Circulation* 2001; 103:2323.
- Danesh, J, Youngman, L, Clark, S, et al. *Helicobacter pylori* infection and early onset myocardial infarction: case-control and sibling pairs study. *BMJ* 1999; 319:1157.
- Dannenberg, AL, Keller, JB, Wilson, PWF, et al. Leisure time physical activity in the Framingham Offspring Study. Description, seasonal variation, and risk factor correlates. *Am J Epidemiol* 1989; 129:76.
- Dart, AM, Lacombe, F, Yeoh, JK, et al. Aortic distensibility in patients with isolated hypercholesterolaemia, coronary artery disease, or cardiac transplant. *Lancet* 1991; 338:270.
- Daugherty, A, Manning, MW, Cassis, LA. Angiotensin II promotes atherosclerotic lesions and aneurysms in apolipoprotein E-deficient mice. *J Clin Invest* 2000; 105:1605.
- David W, Guy De Backer, Ole F, Ian G, Giuseppe M and Kalevi P: Prevention of coronary heart disease in clinical practice; Recommendations of the Second Joint Task Force of European and other Societies on Coronary Prevention. *European Heart Journal* (1998) 19, 1434–1503
- Davies MJ, Woolf, N, Rowles, PM, Pepper, J. Morphology of the endothelium over atherosclerotic plaques in human coronary arteries. *Br Heart J* 1988; 60:459.
- Davis, PH, Dawson, JD, Riley, WA, Lauer, RM. Carotid intimal-medial thickness is related to cardiovascular risk factors measured from childhood through middle age: the Muscatine study. *Circulation* 2001; 104:2815.
- De Bacquer, D, De Backer, G, Kornitzer, M, et al. Prognostic value of ischemic electrocardiographic findings for cardiovascular mortality in men and women. *J Am Coll Cardiol* 1998; 32:680.
- De Bacquer, D, De Backer, G, Kornitzer, M, et al. Prognostic value of ischemic electrocardiographic findings for cardiovascular mortality in men and women. *J Am Coll Cardiol* 1998; 32:680.

- del Sol, AI, Moons, KG, Hollander, M, et al. Is carotid intima-media thickness useful in cardiovascular disease risk assessment? The Rotterdam study. *Stroke* 2001; 32:1532.
- Diez Roux, AV, Merkin, SS, Arnett, D, et al. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med* 2001; 345:99.
- Doll, R, Peto, R, Boreham, J, Sutherland, I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ* 2004; 328:1519.
- Donahue, RP, Abbott, RD, Reed, DM, et al. Alcohol and hemorrhagic stroke: The Honolulu Heart Program. *JAMA* 1986; 255:2311.
- Dong, ZM, Brown, AA, Wagner, DD. Prominent role of P-selectin in the development of advanced atherosclerosis in apoE-deficient mice. *Circulation* 2000; 101:2290.
- Downs, JR, Clearfield, M, Weis, S, et al for the AFCAPS/TexCAPS Research Group. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: Results of AFCAPS/TexCAPS. *JAMA* 1998; 279:1615.
- Duffy, SJ, Biegelsen, ES, Holbrook, M, et al. Iron chelation improves endothelial function in patients with coronary artery disease. *Circulation* 2001; 103:2799.
- Effects of estrogen or estrogen/progestin regimens on heart disease risk factors in postmenopausal women. The Postmenopausal Estrogen/Progestin Interventions (PEPI) Trial. The Writing Group for the PEPI Trial. *JAMA* 1995; 273:199.
- Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. The Trials of Hypertension Prevention, phase II. The Trials of Hypertension Prevention Collaborative Research Group. *Arch Intern Med* 1997; 157:657.
- Eidelman, RS, Hebert, PR, Weisman, SM, Hennekens, CH. An update on aspirin in the primary prevention of cardiovascular disease. *Arch Intern Med* 2003; 163:2006.
- Ekelund LG, Haskell WL, Johnson JL, Whaley FS, Criqui MH, Sheps DS. Physical fitness as a predictor of cardiovascular mortality in

asymptomatic North American men. The Lipid Research Clinics Mortality Follow-up Study. *N Engl J Med* 1988;319(21):1379–84.

- Ekelund, LG, Haskell, WL, Johnson, JL, et al. Physical fitness as a predictor of cardiovascular mortality in asymptomatic North American men. The Lipid Research Clinics Mortality Follow-up Study. *N Engl J Med* 1988; 319:1379.
- El-Khairi, L, Ueland, PM, Refsum, H, et al. Plasma total cysteine as a risk factor for vascular disease: the European Concerted Action Project. *Circulation* 2001; 103:2544.
- Ellison, RC. Cheers! [editorial; comment]. *Epidemiology* 1990; 1:337.
- Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001; 285:2486.
- Falk, E. Plaque rupture with severe pre-existing stenosis precipitating coronary thrombosis: Characteristics of coronary atherosclerotic plaques underlying fatal occlusive thrombi. *Br Heart J* 1983; 50:127.
- Fang, J, Alderman, MH. Serum uric acid and cardiovascular mortality the NHANES I epidemiologic follow-up study, 1971-1992. National Health and Nutrition Examination Survey. *JAMA* 2000; 283:2404.
- Febbraio, M, Hajjar, DP, Silverstein, RL. CD36: a class B scavenger receptor involved in angiogenesis, atherosclerosis, inflammation, and lipid metabolism. *J Clin Invest* 2001; 108:785.
- Final report on the aspirin component of the ongoing Physicians' Health Study. Steering Committee of the Physicians' Health Study Research Group. *N Engl J Med* 1989; 321:129.
- Folsom, AR, Nieto, J, Sorlie, P, et al, for the Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Helicobacter pylori* seropositivity and coronary heart disease incidence. *Circulation* 1998; 98:845.
- Franklin, SS, Khan, SA, Wong, ND, et al. Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham Heart Study. *Circulation* 1999; 100:354.



- Franklin, SS, Larson, MG, Khan, SA, et al. Does the relation of blood pressure to coronary heart disease risk change with aging?: the Framingham Heart Study. *Circulation* 2001; 103:1245.
- Freedman, DS, Williamson, DF, Gunter, EW, Byers, T. Relation of serum uric acid to mortality and ischemic heart disease. The NHANES I Epidemiologic Follow-up Study. *Am J Epidemiol* 1995; 141:637.
- Fuchs, CS, Stampfer, MJ, Colditz, GA, et al. Alcohol consumption and mortality among women. *N Engl J Med* 1995; 332:1245.
- Fung, TT, Stampfer, MJ, Manson, JE, et al. Prospective study of major dietary patterns and stroke risk in women. *Stroke* 2004; 35:2014.
- Fuster V, Pearson TA. 27th Bethesda Conference: Matching the Intensity of Risk Factor Management with the Hazard for Coronary Disease Events. *J Am Coll Cardiol* 1996; 27: 957–1047.
- Fuster, V, Badimon, L, Badimon, JJ, Chesebro, JH. The pathogenesis of coronary artery disease and the acute coronary syndromes. *N Engl J Med* 1992; 326:242.
- Gaede, P, Vedel, P, Larsen, N, et al. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003; 348:383.
- Garcia Rodriguez, LA, Hernandez-Diaz, S, de Abajo, FJ. Association between aspirin and upper gastrointestinal complications: systematic review of epidemiologic studies. *Br J Clin Pharmacol* 2001; 52:563.
- Gardin, JM, McClelland, R, Kitzman, D, et al. M-mode echocardiographic predictors of six- to seven-year incidence of coronary heart disease, stroke, congestive heart failure, and mortality in an elderly cohort (the Cardiovascular Health Study). *Am J Cardiol* 2001; 87:1051.
- Gaw S, Yudkin JS, Blauth C, Drury P *et al.* Prevention and management of cardiovascular disease in patients with diabetes mellitus: an evidence base. *Diabet Med* 2002; 18: T110–32.
- Gawaz, M, Neumann, F-J, Dickfeld, T, et al. Activated platelets induce monocyte chemotactic protein-1 secretion and surface expression of intercellular adhesion molecule-1 on endothelial cells. *Circulation* 1998; 98:1164.

- Genest Jr JJ, Martin-Munley SS, McNamara JR *et al.* Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* 1992; 85: 2025–33.
- Genest, JJ, Martin-Munley, SS, McNamara, JR, et al. Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* 1992; 85:2025.
- Gerstein, HC, Pais, P, Pogue, J, et al. Relationship of glucose and insulin level to the risk of myocardial infarction: A case-control study. *J Am Coll Cardiol* 1999; 33:612.
- Gill, JS, Shipley, MJ, Tsementzis, SA, et al. Alcohol consumption--a risk factor for hemorrhagic and non-hemorrhagic stroke. *Am J Med* 1991; 90:489.
- Gillman, MW, Cupples, LA, Gagnon, D, et al. Protective effect of fruits and vegetables on development of stroke in men. *JAMA* 1995; 273:1113.
- Gnasso, A, Carallo, C, Irace, C, et al. Association between intima-media thickness and wall shear stress in common carotid arteries in healthy male subjects. *Circulation* 1996; 94:3257.
- Gordon, T, Castelli, WP, Hjartland, MC, et al. High density lipoprotein as a protective factor against coronary artery disease. The Framingham Study. *Am J Med* 1977; 62:707.
- Greenland, P, Daviglius, ML, Dyer, AR, et al. Resting heart rate is a risk factor for cardiovascular and noncardiovascular mortality: the Chicago Heart Association Detection Project in Industry. *Am J Epidemiol* 1999; 149:853.
- Greenland, P, Knoll, MD, Stamler, J, et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 2003; 290:891.
- Gregg, EW, Cheng, YJ, Cadwell, BL, et al. Secular trends in cardiovascular disease risk factors according to body mass index in US adults. *JAMA* 2005; 293:1868.
- Gunn, M, Stephens, JC, Thompson, JR, et al. Significant association of cagA positive *Helicobacter pylori* strains with risk of premature myocardial infarction. *Heart* 2000; 84:267.

- Hagiwara, H, Mitsumata, M, Yamane, T, et al. Laminar shear stress-induced GRO mRNA and protein expression in endothelial cells. *Circulation* 1998; 98:2584.
- Harrington, RA, Becker, RC, Ezekowitz, M, et al. Antithrombotic therapy for coronary artery disease: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126:513S.
- Harris, TB, Cook, EF, Kannel, WB, et al. Proportional hazards analysis of risk factors for coronary heart disease in individuals aged 65 or older. *J Am Geriatr Soc* 1988; 36:1023.
- Harrison, DH, Armstrong, ML, Freiman, PC, Heistad, DD. Restoration of endothelium dependent relaxation by dietary treatment of atherosclerosis. *J Clin Invest* 1987; 80:1808.
- Hasdai, D, Holmes, DR, Carratt, KN, et al. Mechanical pressure and stretch release endothelin-1 from human atherosclerotic coronary arteries in vivo. *Circulation* 1997; 95:357.
- Hasenstab, D, Lea, H, Hart, CE, et al. Tissue factor overexpression in rat arterial neointima models thrombosis and progression of advanced atherosclerosis. *Circulation* 2000; 101:2651.
- He, FJ, Nowson, CA, MacGregor, GA. Fruit and vegetable consumption and stroke: meta-analysis of cohort studies. *Lancet* 2006; 367:320.
- He, K, Song, Y, Daviglius, ML, et al. Accumulated evidence on fish consumption and coronary heart disease mortality: a meta-analysis of cohort studies. *Circulation* 2004; 109:2705.
- Hebert, PR, Moser, M, Mayer, J, et al. Recent evidence on drug therapy of mild to moderate hypertension and decreased incidence of coronary heart disease. *Arch Intern Med* 1993; 153:578.
- Heitzer, T, Schlinzig, T, Krohn, K, et al. Endothelial dysfunction, oxidative stress, and risk of cardiovascular events in patients with coronary artery disease. *Circulation* 2001; 104:2673.
- Hennekens, CH, Hollar, D, Baigent, C. Sex-related differences in response to aspirin in cardiovascular disease: an untested hypothesis. *Nat Clin Pract Cardiovasc Med* 2006; 3:4.
- Hennekens, CH, Sacks, FM, Tonkin, A, et al. Additive benefits of pravastatin and aspirin to decrease risks of cardiovascular disease:

randomized and observational comparisons of secondary prevention trials and their meta-analyses. *Arch Intern Med* 2004; 164:40.

- Higazi, AA, Nassar, T, Ganz, T, et al. The alpha-defensins stimulate proteoglycan-dependent catabolism of low-density lipoprotein by vascular cells: a new class of inflammatory apolipoprotein and a possible contributor to atherogenesis. *Blood* 2000; 96:1393.
- Holvoet, P, Perez, G, Zhian, Z, et al. Malondialdehyde-modified LDL in patients with atherosclerotic disease. *J Clin Invest* 1995; 95:2611.
- Howard, BV, Van Horn, L, Hsia, J, et al. Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006; 295:655.
- Hubert, HB, Feinleib, M, McNamara, PM, et al. Obesity as an independent risk factor for cardiovascular disease: A 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 1983; 67:968.
- Hubert, HB, Feinleib, M, McNamara, PM, et al. Obesity as an independent risk factor for cardiovascular disease: a 26-year follow-up of participants in the Framingham Heart Study. *Circulation* 1983; 67:968.
- Hubinette, A, Cnattingius, S, Ekbom, A, et al. Birthweight, early environment, and genetics: a study of twins discordant for acute myocardial infarction. *Lancet* 2001; 357:1997.
- Hung, J, Whitford, EG, Parsons, RW, et al. Association of sleep apnea with myocardial infarction in men. *Lancet* 1990; 336:261.
- Hunt, ME, O'Malley, PG, Vernalis, MN, et al. C-reactive protein is not associated with the presence or extent of calcified subclinical atherosclerosis. *Am Heart J* 2001; 141:206.
- Hypertension prevalence and the status of awareness, treatment, and control in the United States. Final report of the Subcommittee on Definition and Prevalence of the 1984 Joint National Committee. *Hypertension* 1985;7(3 Pt 1):457–68.
- Ihling, C, Szombathy, T, Bohrmann, B, et al. Coexpression of endothelin-converting enzyme-1 and endothelin-1 in different stages of human atherosclerosis. *Circulation* 2001; 104:864.

- Iiyama, K, Hajra, L, Iiyama, M, et al. Patterns of vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 expression in rabbit and mouse atherosclerotic lesions and at sites predisposed to lesion formation. *Circ Res* 1999; 85:199.
- Inoue, T, Uchida, T, Kamishirado, H, et al. Antibody against oxidized low density lipoprotein may predict progression or regression of atherosclerotic coronary artery disease. *J Am Coll Cardiol* 2001; 37:1871.
- Inoue, T, Uchida, T, Kamishirado, H, et al. Clinical significance of antibody against oxidized low density lipoprotein in patients with atherosclerotic coronary artery disease. *J Am Coll Cardiol* 2001; 37:775.
- Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 1998; 352:837.
- Islam, KN, Devaraj, S, Jialal, I. Alpha-tocopherol enrichment of monocytes decreases agonist-induced adhesion to human endothelial cells. *Circulation* 1998; 98:2255.
- Iuliano, L, Mauriello, A, Sbarigia, E, et al. Radiolabeled native low-density lipoprotein injected into patients with carotid stenosis accumulates in macrophages of atherosclerotic plaque: effect of vitamin E supplementation. *Circulation* 2000; 101:1249.
- Jarvisalo, MJ, Jartti, L, Nanto-Salonen, K, et al. Increased Aortic Intima-Media Thickness: A Marker of Preclinical Atherosclerosis in High-Risk Children. *Circulation* 2001; 104:2943.
- Jha, P, Flather, M, Lonn, E, et al. The antioxidant vitamins and cardiovascular disease. A critical review of epidemiologic and clinical trial data. *Ann Intern Med* 1995; 123:860.
- John, S, Schlaich, M, Langenfeld, M, et al. Increased bioavailability of nitric oxide after lipid-lowering therapy in hypercholesterolemic patients. A randomized, placebo-controlled, double-blind study. *Circulation* 1998; 98:211.
- Johnson, RC, Chapman, SM, Dong, ZM, et al. Absence of P-selectin delays fatty streak formation in mice. *J Clin Invest* 1997; 99:1037.

- Joshipura, KJ, Ascherio, A, Manson, JE, et al. Fruit and vegetable intake in relation to risk of ischemic stroke. *JAMA* 1999; 282:1233.
- Jousilahti, P, Vartiainen, E, Tuomilehto, J, et al. Sex, age, cardiovascular risk factors, and coronary heart disease: A prospective follow-up study of 14,786 middle-aged men and women in Finland. *Circulation* 1999; 99:1165.
- Juan, SH, Lee, TS, Tseng, KW, et al. Adenovirus-mediated heme oxygenase-1 gene transfer inhibits the development of atherosclerosis in apolipoprotein e-deficient mice. *Circulation* 2001; 104:1519.
- Julius, S, Nesbitt, SD, Egan, BM, et al. Feasibility of treating prehypertension with an angiotensin-receptor blocker. *N Engl J Med* 2006; 354:1685.
- Kaartinen, M, Penttila, A, Kovanen, PT. Mast cells in rupture prone areas of human coronary atheromas produce and store TNF-alpha. *Circulation* 1996; 94:2787.
- Kannel, WB, Dannenberg, AL, Levy, D. Population implications of electrocardiographic left ventricular hypertrophy. *Am J Cardiol* 1987; 60:85I.
- Kannel, WB, Gordon ,T, Schwartz, MJ. Systolic versus diastolic blood pressure and risk of coronary heart disease: The Framingham Study. *Am J Cardiol* 1971; 27:335.
- Kannel, WB, Kannel, C, Paffenbarger, RS Jr, Cupples, LA. Heart rate and cardiovascular mortality: the Framingham Study. *Am Heart J* 1987; 113:1489.
- Kannel, WB, McGee, DL. Diabetes and cardiovascular risk factors: The Framingham Study. *Circulation* 1979; 59:8.
- Kannel, WB, McGee, DL. Diabetes and glucose tolerance as risk factors for cardiovascular disease: The Framingham Study. *Diabetes Care* 1979; 2:120.
- Kannel, WB. Left ventricular hypertrophy as a risk factor: the Framingham experience. *J Hypertens Suppl* 1991; 9:S3.
- Kenchaiah, S, Evans, JC, Levy, D, et al. Obesity and the risk of heart failure. *N Engl J Med* 2002; 347:305.

- King, AC, Haskell, WL, Young, DR, et al. Long-term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50-65 years. *Circulation* 1995; 91:2596.
- Knoops, KT, de Groot, LC, Kromhout, D, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* 2004; 292:1433.
- Knutsen, R, Knutsen, SF, Curb, JD, et al. The predictive value of resting electrocardiograms for 12-year incidence of coronary heart disease in the Honolulu Heart Program. *J Clin Epidemiol* 1988; 41:293.
- Kockx, MM, De Meyer, GR, Muhring, J, et al. Apoptosis and related proteins in different stages of human atherosclerotic plaques. *Circulation* 1998; 97:2307.
- Koenig, W, Rothenbacher, D, Hoffmeister, A, et al. Infection with helicobacter pylori is not a major independent risk factor for stable coronary heart disease: lack of a role of cytotoxin- associated protein A-positive strains and absence of a systemic inflammatory response. *Circulation* 1999; 100:2326.
- Koenig, W, Sund, M, Frohlich, M, et al. C-reactive protein, a sensitive marker of inflammation, predicts future risk of coronary heart disease in initially healthy middle-aged men. Results from the MONICA (Monitoring Trends and Determinants in Cardiovascular Disease) Augsburg Cohort study, 1984 to 1992. *Circulation* 1999; 99:237.
- Krauss, RM, Winston, M. Obesity: Impact on cardiovascular disease. *Circulation* 1998; 98:1472.
- Kristal-Boneh, E, Silber, H, Harari, G, Froom, P. The association of resting heart rate with cardiovascular, cancer and all-cause mortality. Eight year follow-up of 3527 male israeli employees (the CORDIS study). *Eur Heart J* 2000; 21:116.
- Kuczmarski, R.J., Johnson, C.L., Flegal, K.M., Campbell, S.M. Increasing prevalence of overweight among US adults. *Journal of the American Medical Association*. 1994; 272:205-211.
- Kurth, T, Moore, SC, Gaziano, JM, et al. Healthy lifestyle and the risk of stroke in women. *Arch Intern Med* 2006; 166:1403.

- Kuulasmaa, K, Tunstall-Pedoe, H, Dobson, A, et al. Estimation of contribution of changes in classic risk factors to trends in coronary-event rates across the WHO MONICA Project populations. *Lancet* 2000; 355:675.
- Lamarche, B, Moorjani, S, Lupien, PJ, et al. Apolipoprotein A-I and B levels and the risk of ischemic heart disease during a five-year follow-up of men in the Quebec cardiovascular study. *Circulation* 1996; 94:273.
- LaMonte, MJ, Eisenman, PA, Adams, TD, et al. Cardiorespiratory fitness and coronary heart disease risk factors: the LDS hospital fitness institute cohort. *Circulation* 2000; 102:1623.
- LaMonte, MJ, Eisenman, PA, Adams, TD, et al. Cardiorespiratory fitness and coronary heart disease risk factors: the LDS hospital fitness institute cohort. *Circulation* 2000; 102:1623.
- Lauer, MS, Okin, PM, Larson, MG, et al. Impaired heart rate response to graded exercise. Prognostic implications of chronotropic incompetence in the Framingham Heart Study. *Circulation* 1996; 93:1520.
- Laukkanen, JA, Lakka, TA, Rauramaa, R, et al. Cardiovascular fitness as a predictor of mortality in men. *Arch Intern Med* 2001; 161:825.
- Law, MR, Morris, JK. By how much does fruit and vegetable consumption reduce the risk of ischaemic heart disease? *Eur J Clin Nutr* 1998; 52:549.
- Leeson, CP, Whincup, PH, Cook, DG, et al. Cholesterol and arterial distensibility in the first decade of life: A population-based study. *Circulation* 2000; 101:1533.
- Lehr, HA, Sagban, TA, Ihling, C, et al. Immunopathogenesis of atherosclerosis: Endotoxin accelerates atherosclerosis in rabbits on hypercholesterolemic diet. *Circulation* 2001; 104:914.
- Lekakis, JP, Papamichael, CM, Cimponeriu, AT, et al. Atherosclerotic changes of extracoronary arteries are associated with the extent of coronary atherosclerosis. *Am J Cardiol* 2000; 85:949.
- Leon, AS, Connett, J, Jacobs, DR, et al. Leisure-time physical activity levels and risk of coronary heart disease and death. The Multiple Risk Factor Intervention Trial. *JAMA* 1987; 258:2388.



- Lerman, A, Edwards, BS, Hallett, JW, et al. Circulating and tissue endothelin immunoreactivity in advanced atherosclerosis. *N Engl J Med* 1991; 325:997.
- Levin, RI. The puzzle of aspirin and sex. *N Engl J Med* 2005; 352:1366.
- Levine, GN, Frei, B, Koulouris, SN, et al. Ascorbic acid reverses endothelial vasomotor dysfunction in patients with coronary artery disease. *Circulation* 1996; 93:1107.
- Levy, D. Echographically detected left ventricular hypertrophy: Prevalence and risk factors. The Framingham Heart Study. *Ann Intern Med* 1988; 108:7.
- Li, H, Freeman, MW, Libby, P. Regulation of smooth muscle cell scavenger receptor expression in vivo by atherogenic diets and in vitro by cytokines. *J Clin Invest* 1995; 95:122.
- Little, WC, Constantinescu, M, Applegate, RJ, et al. Can coronary angiography predict the site of a subsequent myocardial infarction in patients with mild to moderate coronary artery disease? *Circulation* 1988; 78:1157.
- Lloyd-Jones, DM, Larson, MG, Beiser, A, et al. Lifetime risk of developing coronary heart disease. *Lancet* 1999; 353:89.
- Luepker RV, Rosamond WD, Murphy R *et al.* Socioeconomic status and coronary heart disease risk factor trends. The Minnesota Heart Survey. *Circulation* 1993; 88 (part 1): 2172–9.
- Lynch JW, Kaplan GA, Cohen RD *et al.* Do cardiovascular risk factors explain the relation between socio-economic status, risk of all cause mortality, cardiovascular mortality, and acute myocardial infarction? *Am J Epidemiol* 1996; 144: 934–42.
- Major outcomes in moderately hypercholesterolemic, hypertensive patients randomized to pravastatin vs usual care: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT-LLT). *JAMA* 2002; 288:2998.
- Makimattila, S, Mantysaari, M, Groop, PH, et al. Hyperreactivity to nitrovasodilators in forearm vasculature is related to autonomic dysfunction in insulin-dependent diabetes mellitus. *Circulation* 1997; 95:618.

- Makimattila, S, Virkamaki, A, Groop, PH, et al. Chronic hyperglycemia impairs endothelial function and insulin sensitivity via different mechanisms in insulin-dependent diabetes mellitus. *Circulation* 1996; 94:1276.
- Malik, I, Danesh, J, Whincup, P, et al. Soluble adhesion molecules and prediction of coronary heart disease: a prospective study and meta-analysis. *Lancet* 2001; 358:971.
- Mancini, GBJ, Henry GC, Macaya, C, et al. Angiotensin-converting enzyme inhibition with quinapril improves endothelial vasomotor dysfunction in patients with coronary artery disease. The TREND (Trial on Reversing Endothelial Dysfunction) Study. *Circulation* 1996; 94:258.
- Manninen, V, Tenkanen, L, Koskinn, P, et al. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary artery disease in the Helsinki Health Study: Implications for treatment. *Circulation* 1992; 85:37.
- Manson, JE, Colditz, GA, Stampfer, MJ, et al. A prospective study of obesity and risk of coronary heart disease in women. *N Engl J Med* 1990; 322:882.
- Manson, JE, Hu, FB, Rich-Edwards, JW, et al. A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women. *N Engl J Med* 1999; 341:650.
- Marmot MG, Kogevinas M, Elston MA. Socioeconomic status and disease. *WHO Reg Publ Eur Ser* 1991; 37: 113–46.
- Mathew, V, Cannan, CR, Miller, VM, et al. Enhanced endothelin-mediated coronary vasoconstriction and attenuated basal nitric oxide activity in experimental hypercholesterolemia. *Circulation* 1997; 96:1930.
- Maxwell, AJ, Bruinsma, KA. Uric acid is closely linked to vascular nitric oxide activity. Evidence for mechanism of association with cardiovascular disease. *J Am Coll Cardiol* 2001; 38:1850.
- Mayers, DG. The iron hypothesis: does iron play a role in atherosclerosis?. *Transfusion* 1996; 40:1023.
- Mayr, M, Kiechl, S, Willeit, J, et al. Infections, immunity, and atherosclerosis: associations of antibodies to *Chlamydia pneumoniae*, *Helicobacter pylori*, and cytomegalovirus with immune reactions to heat-

- shock protein 60 and carotid or femoral atherosclerosis. *Circulation* 2000; 102:833.
- McCrohon, JA, Jessup, W, Handelsman, DJ, et al. Androgen exposure increases human monocyte adhesion to vascular endothelium and endothelial cell expression of vascular cell adhesion molecule-1. *Circulation* 1999; 99:2317.
  - McGill, HC Jr, McMahan, CA, Herderick, EE, et al. Obesity accelerates the progression of coronary atherosclerosis in young men. *Circulation* 2002; 105:2712.
  - McGill, HC, McMahan, A, Zieske, AW, et al. Association of coronary heart disease risk factors with microscopic qualities of atherosclerosis in youth. *Circulation* 2000; 102:374.
  - Melnick, JL, Hu, C, Burek, J, et al. Cytomegalovirus DNA in arterial walls of patients with atherosclerosis. *J Med Virol* 1994; 42:170.
  - Minamino, T, Kurihara, H, Takahashi, M, et al. Endothelin-converting enzyme expression in the rat vascular injury model and human coronary atherosclerosis. *Circulation* 1997; 95:221.
  - Miura, K, Daviglus, ML, Dyer, AR, et al. Relationship of blood pressure to 25-year mortality due to coronary heart disease, cardiovascular diseases, and all causes in young adult men. The Chicago Heart Association detection project in industry.
  - Mogensen, CE. Combined high blood pressure and glucose in type 2 diabetes: double jeopardy (editorial). *BMJ* 1998; 317:693.
  - Morrison, JA, Jacobsen, DW, Sprecher, DL, et al. Serum glutathione in adolescent males predicts parental coronary heart disease. *Circulation* 1999; 100:2244.
  - Muhlestein, JB, Hammond, EH, Carlquist, GF, et al. Increased incidence of Chlamydia species within the coronary arteries of patients with severe atherosclerosis versus other forms of cardiovascular disease. *J Am Coll Cardiol* 1995; 27:1555.
  - Muhlestein, JB, Horne, BD, Carlquist, JF, et al. Cytomegalovirus seropositivity and C-reactive protein have independent and combined predictive value for mortality in patients with angiographically demonstrated coronary artery disease. *Circulation* 2000; 102:1917.

- Mustafa, A, Nityanand, S, Berglund, L, et al. Circulating immune complexes in 50-year-Old men as a strong and independent risk factor for myocardial infarction. *Circulation* 2000; 102:2576.
- Naftilan AJ, Pratt RE, Dzau VJ. Induction of platelet-derived growth factor A-chain and c-myc gene expressions by angiotensin II in cultured rat vascular smooth muscle cells. *J Clin Invest* 1999;83(4):1419–24.
- Nathan, DM, Cleary, PA, Backlund, JY, et al. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med* 2005; 353:2643.
- Nelson, M, Reid, C, Beilin, L, et al. Rationale for a trial of low-dose aspirin for the primary prevention of major adverse cardiovascular events and vascular dementia in the elderly: Aspirin in Reducing Events in the Elderly (ASPREE). *Drugs Aging* 2003; 20:897.
- Nieto, FJ, Adam, E, Sorlie, P, et al. Cohort study of cytomegalovirus infection as a risk factor for carotid intimal-medial thickening. A measure of subclinical atherosclerosis. *Circulation* 1996; 94:922.
- Njolstad, I, Arnesen, E, Lund-Larsen, PG. Smoking, serum lipids, blood pressure, and sex differences in myocardial infarction. A 12-year follow-up of the Finnmark Study. *Circulation* 1996; 93:450.
- Nwasokwa, ON, Weiss, M, Glasdstone, C, et al. Effect of coronary artery size on the prevalence of atherosclerosis. *Am J Cardiol* 1996; 78:741.
- O'Brien, KD, Olin, KL, Alpers, CE, et al. Comparison of apolipoprotein and proteoglycan deposits in human coronary atherosclerotic plaques: Colocalization of biglycan with apolipoproteins. *Circulation* 1998; 98:519.
- Packard, CJ, O'Reilly, DS, Caslake, MJ, et al. Lipoprotein-associated phospholipase A2 as an independent predictor of coronary heart disease. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 2000; 343:1148.
- Paffenbarger, RS Jr, Hyde, RT, Wing, AL, et al. The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *N Engl J Med* 1993; 328:538.
- Panza JA, Quyyumi AA, Brush JE, Jr, Epstein SE. Abnormal endothelium-dependent vascular relaxation in patients with essential hypertension. *N Engl J Med* 1990;323(1):22–7.

- Pasceri, V, Cammarota, G, Patti, G, et al. Association of virulent *Helicobacter pylori* strains with ischemic heart disease. *Circulation* 1998; 97:1675.
- Pasceri, V, Willerson, JT, Yeh, ETH. Direct proinflammatory effect of C-reactive protein on human endothelial cells. *Circulation* 2000; 102:2165.
- Patel, P, Mendall, MA, Carrington, D, et al. Association of *Helicobacter pylori* and *Chlamydia pneumoniae* infections with coronary heart disease and cardiovascular risk factors. *BMJ* 1995; 311:711.
- Patel, SS, Thiagarajan, R, Willerson, JT, et al. Inhibition of  $\alpha_4$  integrin and ICAM-1 markedly attenuate macrophage homing to atherosclerotic plaques in apoE-deficient mice. *Circulation* 1998; 97:75.
- Pearson, TA, Blair, SN, Daniels, SR, et al. AHA Guidelines for Primary Prevention of Cardiovascular Disease and Stroke: 2002 Update: Consensus Panel Guide to Comprehensive Risk Reduction for Adult Patients Without Coronary or Other Atherosclerotic Vascular Diseases. American Heart Association Science Advisory and Coordinating Committee. *Circulation* 2002; 106:388.
- Podrez, EA, Febbraio, M, Sheibani, N, et al. Macrophage scavenger receptor CD36 is the major receptor for LDL modified by monocyte-generated reactive nitrogen species. *J Clin Invest* 2000; 105:1095.
- Potter, DD, Sobey, CG, Tompkins, PK, et al. Evidence that macrophages in atherosclerotic lesions contain angiotensin II. *Circulation* 1998; 98:800.
- Powell, KE, Thompson, PD, Caspersen, CJ, et al. Physical activity and the incidence of coronary heart disease. *Annu Rev Public Health* 1987; 8:253.
- Prescott, E, Hippe, M, Schnohr, P, et al. Smoking and the risk of myocardial infarction in women and men: Longitudinal population study. *BMJ* 1998; 316:1043.
- Prescott, E, Hippe, M, Schnohr, P, et al. Smoking and the risk of myocardial infarction in women and men: Longitudinal population study. *BMJ* 1998; 316:1043.

- Rader DJ, Hoeg JM, Brewer HB Jr. Quantitation of plasma apolipoproteins in the primary and secondary prevention of coronary artery disease. *Ann Intern Med* 1998;120(12):1012–25.
- Rajavashisth, T, Qiao, J-H, Tripathi, S, et al. Heterozygous osteopetrotic (op) mutation reduces atherosclerosis in LDL receptor-deficient mice. *J Clin Invest* 1998; 101:2702.
- Reaven, GM. Banting Lecture 1988: Role of insulin resistance in human disease. *Diabetes* 1988; 37:1595.
- Rectenwald, JE, Moldawer, LL, Huber, TS, et al. Direct evidence for cytokine involvement in neointimal hyperplasia. *Circulation* 2000; 102:1697.
- Redberg, RF, Rifai, N, Gee, L, Ridker, PM. Lack of association of C-reactive protein and coronary calcium by electron beam computed tomography in postmenopausal women: implications for coronary artery disease screening. *J Am Coll Cardiol* 2000; 36:39.
- Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. The Expert Panel. *Arch Intern Med* 1988; 148:36.
- Rexrode, KM, Hennekens, CH, Willett, WC, et al. A prospective study of body mass index, weight change, and risk of stroke in women. *JAMA* 1997; 277:1539.
- Rich-Edwards, JW, Manson, JE, Hennekens, CH, Buring, JE. The primary prevention of coronary heart disease in women. *N Engl J Med* 1995; 332:1758.
- Ridker, PM, Buring, JE, Rifai, N. Soluble P-Selectin and the Risk of Future Cardiovascular Events. *Circulation* 2001; 103:491.
- Ridker, PM, Buring, JE, Shih, J, et al. Prospective study of C-reactive protein and the risk of future cardiovascular events among apparently healthy women. *Circulation* 1998; 98:731.
- Ridker, PM, Cook, NR, Lee, IM, et al. A randomized trial of low-dose aspirin in the primary prevention of cardiovascular disease in women. *N Engl J Med* 2005; 352:1293.

- Ridker, PM, Cushman, M, Stampfer, MJ, et al. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997; 336:973.
- Ridker, PM, Glynn, RJ, Hennekens, CH. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* 1998; 97:2007.
- Ridker, PM, Glynn, RJ, Hennekens, CH. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* 1998; 97:2007.
- Ridker, PM, Hennekens, CH, Buring, JE, Rifai, N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 2000; 342:836.
- Ridker, PM, Hennekens, CH, Stampfer, MJ, et al. Prospective study of herpes simplex virus, cytomegalovirus, and the risk of future myocardial infarction and stroke. *Circulation* 1998; 98:2796.
- Ridker, PM, Rifai, N, Stampfer, MJ, Hennekens, CH. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy Men. *Circulation* 2000; 101:1767.
- Ridker, PM. Evaluating novel cardiovascular risk factors: Can we better predict heart attacks? *Ann Intern Med* 1999; 130:933.
- Rimm, EB, Ascherio, A, Giovannucci, E, et al. Vegetable, fruit, and cereal fiber intake and risk of coronary heart disease among men. *JAMA* 1996; 275:447.
- Rimm, EB, Giovannucci, EL, Willett, WC, et al. Prospective study of alcohol consumption and risk of coronary disease in men. *Lancet* 1991; 338:464.
- Rimm, EB, Stampfer, MJ, Ascherio, A, et al. Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 1993; 328:1450.
- Roest, M, van der, Schouw YT, de Valk, B, et al. Heterozygosity for a Hereditary Hemochromatosis Gene Is Associated With Cardiovascular Death in Women. *Circulation* 1999; 100:1268.

- Romm, PA, Green, CE, Regan, K, Rackley, CE. Relation of serum lipoprotein cholesterol levels to the presence and severity of angiographic coronary artery disease. *Am J Cardiol* 1991; 67:479.
- Roncaglioni, MC, Santoro, L, D'Avanzo, B, et al. Role of family history in patients with myocardial infarction: An Italian case-control study. GISSI-EFRIM Investigators. *Circulation* 1992; 85:2065.
- Rosenberg, L, Kaufman, DW, Helmrich, SP, Shapiro, S. The risk of myocardial infarction after quitting smoking in men under 55 years of age. *N Engl J Med* 1985; 313:1511.
- Rosengren, A, Wedel, H, Wilhelmsen, L. Body weight and weight gain during adult life in men in relation to coronary heart disease and mortality. A prospective population study. *Eur Heart J* 1999; 20:269.
- Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993;362(6423):801–9.
- Ross, R. The pathogenesis of atherosclerosis: A perspective for the 1990s. *Nature* 1993; 362:801.
- Rupprecht, HJ, Blankenberg, S, Bickel, C, et al. Impact of viral and bacterial infectious burden on long-term prognosis in patients with coronary artery disease. *Circulation* 2001; 104:25.
- Sacks, FM, Pfeffer, MA, Moye, LA, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* 1996; 335:1001.
- Salomaa, V, Matei, C, Alesksic, N, et al. Soluble thrombomodulin as a predictor of incidence coronary heart disease and symptomless carotid artery atherosclerosis in the Atherosclerosis Risk in Communities (ARIC) study: A case-cohort study. *Lancet* 1999; 353:1729.
- Sandvik, L, Erikssen, J, Thaulow, E, et al. Physical fitness as a predictor of mortality among healthy, middle-aged Norwegian men. *N Engl J Med* 1993; 328:533.
- Scandinavian Simvastatin Survival Study Group. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet* 1994; 344: 1383–9.



- Schachinger, V, Britten, MB, Zeiher, AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation* 2000; 101:1899.
- Schchinger, V, Britten, MB, Elsner, M, et al. A positive family history of premature coronary artery disease Is associated with impaired endothelium-dependent coronary blood flow regulation. *Circulation* 1999; 100:1502.
- Schoenhagen, P, Ziada, KM, Kapadia, SR, et al. Extent and direction of arterial remodeling in stable versus unstable coronary syndromes: an intravascular ultrasound study. *Circulation* 2000; 101:598.
- Schoenhagen, P, Ziada, KM, Vince, DG, et al. Arterial remodeling and coronary artery disease: the concept of "dilated" versus "obstructive" coronary atherosclerosis. *J Am Coll Cardiol* 2001; 38:297.
- Schwartz CJ, Valente AJ, Sprague EA, Kelley JL, Cayatte AJ, Rozek MM. Pathogenesis of the atherosclerotic lesion. Implications for diabetes mellitus. *Circulation* 1992;15(9):1156–67.
- Schwartz O, Gardemann A, Haberbosch W *et al*. Positive association of the  $\alpha$  fibrinogen H1/H2 gene variation to basal fibrinogen levels and to the increase in fibrinogen concentration during acute phase reaction but not to coronary artery disease and myocardial infarction. *Thromb Haemost* 1999; 77: 1120–6.
- Sdringola, S, Patel, D, Gould, KL. High Prevalence of Myocardial Perfusion Abnormalities on Positron Emission Tomography in Asymptomatic Persons With a Parent or Sibling With Coronary Artery Disease. *Circulation* 2001; 103:496.
- Selhub, J, Jacques, PF, Wilson, PWF, et al. Vitamin status and intake as primary determinants of homocysteinemia in the elderly. *JAMA* 1993; 270:2693.
- Selhub, J, Jacques, PF, Wilson, PWF, et al. Vitamin status and intake as primary determinants of homocysteinemia in the elderly. *JAMA* 1993; 270:2693.
- Sesso, HD, Lee, IM, Gaziano, JM, et al. Maternal and paternal history of myocardial infarction and risk of cardiovascular disease in men and women. *Circulation* 2001; 104:393.

## References

---

- Sesso, HD, Stampfer, MJ, Rosner, B, et al. Two-year changes in blood pressure and subsequent risk of cardiovascular disease in Men. *Circulation* 2000; 102:307.
- Sever, PS, Dahlof, B, Poulter, NR, et al. Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial-Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet* 2003; 361:1149.
- Shahar, E, Chambless, LE, Rosamond, WD, et al. Plasma lipid profile and incident ischemic stroke: the Atherosclerosis Risk in Communities (ARIC) study. *Stroke* 2003; 34:623.
- Shepherd, J, Cobbe, SM, Ford, I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 1995; 333:1301.
- Shepherd, J, Cobbe, SM, Ford, I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland Coronary Prevention Study Group. *N Engl J Med* 1995; 333:1301.
- Shepherd, J, Cobbe, SM, Ford, I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med* 1995; 333:1301.
- Simonini, A, Moscucci, M, Muller, DW, et al. IL-8 is an angiogenic factor in human coronary atherectomy tissue. *Circulation* 2000; 101:1519.
- Singer, DE, Nathan, DM, Anderson, KM, et al. Association of HbA1c with prevalent cardiovascular disease in the original cohort of the Framingham Heart Study. *Diabetes* 1992; 41:202.
- Siscovick, DS, Ekelund, LG, Johnson, JL, et al. Sensitivity of exercise electrocardiography for acute cardiac events during moderate and strenuous physical activity. The Lipid Research Clinics Coronary Primary Prevention Trial. *Arch Intern Med* 1991; 151:325.
- Sorlie, PD, Nieto, FJ, Adam, E, et al. A prospective study of cytomegalovirus, herpes simplex virus 1, and coronary heart disease: the atherosclerosis risk in communities (ARIC) study. *Arch Intern Med* 2000; 160:2027.

- Srinivasan SR, Wattigney W, Webber LS, Berenson GS. Race and gender differences in serum lipoproteins of children, adolescents, and young adults—emergence of an adverse lipoprotein pattern in white males: the Bogalusa Heart Study. *Prev Med* 1991;29(6):671–84.
- Staessen, JA, Wang, JG, Thijs, L. Cardiovascular protection and blood pressure reduction: a meta-analysis. *Lancet* 2001; 358:1305.
- Stamler, J, Stamler, R, Neaton, JD. Blood pressure, systolic and diastolic, and cardiovascular risks: U.S. population data. *Arch Intern Med* 1993; 153:598.
- Stampfer, MJ, Colditz, GA, Willett, WC, et al. Postmenopausal estrogen therapy and cardiovascular disease: Ten-year follow-up from the Nurses' Health Study. *N Engl J Med* 1991; 325:756.
- Stampfer, MJ, Hennekens, CH, Manson, JE, et al. Vitamin E consumption and risk of coronary heart disease in women. *N Engl J Med* 1993; 328:1444.
- Stampfer, MJ, Malinow, MR, Willett, WC, et al. A prospective study of plasma homocysteine and risk of myocardial infarction in US physicians. *JAMA* 1992; 268:877.
- Stampfer, MJ, Malinow, MR, Willett, WC, et al. A prospective study of plasma homocysteine and risk of myocardial infarction in US physicians. *JAMA* 1992; 268:877.
- Strydom, HC, Chandler, AB, Dinsmore, RE, et al. Definition of advanced types of atherosclerotic lesions in a histological classification of atherosclerosis: A report from the Committee on Vascular Lesions of the Council on Arteriosclerosis, American Heart Association. *Circulation* 1995; 92:1355.
- Stec, JJ, Silbershatz, H, Ofler, GH, et al. Association of fibrinogen with cardiovascular risk factors and cardiovascular disease in the Framingham offspring population. *Circulation* 2000; 102:1634.
- Stein, JH, Keevil, JG, Wiebe, DA, et al. Purple grape juice improves endothelial function and reduces the susceptibility of LDL cholesterol to oxidation in patients with coronary artery disease. *Circulation* 1999; 100:1050.

- Steinberg, D, Parthasarathy, S, Carew, TE, et al. Beyond cholesterol: Modifications of low-density lipoproteins that increase its atherogenicity. *N Engl J Med* 1989; 320:915.
- Strachan, DP, Mendall, MA, Carrington, D, et al. Relation of *Helicobacter pylori* infection to 13-year mortality and incident ischemic heart disease in the Caerphilly Prospective Heart Disease Study. *Circulation* 1998; 98:1286.
- Strawn, WB, Chappell, MC, Dean, RH, et al. Inhibition of early atherogenesis by losartan in monkeys with diet-induced hypercholesterolemia. *Circulation* 2000; 101:1586.
- Stroke and coronary heart disease in mild hypertension: risk factors and the value of treatment. Medical Research Council Working Party. *Br Med J (Clin Res Ed)* 1988; 296:1565.
- Strong, JP, Malcom, GT, McMahan, CA, et al. Prevalence and extent of atherosclerosis in adolescents and young adults. Implications for prevention from the pathobiological determinants of atherosclerosis in youth study. *JAMA* 1999; 281:727.
- Suh, I, Shaten, JB, Cutler, JA, Kuller, LH. Alcohol use and mortality from coronary heart disease: The role of high-density lipoprotein cholesterol. *Ann Intern Med* 1992; 116:881.
- Summary of the second report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. *JAMA* 1993; 269:3015.
- Suwaidi, JA, Hamasaki, S, Higano, ST, et al. Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction. *Circulation* 2000; 101:948.
- Suwaidi, JA, Hamasaki, S, Higano, ST, et al. Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction. *Circulation* 2000; 101:948.
- Takahashi, M, Kitagawa, S, Masuyama, JI, et al. Human monocyte-endothelial cell interaction induces synthesis of granulocyte-macrophage colony-stimulating factor. *Circulation* 1996; 93:1185.
- The Coronary Drug Project. Initial findings leading to modifications of its research protocol. *JAMA* 1970;214(7):1303–13.

- The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. Arch Intern Med 1997; 157:2413.
- Theorell T, Tsutsumi A, Hallqvist J *et al.* Decision latitude, job strain and myocardial infarction: A study of working men in Stockholm. Am J Pub Health 1998; 88: 382–8.
- Third report of the National Cholesterol Education Program (NCEP) Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). Circulation 2002; 106:3143.
- Thun, MJ, Peto, R, Lopez, AD, et al. Alcohol consumption and mortality among middle-aged and elderly US adults. N Engl J Med 1997; 337:1705.
- Tintut, Y, Patel, J, Parhami, F, Demer, LL. Tumor necrosis factor- $\alpha$  promotes In vitro calcification of vascular cells via the cAMP pathway. Circulation 2000; 102:2636.
- Touboul, PJ, Elbaz, A, Koller, C, et al. Common carotid artery intima-media thickness and brain infarction: the etude du profil Genetique de l'Infarctus Cerebral (GENIC) case-control study. Circulation 2000; 102:313.
- Tsao, PS, Buitrago, R, Chan, JR, et al. Fluid flow inhibits endothelial adhesiveness. Nitric oxide and transcriptional regulation of VCAM-1. Circulation 1996; 94:1682.
- Tsao, PS, Wang, B, Buitrago, R, et al. Nitric oxide regulates monocyte chemotactic protein-1. Circulation 1997; 96:934.
- Tuomainen, TP, Kontula, K, Nyyssönen, K, et al. Increased Risk of Acute Myocardial Infarction in Carriers of the Hemochromatosis Gene Cys282Tyr Mutation: A Prospective Cohort Study in Men in Eastern Finland. Circulation 1999; 100:1274.
- Tuzcu, EM, Kapadia, SR, Tutar, E, et al. High prevalence of coronary atherosclerosis in asymptomatic teenagers and young adults: evidence from intravascular ultrasound. Circulation 2001; 103:2705.
- U.S. Department of Agriculture, U. S. Department of Health and Human Services. Nutrition and Your Health: Dietary Guidelines for Americans.

- Washington, DC: U.S. Government Printing Office; 1995. Report No.: Home and Garden Bulletin No. 232.
- US Preventive Services Task Force. Guide to clinical preventive services, 3rd ed, 2000-2002. [www.ahrq.gov/clinic/prevnew.htm](http://www.ahrq.gov/clinic/prevnew.htm) (Accessed March 7, 2005).
  - Vaccarino, V, Berger, AK, Abramson, J, et al. Pulse pressure and risk of cardiovascular events in the systolic hypertension in the elderly program. *Am J Cardiol* 2001; 88:980.
  - van der Wal AC, Becker, AE, van der, Loos CM, Das, PK. Site of intimal rupture or erosion of thrombosed coronary atherosclerotic plaques is characterized by an inflammatory process irrespective of the dominant plaque morphology. *Circulation* 1994; 89:36.
  - Vanharanta, M, Voutilainen, S, Lakka, TA, et al. Risk of acute coronary events according to serum concentrations of enterolactone: a prospective population-based case-control study. *Lancet* 1999; 354:2112.
  - Vasan, RS, Larson, MG, Benjamin, EJ, et al. Left ventricular dilatation and the risk of congestive heart failure in people without myocardial infarction. *N Engl J Med* 1997; 336:1350.
  - Vasan, RS, Larson, MG, Leip, EP, et al. Impact of high-normal blood pressure on the risk of cardiovascular disease. *N Engl J Med* 2001; 345:1291.
  - Vink, H, Constantinescu, AA, Spaan, JA. Oxidized lipoproteins degrade the endothelial surface layer: implications for platelet-endothelial cell adhesion. *Circulation* 2000; 101:1500.
  - Waldius, G, Jungén, I, Holme, I, et al. High apolipoprotein B, low apolipoprotein A-I, and improvement in the prediction of fatal myocardial infarction (AMORIS study): a prospective study. *Lancet* 2001; 358:2026.
  - Wallace, AM, McMahon, AD, Packard, CJ, et al. Plasma Leptin and the Risk of Cardiovascular Disease in the West of Scotland Coronary Prevention Study (WOSCOPS). *Circulation* 2001.
  - Wannamethee, SG, Shaper, AG, Whincup, PH, Walker, M. Smoking cessation and the risk of stroke in middle-aged men. *JAMA* 1995; 274:155.

- Wannamethee, SG, Shaper, AG, Whincup, PH. Serum urate and the risk of major coronary heart disease events. *Heart* 1997; 78:147.
- Wei, M, Gibbons, LW, Mitchell, TL, et al. Low fasting plasma glucose level as a predictor of cardiovascular disease and all-cause mortality. *Circulation* 2000; 101:2047.
- Whincup, P, Danesh, J, Walker, M, et al. Prospective study of potentially virulent strains of *Helicobacter pylori* and coronary heart disease in middle-aged men. *Circulation* 2000; 101:1647.
- Wiedermann, CJ, Kiechl, S, Dunzendorfer, S, et al. Association of endotoxemia with carotid atherosclerosis and cardiovascular disease: prospective results from the Bruneck Study. *J Am Coll Cardiol* 1999; 34:1975.
- Wilhelmsson, C, Vedin, JA, Elmfeldt, D, et al. Smoking and myocardial infarction. *Lancet* 1975; 1:415.
- Wilhelmsson, C, Vedin, JA, Elmfeldt, D, et al. Smoking and myocardial infarction. *Lancet* 1975; 1:415.
- Wilking, SVB, Belanger, AJ, Kannel, WB, et al. Determinants of isolated systolic hypertension. *JAMA* 1988; 260:3451.
- Williams, PT. Relationship of distance run per week to coronary heart disease risk factors in 8283 male runners. The National Runners' Health Study. *Arch Intern Med* 1997; 157:191.
- Wilson, PW, D'Agostino, RB, Levy, D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* 1998; 97:1837.
- Wilson, PW, Kannel, WB. Hypercholesterolemia and Coronary Risk in the Elderly: The Framingham Study. *Am J Geriatr Cardiol* 1993; 2:56.
- Wilson, PW, Kauppila, LI, O'Donnell, CJ, et al. Abdominal Aortic Calcific Deposits Are an Important Predictor of Vascular Morbidity and Mortality. *Circulation* 2001; 103:1529.
- Wilson, PW. Established risk factors and coronary artery disease: The Framingham Study. *Am J Hypertens* 1994; 7:7S.
- Winkleby MA, Fortmann SP, Barrett DC. Social class disparities in risk factors for disease. Eight-year prevalence patterns by level of education. *Prev Med* 1990; 19: 1–12.

- Witztum, JL. The oxidation hypothesis of atherosclerosis. *Lancet* 1994; 344:793.
- Wolk, A, Manson, JE, Stampfer, MJ, et al. Long-term intake of dietary fiber and decreased risk of coronary heart disease among women. *JAMA* 1999; 281:1998.
- Woods, A, Brull, DJ, Humphries, SE, Montgomery, HE. Genetics of inflammation and risk of coronary artery disease: the central role of interleukin-6. *Eur Heart J* 2000; 21:1574.
- Wu, KK, Aleksic, N, Ahn, C, et al. Thrombomodulin Ala455Val polymorphism and risk of coronary heart disease. *Circulation* 2001; 103:1386.
- Xu, Q, Schett, G, Perschinka, H, et al. Serum soluble heat shock protein 60 is elevated in subjects with atherosclerosis in a general population. *Circulation* 2000; 102:14.
- Yusuf, S, Hawken, S, Ounpuu, S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004; 364:937.
- Zavaroni, I, Bonora, E, Pagliara, M, et al. Risk factors for coronary artery disease in healthy persons with hyperinsulinemia and normal glucose tolerance. *N Engl J Med* 1989; 320:702.
- Zhang, R, Brennan, ML, Fu, X, et al. Association between myeloperoxidase levels and risk of coronary artery disease. *JAMA* 2001; 286:2136.
- Zhou, YF, Guetta, E, Yu, ZX, et al. Human cytomegalovirus increases modified low density lipoprotein uptake and scavenger receptor mRNA expression in vascular smooth muscle cells. *J Clin Invest* 1996; 98:2129.
- Zhou, YF, Shou, M, Guetta, E, et al. Cytomegalovirus infection of rats increases the neointimal response to vascular injury without consistent evidence of direct infection of the vascular wall. *Circulation* 1999; 100:1569.
- Zhu, J, Nieto, FJ, Horne, BD, et al. Prospective study of pathogen burden and risk of myocardial infarction or death. *Circulation* (Online) 2001; 103:45.



## *References*

---

- Zhu, J, Quyyumi, AA, Norman, JE, et al. Cytomegalovirus in the pathogenesis of atherosclerosis: the role of inflammation as reflected by elevated C-reactive protein levels. *J Am Coll Cardiol* 1999; 34:1738.
- Zhu, J, Quyyumi, AA, Norman, JE, et al. Effects of total pathogen burden on coronary artery disease risk and C-reactive protein levels. *Am J Cardiol* 2000; 85:140.
- Zhu, J, Quyyumi, AA, Rott, D, et al. Antibodies to human heat-shock protein 60 are associated with the presence and severity of coronary artery disease: evidence for an autoimmune component of atherogenesis. *Circulation* 2001; 103:1071.
- Zioncheck, TF, Powell, LM, Rice, GC, et al. Interaction of recombinant apolipoprotein(a) and lipoprotein(a) with macrophages. *J Clin Invest* 1991; 87:767.