

REFERENCES

References

Akiyama, S.K. and Yamada K.M (1997): Fibronectin
Adv – Enzymol, 59: 1-57.

Akiyama, s.k. and Yamada, K.M (1993): Fibronectin in connective tissue
diseases.

Edited by Wagner, B.M, Flischrmajor, R. and Kaufman N.

Published by william and Willkins, p 55.

Andern-Sandberg and Ihse (1983): Management of proximal
cholangiocarcinomas by surgical resection and radiotherapy
Am.J.Surg ; 159:91

Aranha GV, Prinz' RA, Freeark RJ (1994): The spectrum of biliary tract
obstruction from chronic pancreatitis
Arch. Surg 119:595

Asch, B.B., Kamat, B.R., and Burstein, N.A. (1991): Interaction of normal
dysplastic and malignant mammary epithelial cells with FN in vivo and
vitro
Cancer Res, 41: 215 – 225.

Baranov V, Rescan PY, Lissel M. et al (1990): Ultrastructural localization of the major components of the extracellular matrix in normal mouse liver
J submicrosc. Cytol Pathol 22:497 – 506

Behar , Jose (1995): Physiology of the biliary tract
Bokus Gastroentrology 5th edition Vol.3. 2554 – 72 saunders

Boyer JL, Soroka CJ (1995): Vesicle targeting to the apical domain regulates bile excretory function in isolated rate hepatocyte couplets
Gastroenterology;109:1600

Broome V, Lofberg R, veress B et al (1995) : Primary sclerosing cholangitis and ulcerative colits. Evidence for increased neoplastic potential
Hepatology;22 :1404

Brumfeld, V. and Weber, M.W (1998) : Studies on fibronectin and its domains
Arch – Biochem. Biophys, 302:134 – 143.

Carson S., Mosesson. H.W. and Diamond H.S. (1991) : Detection and quantitation of fibronectin in synovial fluid from patients with Rheumatic disease.
Arthritis Rheum. 24:1261 – 1267.

Carter, W.G. and Hakomori, T. (1979): Isolation of galactoprotein from hamster embryofibroblasts and the carbohydrate unit.
Biochemistry, 18:730-736.

Chijiwa K, Koga A, Yamazaki T, et al (1991) : Fibronectin: a possible factor promoting cholesterol monohydrate crystallization in bile
Biochim Biophys Acta 1086:48

Cohen, M.P., Saini, R., Klepser, H., et al (1996): Fibronectin binding to glomerular basement membrane is altered in diabetes
Diabetes, 36:758 – 769.

Cohn JA, Strong TV, Picciotto MR et al; (1993): Localization of the cystic fibrosis membrane conductance regulator in human bile duct epithelial cells.
Gastroenterology, 105:1857

Colli A, Buccino G, Cocciolo M, et al (1996): Diagnostic accuracy of fibronectin in differential diagnosis of ascites.
Cancer 58: 2489 – 2493

Cyriel I & Guido N.J. (1998): Primary sclerosing cholangitis: A clinical review
Am. J Gastroenterol 93:512 – 523

Davidson B, varsamidakis N, Dooley J (1992) : The value of exfoliative cytology for investigating bile duct strictures.

GUT 15:108-110

Delpuech,P., Desch,G. and frautus F. (1999) : Fibronectin is unsuitable as a tumour marker in pleural effusion.

Clin. Chem. 35(1): 166 – 168.

De-Russe, J., colombat, P., Lavoix,X. et al . (1995) : Plasma fibronectin in various homeopathic diseases.

Clin.Chem.Acta, 145:49 – 54.

Desa LA, Akosa AB, Lazzara S et al (1991): Cytodiagnosis in the management of extrahepatic biliary strictures

Gut, 32:1188

Domula , M. , Byknowska , K. , and Kopic , M. (1995) : Plasma fibronectin concentration in health and septic infants

J.Pediat. , 144 : 49-54 .

Donohue JH, Nagorney DM, Grant CS (1990): Carcinoma of the gall bladder, does radical resection improve outcome?

Arch. Surg 125:237

Dowling R M (1992): The enterohepatic circulation

Gastroentrology, 62:122

Drapiewski JF. (1984): Carcinoma of the pancreas: possible clinical presentations

Am. J.clin. pathology 14:549

Edsall, J.T., Gillbert, G.A. and scherage, H.A. (1955): The non – clotting component of human plasma fractin I (cold – insoluble globulin).

J – Am. Chem. Soc. 77:157.

Ekbom A, Hsieh C, Yuen J et al (1993): Risk of extrahepatic bile duct cancer after cholecystectomy

Lancet 342:1262.

Engvall , E. , Ruoslathi , E. and Miller , E.J. (1998): Affinity of fibronectin to collagen of different genetic types and to fibrinogen

J. Exp.Med. , 187:1584-1593.

Eric , J.B. and Jennifer , L.G. (1998) : Fibronectin receptors of phagocytes.

J.Exp. Med., 167 : 777-783.

Fateen A, Sherif A, Arafa M et al (1994): Significance of measuring fibronectin in bile as a malignant predictor

Journal of biomedical sciences and therapeutics (Egypt) 10(11) 1

Nov. suppl.

Feith MP, Klompmaker J, Maring JK (1997): Biliary reconstruction during liver transplantation in patients with primary sclerosing cholangitis
Transplant Proc 29:560 – 561

Feldman, B.F., Thomson, D.B, and D’Niell,s. (1995): Plasma fibronectin concentration in dogs with disseminated intravascular coagulation
Am. J. Vet. Res 46:1171 – 1174

Fernando, G.C., Allison, E,Mark. I.M. and Jerry, M.B. (1995): Decreased plasma fibronectin level in children with hemolytic uremic syndrome
Am. J. Med. 78:549 – 554.

Fleisch. R., Perlish J.S., Kieg T. et al (1991): Variability in collagen and fibronectin synthesis by scleroderma fibroblasts in primary cultures
J.invest. Dermatol 70: 400 – 403.

Fukuda , H. and Hakomori , S. (1979): Carbohydrate structure of galactoprotein a , a major transformation sensitive glycoprotein released from embryo fibroblasts.
J.Biol . Chem. , 254:5451-5463.

Furcht , L.T. and Wendelscharfer – Crabb (1989): The role of fibronectin in normal platelets aggregation .
J.Cell. Biol. , 83:61-67.

Garcia Pardo (1997): Primary structure of human plasma fibronectin.
Biochem.J , 241:923-926

Ginsberg , M.H , pointerm R.G. and Plow , E.F.(1990): Thrombin increases expression of fibronectin antigen on the platelet surface .
Proc.Nat .Acad.Sci.USA . 77:1049-1054

G.J. Romanes (1998): Ducts of Liver
Cunningham's maual of practical anatomy , 16th edition 1998
volume 2: 129 – 130 .Oxford

Geerts A, Geuze HJ, Slot JW, et al (1996): Immuno localization of procollagen III Fibronectin and heparin sulfate Proteoglycan on ultrathin frozen sections of the normal fat liver.
Histochemistry;84 :355-362.

Gonzalez – Calvin, J., Scully, M.F., Sanger. Y et al (1992): Fibronectin in fulminate hepatic failure
British Med.J. 285:1231 – 1234

Grinnell , F. , Feld , M. and Miner , D.(1990): Fibroblast adhesion to fibrinogen and fibrin substrat requirement for cold insoluble globulin (plasma fibronectin)
Cell , 19:517-522.

Harry , R.H. , Nancy , H.A and John , F.B.(1993): Mechanism of fibronectin enhancement of group B streptococcal phagocytosis by human neutrophils and culture – derived microphages.
Infect.Immun., (61) : 2334-2339.

Hoffbrand, A.V.(1998): Congenital and acquired coagulation disorders in:
Recent Advances of haematology
Churchill livingstone, Edinburgh, London, Melbourne and New York
1st ed., p:319.

Hormann, M. Seidl, M. (1990): Affinity chromatography on immobilized fibrin monomer III The fibrin affinity center of fibronectin
Physiol. Chem., 361: 1449-1454

Hynes Ro (1990): Fibronectins,
Berlin – Heidelberg – New York springer 300 – 334

Hynes, R.O. and Destree A.T (1998): Relationship between fibronectin (LETS) and actin
Cell, 15: 875 – 877.

Jane, A.D, Ardenne and Me Gee J.O.O (1999): Fibronectin in disease
J. pathol , 142:235 – 251.

Juvonen T, Kaira luoma Mi, Malinen H, et al (1993): extracellular matrix protein in bile and serum of patients with gall stones diseases
Connect tissue Res 29:171 – 181

Kaw M, Silverman WB, Rabinovitz M (1995): Biliary tract calculi in primary sclerosing cholangitis
Am J Gastroenterol 90:72 – 75.

Keski – Oja, J., Sen, A., and Todarce, G.J. (1990): Direct association of fibronectin and actin Molecules in vitro.
J.cell. Biol.,85:527 – 532

Kinsella, M.B., Smith E.A, Miller K.S. et al (1999): Spontaneous production of fibronectin by alveolar macrophages in patients with scleroderma
Arthritis Rheum 32: 577 – 583.

Klatskin G (1965): Adenocarcinoma of the hepatic duct at its bifurcation within the portahepatis. An unusual tumour with distinctive clinical and pathological features
Am J med 38:24

Kor,L, Lai, C.S., Nettesheim, D. et al (1993): IH NMR. Based determination of the three – dimensional structure of the human plasma fibronectin fragmentcontaining inter – chain disulfide bonds.
J. Biol – chem. Apr 25:8580 – 8589.

Korner.T, Kropf. J, Hackier R, et al (1996): Fibronectin in human bile fluid for diagnosis of malignant biliary disease.

Hepatology: 23 (3): 423 – 428.

Kullak-Ublick G-A, Magenbuch B, Stieger B et al (1995): Molecular and functional characterization of an organic anion transporting polypeptide cloned from human liver .

Gastroenterology;109:1274

Kumar A, Aggrwal S, Berry M (1990): Ultrasonography of carcinoma of the gall bladder : on analysis of 80 cases.

J clin. Ultrasound 18:715

Kurothong TR, Deery A, Dooley JS et al (1993): A prospective study of biliary cytology in 100 patients with bile duct strictures

Hepatology 18:1399

Kuusela, P. (1998): fibronectin binds to staphylococcus aureus

Nature, 276:718 – 721

Kuusela, P. vaheri, A., Palo, J et al, (1978): Demonstration of fibronectin in human CSF.

J.lab. clin. Med 92:595 – 599.

Lafleur , M. , Beaulie , A.D., and Poubelle , P.(1997): Fibronectin gene expression in PNL .
J.Biol.Chem. , 262:2111 – 2120 .

Lai ECS, chu KM, Loc – Y et al (1992): Choice of palliation for malignant hilar biliary obstruction
Am.J.surg 163:208

Lee JG, Schutz SM, leung JW (1995): Endoscopic therapy of sclerosing cholangitis
Hepatology 21:661 – 667

Lillemoe KD (1997): Benign post operative bile duct stricture
Baillieres clin Gastroenterol, 11:4, 749 – 779

Lindor KD, Wiesner RM, colwell LL (1998): The combination of prednisone and colchicine in patient with primary sclerosing cholangitis
Am J gastroenterol 86:57 – 61

Low VH (1997): Retrograde cholangiography of malignant biliary strictures: spectrum of appearance and pitfalls.
Abdom imaging Jul, 22:4,421 – 5

Maccarty, RL, La Russo NF, Wiesner RH (1999): Primary sclerosing cholangitis, finding on cholangiography and pancreaticography
Radiology 149 : 39 – 44

Maglund C, Y latupa S, Mertaniemi p et al (1997): Cellular fibronectin concentration in the plasma of patients with malignant and benign diseases: a comparison with CA 19-9 and CEA.

Br.J. cancer 76(6) : 777 – 783

Majoie CB, Reeder JW, Sanders JB (1991): Primary sclerosing cholangitis: A modified classification of cholangiographic findings.

Am J Roentgenol 157:495 – 497

Matsuda, n., Yamanaka, T. and Mastude A (1992): Distribution of fibronectin in plasma and liver diseases

Clin. Chem. Acta, 18:191 – 196.

May GR, Bender CE, La Russon F. (1995): Non operative dilatation of dominant stricture in primary sclerosing cholangitis

Am J Radiol 145:1061 – 1064

Mc Donagh , J.(1998): A molecular glue

Arch.Pathol.Lab.Med. , 105:393-396.

Merier PJ (1995): Molecular mechanisms of hepatic bile salt transport from sinusoidal blood into bile.

AM. J. physiol ; 269:G801

Miquel JF, von-Ritter C, Del-pozo et al (1994): Fibronectin in human gall bladder bile: cholesterol pronucleating and/or mucin (link) protein?
Am – J – physiol 267: 393 – 400

Moriwaki Y., Kohjiro N., Itoh M., et al (1999): Discrimination of tuberculous from carcinomatous pleural effusion by biochemical marker: adenosine deaminase, lysozyme, fibronectin and carcinoembryonic antigen.
Jpn. J. Med 28(4): 478 – 487.

Morrison, P.R. , Edsall, J.T and Miller, S.G. (1948): Preparation and properties of serum and plasma proteins The separation of purified fibrinogen from fraction I of human plasma
J.Am chem. Soc.,76:3103.

Mosesson, M.W. and Amrani, D.L. (1980): The structure and biological activity of plasma fibronectin
Blood,56;145-158.

Mosesson, M.W., Coleman, R.W. and Sherry, S. (1968): chronic intravascular coagulation syndrome report of cases with special studies of an associated plasma cryoprecipitate (Cryofibrinogen)
N.Engl. J.Med. 278:815-821.

Mosher, D.F. (1975): Cross – linking of cold – insoluble Globulin by fibrin stabilizing factor
J.Biol. Chem. 250:6614 – 6621.

Mosher, D.F. (1980): Fibronectin concentration in progressive homeostasis and thrombosis.

J clin invest, 69: 450-456.

Mosher, D.F. and proctor, R.A. (1990): Binding and factor XIIIa mediated cross – linking of a 27-kilodalton fragment of fibronectin to staph-aureus
Science, 209:927-931.

Mosher, D.F. and William E.M. (1997): Fibronectin concentration is decreased in plasma of severely ill patients with disseminated intravascular coagulation

J.Lab.Clin.Med., 91:729 – 735.

Moto Y, okamoto K, Uchimura M (1994): Composite tumour(ordinary adenocarcinoma typical carcinoid and goblet cell adenocarcinoid) of the gall bladder: a variety of composite tumour

Am J Gastroenterol 79:645

Myhre , E.B. and Kuusela, P. (1993): Binding of human fibronectin to group A, C and G streptococci Infect.

Immun. 40:29 – 34.

Nagorney DM, Donahue JH, Farnell JH et al (1993): Outcome after curative resections of cholangiocarcinoma

Arch. Surg 128:871

Nagy, B., Katona E., Erdei J. et al (1998): Fibronectin in broncho alveolar lavage fluid and plasma from children with chronic inflammation of the lung
Scand.Pediatr, Acta 77:727 – 733.

Nathanson MH, Boyer JL (1991): Mechanisms and regulation of bile secretion
Hepatology.14:55

Niederau C, Sonnenberg A, Mueller J (1984): Comparison of the extrahepatic bile duct size measured by ultrasound and different radiographic methods.
Gastroenterology , 87:615

Nix GAJJ, van overbeeke ic, Wilson JHP (1998): ERCP diagnosis of tumours in the region of the head of pancreas. Analysis of criteria and computer aided diagnosis
Dig. Dis. Sci 33:577

Northorer JMA, Terblanche JA. (1979): A new look at the arterial supply of the bile duct in man and its surgical implication
Br.J.surg 66 : 379

Obefield and Rossi (1988): Intrahepatic cholangiocarcinoma radiologic – pathologic correlation .
Radiology ; 167:689.

Olden, K., Pratt, R.M., and Yamada, K.M. (1998): Role of carbohydrate secretion and turnover effects of tunicamycin on the major cell surface glycoprotein of chick embryo fibroblasts.

Cell, 13:461 – 467.

Parkin DM, Ohshima H, Srivatanakul P et al (1993): Cholangiocarcinoma. Epidemiology, mechanism of carcinogenesis and prevention Cancer

Epidemiol – Biomarker prevention 2:537

Pearlstein, E. (1986): Plasma membrane glycoprotein which mediates adhesion of fibroblasts to collagen.

Nature, 62:497 – 503.

Pearlstein, E., Gold, L.i and Garcia Parado, A. (1990): Fibronectin: A review of its structure and biological activity

Mol. cell. Biochem. 29: 103 – 108.

Perri , R.T., Kay , N.E. Mc carthy , J, et al (1992): Fibronectin enhances in vitro monocyte – macrophage mediated tumouricidal activity .

Blood , 60 : 430-435.

Plow , E.G , Brdwell , C.R. and Ginsberg , M (1989): Identification and quantitation of platelet associated fibronectin antigen .

J.Clin.Invest. , 63 : 540-544

Polydorou AA, Cairns SR, Dowsett JF et al (1991): Palliation of proximal malignant biliary obstruction by endoscopic endoprosthesis insertion

Gut, 32:685

Poroyko MK, la russo NF, Wiesner RH (1991): Primary sclerosing cholangitis: A progressive disease ?

[Review] Semin liver disease 11:18-25.

Ramage JK, Donaghy A, Farrant JM et al (1995): Serum tumour markers for the diagnosis of cholangiocarcinoma in primary sclerosing cholangitis

Gastroenterology 108:865

Richard , A. , Proctor and Vann (1995): Role of fibronectin in human monocyte and macrophage bacterial activity .

Infect . Immun . 47 : 629 – 637 .

Rosen CB, Nagorrey DM, Wiesner RH (1991): Cholangiocarcinoma complicating primary sclerosing cholangitis .

Ann surg 213:21 – 25.

Ruoslahti, E. and Engvall, E. (1990): Complexing of fibronectin glycosamin – oglycans and collagen.

Biochem – Biophys, Acta, 631:350 – 356.

Ruoslahti, E., and vaheri. A. (1979): Novel human serum protein from fibroblust plama membrane.

Nature, 248:790-799.

Ruoslahti; E., Hayman, E.G., Pierschbacher, M. et al (1992): Fibronectin: purification, Immunochemical properties and biological activities.

Methods in enzymology 82:303 – 308

Saba , T.M. (1990): Physiology and physiopathology of reticuloendothelil system .

Arch . Intern . Med . , 126 : 1091-1102 .

Scovill , W.A , Saba , T.M. and Blumen stock , F.A. (1988): Opsonic as surface binding glycoprotein therapy during sepsis .

Ann . Surg . 188 : 521-524.

Sekiguchi, K. and Hakomori, S.I (1990): Functional domain structure of fibronectin

Proc. Natl. Acad. Sci. USA 77:2661-2663.

Sellinger M, Barrett C, Malle P et al (1998): Cryptic Na⁺,K⁺ - ATPase activity in rat liver canalicular plasma membranes; evidence for its basolateral origin

Hepatology; 11: 223

Sherman , L.A , Harwing , S. and lee , J. (1995): In vitro formation and in vivo clearance of fibrinogen fibrin complexes .

J.Lab.Clin.Med. , 80 : 100-104

Simpson , W.A , Courtney , II and Beachey , E.I. (1992): Laboratory diagnosis of bacterial meningitis .

Trans , Roy. Soc.Trop. Med.Hyg. (Suppl 1) 85 : 6-8

Smit JJM, Schinkel AH, Elferink RPJ et al; (1993): Homozygous disruption of the murine *mdr2* P-glycoprotein gene leads to a complete absence of phospholipid from bile and to liver disease

Cell, 75;451.

Smith R.T and von korff (1957): A heparin perceptible fraction of human plasma Isolation and characterization of the fraction

J., clin. Invest. , 36:596-599.

Stabile BE, Calabria R, Wilson SE (1997): Stricture of the common bile duct from chronic pancreatitis

Surgery 165:121

Stahl TJ, Allen MOC, Ansel HJ (1988): Partial biliary obstruction caused by chronic pancreatitis, an appraisal of indications for surgical biliary drainage

Am surg 207:26.

Stathakis, N.E., Fountas, A. and Tsianos, E. (1991): Plasma fibronectin in normal subject in various disease states

J.clin.pathol, 34:504 – 508.

Stenman, S, and vaheiri, A (1998): Distribution of a major connective tissue protein, fibronectin in normal human tissue.

J.exp – Med , 401:1054 – 1064

Sugwo CA (1983): Iatrogenic injury to the bile duct.

Arch . Surg. 125 : 1028.

Tamkun Jw, Hynes RO (1993): Plasma fibronectin is synthesized and secreted by hepatocytes.

J.Biol Chem, 258:4641 – 4647.

Terblanche J, Allison HF, Northover JMA (1993): An ischemic basis for biliary stricturs

Surgery 94:52

Tompkins PK, Saunders KD, Roslyn JJ et al (1990): Changing patterns in diagnosis and management of bile duct cancer

Ann. Surg 211:614

Torikata, C., Villiger B, Kahn C. et al (1992): Ultrastructural distribution of fibronectin in normal and fibrotic human lung.

Am. J. pathol. 96:652 – 657

Torre , D. , Zeroü , C. , and Speraza , F. (1991): Cerebrospinal fluid concentration of fibronectin in meningitis .

J . Clin . Pathol . , 44 : 783-784.

Tsukada N, Ackerley CA, phillips MJ (1995): The structure and organization of the bile canalicular cytoskeleton with special reference to actin and actin-binding proteins

Hepatology, 21:1106.

Tsukamoto Y . , Helsel , W. E. and Wahi , S. M. (1991): Macrophage production of Fibronectin , a chemoattractant for fibroblasts .

J . Immunol . 127 : 675-678.

Vauthey J.N, Blumgart LH (1994): Recent advances in the management of cholangiocarcinomas

Semin. Liver dis 14:109

Wagner, D.D. and Hynes, R.O.(1990): Topological arrangement of the major structural features of fibronectin

J.Biol. chem. 235:4304-4312

Wagner,D.D and Hynes, R.O. (1999): Domain structure of fibronectin and its relation to function. Disulfide sulfhydryl groups.

J. Biol. Chem , 254:6746 – 6749.

Warshaw AL, Fernandez .Del.Castillo C (1992): Pancreatic carcinoma

N eng J med 326:455

Wash burn WK, Lewis WD,Jenkins RL (1996): Aggressive surgical resection for cholangiocarcinoma

Arch. Surg 130:270

Weller, M., Sommer, N., and stevens, A. (1990): Increased intrathecal synthesis of fibronectin in bacterial and carcinomatous meningitis

Med clin. N. Am. 69(2) :219 – 229

Wilson JA (1989): Chronic pancreatitis ; a cause of cholestasis .

Gut ; 18 : 196.

Yamada, S, Willingham.Y.and Yamada, K.M., M.C. (1990): Intracellular localization of fibronectin by immunoelectron microscopy

J.Histochem. Cytochem.,28:953-957.

Yokota T, Oritani K, Mitsui M et al (1998): Growth – supporting activities of fibronectin on hematopoietic stem progenitor cell in vitro and in vivo Blood. 91(9): 3263 – 3272.

Zucke , M. b . Mosesson , M.W. , Brochman , M.J. et al (1989): Release of platelet fibronectin from alpha granules induced by thrombi by or collagen .

Blood , 54 : 8-12.