

REFERENCES

References

- Adler A.I.; Neil H.A.; Manley S.E.; Holman R.R. and Turner R.C. (1999):*** Hyperglycemia and hyperinsulinemia at diagnosis of diabetes and their association with subsequent cardiovascular disease in the United Kingdom prospective diabetes study (UKPDS 47). *Am. Heart. J.* 138: 5353-59.
- Aitman T.J.; Glazier A.M.; and Wallace C.A. (1999) :*** Identification of Cd36 (Fat) as an insulin-resistance gene causing defective fatty acid and glucose metabolism in hypertensive rats. *Nature Genetics.* 21: 76-83.
- Alemzadeh R; Holshouser S; Massey P and Koontz L; (2002):*** Chronic suppression of insulin by diazoxide alters the activities of key enzymes regulating hepatic gluconeogenesis in Zucker rats. *Eur. J. Endocrinol.* 146(6): 871-9.
- Altshuler D.; Hirschorn J.N.; and Klannemark M., (2000) :*** The common PPARGPro12Ala polymorphism is associated with decreased risk of type 2 diabetes. *Nat. Genet.* 26: 76-80.
- Anada B; (2000):*** Effects of type 2 diabetes on the ability of insulin and glucose to regulate splanchnic and muscle glucose metabolism. *Diabetes.* 49: 272-83.

Alwan AS; and King H; (1993): Diabetes in the Eastern Mediterranean Region WH statistics quart. 43(4) 355-9

American Diabetes Association (2002) : Gestational diabetes mellitus (Position Statement). Diabetes Care 25 (Suppl. 1). 593-596.

Andersson D.K.G. and Svaardsudd K. (1995) : Longterm glycemic control relates to mortality in type II diabetes. Diabetes Care. 18: 1534-1543.

Anderson E.A.; Balon T.W.; and Hoffman R.P. (1992) : Insulin increases sympathetic activity but not blood pressure in borderline hypertensive humans. Hypertension 19: 621-627.

Anderson R.L.; Hamman R.F.; and Savage P.J. (1995) : Exploration of simple insulin sensitivity measures derived from frequently sampled intravenous glucose tolerance (FSIGIT) tests. The insulin resistance atherosclerosis study. Am. J. Epidemiol. 142: 724-32.

Andrew TH; (2001): MODY, from clinical description to molecular genetic characterization. Clin. Endocrinol. and Metabolism. 15(3) 309-23.

Arab MM; (1992): Diabetes mellitus in Egypt. Wld. Hlth. Statist. Quart. 45: 334-7.

Arab MM; and Abd Al-Moneim NA; (1994): A socio-economic background for the epidemiology of DM 2nd March 1993, Medico-Graphia 16(1) 84-90..

Arab MM; and El-Sewy FZ; (1996): Diabetes in the Egyptian Deserts. Diabetes Care. 19: 92.

Arslanian S; and Suprasongsin; (1996): Differences in vivo insulin secretion and sensitivity of healthy black versus white adolescents. J. Pediatr. 129: 440-3.

Ashcroft F.M. and Rorsman P. (1990) : ATP-sensitive K⁺ channels : a link between B-cell metabolism and insulin secretion. Biochem. Soc. Trans. 18: 109-11.

Atkinson M.A. and Maclaren N.K. (1994) : The pathogenesis of insulin dependent diabetes. N. Engl. J. Med. 331: 1428-1436.

Avicena ; 980-1037.

Aziz A. Ismail & G.V. Gilli (1999): The epidemiology of type 2 diabetes and its current measurement. Bailliere's Clinical Endocrinology and Metabolism. 13:No. 2 pp. 197-220.

Bak J. Moller N.; and Schmitz O. (1992) : IN vivo action and muscle glycogen synthase activity in type II (noninsulin dependent)

diabetes mellitus : effects of diet treatment. Diabetologia. 35: 777-84.

Barnard H; Wolffers IN and Arab MM; (1996): Could the low prevalence of diabetes mellitus in the Egyptian deserts be the result of protective element in the Bedouin diet?. The Egyptian. J. of Diabetes. 1(1): 23-25.

Banerji M. and Lebovitz H. (1989) : Insulin sensitive and insulin resistant variants in IDDM. Diabetes. 38: 784-792.

Barnett A.H.; Eff C.; and Leslie R.D.G. (1981) : Diabetes in indication twins. Diabetologia. 20: 87-93.

Baron AD and Brechtel G; (1993): Insulin differentially regulate systemic and skeletal muscular vascular resistance. Am. J. phsiol. 265: E61-E67.

Barroso I; Gurnell M.; and Crowley V.E.F. (1999) : Dominant negative mutations in human PPAR γ associated with severe insulin resistance, diabetes mellitus and hypertension. Nature. 402: 880-3.

Beatty OL; Harper R; Sheridan B; Atkinson AB and Bell OM; (1993): Insulin resistance in offspring of hypertensive patients. BMJ. 307: 92-96.

Beischer W. (1976) : Human C-peptide. Part I Radioimmunoassay. Klin. Wscr. 54: 709.

Beischer W. (1976) : Human C-peptide. Part II clinical studies. Klin. Wscr. 54: 717.

Bell P.; Firth R. and Rizza R. (1989) : Assessment of the postprandial pattern of glucose metabolism in nondiabetic subjects and patients with noninsulin-dependent diabetes mellitus using a simultaneous infusion of (2-³H) and (3-³H)- glucose. Metabolism. 38: 38-45.

Bell PM. (1996): Clinical significance of insulin resistance. Diabet Med, 15: 132-48.

Bennet PH; (1990): Epidemiology of diabetes mellitus. In Rifkins M & Porte D (acts.). diabetes Theory and Practice. 4th. Ed. El-Sovier. 337-77.

Bennett S.T. and Todd J.A. (1996) : Human type I diabetes and the insulin gene : principles of mapping polygenes. Annual Reviews in Genetics. 30: 343-370.

Bergman R.; Phillips J. and Cobelli C. (1981) : Physiologic evaluation of factors controlling glucose tolerance in man : measurement of insulin sensitivity and cell glucose sensitivity from the response to intravenous glucose. J. Clin. Invest. 68: 1456-67.

Bernard DJ; Rifkin H; Omara MH; and Zaied MA; (1992): Standards of case of diabetic patients in a typical Egyptian Community. British. J. of General Practice. 46: 401-5.

Bjorntorp P. (1999) : Neuro-endocrine perturbations as a cause of insulin resistance. Diabetes. Metab. Res. Rev. 15: 427-441.

Bjorntorp P; (1990): "Portal" adipose tissue as a generator of risk factors for cardiovascular disease and diabetes. Arteriosclerosis. 10: 493-97.

Blanchard J.F.; Ludwig S.; and Wajda A. (1996) : Incidence and prevalence of diabetes in Manitoba 1986-1991. Diabetes Care 19: 807-811.

Blix P.M. (1982) : Urinary C-Peptide : an indicator of B-cell secretion under different metabolic conditions. J. Clin. Endocrinol. Metab. 54:3,574.

Boden G. (1997) : Role of fatty acids in the pathogenesis of insulin resistance and NIDDM. Diabetes. 46: 3-10.

Bogardus C.; Lillioja S.; Mott D.M.; Hollenbeck C. and Reaven G. (1985) : Relationship between degree of obesity and in vivo insulin action in man. Am. J. Physiol. 248: E286-E291.

Bogardus C; Lilloja S.; Stone K., and. Mott D; (1984): Correlation between muscle glycogen synthase activity and in vivo insulin action in man. J. Clin. Invest. 73: 1185-90.

Bonadonna R.C. and Bonora E. (1997) : Glucose and free fatty acid metabolism in human obesity. Relationship with insulin resistance. Diabetes Rev. 5: 21-51.

Bonadonna R.C.; Croop L.C.; and Zych K. (1990) : Dose-dependent effect of insulin of plasma free fatty acid turnover oxidation in humans. American Journal of Physiology. 259: E736-E750.

Bonadonna R.C.; Del-Patro S; Borona E; Succomani MP; Guilli G; Natali A; Frascerra S; Pecori N; Ferrannini E; Bier D; Cobelli C; and DeFronzo RA; (1996): Roles of glucose transport and glucose phosphorylation in muscle insulin resistance of NIDDM. Diabetes. 45: 915-25.

Bonora E.; Targher G.; Alberiche M.; bonadonna R.C.; and Saggiani F.; Zenere M.B., (2002) : Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity. Studies in subjects with various degrees of glucose tolerance and insulin sensitivity. Diabetes Care. 23: 57-63.

Borkman M; Storlien LH; Pan DA; Jenkins AB; Chisholm DJ and Campbell IV; (1993): The relationship between insulin

sensitivity and the fatty acid composition of skeletal muscle phospholipids. *N. Engl. J. Med.* 328: 238-44.

Bouchard C. (1995): Genetics and the metabolic syndrome. *Int. J. Obes.* 19: 552-9.

Bouchard C. (1997) : Genetics of human obesity : recent results from linkage studies. *Journal of Nutrition.* 127: 18875-18905.

Bouchard C; Perusse L; Deriaz O; Despres JP; and Tremblay A; (1993): Genetic influences on energy expenditure in humans. *Crit. Rev. Food Sci. Nutr.* 33: 345-50.

Boulay MR; Ama PFM; and Bouchard C; (1988): Racial variation in work capacities and powers. *Canadian J. of Sports Science.* 13: 127-35.

Boulton TGC; Margarey AM and Cockington RA; (1995): Serum lipids and apolipoprotein from 1 to 15 years: changes with age and puberty, and relationships with diet, parental cholesterol and family history of ischaemic heart disease. *Acta. Paediatr.* 48: 1113-8.

Bressler R. and Johnson D.G. (1997) : Pharmacological regulation of blood glucose levels in non-insulin- dependent diabetes mellitus. *Arch. Intern. Med.* 157: 836-848.

- Brown S.D.M. and Nolan P.M. (1998) :** Mouse mutagenesis- systematic studies of mammalian gene function. *Human Molecular Genetics*. 7: 1627-1633.
- Brutis CA and Ashwood ER; (1994):** Carbohydrates. *Clin. Chemistry*. W.B Saunders Comp. 2nd. Edition. Vol. (2). Pages: 936-39.
- Buffington CK and Kitabchi AE; (1994):** Evidence for a defect in insulin metabolism in hyper-androgenic women with polycystic ovarian syndrome. *Metabolism*. 43: 1367-72.
- Burke CM; Bullough RC; and Melby CL; (1993):** Resting metabolic rate and post-prandial thermogenesis by level of aerobic fitness in young women. *Eur. J. Clin. Nutr.* 47: 575-85.
- Buthlezi HE; Valsania P; Halter JB; and Lin X; (1998):** Differential effects of BMI on diabetes risk among black and white Americans. *Diabetes Care*. 21: 1828-35.
- Butkiewicz EK; Leibson C; O'Brien PC; Palumbo PJ; and Rizza RA. (1995):** Insulin therapy for diabetic keto-acidosis. *Diabetes Care*. 18:1187-90.
- Byrne C.D.; Wareham N.J.; and Day N.E. (1995) :** Decreased non-esterified fatty acids suppression and features of the insulin resistance syndrome occur in a sub-group of individuals with normal glucose tolerance. *Diabetologia*. 38: 1358-66.

Cambell M.J. & Machin D. (1993): In: Medical Statistics. A common sense approach. 2nd. Ed. Lohm Willey & sons. New- york; Chichester; Brisbane; Toronto & Singapore. Page 31-35.

Carpenter S.M. & Coustan R.C. (1982): Diagnosis of diabetes mellitus: New criteria. J. Clin Endocrinol Metab 61: 1-6.

Ceesay S.M.; Prentice A.M.; and Day K.C.(1989) : The use of heart rate monitoring in the estimation of energy expenditure : a validation study using indirect whole-body calorimetry. Br. J. Nutr. 61: 175-86.

Chan J.M; Rimm E.B. and Colditz G.A; (1994): Obesity, fat distribution and weight gain as risk factors for clinical diabetes in men. Diabetes Care. 17: 961-69.

Charles M.A.; Fontbonne A.; and Thibult N.(1991): Risk factors for NIDDM in white population. Paris prospective study. Diabetes. 40: 796-799.

Chisholm DJ; Campbell LV; and Kraeger EW.; (1997): Pathogenesis of insulin resistance syndrome (syndrome X). Clin Exp Pharmacol Phsiol. 24:792-4.

Chou CJ; Haluzik M; Gregory C; Dietz KR; Vinson C; Gavrilova O; and Reitman ML; (2002): WY14,643, a peroxisome proliferator-activated receptor alpha agonist, improves hepatic

and muscle steatosis and reverses insulin resistance in lipotrophic A-ZIP/F-1 mice. *J. Biochem.* 5;277(27):24484-9.

Christine HL; Barbara AG; and Michael IG; (2000): Role of dietary factors in ethnic differences in early risk of cardiovascular disease and type 2 diabetes. *Am. J. of Clin. Nutr.* 71: 725-32.

Clark P. (1993) : Evaluation of medgenix insulin-irma. Addenbrooke's hospital Cambridge. UK. *Clin. Endocrinol.* 32: 689-93.

Clement K.; Hereberg S.; and Passinge B.(2000) : The pro115 Gln and Pro 12Ala PPAR γ mutations in obesity and type 2 diabetes. *Int. J.Obes. Relat. Metab. Disord.* 24: 391-3.

Considine R.V.; Sinha M.K.; and Heiman M.L.(1996) : Serum immunoreactive-leptin concentrations in normal weight and obese humans. *New England Journal of medicine.* 334: 292-295.

Cooper R.S.; Rotimi C.N.; and Kaufman J.S. (1997) : Prevalence of NIDDM among population of the African Diaspora. *Diabetes Care.* 20: 343-348.

Cowie CC; Harris MI; Silverman RE; Johnson EW; and Rust KF; (1993): Effect of multiple risk factors on differences between blacks and whites in the prevalence of NIDDM subjects in the US. *Am. J. Epidemiol.* 137: 719-32.

Cruickshank GK; Cooper J; Burnett M; MacDuff J; and Drubra U;
(1991): Ethnic differences in fasting plasma C-Peptide and
 insulin in relation to glucose tolerance and blood pressure.
 Lancet. 5;338(8771) 842-7.

Damsbo PA; Vaag O; Holter-Nielsen and H. Beck-Nielsen (1991):
 Reduced glycogen synthase activity in skeletal muscle from
 obese patients with and without diabetes mellitus. Diabetologia.
 34:239-45.

Daniel W.W. (1987): In: Biostatistics, A foundation for analysis in the
 health sciences. 4th. Ed. John Willey & Sons. New-york;
 Chichester; Brisbane; Toronto & Singapore. Page 110-12.

Daughaday W.H; (1990): The possible autocrine/paracrine and
 endocrine roles of insulin-like growth factors of human tumors.
 Endocrinology. 127:1-4.

Davidson M.B. and Peters A.L. (1997) : An overview of metformin in
 the treatment of type 2 diabetes mellitus. Am.J. Med. 102: 99-
 110.

Davis TME, Jackson D; Balme M and Bruce DG. (1999): . Prevalence
 of, and risk factors for, hyperlactaemia in metformin-treated
 patients: the Fremantle Diabetes Study. Diabetologia 42 (Suppl.
 1): 225.

Day CP; Grove J; Daly AK; Stewart MW; Aveny PJ; and Walker M. (1998): Tumor necrosis factor-alpha gene promoter polymorphism and decreased insulin resistance. *Diabetologia*, 430-4

Deeb S.S.; Fajas L.; and Nemoto M. (1998) : A Pro 12Ala substitution in PPAR γ associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. *Nat. Genet.* 20: 284-7.

DeFronzo R.A.; Bonadonna R. and Ferrannini E. (1992) : Pathogenesis of NIDDM : a balanced overview. *Diabetes Care* 15: 318-68

DeFronzo R.A; Ferrannini E. and Simonson D. (1989) : Fasting hyperglycemia in noninsulin-dependent diabetes mellitus : contributions of excessive hepatic glucose production and impaired tissue glucose uptake. *Metabolism*. 38: 387-95.

DeFronzo R.A; Gunnarsson R.; and Bjorkman O. (1985) : Effects of insulin on peripheral and splanchnic glucose metabolism in noninsulin-dependent (type II) diabetes mellitus. *J. Clin. Invest.* 76: 149-55.

DeFronzo RA; Jacot E; Jequier E; Maeder E; Wahren J. and Felber JP; (1981): The effect of insulin on the disposal of intravenous glucose. *Diabetes*. 30: 1000-1007.

Del- Prato S; Leonetti F; Simonson DC; Sheehan P; Matsuda M and DeFronzo RA; (1994): Effect of sustained physiologic hyperinsulinemia on insulin secretion and insulin sensitivity in man. *Diabetologia*. 37: 1025-35.

Department of Health (1994) : The health of the nation. A consultative Department of health in England. Report. London. : HMSO.

Desvergne B. and Wahli W. (1999) : Peroxisome proliferator activated receptors : nuclear control of metabolism. *Endocr. Rev.* 20: 649-88.

Diabetes Prevention Program Research Group (1999) : The diabetes prevention program : design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care* 22: 623-634.

Dietz WH; (1996): Is reduced metabolic rate associated with obesity?. *J. Pediatr.* 129: 621-23.

Dobson A.J.; Evans A.; and Ferrario M. (1998) : Changes in estimated coronary risk in the 1980s : data from 38 populations in the WHO Monica project. *Annals of Medicine* 30: 199-205.

Doumas B;(1971). Cholorimetric method using bromocresal green. *Clin. Chem. Acta.* 31: 87.

Duckworth CD; (1988): Insulin degradation: mechanisms, products and significance. *Endocr. Rev.* 9: 319-45.

Eisenbarth G.S; (1986): Type I diabetes mellitus: A chronic autoimmune disease. *N. Engl. J. Med.* 314:1360-86.

Elchebly M.; Payette P.; and Michaliszyn E. (1999) : Increased insulin sensitivity and obesity resistance in mice lacking the protein tyrosine phosphatase- IB gene. *Science.* 283: 1544-1548.

El-Dib (1991): Prevalence of DM. In Northern Sinai. Personal communication.

Emilsson V.; Summers R.J.; and Hamilton S. (1998) : The effects of the beta-3 adrenoceptor agonist BRL 35135 on UCP isoform mRNA expression. *Biochemical and Biophysical Research Communications.* 252: 450-454.

Emoto M.; Nishizawa Y.; Maekawa K.; Hiura Y.; Kanda H.; and Kawagishi T. (1999) : Homeostasis model assessment as a clinical index of insulin resistance in type 2 diabetic patients treated with sulphonylureas. *Diabetes. Care.* 22: 818-822.

Endre T; Mattiasson I; Hulten UL; Lindegarde F and Berglund G; (1994): Insulin resistance is coupled to low physical fitness in

normotensive men with a family history of hypertension. *J. Hypertension*. 12: 81-88.

Engelgau T.Z; Homer R.W; Sirlud L. and Safey D. (1993): incidence of retinopathy in different ethnic populations. *Diabetes Care*. 24: 419-25.

Engelgau T.Z., Finch R.Q., Abter S., and Nael K.J. (1997): insulin secretion and action. *N.Engl L. Med* 334: 777-84.

Engelgau. T.Z.; Narayan k.M.V. and Herman W.H. (2000) : Screening for type 2 diabetes (Technical Review). *Diabetes Care* 10: 1563-1580.

Erikson J; A. Franssila-Kallunki; A. Ekstrand; C. Saloranta; E. Widen; C. Schalin and L. Groop;(1989): Early metabolic defects in persons at increased risk for NIDDM. *N. Engel. J. Med.* 318: 1217-25.

Evans J.C.; Frayling T.M.; and Cassell P.G. (2001) : Studies of association between the gene for calpain-10 and type 2 diabetes mellitus in the United Kingdom. *Am.J. Hum. Genet.* 69: 544-52.

Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (1997) : Report of the Expert Committee on the

Diagnosis and Classification of Diabetes Mellitus. Diabetes Care. 20: 1183-1197.

Expert Committee on the Diagnosis and Classification of Diabetes Mellitus (2002) : Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Diabetes Care. 25: S5-S16.

Expert panel of the IFCC., 1976.

Fagius J. and Berne C. (1994) : Increase in muscle nerve sympathetic activity in humans after food intake. Clinical Science. 86: 159-167.

Fajans S. (1989) : Maturity-onset diabetes of the young (MODY). Diabetes Metab. Rev. 5: 579-606.

Fajans S.S.; Bell G.I.; and Polonsky K.S. (2001) : Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. New Engl. J. Med. 345: 971-80.

Fatani HH; Mira SA; and El-Lubiee (1987): Prevalence of DM in Rural Saudi Arabia. Diabetes Care. 10 (2).

Felber JP; E. Ferrannini A. Golay; HU Meyer; D. Thiebaud; B. Curchod; E. Maeder; E. Jequier and RA DeFronzo; (1987):

Role of lipid oxidation in pathogenesis of insulin resistance of obesity and type-2 diabetes. *Diabetes*. 36: 1341-50.

Felber JP; Thiebaud D; Maeder E; Jequier E; Hendler R. and DeFronzo RA; (1983): Effect of somatostatin-induced insulinopenia on glucose oxidation in man. *Diabetologia*. 25: 325-30.

Feldman RD and Bierbrier GS; (1993): Insulin-mediated vasodilatation: impairment with increased blood pressure and body mass. *Lancet*. 342: 707-9.

Ferrannini E; Wahren J; and Felig P. (1980) : The role of fractional glucose extraction in the regulation of splanchnic glucose metabolism in normal and diabetic man. *Metabolism*. 29: 28-35.

Ferrannini E; (1998): Insulin resistance versus insulin deficiency in NIDDM patients. Problems and prospects. *Endocrinol. Rev.* 19: 477-90.

Ferrannini E; and Cobelli C; (1987): The kinetics of insulin in man. II. Role of the liver. *Diabetes. Metab. Rev.* 3: 365-97.

Ferrannini E; Wahren J; Faber OK; Felig P; Binder C; and DeFronzo RA; (1983): Splanchnic and renal metabolism of insulin in human subjects: a dose response study. *Am. J. Physiol.* 244: E517-E527.

- Finch P.M; Aiden C; Morris B.J. and Lateman D. (1999):*** Diagnosis of diabetes mellitus: Correlation of FPG with 2-h plasma glucose. Diabetes Care. 35: 1118-23.
- Fitzgerald A.P. and Jarrett R.J. (1991) :*** Are conventional risk factors for mortality relevant in type 2 diabetes. Diabetic Medicine. 8: 475-480.
- Flegal K; Ezzat T; and Hanis H; (1991):*** Prevalence of diabetes in Mexicans , Cubans and Puerto-Ricans from the Hispanic Health and Nutritional Epidemiology Survey 1992-1994. Diabetes Care. 14 (suppl 3) 626-56.
- Flier J.S. (1997) :*** Leptin expression and action : new experimental paradigms. Proceedings of the National Academy of Sciences of the US. 94: 4242-4245.
- Flier J.S.; Kahn C.R. and Roth J. (1979) :*** Receptors antireceptor antibodies and mechanisms of insulin resistance. N.Engl. J. Med. 300. 8: 413-419.
- Food and Nutrition Board (1995) :*** Criteria for evaluating weight management programs. In Thomas PR (Ed.). Nature and Problem of Obesity. Pp. 10-11. Washington. D.C. : National Academy Press.

- Foppa I. and Spiegelman D. (1997) :*** Power and sample size calculations for case-control studies of gene-environment interactions with a polytomous exposure variable. *Am. J. Epidemiol.* 146: 596-604.
- Foresch E.R & Zapf J. (1985):*** Insulin-like growth factors and insulin: comparative aspects. *Diabetologia.* 28:485-93.
- Fraser GE; (1994):*** Diet and coronary heart disease: beyond dietary fats and low-density-lipoprotein cholesterol. *Am. J. Clin. Nutr.* 59(suppl): 1117S-23S.
- Frier B.M.; Ashby J.P.; Nairn I.M. and Bairs J.D. (1981) :*** Plasma insulin, C-peptide and glucagon concentrations in patients with insulin-dependent diabetes treated with chlorpropamide. *Diab. Metab.* 7. 1: 45-49.
- Froguel P. (1997) :*** Tracking down genes to cure diabetes : an achievable task for the 21st century? *Diabetes and Metabolism.* 23(Supplement 2) : 8-13.
- Froguel P.; zouali H.; and Vionnet M., (1993) :*** Familiar hyperglycemia due to mutations in glucokinase. Definition of a subtype of diabetes mellitus. *N. Engl. J. Med.* 328: 687-702.

- Fujimoto W.Y. (1996) : Overview of non-insulin dependent diabetes mellitus (NIDDM) in different population groups. Diabetic Medicine. 13: 57-510.***
- Fujimoto W.Y.; Leonetti D.L.; and Kinyour I.L. (1987) : Prevalence of diabetes mellitus and impaired glucose tolerance among second generation lapanser- American Men. Diabetes. 36: 721-9.***
- Fukagama NK; Anderson JW; Hageman G; Young VR; and Minaker KL; (1990): High-carbohydrate, high-fiber diets increase peripheral insulin sensitivity in healthy young and old adults. Am. J. Clin. Nutr. 52: 524-8.***
- Furnsinn C.; Brunmair B.; and Furtmullar R. (1998) : Failure of leptin to affect basal and insulin-stimulated glucose metabolism of rat skeletal muscle in vitro. Diabetologia. 41: 524-529.***
- Galli J.; Li L.S.; and Glaser A. (1996) : Genetic analysis of non-insulin dependent diabetes mellitus in the GK rat. Nature Genetic. 12: 31-37.***
- Gauguier D.; Froguel P.; and Parent V. (1996) : Chromosomal mapping of genetic loci associated with non-insulin dependent diabetes in the GK rat. Nature Genetics. 12: 38-43.***

Gerich E; (2000): Contributions of impaired insulin secretions and insulin sensitivity in type 2 diabetes mellitus: current therapeutic implications. Dialogue. 3 page 17.

Gerich J. (1988): Glucose counter-regulation and its impact on diabetes mellitus. Diabetes. 37:1608-17.

Gerich J. (1991): Is muscle the major site of insulin resistance in type 2 (non insulin-dependent) diabetes mellitus? Diabetologia. 34: 607-10.

Gerich J. (1993): Control of glycaemia. Bailliere's Clin. Endocrinol. Metab. 7: 551-86.

Gerich J. (1999) : Is insulin resistance the principal cause of type 2 diabetes? Diabetes, Obesity and Metab. 1: 1-7.

Gerich J; (2000): Contribution of impaired insulin secretion and insulin sensitivity in type 2 diabetes mellitus: current evidence and therapeutic implications. Dialogue 23: 19-22.

Gerich J.; Mitrakou A.; Kelley D.(1990) : Contribution of impaired muscle glucose clearance to reduced post-absorptive systemic glucose clearance in NIDDM. Diabetes. 39: 211-16.

Ghosh S.; Watanabe R.M.; and Valle T.T (2000) : The Finland-United States Investigation of non-insulin-dependent diabetes mellitus genetics (FUSION) study. I. An. Autosomal genome scan for genes that predispose to type 2 diabetes. American Journal of Human Genetics. 67: 1174-1185.

Glombitza KW; Mahran GH; and Mirhom YM; (1994): Hypoglycemic effect of Zizyphus Spinachristi in rats. Plant Med. 60: 244-7.

Goran MI; Nagy TR and Treuth MS; (1997): Visceral fat in white and African-American prepubertal children. Am. J. Clin. Nutr. 65: 1703-8.

Gordon TM; (1977). Determination of cholesterol after precipitation of chylomicrons, VLDL and LDL by phosphotungestic acid and magnisium chloride. Amer. J. Med.62: 707.

Gower BA; Granger WM; Franklin F; Shewchuk RM; and Goran NI; (2002): Contribution of insulin secretion and clearance to glucose-induced insulin concentration in African-American and Caucasian children. J. Clin. Endocrinol Metab. 87(5) 2218-24.

Gower BA; Nagy TR; and Goran MI; (1999): Visceral fat, insulin sensitivity, and lipids in prepubertal children. Diabetes. 48: 1515-21.

Granberry MC; Schneider EF; and Fonseca VA. (1998): The role of triglitazone in treating the insulin resistance syndrome. Pharmacotherapy, 18:937-87.

Grassi G.; Cattaneo B.M.; and Seravalle G. (1996) : Obesity and the sympathetic nervous system. Blood Pressure. 5 (Supplement 1): 43-46.

Groop L; Ekkstrand A. and Forsblom C; (1993): Insulin resistance, hypertension and microalbuminuria in patients with type 2 diabetes mellitus. Diabetologia. 36: 642-47.

Guarino MP; Afonso RA; Raimundo N; Raposo JF; and Macedo MP; (2003): Hepatic glutathione and nitric oxide are critical for hepatic insulin-sensitizing substance action. Am. J. Physiol. Gastrointest. Liver. Physiol. 284(4): G588-94.

Guo S.W. (2000) : Gene environment interaction and the mapping of complex traits : some statistical models and their implications. Hum. Hered. 50: 286-303.

Haffner S.M.; Stern M.P.; and Mitchell B.D. (1990) : Incidence of type II diabetes in Mexican Americans predicted by fasting insulin and glucose levels, obesity, and body-fat distribution. Diabetes. 39: 283-288.

Haffner SM.; (1997): Progress in population analysis of the insulin resistance syndrome. Ann NY Acad Sci, 827:1-12.

Haffner SM; (1998) : Management of dyslipidemia in adults with diabetes (technical review). *Diabetes Care* 21: 160-178.

Haffner SM; Stern MP; Watanabe RM; and Bergman RN; (1992): Relationship of insulin clearance and secretion to insulin sensitivity in non-diabetic Mexican Americans. *Eur. J. Clin. Invest.* 22: 147-53.

Haffner SM; Valdez R.A.; Hazuda H.H.; Mitchell B.D.; Morales P.A.; and Stern M.P. (1992) : Prospective analysis of the insulin resistance syndrome (syndrome X). *Diabetes.* 41: 715-722.

Hager J.; Hansen L.; and Vaisse C. (1995) : A missense mutation in the glucagon receptor gene is associated with non-insulin-dependent diabetes mellitus. *Nature Genetics.* 9: 299-304.

Haggag K.M.S., (1990): Prevalence of diabetes mellitus among a section of the Egyptian population at the governorate of the Red –Sea; a part of multinational Center Study in the Mediterranean Countries. Thesis, University of Alexandria. P.23-26.

Hales C.N. and Barker D.J. (1992) : Type 2 (non-insulin-dependent) diabetes mellitus : the thrifty phenotype hypothesis. *Diabetologia.* 35: 595-601.

Hales C.N.; Barker D.J.; and Clark P.M. (1991): Fetal and infant growth and impaired glucose tolerance at age 64. British Medical Journal. 303: 1019-1022.

Hamman R.F. (1992): Genetic and environmental determinants of non-insulin-dependent diabetes mellitus (NIDDM). Diabetes Metab. Rev. 8: 287-338.

Han T.S.; Van Ler E.M.; Seidell J.C. and Lean M.E.J. (1995) : Waist circumference action levels in the identification of cardiovascular risk factors : prevalence study in a random sample. British Medical Journal. 311: 1401-1405.

Handberg A; Vaag A; Vinten J. and Beck-Nielsen H; (1993): Decreased tyrosine kinase activity in partially purified insulin receptors from muscle of young, non-obese first degree relatives of patients with type 2 diabetes mellitus. Diabetologia. 36: 668-47.

Hani E.H.; Boutin P.; and Durand E. (1998) : Missense mutations of the pancreatic islet beta cell inwardly rectifying K⁺ channel gene (KIR6.2/BIR): a meta-analysis suggests a role in the polygenic basis of type II diabetes mellitus in Caucasians. Diabetologia. 41: 1511-1515.

Hanis C.L.; Boerwinkle E. and Chakraborty R. (1996) : A genome-wide search for human non-insulin-dependent type 2 diabetes genes reveals a major susceptibility locus on chromosome 2. Nat. Genet. 13: 161-71.

Hanson R.L.; Ehm M.G.; and Pettitt D.J. (1998) : An autosomal genomic scan for loci linker to type II diabetes mellitus and body-mass index in Pima Indians. Am. J. Hum. Genet. 63: 1124-32.

Hanson R.L.; Pettitt D.J.; and Bennett P.H. (1995) : Familial relationships between obesity and NIDDM. Diabetes. 44: 418-22.

Hanson R.L.; Pratley R.E.; and Bogardus C. (2000) : Evaluation of simple indices of insulin sensitivity and insulin secretion for use in epidemiologic studies. Am. J. Epidemiol. 151: 190-8.

Hara K.; Okada T.; and Tobe K. (2000) : the Pro12Ala polymorphism in PPAR γ 2 may confer resistance to type 2 diabetes. Biochem. Biophys. Res. Commun. 271: 212-6.

Harding A.H.; Sargeant L.A.; and Welch L.A. (2001) : Fat consumption and HbA $_{1c}$ levels: The EPIC-Norfolk study. Diabetes Care. 24: 1911-6.

Harris M.I. (1989) : Impaired glucose tolerance in the U.S. population.
Diabetes Care. 12: 464-474.

Harris M.I; Cowie CC; Gu K; Francis ME; Flegal K; and Eberhardt MS; (2002): Higher fasting insulin but lower fasting C-Peptide levels in African-Americans in the US population. Diabetes. Metab. Res. Rev. 18(2): 149-55.

Hattersley AT and Tooke JE; (1999) : The fetal insulin hypothesis an alternative explanation of the association of low birth weight with diabetes and vascular disease. Lancet. 353: 1789-1792.

Haynes W.G.; Sivitz W.I.; and Morgan D.A. (1997) : Sympathetic and cardiorenal actions of leptin. Hypertension. 30: 619-623.

Helmrich SP; Ragland DR; Leung RW and Paffenbarger RS; (1991):
Physical exercise and reduced occurrence of NIDDM. N. Engl. Med. 325: 147-52.

Hemdan; G; (1993): Sinai. Kitab Al-Hilal. No. 311. Dar Al-Hilal. No. 309. (published). Cairo. (arabic). **P: 1213-17.**

Hemdan; G; (1993): The personality of Egypt. Kitab Al-Hilal (publisher). Cairo. (arabic). **P: 2145-66.**

Hennis MMI; Shrago E; and Kissebah AH; (1990): Receptor and post-receptor effects of free fatty acids on hepatic insulin dynamics. Int. J. Obesity. 14: 831- 41.

Henriksen JE; Alford F; and Handelsberg A; (1994): Increased glucose effectiveness in normoglycaemic but insulin resistant relatives of patients with NIDDM. J. Clin. Invest. 94: 1196-1204.

Hoekstra J.B.L; van Rijn H.J.M; and Erkelens D.W. (1982): C-Peptide. Diabetes Care. 5:438-46.

Hoet J.J.; Tripathy B.B.; Rao R.H. and Yajnik C.S. (1996) : Malnutrition and diabetes in the tropics. Diabetes Care. 19: 1014-1017.

Holmang A; Brezezinka Z and and Bjorntorp P; (1993): Effects of hyperinsulinemia on muscle fiber composition and capillarization in rats Diabetes. 42:1073-91.

Horikawa Y.; Oda N.; and Cox N.J. (2000) : Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. Nat. Genet. 26: 163-75.

Hosker JP; Burnett MA; Davies EG; Matthews DR; Rudenski AS and Turner RC; (1984): loss of normal sigmoidal beta cell response curve to glucose in NIDDM patients. Diabetologia. 27: 289 A.

Hotamisligil G.S.; Arner P.; and Caro J.F. (1995) : Increase adipose tissue expression of tumor necrosis factor- alpha in human obesity and insulin resistance. *Journal of Clinical Investigation.* 95: 2409-2415.

Hotamisligil G.S.; Budavari A.; Murray D. and Spiegelman B.M. (1994) : Reduced tyrosine kinase activity of the insulin receptor in obesity-diabetes. Central role of tumor necrosis factor-alpha. *Journal of Clinical Investigation.* 94: 1543-1549.

Hotamisligil G.S.; Shargill N.S. and Spiegelman B.M. (1993) : Adipose expression of tumor necrosis factor- alpha : direct role in obesity-linked insulin resistance. *Science.* 259: 87-91.

Hother-Nielsen O.; Henriksen J.; and Holst J. (1996) : Effects of insulin on glucose turnover rates in vivo : isotope dilution versus constant specific activity technique. *Metabolism.* 45: 82-91.

Houghton P. (1991) : Selective influences an morphological variation amongst . Pacific Homo Sapiens. *J. Hum. Evol.* 41-51.

Howard B.V. (1999) : Insulin resistance and lipid metabolism. *Am.J. Cardiol.* 84: 28-32.

- Howitt A.J. and Cheales N.A. (1993) :** Diabetes registers : a grassroots approach. British Medical Journal. 307: 1046-1048.
- Huang W.; Connor E.; DelaRosa T.; Muir A.; Schatz D.; Silverstein J.; Crockett S.; She J.X. and Maclaren N.K. (1996) :** Although DR3-DQB1* may be associated with multiple component diseases of the autoimmune polyglandular syndromes, the human leukocyte antigen DR4-DQB 110302 haplotype is implicated only in beta cell autoimmunity. J. Clin. Endocrinol. Metab. 81: 1-5.
- Humphriss DB; Stewart MW; Berrish TS; Barriocanal LA; Trajano LR; Ashworth LA; Brown MD; Miller M; Avery PJ; Alberti KGMM and Walker M; (1997):** Multiple metabolic abnormalities in normal glucose tolerant relatives of NIDDM families. Diabetologia. 40: 1185-90.
- Hwang S.J.; Beaty T.H.; and Liang K.Y. (1994) :** Minimum sample-size estimation to detect gene environment interaction in case-control designs. Am. J. Epidemiol. 140: 1029-37.
- Ismail A.A. (1998) :** The Burden of diabetes in North Liverpool : Epidemiology. Complications and coronary artery disease risk factors. Ph.D. thesis. University of Liverpool. P. 144-51.

IStock M. (1998) : Energy balance and animal models of obesity. In Kopelman PG and Stock M (eds). Clinical Obesity. Pp. 50-72. Oxford : Blackwell Science.

Jaspan J; and Polonsky k; (1982): Glucose ingestion in dogs alters the hepatic extraction of insulin. In vivo evidence for a relationship between the biological action and extraction of insulin. J. Clin. Invest. 69: 516-25.

Jemenez F; Gumbiner B; Flynn T; and Thorburn AW; (1987): Effect of weight loss in massive obesity on insulin and C-Peptide dynamics: sequential changes in insulin production, clearance, and sensitivity. J. Clin. Endocrinol. Metab. 64: 661-68.

Jendrassik J, and Grof P; (1983). Photometric method using Jendrassik and Grof technique. Biochem. 297: 81.

Jervell J. (1997) : Type 2 diabetes (NIDDM) can be prevented. IDF Bulletin. 42: 1-3.

Jones CNO; Pei D; Staris P; Polonsky KS; Chen Y-D. I and Reaven GM; (1997): Alterations in the glucose-stimulated secretory dose response curve and in insulin clearance in non-diabetic insulin resistant individuals. J. Clin. Endocrinol. Metab. 82: 1834-38.

Jones CNO; Abbasi F; Carantoni M; Polonsky KS; and Reaven GM; (2000): Roles of insulin resistance and obesity in regulation of plasma insulin concentrations. *Am. J. Endocrinol. and Metab.* 278(3): E501.

Judzewitsch R.G.; Pfeifer M.A.; Best J.D.; Beard J.C.; Halter J.B. and Porte D. Jr. (1982) : Chronic chlorpropamide therapy of non-insulin dependent diabetes augments basal and stimulated insulin secretion by increasing islet sensitivity to glucose. *J.Clin. End. And Metab.* 55. 2: 321-328.

Kageyama T.; Nishikido N.; and Honda Y. (1997) : Effects of obesity, current smoking status, and alcohol consumption on heart rate variability in male white-collar workers. *International Archives of Occupational and Environmental Health.* 68: 447-454.

Kahn C.R.; Vincent C. and Doria A. (1996) : A genetics of non-insulin-dependent (type II) diabetes mellitus. *Annual Review in Medicine.* 47: 509-531.

Kaprio J.; Tuomilehto J.; and Koskenvuo M. (1992) : Concordance for type I (insulin-dependent) and type 2 (non-insulin dependent) diabetes mellitus in a population based cohort of twins in Finland. *Diabetologia* 35: 1060-1067.

Karakash C; Assimacopoulos-Jeannet; and Jeanrenaud B; (1976): An anatomy of insulin removal in perfused livers of obese-hyperglycemic mice. *J. Clin. Invest.* 57: 1117-24.

Karam JH. (1996): Reversible insulin resistance in non-insulin-dependant diabetes mellitus. *Horm. Metab. Res.* 28: 440-4.

Karakas SEK; (2000): Ethnic differences in the insulin resistance syndrome. *Am. J. of Clin. Nutrition.* 71: 670-71.

Kelley D. and Simoneau J. (1994) : Impaired free fatty acid utilization by skeletal muscle in non-insulin dependent diabetes mellitus. *J. Clin. Invest.* 94: 2349-56.

Kelley D.; Mitrakou A.; and Marsh H. (1988) : Skeletal muscle glycolysis oxidation and storage of an oral glucose load. *J. Clin. Invest.* 81: 1563-71.

Kelley D.; Mookan M. and Veneman T. (1994) : Impaired post-prandial glucose utilization in non-insulin dependent diabetes mellitus. *Metabolism.* 43: 1549-57.

Kelley DE and Mandarino LJ; (1990): Hyper-glycemia normalizes insulin-stimulated skeletal muscle glucose oxidation and storage in type-2 diabetes mellitus. *J. clin. Invest.* 86: 1999-2007.

Kenny S.J. (1995): Prevalence and incidence of NIDDM. In Haris MI, Cowie CC, Reiber G et al. (eds). Diabetes in America. 2nd edition. NIH publication NO. 95-1468. Pp. 47-68. Washington, DC : US Government Printing Office.

Khan S.E. and Porte D.Jr; (1996): Patho-physiology of type 2 diabetes: implications for treatment. In Porte D.Jr & Sherwin R.S (eds). Diabetes mellitus; theory and practice pp 478-512. New-york: Appleton and Lange.

Khaw K.T. (1994) : Genetics and environment : Geoffrey rose revisits. Lancet. 343: 838-839.

Khaw K.T.; Wareham N.J.; and Luben R. (2001) : Glycated haemoglobin, diabetes, and mortality in men in Norfolk cohort of European Prospective Investigation of Cancer and Nutrition (EpiC-Norfolk). BMJ. 322: 15-8.

Khoury M.J. and Flanders W.D. (1996) : Nontraditional epidemiologic approaches in the analysis of gene-environment interaction : case control studies with no controls! Am.J. Epidemiol. 144: 207-13.

Kim KS; Owen WL; Williams D; and Adams-Campbell; (2000): A comparison between BMI and Conicity index on predicting

coronary heart disease: Framingham Heart Study. *Ann. Epidemiol.* 10: 424-31.

King H. and Rewers M. (1993) : Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults. *Diabetes Care.* 1: 157-77.

King H. and Rewers M. on behalf of the WHO Ad Hoc Diabetes Reporting Group (1991) : Diabetes in adults in now and third world problem. *Bulletin of the World Health Organization.* 60: 642-648.

King S.L and John P. W. (1999): Obesity and diabetes. *Balliere Clinical Endocrinology and Metabolism.* 13: 221-37

Kissebah A.H.; Vydellingum N.; Murray R.; Evans D.F.; Hartz A.J.; Kalkhoff R.K. and Adams P.W. (1982) : Relationship of body fat distribution to metabolic complications of obesity. *J. Clin. Endocrinol. Metab.* 54: 254-260.

Kliwer S.A.; Sundseth S.S.; and Jones S.A. (1997) : Fatty acids and eicosanodis regulate gene expression through direct interactions with peroxisome proliferator-activated receptors alpha and gamma. *Proc. Natl. Acad. Sci. USA.* 94: 4318-23.

Knowler W.C.; Pettitt D.J.; and Saad M.F. (1990) : Diabetes mellitus in the Pima Indians : incidence risk factors and pathogenesis. Diabetes Metab. Rev. 6: 1-27.

Knowler W.C.; Pettitt D.J.; Bennett P.H. and Williams R.C. (1983): Diabetes mellitus in the Pima Indians : genetic and evolutionary considerations. American Journal of Physical Anthropology. 62: 107-114.

Koivisto VA; Yki-Jarvinen H and DeFronzo RA; (1986): Phsiological training and insulin sensitivity. Diabetes. Metab. Rev. 1: 445-81.

Komers R; and Vr'ana A. (1998): Thiazolidinediones—tools for the research of metabolic syndrome X. Physiol Res, 47:215-25.

Kosaka K.; Hagura R. and Kuzuya T. (1977) : Insulin responses in equivocal and definite diabetes, with special reference to subjects who had mild glucose intolerance but later developed definite diabetes. Diabetes. 26. 10: 944-952.

Kostir JV and Sonka JA. (1952). Jaffe method (kinetic test without deproteinization). New Method Biochim Biophys. Acta. 8: 86.

Krentz AJ and Natrass M.. (1996): Insulin resistance: a multifaceted syndrome. Insights gained using low-dose insulin infusion technique. Diabet Med, 13:30-9.

Kruszynska Y.T. (1997) : Normal metabolism : the physiology of fuel homeostasis. In Pickup J and Williams G (eds) Textbook of Diabetes. 2nd edn. Ch. 11. Oxford : Blackwell Science.

Kaufman JS; and Cooper RS; (2001): Considerations for use of racial/ethnic classification in etiologic research (commentary). Am. J. Epidemiol. 145: 291-98.

Ku CY; Gower BA; Nagy TR and Goran MI; (1998): Relationships between dietary fat, and serum lipid profile in prepubertal children. Obes. Res. 6: 400-7.

Laakso M. (1993) : How good a marker is insulin level for insulin resistance? Am. J. Epidemiol. 137: 959-65.

Laakso M. (1999) : Hyperglycemia and cardiovascular disease in type 2 diabetes. Diabetes. 48: 973-42.

Laasko M; Edelman SV; Brechtel G; and Baron AD; (1990): Decreased effect of insulin to stimulate skeletal muscle blood flow in obese man. A novel mechanism for insulin resistance. J. Clin. Invest. 85: 1844-52.

Lan M.S.; Wasserfall C.; maclaren N.K. and Notkins A.L. (1996) : IA-2, a transmembrane protein of the protein tyrosine phosphatase

family, is a major autoantigen in insulindependent diabetes Mellitus. Proc. Natl. Acad. Sci. USA. 93: 6367-6370.

Lang D.M.; Matthews D.R.; Peto J. and Turner R.C. (1979) : Cyclic oscillations of basal plasma glucose and insulin concentrations in human beings. N. Engl. J. Med. 301: 1023-1027.

Larsson B.; Svarsudd K.; and Welin L. (1984) : Abdominal adipose tissue distribution, obesity and risk of cardiovascular disease and death ; 13 year follow up of participants in the study of men born in 1913. British Medical Journal. 288: 1401-1404.

Latha PP; Mercedes RC; and Stephen PF; (2002): Heterogenicity in the relationship between ethnicity, BMI, and fasting insulin. Diabetes Care. 25(8): 1531-57.

Laws A; Stefanic ML and Reaven GM; (1989): Insulin resistance and hyper-triglyceridemia in non-diabetic relatives of patients with NIDDM. J. Clin. Endocrinol. Metab. 69: 343-47.

Lee J.H.; Reed D.R. and Price R.A. (1997) : Familial risk ratios for extreme obesity; implications for mapping human obesity genes. International Journal of Obesity. 21: 935-940.

Leteixhe JM; Yanovski SZ; Avila NA; and Hubbard VS; (1994): Insulin secretion , clearance and action before and after gastroplasty in severely obese subjects. Int. J. Obestet. Relat. Metab. Disord. 18: 295-300.

- Levy J.; Atkinson A.; and Bell P. (1998) :** Beta-cell deterioration determines the onset and rate of progression of secondary dietary failure in type 2 diabetes mellitus : the 10 year follow-up of the Belfast Diet Study. *Diabet. Med.* 15: 290-6.
- Lewis G.F. and Steiner G. (1996):** Acute effects of insulin in the control of VLDL production in humans. Implications for the insulin-resistant state. *Diabetes. Care.* 19: 390-393.
- Lewis T.V.; Dart A.M. and Chen-Dusting J.P. (1999):** Endothelium-dependent relaxation by acetylcholine is impaired in hypertriglyceridemic humans with normal levels of plasma LDL cholesterol. *J. Am. Coll. Cardiol.* 33: 805-812.
- Lillioja S.; Mott D.M.; and Zawadzki J.K. (1987) :** In vivo insulin action is a familial characteristic in non-diabetic Pima Indians. *Diabetes.* 36: 1329-35.
- Lillioja S.; Mott D.M.; Spraul M.; Ferraro R.; Foley J.E.; and Ravussin E. (1995) :** Insulin resistance and insulin secretory dysfunction as precursors of non-insulin- dependent diabetes mellitus. Prospective studies of Pima Indians. *N. Engl. J. Med.* 329: 1988-1995.
- Lillioja S.; Mott DM.; and Spraul M. (1993) :** Insulin resistance and insulin secretory dysfunction as precursors of non-insulin

dependent diabetes mellitus : prospective studies of pima Indians. N. Engl. J. Med. 329: 1988-92.

Lillioja S.; Nyomba B.L.; and Saad M.F. (1991) : Exaggerated early insulin resistance in a diabetes-prone population : a metabolic comparison of pima Indians and Caucasians. Journal of Clinical Endocrinology and Metabolism. 73: 866-876.

Lillioja S; Young AA and Culter CL; (1987): Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. J. Clin. Invest. 80: 415-24.

Lilloja S; Mott DM; BV Howard; PH Bennett; H. Yki-Jarvinen; D. Freymond; BL Nyomba; F. Zurlo; B. Swinburn and C. Bogardus; (1988): Impaired glucose tolerance as a disorder of insulin action . longitudinal and cross-sectional studies in Pima-Indians. N. Engl. J. Med. 318: 1217-25.

Lindquist CH; Gower BA; and Goran MI; (2000): Role of dietary factors in ethnic differences in early risk of cardiovascular disease and type 2 diabetes. Am. J. Clin. Nutr. 71: 725-32.

Lithell H; Lindegrade F; Hellsing K; Lundquist G; Nygaard E and Saltin B; (1981): Body weight, skeletal muscle morphology and enzyme activities in relation to fasting serum insulin concentration and glucose tolerance in 48-years-old men. Diabetes. 30: 19-25.

Lloyd T; Chinchilli VM; and Rollings N; (1998): Fruit consumption, fitness, and cardiovascular health in female adolescents: The Penn State Young Women's Health Study. *Am. J. Clin. Nutr.* 67: 624-30.

Luan J.; Browne P.O.; and Harding A.H. (2001) : Evidence for gene-nutrient interaction at the PPAR γ locus. *Diabetes.* 50: 686-9.

Luan J.A.; Wong M.Y.; and Day N.E. (2001) : Sample size determination for studies of gene environment interaction. *Int. J. Epidemiol.* 30: 1035-40.

Ludvik B; Nolan JJ; Roberts A; Baloga J; Joyce M; Bell GM; and Olefsky JM; (1997): Evidence for decreased splanchnic glucose uptake after oral glucose administration in NIDDM. *J. Clin. Invest.* 100: 2345-61.

Macdonald I.A. (1995) : Advances in our understanding of the role of the sympathetic nervous system. *International Journal of Obesity.* 19 (Supplement 7) : 52-57.

Maheux P.; Azhar S.; Kern P.A.; Chen Y.D. and Reaven G.M. (1997) : Relationship between insulin-mediated glucose disposal and regulation of plasma and adipose tissue lipoprotein lipase. *Diabetologia.* 40: 850-858.

Mahgoub MA; (2000): Diabetes. The Egyptian Society of Pancreatic Islet Transplantation for Diabetic Patients. First Edition. P: 5-7.

Mancini F.P.; Vaccaro O.; and Sabatino L. (1999) : Pro 12Ala substitution in the peroxisome proliferator- activated receptor- γ is not associated with type 2 diabetes. Diabetes. 48: 1466-8.

Mandarino L; K. Wright L; Verity J; Nicholas J; Bell O; Kolterman and Beck-Nielsen; (1987): Effects of insulin infusion on human skeletal muscle pyruvate dehydrogenase, phosphofructokinase, and glycogen synthase. J. Clin. Invest . 80: 655-63.

Mangalolu L; Cheung RC; Van Iderstine SC; Taghibiglou C; Pontrelli L; and AdeliK; (2002): Treatment with atorvastatin ameliorates hepatic very-low-density lipoprotein overproduction in an animal model of insulin resistance, the fructose-fed Syrian golden hamster: evidence that reduced hypertriglyceridemia is accompanied by improved hepatic insulin sensitivity. Metabolism. 51(4): 409-18.

Mantzoros C.S.; Moschos S.; and Avramopoulos L. (1997) : Leptin concentrations in relation to body mass index and the tumor necrosis factor- α system in humans. Journal of Clinical Endocrinology and Metabolism. 82: 3408-3413.

- Marin P; Anderson B; Krotkiewski M; and Bjorntorp P; (1994):***
Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care*. 17: 382-86.
- Marshall J.A.; Hamman R.F. and Baxter J. (1991) :*** High-fat, low-carbohydrate diet and the etiology of non-insulin-dependent diabetes mellitus : the San Luis Valley diabetes study. *Am. J. Epidemiol.* 134: 590-603.
- Masson E.; MacFarlane I.A; Power E; and Wallymahmed M. (1992) :***
An audit of the management and outcome of hospital in-patients with diabetes : resource planning implications for the diabetes care team. *Diabetic Medicine*. 9: 753-755.
- Mather H.M. and Keen H. (1995) :*** The Southall diabetes survey: prevalence of known diabetes in Asians and Europeans. *British Medical Journal*. 291: 1081-1084.
- Matsuda J.; Hosoda K.; and Itoh H. (1998) :*** Increased adipose expression of the uncoupling protein-3 gene by thiazolidinediones in Wistar fatty rats and in cultured adipocytes. *Diabetes*. 47: 1809-1814.
- Matthews DR; Hosker JP; Rudenski AS; Naylor BA; Treacher DF and Turner RC; (1985):*** Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 28: 412-19.

- Mayer E.J.; Newman B.; and Austin M.A. (1996):*** Genetic and environmental influences on insulin levels and the insulin resistance syndrome : an analysis of women twins. Am. J. Epidemiol. 143: 323-32.
- McCance D.R.; Hanson R.L.; and Charles M.A. (1994) :*** Comparison of tests for glycated haemoglobin and fasting and two hour plasma glucose concentrations as diagnostic methods for diabetes. BMJ. 308: 1323-8.
- McCane T., Siruda R.L.; and Handerson U.S. (1991):*** Sugar in urine Vs. blood sugar for diagnosis of Diabetes. Diabet Med 11: 299-03.
- McCane T., Yasem E.T., Bakhdah K.M., and Fainenta I. (1993):*** Retinopathy in relation to FPG, 2-h PPG and HbA_{1c}. Diabetes 36:730-39.
- McCarthy MI and Kneel AL; (2001):*** The genetics of type 2 diabetes. Clin. Endocrinol. and Metab. 15: No.3 293-308.
- McGinnis J.M. and Foege W.H. (1993) :*** Actual causes of death in the United States. JAMA. 270: 2207-2212.
- McMahon M.; Marsh H. and Rizza R. (1989) :*** Effects of basal insulin supplementation of disposition of a mixed meal in obese patients with NIDDM. Diabetes. 38: 291-303.

- Meirhaeghe A.; Fajas L.; and Helbecque N. (2000) :*** Impact of the peroxisome proliferator activated receptor γ 2 pro12Ala polymorphism on adiposity, lipids and non-insulin- dependent diabetes mellitus. *Int. J.Obes. Relat. Metab. Disord.* 24: 195-9.
- Meirhaeghe A.; Helbecque N.; and Cottel D. (1999) :*** β 2-adrenoceptor gene polymorphism. Body weight, and physical activity. *Lancet.* 353: 896.
- Meirhaeghe A.; Luan J.; Selberg-Franks P. (2001):*** The effect of the Gly16Arg polymorphism of the beta2 adrenergic receptor gene on plasma free fatty acid levels in modulated by physical activity. *J. Clin. Endocrinol. Metab.* 86: 5881-7.
- Mikins KJ; Sonne B; Farrell PA; Tronier PA and Galbo H; (1988):*** Effects of physical exercise on sensitivity and responsiveness to insulin in humans. *Am. J. Physiol.* 254: E248-E259.
- Miller M.I; (1998):*** Ethno-ophthalmology in the Egyptian Delta; a historical system approach to ethno-medicine in the Middle-East. *Sci Med.* 26:651-7
- Mitrakou A.; Kelley D.; and Veneman T. (1990) :*** Contribution of abnormal muscle and liver glucose metabolism in postprandial hyperglycemia in noninsulin-dependent diabetes mellitus. *Diabetes.* 39: 1381-90.

Mitrakou A.; Vuorinen-Makkola H; and Raptis G; (1992):

Simultaneous assessment of insulin secretion and insulin sensitivity using a hyper-glycemic clamp. J. Clin. Endocrinol. Metab. 75: 379-82.

Mokdad A.H.; Serdula M.K.; Dietz W.H.; Bowman B.A.; Marks J.S.

and Koplan J.P. (1999): The spread of obesity epidemic in the United States. 1991-1998. JAMA. 282: 1519-1522.

Morris R.D.; Rimm D.L.; and Hartz A.J. (1988) : Obesity and heredity

in the etiology of non-insulin dependent diabetes mellitus in 32, 662 adult white women. Amer. J. Epid. 130: 112-21.

Munoz KA; Krebs-Smith SM; Ballard-Barbash R; and Cleavland LE;

(1997): Food intakes of US. Children and adolescents compared with recommendations. Pediatrics. 100: 323-9.

Murata C.; Suzuki Y.; and Muramatsu T. (2000) : Inactive aldehyde

dehydrogenase 2 worsens glycemic control in patients with type 2 diabetes mellitus who drink low to moderate amounts of alcohol. Alcohol. Clin. Exp. Res. 24: 5s-11s.

Nagy T; Gower B; Trowbridge C; Desenberg C; and Goran m; (1998):

Fat distribution and insulin response in prepubertal African-American and white children. Am. J. Clin. Nutr. 67: 821-27.

National Diabetes Data Group (1979) : Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. Diabetes. 28: 1039-1057.

National Health and Medical Research Council (1997) : Economic issues in the prevention and treatment of overweight and obesity. Acting on Australia's Weight : A strategic Plan for the prevention of Overweight and Obesity. Pp. 85-95. Canberra : Australian Government Publishing Service.

Neel J.V. (1962) : Diabetes mellitus : a thrifty genotype rendered detrimental by "progress". Am. J. Human Genet. 14: 353-62.

Nicholas J. W; Paul W.F and Anne H.H. (2002): Establishing the role of gene-environment interactions in the etiology of type 2 diabetes. Endocrinol. Metab. Clin. N. am. 31:553-566.

Nolte L.A.; Hansen P.A.; and Chen M.M. (1998) : Short-term exposure to tumor necrosis factor-alpha dose not affect insulin-stimulated glucose uptake in skeletal muscle. Diabetes. 47: 721-726.

Nyholm B; Mengel S and Nielsen S; (1996): Insulin resistance in relatives of NIDDM patients: the role of physical fitness and muscle metabolism. Diabetologia. 39: 813-22.

O'Rahilly S.P.; Nugent Z.; Rudenski A.S. (1986) : Beta-cell dysfunction, rather than insulin insensitivity, is the primary defect in familial type 2 diabetes. *Lancet.* 2: 360-364.

O'Rahilly SP; Turner R.C; and Matthews D.R. (1988): Impaired pulsatile secretion of insulin in relatives of patients with non-insulin-dependent patients. *N. Engl. J. Med.* 381: 1225-1230.

O'Sullivan J.B. & Muhen C.M; (1964): Criteria for the oral glucose tolerance test in pregnancy. *Diabetes,* 13: 227-85.

Oakuma M; Oakuma K. and York D.A; (1998): Effects of leptin on insulin secretion from isolated pancreatic islets. *Diabetes.* 47: 219-23.

Okhubo Y.; Kishikawa H.; Araki E.; Miyata T.; Isami S.; Motoyoshi S.; Kojima Y.; Furuyoshi N. and Shichiri M. (1995) : Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus : a randomized prospective 6- year study. *Diabetes Res. Clin. Pract.* 28: 103-117.

Olefsky J.M.; Kolterman O.G. and Scarlett J.A. (1982) : Insulin action and resistance in obesity and nonisnulin-dependent type II diabetes mellitus. *Am. J. Physiol.* 243: E15-E30.

Ontko J; (1972): Metabolism of free fatty acids in isolated liver cells. Factors affecting the partition between esterification and oxidation. J. Biol. Chem. 247: 1788-1800.

Ookuma M.; Ookuma K. and York D.A. (1998) : Effects of leptin on insulin secretion from isolated rat pancreatic islets. Diabetes. 47: 219-223.

Page R.C.; Hattersley A.T.; and Levy J.C. (1995) : Clinical characteristics of subjects with a missense mutation in glucokinase. Diabetic Med. 12: 209-17.

Pan X.R.; Cao H.B.; and Li G.W. (1997): Effect of dietary and exercise in preventing NIDDM in people with impaired glucose tolerance : the Da Qing IGT and diabetes study. Diabetes Care. 20: 537-44.

Panza J.A.; Quyyumi A.A.; Brush J.E.Jr. and Epstein S.E. (1990) : Abnormal endothelium-dependent relaxation in patients with essential hypertension. N. Engl. J. med. 323: 22-27.

Papoz L. and the EURODIAB Subarea C Study Group (1993) : Utilization of drug sales data for the epidemiology of chronic disease : the example of diabetes. Epidemiology. 4: 421-427.

Peavy DE; Edmondson JW, and Duckworth WC; (1984): Selective effects of inhibitors of hormone processing on insulin action in isolated hepatocytes. *Endocrinology*. 114: 753-60.

Pedersen O. (1989): The impact of obesity on the pathogenesis of non-insulin-dependent mellitus: A review of current hypotheses. *Diabetes Metab. Rev.* 5:495-509.

Pedersen O. (1999) : Genetics of insulin resistance. *Experimental and Clinical Endocrinology and Diabetes*. 107: 113-118.

Peiris AN; Mueller RA; Smith GA; Struve MF; and Kissebah AH; (1986): Splanchnic insulin metabolism in obesity. Influence of body fat distribution. *J. Clin. Invest.* 78: 1648-57.

Peiris A.N.; Struve M.F.; and Mueller R.A. (1988) : Glucose metabolism in obesity : influence of body fat distribution. *Journal of Clinical Endocrinology and Metabolism*. 67: 760-767.

Perusse L. and Chagnon Y.C. (1997) : Summary of human linkage and association studies. *Behavior Genetics*. 27: 359-372.

Petersen KF; Laurent D; Rothman DL; Cline GW; and Shulman GL; (1998): Mechanism by which glucose and insulin inhibit net

hepatic glycogenolysis in humans. *J. Clin. Invest.* 101: 1203-09.

Petersen KF; Oral EA; Dufour S; Befroy D; Ariyan C; Yu C; Cline GW; DelPaoli AM; Tylor SI; Gorden P; and Shulman GI; (2002): Leptin reverses insulin resistance and hepatic steatosis in patients with severe lipodystrophy. *J. Clin. Invest.* 109(10): 1345-50.

Petty R.H.K.; Harding A.E. and Morgan-Hughes I.A. (1986) : The clinical features of mitochondrial myopathy. *Brain.* 109: 915-938.

Pfeifer M,A; Halter J.B; and Porte D. (1981): Insulin secretion in diabetes mellitus. *Am. J. Med.* 70:579-588.

Phillips A.N. Davey and Smith G. (1993): The design of prospective epidemiological studies : more subjects of better measurements? *J. Clin. Epidemiol.* 46: 1203-11.

Phillips D.I.; Hirst S.; and Clark P.M. (1994) : Fetal growth and insulin secretion in adult life. *Diabetologia.* 37: 592-596.

Phillips D.I.W. (1993) : Twin studies in medical research : can they tell us whether diseases are genetically determined? *Lancet.* 341: 1008-1010.

Phillips D.L.; Barker D.J.; and Hales C.N. (1994) : Thinness at birth and insulin resistance in adult life. Diabetologia. 37: 150-154.

Pierce M.; Keen H. and Bradley C. (1995) : Risk of diabetes in offspring of parents with non-insulin dependent diabetes. Diabetic Med. 12: 6-13.

Pinkney J.H.; Stehouver C.D.A.; Coppack S.W. and Yudkin J.S. (1997): Endothelial dysfunctions : cause of the insulin resistance syndrome. Diabetes. 46: 59-513.

Polonsky K. Frank B. and Pugh W; (1986): the limitation to and valid use of C-Peptide as a marker of the secretion of insulin. Diabetes. 35:379-86.

Polonsky K.S.; Sturis J. and Bell G.I. (1996) : Non-insulin-dependent diabetes mellitus : a genetically programmed failure of the beta cell to compensate for insulin resistance. N. Engl. J. Med. 334: 777-784.

Polonsky KS; Given BD; Hirsch L; Shaprio FT; Tillil H; Beebe C; Galloway JA; Frank BH; Karrison T and Van Cauter E; (1988): Quantitative study of insulin secretion and clearance in normal and obese subjects. J. Clin. Invest. 81: 435-41.

Pospisilik JA; Stafford SG; Demuth HU; McIntosh CH; and Pederson RA; (2002): Long-term treatment with dipeptidyl peptidase IV inhibitor improves hepatic and peripheral insulin sensitivity in the Vancouver diabetic fatty rat: an euglycemic-hyperinsulinemic clamp study. *Diabetes*. 51(9): 2677-83.

Poulsen P.; Kyvik K.O.; Vaag A. and Beck-Nielsen H. (1999) : Heritability of type II (non-insulin dependent) diabetes mellitus and abnormal glucose tolerance- apopulation based twin study. *Diaetologia*. 42: 139-145.

Poulton J.; Brown M.S.M.; and Copper A. (1998) : A common mitochondrial DNA variant is associated with insulin resistance in adult life. *Diabetologia*. 41: 54-58.

Quinones-Galvan A. and Ferrannini E. (1997) : Renal effects of insulin in man. *J. Nephrol*. 10: 188-191.

Ramachandran A.; Snehalatha C.; and Dharmaraj A. (1992) : Prevalence of glucose intolerance in Asian Indians. *Diabetes Care*. 15: 1348-1355.

Randle P.J.; Garland P.B.; Hales C.N. and Newholme C.A. (1963) : The glucose fatty acid cycle. Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet*. 1: 785-789.

Ravelli A.C.J.;van der Meulen J.H.P.; and Michels P.P.J. (1998) :

Glucose tolerance in adults after prenatal exposure to famine.
Lancet. 351: 173-177.

Reaven GM and Greenfield MS; (1981): Diabetic hypertriglyceridemia:

evidence for three clinical syndromes. Diabetes. 30: 66-75.

Reaven GM and Laws A; (1994): Insulin resistance, compensatory

hyperinsulinemia and coronary heart disease. Diabetologia 37:
948-52.

Reaven GM; (1988): Role of insulin resistance in human disease.

Diabetes. 37: 1595-1607.

Rebuffe-Scrive M; Andersson B; Olbe I; and Bjorntorp P; (1989):

Metabolism of adipose tissue in intra-abdominal depots in
severely obese men and women. Metabolism. 39: 1021-25.

Richard W; (1973). Enzymatic colorimetric method with lipid clearing

factor. Clin. Chem; 19: 1350.

Ricquier D. and Bouillaud F. (1998) : The mitochondrial uncoupling

proteins. Medicine Science. 14: 889-897.

Ringel J.; Engeli S.; and Distler A. (1999) : Pro 12 Ala missense

mutation of the peroxisome proliferator activated receptor γ
and diabetes mellitus. Biochem. Biophys Res. Commun. 254:
450-3.

Ristow M.; Giannakidou W.; and Hebinck J. (1998) : An association between NIDDM and a GAA trinucleotide repeat in the X25/Frataxin (FA) gene. *Diabetes*. 47: 851-854.

Rheeder PJ; Gerald PB; Hales CN; and Newsholm EA; (2001): Role of dietary factors in ethnic differences in early risk of cardiovascular disease and type 2 diabetes. *Am. J. Clin. Nutr.* 71: 725-32.

Robbins D.C; Tager H.S; and Rubenstein A.H. (1984): Biologic and clinical importance of proinsulin. *N. Engl. J. Med.* 310:1165-75.

Roder ME; Knip M; Hartling SG; Karjalainen J; Akerblom HK; and Binder C; (1994): The Childhood Diabetes in Finland Study Group: Disproportionately elevated proinsulin levels precede the onset of IDDM in sibilings with low first phase insulin responses. *J. Clin. Endocrinol. Metab.* 79: 1570-75.

Rodriguez BL; Lau NI; and Burchfiel CM; (1999): Glucose intolerance and 23-year risk of coronary heart disease and total mortality: the Honolulu Heart Program. *Diabetes Care*. 22: 1262-5

Rojdmark S; Bloom G; Chou MCY; Jaspan J; and Field JB; (1978): Hepatic insulin and glucagon extraction after their augmented secretion in dogs. *Am. J. Physiol.* 235: E88-E96.

Rossell R; Comis R; Casamitjana R; Segura R; Vilardell E; and Rivera F; (1983): Reduced hepatic insulin clearance in obesity: relationship with plasma insulin levels. . J. Clin. Endocrinol. & Metab. 56: 608-11.

Saad M.F.; Knowler W.C.; and Pettitt D.J. (1991) : A two-step model for development of non-insulin-dependent diabetes. Am. J. Med. 90: 229-35.

Saghizadeh M.; Ong J.M.; and Garvey W.T. (1996) : The expression of TNF alpha by human muscle. Relationship to insulin resistance. Journal of Clinical Investigation. 97: 1111-1116.

Salas J.; Jansen S.; and Lopez-Miranda J. (1998) : The SstI polymorphism of the apolipoprotein C-III gene determines the insulin response to an oral-glucose-tolerance test after consumption of a diet rich in saturated fats. Am. J. Clin. Nutr. 68: 396-401.

Sargeant L.A.; Khaw K.T.; and Bingham S. (2001) : Fruit and vegetable intake and population glycosylated haemoglobin levels : the EPIC-Norfolk Study. Eur. J. Clin. Nutr. 55: 342-8.

Sargeant L.A.; Wareham N.J. and Khaw K.T. (2000) : Family history of diabetes identifies a group at increased risk for the metabolic

consequences of obesity and physical inactivity in EPIC-Norfolk : a population-based study. *Int. J. Obes.* 24: 1333-9.

Savontaus E.; Rouru J.; and Boss O. (1998) : Differential regulation of uncoupling proteins by chronic treatments with beta3 adrenergic agonist BRL 35135 and metformin in obese fa/fa Zucker rats. *Biochemical and Biophysical Research Communications.* 246: 899-904.

Scarlett J.A.; Gray R.S.; Griffin J.; Olefsky J.M. and Kolterman O.G. (1982) : Insulin treatment reverses the insulin resistance of type II diabetes mellitus. *Diabetes Care.* 5: 353-363.

Scherbaum W.A. (1998) : The role of amylin in the physiology of glycemic control. *Experimental and Clinical Endocrinology and Diabetes.* 106: 97-102.

Schichiri M.; Kishikawa H.; Ohkubo Y. and Wake N. (2000) : Long-term results of the Kumamoto study on optimal diabetes control in type 2 diabetes control in type 2 diabetic patients. *Diabetes Care* 23 (Suppl. 2) : B21- B29.

Schrezenmier J. (1996): Hyper-insulinaemia, hyper-proinsulinaemia and insulin resistance in the metabolic syndrome. *Experienta* 426-32.

Schulman DDL; Rothman T; Jue P; Stein RA; DeFronzo and Schulman RG; (1990): Quantitation of muscle glycogen

synthesis in normal subjects and subjects with NIDDM by C_{13} - nuclear magnetic resonance spectroscopy. N. engl. Med. 322: 223-228.

Schumacher M.C.; Hasstedt S.J.; and Hani S.J. (1992) : Major gene effect for insulin levels in familial NIDDM pedigrees. Diabetes. 41: 416-23.

Schwartz M.W. and Seeley R.J. (1997) : Neuroendocrine responses to starvation and weight loss. New England. Journal of Medicine. 336: 1803-1811.

Segal K.R.; Landt M. and Klein S. (1996) : Relationship between insulin sensitivity and plasma leptin concentration in lean and obese men. Diabetes. 45: 988-991.

Seidell JC; Oosterlee A; Deurenberg P; Hautvast JGAJ; and Ruijs JIJ; (1988): Abdominal fat depots measured with computed tomography: effects of degree of obesity, sex, and age Eur. J. Clin. Nutr. 42: 805-15.

Shapiro ET; Polonsky KS; Copinschi G; Bosson D; Tillil H; Blackman J; Lewis G; and Vaccaro E; (1991): Nocturnal elevation of glucose levels during fasting in NIDDM. J. Clin. Endocrinol. Metab. 72: 444-54.

Shapiro ET; Tillil H; Rubenstein AH and Polonsky KS; (1988):
Peripheral insulin parallels changes in insulin secretion more closely than C-Peptide after bolus intravenous glucose administration. *J. Clin. Endocrinol. & Metab.* 67: 1094-99.

Smith P.G. and Day N.E. (1984) : The design of case-control studies : the influence of confounding and interaction effects. *Int. J. Epidemiol.* 13: 356-65.

Smith U; (1980): Adrenergic control of human adipose tissue lipolysis. *Eur. J. Clin. Invest.* 10: 343-44.

Snieder I.I.; Boomsma D.I.; and vanDoornen L.J.P. (1999) : Bivariate genetic analysis of fasting insulin and glucose levels. *Genet Epidemiol.* 16: 426-46.

Snitker S.; Pratley R.E.; and Nicolson M. (1997) : Relationship between muscle sympathetic nerve activity and plasma leptin concentration. *Obesity Research.* 5: 338-340.

Sonnenberg GE; Hoffman RG; Mueller RA; and Kissebah AH; (1994):
Splanchnic insulin dynamics and secretion pulsilities in abdominal obesity. *Diabetes.* 43: 468-77.

Sorensen T.I. (1995): The genetics of obesity. *Metabolism : Clinical and Experimental.* 44: 4-6.

Sowers J.R. (1990) : At the cutting edge. Insulin resistance and hypertension. *Mol. Cell. Endocrinol.* 74: C87-C89.

Spurr G.B.; Prentice A.M.; and Murgatrody P.R. (1988) : Energy expenditure from minute-by minute heart rate recording : comparison with indirect calorimetry. *Am. J. Clin. Nutr.* 48: 552-9.

Stanhope J.M. and Prior I.A.M. (1980) : The Tokelau Island migrant study : prevalence and incidence of diabetes mellitus. *New Z Med. J.* 92: 417-21.

Starr J.I.; Mako M.E.; Juhn D. and Rubenstein A.H. (1978) : Measurement of serum proinsulin-like material : cross-reactivity of porcine and human proinsulin in the insulin radioimmunoassay. *J. Lab. Clin. Med.* 91. 4: 691-692.

Stephens J.M.; Lee J. and Pilch P.F. (1997) : Tumor necrosis factor- α -induced insulin resistance in 3T3-L1 adipocytes in accompanied by a loss of insulin receptor substrate-I and GLUT4 expression without a loss of insulin receptor-mediated signal transduction. *Journal of Biological Chemistry.* 272: 971-976.

Steppan C.M.; Bailey S.T.; Bhat S.I.D.; Brown E.J.; Banerjee R.R.; and Wright C.M. (2001) : The hormone resistin links obesity to diabetes. *Nature*. 409: 307-312.

Steven DM; (2000): Longitudinal compensation for fat-induced insulin resistance includes reduced hepatic insulin clearance and enhanced beta-cell response. *Diabetes*. 49: 2116-25.

Stock M; (1998): Energy balance and animal models of obesity. In Kopelman PG & Stock M (eds). *Clinical obesity* pp 50-72. Oxford: Blackwell Science.

Stoffers DA.; Ferrer J.; Clarke W.L. and Habener J.F. (1997): Early-onset type-II diabetes mellitus (MODY4) linked to IPFI. *Nature Genetics*. 17: 138-139.

Strombland G and Bjorntorp P; (1986): Reduced hepatic insulin clearance in rats with dietary induced obesity. *Metabolism*. 35: 323-27.

Stumvoll M.; Meyer C.; and Mitrakou A. (1997) : Renal glucose production and utilization : new aspects in humans. *Diabetologia*. 40: 749-57.

Sumner AE; Kushner H; Sherif KD; Tulenko TN; Falkner B; and Marsh JB; (1999): Sex differences in African-Americans

regarding sensitivity to insulin's gluco-regulatory and antilipolytic actions. *Diabetes Care*. 22: 71-77.

Svedberg J; Bjorntorp P; Lonnroth P; and Smith U; (1991): Etomoxir prevents the inhibitory effect of free fatty acids on insulin binding and action in isolated fat hepatocytes. *Diabetes*.40:783-86.

Svedberg J; Bjorntorp P; Smith U; and Lonnroth P; (1990): Free fatty acid inhibition of insulin binding, degradation and action in isolated fat hepatocytes. *Diabetes*. 39:570-74.

Svedberg J; Gunnar S; Wirth A; Smith U and Bjorntrop P; (1991): Fatty acids in the portal vein of the rat regulate hepatic insulin clearance. *Clin. Invest*. 88: 2054-58.

Swinburn B.A. (1996) : The thrifty genotype hypothesis : how does it look after 30 years? *Diabete. Med*. 13: 695-9.

Tallarigo L., Giampietro O., and Peeno G.(1986): Relations of glucose tolerance to complications of pregnancy in non-diabetic women. *N. Engl. J. Med*. 315:989-92.

Tattersall R.B. and Pyke D.A. (1972) : Diabetes in twins. *Lanct*. 11: 1120-5.

Taylor S. (1992) : Insulin resistance : lessons from patients with mutations in the insulin-receptor gene. Diabetes. 41: 1473-90.

Temple R.C.; Carrington C.A.; Luzio S.D.; Owens D.R.; Schneider A.E.; Sobey W.J. and Hales C.N. (1989) : Insulin deficiency in non-insulin dependent diabetes. The lancet. Feb. 11: 293-295.

Temple R.C.; Clark P.M. and Hales C.N. (1992) : Measurement of insulin secretion in type 2 diabetes : problems and pitfalls. Diabetic medicine. 9: 503-512.

Temple R.C.; Clark P.M.; Nagi D.K.; Schneider A.E.; Yudkin J.S. and Hales C.N. (1990) : Physical exercise and reduced occurrence of NIDDM. N. Engl.Med.325: 147-52.

Terauchi Y.; Iwamoto K. and Tamemoto H. (1997) : Development of non-insulin-dependent diabetes mellitus in the double knockout mice with disruption of insulin receptor substrate-1 and beta cell glucokinase genes. Genetic reconstruction of diabetes as a polygenic disease. Journal of Clinical investigation. 99: 861-866.

Thomas F.; Balkau B. and Vauzelle-Kervroedan F. (1994) : Maternal effect and familial aggregation in NIDDM. The CODIAB study. Diabetes. 43: 63-67.

Thorburn AW; B. Gumbiner; F. Bulacan; G. Brechtel and RR. Henry;
(1991): Multiple defects in muscle glycogen synthase activity contribute to reduced glycogen synthesis in NIDDM. J. Clin Invest. 87: 489-95.

Trider P; (1969). Enzymatic colorimetric method with lipid clearing factor. Ann. Clin. Bioch; 624-27.

Tsaglis H; Ionnidis I; Giakoumaki A; Amantou A and Komitopoulos S,(1999):. The effect of metformin on serum B12 levels: is there any clinical importance? Diabetologia 42 (suppl. 1): 224.

Tuomilehto J. and Golay A; (2000): Sugar and spikes: Revealing the new hidden threat in type 2 diabetes. Endocrinology 53:1-4.

Tuomilehto J.; Lindstrom J.; and Eriksson J.G. (2001) : Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. N. Engl. J. Med. 344: 1343-50.

Tylor S; Accilli D; and Imal Y; (1994): Insulin resistance or insulin deficiency: which is the primary cause of NIDDM? Diabetes. 48: 735-40.

Ukkola O.; Sun G. and Gouchard C. (2001) : Insulin-like growth factor 2 (IGF2) and IGF-binding protein 1 (IGFBP1) gene variants are associated with overfeeding-induced metabolic changes. Diabetologia. 44: 2231-6.

United Kingdom Prospective Diabetes Study (UKPDS) Group. (1995):

UK prospective diabetes study 16. Overview of 6 years therapy of type II diabetes : a progressive disease. Diabetes. 44: 1249-58.

United Kingdom Prospective Diabetes Study Group (1998) : Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). Lancet. 352: 837-853.

Vaag A; Henriksen and Beck-Nielsen H; (1992): Decreased insulin activation of glycogen synthase in skeletal muscles in young non-obese Caucasian first degree relatives of patients with NIDDM. J. clin. Invest. 89:782-88.

Vaag A; Henriksen J and Madsbad S; (1995): Insulin secretion, insulin action and hepatic glucose production in identical twins discordant for NIDDM. J. Clin. Invest. 95: 690-8.

Vaag A; Scott P; Damsbo P; Gall MA; Richer EA; and Beck-Nielsen H; (1991): Effects of the antilipolytic nicotinic acid analogue Acipimox on whole body and skeletal muscle glucose metabolism in patients with NIDDM. J. Clin. Invest. 88: 1282-90.

Valdez R; (1995): Differential impact of obesity in related populations.
Obes. Res. 3 (suppl. 2): 223s- 232s.

Valente AM; Strong W; Sinaiko AR; (2001): Obesity and insulin resistance in young people. Am. Heart. J. 142:440-44.

Van Haeften T.; Pimenta W. and Mitrakou A. (1999) : Different factors regulate glucose tolerance and insulinemia in NGT and IGT subjects. Diabetes. 48 (Suppl. 1: A 436-37. Abstract.

Vanderpump M.P.J.; Tunbridge W.M.G. and French J.M. (1996) : The incidence of diabetes mellitus in an English community : a 20-year follow-up of the Whickham survey. Diabetic Medicine. 13: 741-746.

Varadi A.; Lebel L. and Hashim Y. (1999) : Sequence variants of the sarco (endo) plasmic reticulum Ca^{2+} - transport ATPase 3 gene (SERCA3) in Caucasian type II diabetic patients (UK Prospective Diabetes Study 48). Diabetologia. 42: 1240-1243.

Vasquez B; Reaven GM; and Andrews WG; (1985): Elevated in vivo insulin clearance in Pima Indians with NIDDM. Diabetes. 34(7) 671-6.

Vauhkonen I.; Niskanen L.; Vanninen E.; Kainulainen S.; Uusitupa M. and Laakso M. (1998) : Defects in insulin secretion and

insulin action in non-insulin dependent diabetes mellitus are inherited. Metabolic studies on offspring of diabetic probands. *J. Clin. Invest.* 101: 86-96.

Velho G. and Froguel P. (1997) : Genetic determinants of non-insulin-dependent diabetes mellitus : strategies and recent results. *Diabetes and Metabolism.* 23: 7-17.

Vilbregson S.; Sigurdsson G. and Sigvaldason H. (1997) : Prevalence and incidence of NIDDM in Iceland : evidence for stable incidence among males and females 1967-1991 : The Reykjavik Study. *Diabetic Medicine.* 14: 491-498.

Villar-Palasi C. and J. Larner; (1960): Insulin-mediated effect on the activity of UDPG-glycogen transglucosylase of muscle. *Biochim. Biophys. Acta.* 81: 508-20.

Walder K.; Norman R.A. and Hanson R.L. (1998) : Association between uncoupling protein polymorphisms (UCP2-UCP3) and energy metabolism/obesity in Pima Indians. *Human. Molecular Genetics.* 7: 1431-1435.

Wang Y.; Kuropatwinski K.K. and White D.W. (1997) : Leptin receptor action in hepatic cells. *Journal of Biological Chemistry.* 272: 16216-16223.

Wareham N.J.; Byrne C.D. and Williams R. (1999) : Fasting proinsulin concentrations predict the development of type 2 diabetes. Diabetes Care. 22: 262-70.

Wareham N.J.; Wong M.Y. and Day N.E. (2000) : Glucose intolerance and physical inactivity : the relative importance of low habitual energy expenditure and cardiorespiratory fitness. Am. J. Epidemiol. 152: 132-9.

Warram J.H; Martin B.C; and Krolewski AS. (1990): Slow glucose removal rate and hyper-insulinaemia precede the development of type II diabetes in the offspring of diabetic patients. Ann. Intern. Med. 113:909-915.

Warram JH; Martin BC; Krolewski AS; Soeldner JS and Kahn CR; (1990): Slow glucose removal rate and hyperinsulinemia precede the development of type-2 diabetes in the offspring of diabetic patients. Ann. Intern. Med. 113: 909-15.

Weir G.C.; Nathan D.M. and Singer D.E. (1994) : Standards of care for diabetes (technical review). Diabetes Care. 17: 1514-1522.

Wendorf M. (1989) : Diabetes, the ice free corridor, and the Paleoindian settlement of North America. Am. J. Phys. Anthropol.; 79: 503-20.

White M.F. (1998) : The IRS-signalling system : a network of docking proteins that mediate insulin action. *Molecular and Cellular Biochemistry*. 182: 3-11.

Williams D.R.R.; Wareham N.J. and Brown D.C. (1995) : Glucose intolerance in the community : the Isle of Ely diabetes project. *Diabet. Med.* 12: 30-5.

Wiltshire S.; Hattersley A.T. and Hitman G.A. (2001) : A genome-wide scan for loci predisposing to type 2 diabetes in a UK population (the diabetes [UK] Warren 2 repository) : analysis of 573 pedigrees provides independent replication of a susceptibility locus on chromosome 1q. *Am. J. Hum. Genet.* 69: 553-69.

Wing R.R.; Matthews K.A. and Kuller L.H. (1992) : Environmental and familial contributions to insulin levels and change in insulin levels in middle-age women. *JAMA*. 268: 1890-5.

Withers D.J.; Gutierrez J.S. and Towery H. (1998) : Disruption of IRS-2 causes type 2 diabetes in mice. *Nature*. 391: 900-904.

World Health Organization (1985) : Diabetes Mellitus : Report of a WHO Study Group. WHO Technical Report Series 727. Geneva : WHO.

World Health Organization (1997) : Prevention of diabetes mellitus. 844. Geneva : WHO.

World Health Organization Study Group (1997) : Defining the problem of overweight and obesity. In Report of a WHO consultation on Obesity. Obesity : Preventing and Managing the Global Epidemic. Pp. 7-16. Geneva : WHO.

Wu C; Okar DA; Newgard CB; and Lange AJ; (2002): Increasing fructose 2,6-bisphosphate overcomes hepatic insulin resistance of type 2 diabetes. Am. J. Physiol. Endocrinol. Metab. 282(1): E38-45.

Yki-Jarvinen H. (1998) : Toxicity of hyperglycaemia in type 2 diabetes. Diabetes Metab. Rev. 14: 45-50.

Yu-Poth S; Zhao G; Etherton T; Naglak M; Jonnalagadda S; and Kris-Etherton PM; (1999): Effects of the National Cholesterol Education Programs's Step I and Step II dietary intervention programs on cardiovascular disease risk factors: a meta-analysis. Am. J. Clin. Nutr. 69: 632-46.

Zachary T. Bloomgarden (2002) : New insights in obesity. Diabetes Care. Vol. 25, No. 4, April. 789-795.

- Zhang Y.; Proenca R. and Maffei M. (1994) :** Positional cloning of the mouse obese gene and its human homologue. *Nature*. 372: 425-432.
- Zimmer P. and McCarthy D. (1995) :** The NIDDM epidemic : global estimates and projection – a look into the crystal ball. *Idf. Bulletin*. 40: 8-16.
- Zimmet P., (1992) :** Kelly West Lecture 1991 : Challenges in diabetes epidemiology : from west to the rest. *Diabetes Care*. 15: 232-252.
- Zimmet P.; Dowse G. and Finch C. (1990) :** The epidemiology and natural history of NIDDM- lessons from the South Pacific. *Diabetes and Metabolism. Reviews*. 6: 91-124.
- Zimmet P.; King H. and Taylor R. (1984) :** The high prevalence of diabetes mellitus, impaired glucose tolerance and diabetic retinopathy in Nauru-The 1982 survey. *Diabetes Res*. 1: 13-8.
- Zimmet P; Tuomi T.; Mackay R.; Rowley M.J.; Knowles W.; Cohen M. and Lang D.A. (1994) :** Latent autoimmune diabetes mellitus in adults (LADA) : the role of antibodies to glutamic acid decarboxylase in diagnosis and prediction of insulin dependency. *Diabt. Med*. 11: 299-303.