

CONTENTS

Introduction	1
Aim of the work	4
Review of literatures	5
❖ <i>Anatomy of the venous system of the lower limb</i>	5
❖ <i>The deep veins of the lower limbs</i>	11
❖ <i>Valves of veins of lower limb</i>	19
❖ <i>Venous anatomy of the upper limb</i>	22
❖ <i>Pathogenesis of deep venous thrombosis</i>	28
❖ <i>Risk factors</i>	33
❖ <i>D-dimer test in diagnosis of deep venous thrombosis</i>	38
❖ <i>Contrast venography</i>	45
❖ <i>Duplex imaging in diagnosis of deep vein thrombosis</i>	49
❖ <i>Duplex findings in deep venous thrombosis</i>	60
❖ <i>Advantages and limitations of duplex imaging</i>	63
❖ <i>Duplex study versus other modalities</i>	72
❖ <i>Magnetic resonance venography in diagnosis of deep vein thrombosis.</i>	76
Patients and Methods	84
Results	108
Cases	131
Discussion	145
Summary and conclusion	163
Recommendations	165
References	166
Arabic summary	٢-١

List of Tables

Table (1): Character of our studied patients ,distribution of age, sex their clinical presentation and clinical probability	115
Table (2): Relationship between findings of Contrast Venography and clinical pretest probability of DVT	117
Table (3) Distribution of cases according to their clinical probability and sites of thrombosis	118
Table (4): Relationship between the result of D-dimer test and the clinical probability of our studied patients which revealed a considerable number of patients with a low probability group had negative D- dimer test	119
Table (5): showing the sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of D-dimer test when compared to Contrast Venography . Among the twenty three positive D dimer cases, twenty one patients had a D V T confirmed by C V and among the remaining seven cases with negative D dimer , only one case had D V T confirmed by CV	121
Table (6): sensitivity, specificity, positive predictive value(P P V and negative predictive value(N P V) of D –dimer when a positive D-dimer or medium / high clinical pretest probability combined together and compared to C V	122
Table (7): sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of D –dimer , when a positive D-dimer and medium /high clinical pretest probability combined together and compared to C V	123
Table (8): sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of D-dimer test when positive D-dimer cases or Duplex findings combined together and compared to C V as gold standard, with increased sensitivity and specificity up to (100%)	124

Table (9): sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of D-dimer when positive D dimer and Duplex combined together and compared to C V	125
Table (10) Results of Contract Venography and distribution of cases according to their sites of thrombosis	126
Table (11): Distribution of thrombi as detected by Contarst Venography and Duplex study	127
Table (12): sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of Duplex study when compared to C V	128
Table (13): sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of Magnetic Resonance venography for diagnosis DVT when compared to C V. with complete agreement in both studies	129
Table (14): Distribution of cases and their site of thrombosis in Contrast Venography and MRV which showed complete agreement in both studies as regard to the sites of thrombosis	130

List of Figures

<i>Figure 1: Long saphenous vein and its tributaries</i>	8
<i>Figure 2: Short saphenous vein and its tributaries</i>	10
<i>Figure 3: The principal deep veins of the lower limbs</i>	13
<i>Figure 4: Anatomy of the deep venous system of the lower limb</i>	14
<i>Figure 5: sites of important perforator veins</i>	18
<i>Figure (6) Venous anatomy of the upper limb</i>	23
Figure (7) Thrombosis formation on venous valve cusp (A) Normally, blood flow (straight arrows) proceeds through the valve with the formation of eddy current (curved arrows). (B) A clot nidus may form on the cusp in the setting of stasis. (C) The clot may then extend into the venous lumen	29
<i>Figure(8) Plasmic degradation of crosslinked fibrin</i>	39
<i>Fig (9), Difference in Pitch of an ambulance siren when moving</i>	51
Fig (10) Alignment of the individual magnetic moments with the external field M_0 is the net magnetization	78