

References

- Allen A, Pusey C, Gaskin G, 2002:** Outcome for renal replacement therapy in antineutrophil cytoplasmic antibody-associated vasculitis. *J Am Soc Nephrol.*; 9:1258-1263.
- Amar J, 2000:** Nocturnal blood pressure and 24-hour pulse pressure are potent indicators of mortality in hemodialysis patients. *Kidney Int.*; 57:2485-2491.
- Amato RL, 2005:** Water treatment for hemodialysis updated to include the latest AAMI standards for dialysate (RD52:2004). *Nephrol Nurs J.*; 332: 151-167
- Amerling R, 2005:** Practice guidelines: are they dead on arrival? Yes. *Nephrol News Issues* 2003; 17: 38–39. 3 Amerling R: Practice guidelines: do we really need them? *Perit Dial Int.*; 25: 140–141.
- Amin K, 2002:** Intraperitoneal ondansetron hydrochloride for intractable nausea and vomiting due to diabetic gastroparesis in a patient on peritoneal dialysis. *Perit Dial Int.*;22:539-540.
- Anderson-Haag T, Patel B, 2003:** Safety of colchicine in dialysis patients. *Semin Dial.*; 16:412-413.
- Andress DL, 2007:** Early deposition of aluminum in bone in diabetic patients on hemodialysis. *N Engl J Med.*;316:292-296.
- Andrulli S, 2002:** The role of blood volume reduction in the genesis of intradialytic hypotension. *Am J Kidney Dis.*; 40:1244-1254.
- ANZ (Data Registry Report), 2006:** Australia and New Zealand Dialysis and Transplant Registry..

- Aoki J, 2006:** Coronary revascularization improves long-term prognosis in diabetic and non-diabetic end-stage renal disease. *Circ J*;66:595-599.
- Arora SK, McFarlane SI, 2005:** The case for low carbohydrate diets in diabetes management. *Nutr Metab.*;2:16.
- Ascher E, Gade P, Hingorani A, 2002:** Changes alarm system: Impact of dialysis outcome and quality initiative recommendations. *J Vasc Surg* 31:84-92
- Association for the Advancement of Medical Instrumentation, 2004:** Dialysate for hemodialysis. Arlington, VA: Association for the Advancement of Medical Instrumentation, ANSI/AAMI RD52
- Augustine JJ, 2004:** A randomized controlled trial dialysis in patients with ARF. *Am J Kidney Dis.*;44:1000-1007.
- Avram MM, Goldwasser P, Erroa M, Fein PA, 2001:** Predictors of survival in continuous ambulatory peritoneal dialysis patients: The importance of prealbumin and other nutritional and metabolic markers. *Am J Kidney Dis* 23:91-98, 1994
- Bailey BJR, Briars G, 2002:** Estimating the surface area of the human body. *Stat Med* 15:1325-1332
- Beardsworth SF, 2001:** a protocol for administration during HD and review of published protocols. *Perit Dial Int.*;8:145.
- Bergstrom J, Emerson P, Keshaviah P, 1999:** The complications of peritoneal dialysis, in *Practical Peritoneal Dialysis*. Bristol, England, John Wright & Sons,: 63-69
- Beddhu S, Martin SR, Clouse DF, 2005:** Ramkumar N, Pappas LM. Normalization of protein intake by body weight and the associations of protein intake with nutritional status and survival. *J Ren Nutr.*; 15(4):387-397.

- Bergstrom J, Heimberger O, Lindholm B, 2000:** Calculation of protein equivalent of total nitrogen appearance from urea appearance: Which formulas should be used? *Perit Dial Int* 18:467-473
- Bhatla B, Moore H, Emerson P, Keshaviah P, Prowant B, Nolph KD, Singh A, 1999:** Lean body mass estimation by creatinine kinetics, bioimpedance, and dual energy x-ray absorptiometry in patients on continuous ambulatory peritoneal dialysis. *ASAIO J* 41:M442-M446
- Blake PG, Flowerdew G, Blake RM, Oreopoulos DG, 2001:** Serum albumin in patients on continuous ambulatory peritoneal dialysis: Predictors and correlations with outcomes. *J Am Soc Nephrol* 3:1501-1507
- Blake PG, Sombolos K, Abraham G, Weissgarten J, Pemberton R, Chu GL, Oreopoulos DG, 1999:** Lack of correlation between urea kinetic indices and clinical outcomes in CAPD patients. *Kidney Int* 39:700-706
- Blake PG, Spanner E, McMurray S, Lindsay RM, Ferguson E, 2000:** Comparison of measured and predicted creatinine excretion is an unreliable index of compliance in PD patients. *Perit Dial Int* 16:147-153.
- Bleyer AJ, Hylander B, Sudo S, Nomoto Y, de la Torre E, Chen RA, Burkart JM, 2003:** An international study of patient compliance with hemodialysis. *JAMA* 281:1211-1213
- Bloembergen WE, Stannard DC, Port F, Wolfe RA, Pugh JA, Jones CA, Greer JW, Golper TA, Held PJ, 2003:** Relationship of dose of hemodialysis and cause-specific mortality. *Kidney Int* 50:557-565

- Blumenkrantz MJ, Kopple JD, Moran JK, Grodstein GP, Coburn JW, 1999:** Nitrogen and urea metabolism during continuous ambulatory peritoneal dialysis. *Kidney Int* 20:78-82
- Brandes JC, Piering WF, Beres JA, Blumenthal SS, Fritsche C, 1998:** Clinical outcome of continuous ambulatory peritoneal dialysis predicted by urea and creatinine kinetics. *J Am Soc Nephrol* 2:1430-1435
- Brugnara C, 2003:** Iron deficiency and erythropoiesis: new diagnostic approaches. *Clin Chem.*; 49(10):1573-1578.
- Burkart J, Schreiber M, Tabor T, Korbet S, Stallard R, 2000:** Prospective, multicenter evaluation of patient transfer from PD to HD. *Perit Dial Int* 16:S66
- Burkart JM, Bleyer AJ, Jordan JR, Zeigler NC, 2002:** An elevated ratio of measured to predicted creatinine production in CAPD patients is not a sensitive predictor of noncompliance with the dialysis prescription. *Perit Dial Int* 16:142-146
- Burrowes JD, 2003:** Effects of dietary intake, appetite, and eating habits on dialysis and non-dialysis treatment days in hemodialysis patients: cross-sectional results from the HEMO study. *J Ren Nutr*; 13:191-198.
- Cagney KA, Wu AW, Herbert RJ, Meyer KB, Jenckes MW, Bass EB, Powe NR, 2002:** Formal assessment of quality of life instruments for end-stage renal disease. *Med Decision Making* 15:421-421
- Cai W, 2006:** Association of advanced glycoxidation end products and markers of inflammation with thrombosis of arteriovenous grafts in hemodialysis patients. *Am J Nephrol.*; 26:181-185.

- Canaud B, 2000:** Microbiologic purity of dialysate: rationale and technical aspects. *Blood Purify*;18: 200-213.
- Charytan C, 1999:** Continuous ambulatory peritoneal dialysis after abdominal aortic graft surgery. *Perit Dial Int* 12:227-229
- Charytan C, Spinowitz BS, 2000:** Dialysate Leaks, in Nissenson AR, Fine RN: *Dialysis Therapy* (2 ed). Philadelphia, PA, Hanley and Belfus, pp 188-189
- Che-yi C, 2005:** Acupuncture in haemodialysis patients at the Quchi acupoint for refractory uremic pruritus. *Nephrol Dial Transplant*; 20:1912-1915
- Chobanian AV, 2003:** The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*; 289:2560-2572.
- Chumlea WC, 2003:** Nutritional status assessed from anthropometric measures in the HEMO study. *J Ren Nutr*;13:31-38.
- Churchill D, Thorpe K, Nolph K, Keshaviah P, Oreopoulos P, Page D, 2005:** Increased peritoneal membrane transport is associated with decreased patient and technique survival for continuous peritoneal dialysis patients. *J Am Soc Nephrol* 9:1285-1292
- Casino MB and Lopez LK, 2000:** Prediction of creatinine clearance from serum creatinine. *Nephron* 16:31-41, 1976
- Coyne DW, 2006:** Pernicious anemia: influence of industry on renal guideline development
- Coyne DW, Delmez J, Spence G, Windus DW, 2002:** Impaired delivery of hemodialysis prescriptions: An analysis of causes and an approach to evaluation. *J Am Soc Nephrol* 8:1315-1318

- Cruz DN, Mahnensmith RL, Perazella MA, 2006:** Intradialytic hypotension: Is midodrine beneficial in symptomatic hemodialysis patients? *Am J Kidney Dis* 101-107
- Daugirdas JT, 2003:** Second generation logarithmic estimates of single-pool variable volume Kt/V: An analysis of error. *J Am Soc Nephrol* 4:1205-1213
- Davies SJ, Bryan J, Phillips L, Russell GI, 2003:** The predictive value of KT/V and peritoneal solute transport in CAPD patients is dependent on the type of comorbidity present. *Perit Dial Int* 16:S158-S162
- Dheenan S, 2004:** Continuous peritoneal dialysis in heavyweight individuals: urea and creatinine clearances. *Perit Dial Int* 16:302-306
- Depner TA, 1997:** Single-Compartment Model. in *Prescribing Hemodialysis: A Guide to Urea Modeling*. Norwell, MA, Kluwer Academic Publishers; 81-88
- Depner TA, 2001:** hemodialysis efficiency: an analysis of alarm system. ;8(4):227-235.
- Depner TA, Greene T, Gotch FA, Daugirdas JT, Keshaviah PR, Star RA, 1999:** Hemodialysis Study Group: Imprecision of the hemodialysis dose when measured directly from urea removal. *Kidney Int* 55:635-647
- D'Haese PC, 2000:** Increased bone strontium levels in hemodialysis patients with osteomalacia. *Kidney Int.*; 57:1107-1114.
- D'Haese PC, 2003:** A multi-center study on the effects of lanthanum carbonate (Fosrenol) and calcium carbonate on renal bone disease in dialysis patients. *Kidney Int.*;63: 73-78.

- Dheenan S, 2002:** Effect of sertraline hydrochloride on dialysis hypotension. *Am J Kidney Dis.*; 31:624-630.
- Dhondt AW, Vanholder RC, De Smet RV, Claus SA, Waterloos MA, Glorieux GL, Delanghe JR, Lameire NH, 2003:** Studies on dialysate mixing in the Genius single-pass batch system for hemodialysis therapy. *Kidney Int.*; 63: 1540–1547
- Di Filippo S, 2006:** Reduction in urea distribution volume over time in clinically stable dialysis patients. *Kidney Int.*; 69(4):754-59.
- Diaz-Buxo JA, Crawford-Bonadio TL, St Pierre D, Ingram KM, 2006:** Establishing a successful home dialysis program. *Blood Purify*; 24: 22–27
- Dogan E, 2005:** Relation between Adequacy, some laboratory parameters, and quality-of-life in hemodialysis patients. *Renal Failure* ;27:695-699.
- Don BR, 2001:** Leptin is a negative acute phase protein in chronic hemodialysis patients. *Kidney Int.* ;59:1114-1120.
- Doorenbos CJ, Bosma RJ, Lamberts PJ, 2001:** The slight increase in volume in parallel-plate dialyzers during ultra-filtration. *Nephrol Dial Transplant*; 16:1303–1304.
- Doran T, Fullwood C, Gravelle H, Reeves D, Kontopantelis E, Hiroeh U, Roland M, 2006:** Payfor-performance programs in family practice in the United Kingdom. *N Engl J Med.*; 355: 375–384.
- Duffy R, 2000:** Multistate outbreak of hemolysis in hemodialysis patients traced to faulty blood tubing sets. *Kidney Int.*; 57(4): 1668-1674.
- Edefonti A, Picca M, Damiani B, Loi S, Ghio L, Giani M, Consalvo G, Grassi MR, 2006:** Dietary prescription based on estimated

- nitrogen balance during peritoneal dialysis. *Pediatr Nephrol* 13:253-258
- Eknoyan G, 2002:** Effect of dialysis dose and membrane flux in maintenance hemodialysis. *N Engl J Med*; 19:347
- Emmett M, 2000:** Effect of three bases and a cation exchange resin on serum sodium and potassium.;108:752-760.
- European Pharmacopoeia Supplement, 2001:** Haemodialysis solutions, concentrated, water for diluting. Monograph 1167
- European Renal Association, 2002:** European Dialysis and Transplantation Association. European best practice guidelines for hemodialysis, section IV dialysis fluid purity. *Nephro Dial Transplant* ;17:45-62
- Evanson JA, 1999:** Measurement of the delivery of dialysis in acute renal failure. *Kidney Int.*;55:1501-1508.
- Fagugli RM, 2006:** Effects of short daily hemodialysis and extended standard hemodialysis on blood pressure and cardiac hypertrophy: a comparative study. *J Nephrol.*; 19:77-83.
- Fishbane S, 2001:** A randomized trial of iron deficiency testing strategies in hemodialysis patients. *Kidney Int.*; 60(6):2406-2411.
- Foley RN, 2002:** Blood pressure and long-term mortality in United States hemodialysis patients: USRDS Waves 3 and 4 Study. *Kidney Int*; 62: 1784-1790.
- Frankenfield D, McClellan WM, Helgersen SD, Lowrie EG, Rocco M, Owen WF, 2006:** Relationship between urea reduction ratio, demographic characteristics, and body weight for patients. National ESRD Core Indicators Project. *Am J Kidney Dis* 33:584-591

- Franssen CFM, 2006:** Adenosine and dialysis hypotension. *Kidney Int.*; 69:789-791.
- Friedman EA, 2003:** International comparisons of survival on dialysis: are they reliable? *Hemodialysis Int.*; 7: 59–66.
- Galland R, 2004:** Traeger J. Short daily hemodialysis and nutritional status in patients with chronic renal failure. *Semin Dial.*;17:104-108.
- Galland R, Traeger J, 2004:** Short daily hemodialysis and nutritional status in patients with chronic renal failure. *Semin Dial.*; 17:104-108.
- Genestier S, Hedelin G, Schaffer P, Faller B, 2002:** Prognostic factors in CAPD patients: A retrospective study of a 10-year period. *Nephrol Dial Transplant* 10:1905-1911
- Ghandour H, 2002:** Distribution of plasma folate forms in hemodialysis patients receiving high daily doses of L-folinic or folic acid. *Kidney Int* 2002;62:2246-2249.
- Goel SP, Nolph KD, Moore HL, Prowant BF, Khanna R, Twardowski ZJ, 1999:** A prospective study of the effect of noncompliance on small solute removal in continuous ambulatory peritoneal dialysis. *ASAIO J* 41:452-456
- Goovaerts T, Jadoul M, Goffin E, 2005:** Influence of a pre-dialysis education programme (PDEP) on the mode of renal replacement therapy. *Nephrol Dial Transplant*; 20: 1842-1847
- Gotch FA, 2002:** Kinetic modeling in hemodialysis, in Nissenson AR, Fine RN, Gentile DE (eds): *Clinical Dialysis*. (ed 3). East Norwalk, CT, Appleton & Lange; 156-189
- Goovaerts M, 2005:** The incident of patient cohort study design with uncontrolled dose may result in a substantial overestimation of

mortality (M) as a function of peritoneal dialysis (PD) dose. ASAIO Trans 42:102

Gotch FA, Keen M, 2002: Kinetic modeling in peritoneal dialysis, in Nissenson AR, Fine RN, Gentile DE (eds): Clinical Dialysis (ed 3). Norwalk, CT, Appleton and Lange; 343-375

Gotch FA, Lipps BJ, Keen ML, Panlilio F, 2003: Coinvestigators for the Fresenius Randomised Dialysis Prescriptions and Clinical Outcome Study (RDP/CO): Computerized urea kinetic modeling (UKM) to prescribe and monitor delivered Kt/V (pKt/V, dkT/V) in peritoneal dialysis. Adv Perit Dial 12:43-45

Gregory M and Holley JL, 2000: Assessing compliance with the dialysis prescription in children on peritoneal dialysis. Perit Dial Int 16:69-73

Gulanikar AC, Jindal KK, Hirsch DJ, 1998: Is chronic peritoneal dialysis safe in patients with intrabdominal prosthetic vascular graphs? Nephrol Dial Transplant 6:215-217

Hakim RM, 2005: Assessing the adequacy of dialysis. Kidney Int 37:822-832

Harty J, Boulton H, Heelis N, Uttley L, Venning M, Gokal R, 2000: Limitations of kinetic models as predictors of nutritional and dialysis adequacy in continuous ambulatory peritoneal dialysis patients. Am J Nephrol 13:454-463

Harty J, Faragher B, Venning M, Gokal R, 2002: Urea kinetic modelling exaggerates the relationship between nutrition and dialysis in CAPD patients. (The hazards of cross-sectional analysis). Perit Dial Int 15:105-109

Hays DF, Baird J, Simpson K, Allison ME, 2004: Urea monitoring during dialysis: the wave of the future. ASAIO Trans 37:49-53

- Heaf J, 2002:** CAPD adequacy and dialysis morbidity: Detrimental effect of a high peritoneal equilibration rate. *Ren Fail* 17:575-587
- Health Care Financing Administration, 1999:** Annual Report, End Stage Renal Disease Core Indicators Project. Department of Health and Human Services, Health Care Financing Administration, Office of Clinical Standards and Quality
- Hertel GH, Goldman RS, Murata GH, Malhotra D, 1999:** Volume of distribution and fractional clearance of urea in amputees on continuous ambulatory peritoneal dialysis. *Perit Dial Int* 14:356-361
- Hiroshige K, Yuu K, Soejima M, Takasugi M, Kuroiwa A, 1999:** Rapid decline of residual renal function in patients on automated peritoneal dialysis. *Perit Dial Int* 16:307-315
- Holley JL, 2005:** Palliative care in end-stage renal disease: focus on advance care planning, hospice referral, and bereavement. *Semin Dial*;18:154-156.
- Holley JL, Piraino B, 2001:** Careful patient selection and dialysis prescription are required for effective nightly intermittent peritoneal dialysis. *Perit Dial Int* 14:155-158
- Hood AH, Malhotra D, Harford AM, 1999:** Creatinine clearance in amputees on CPD. *Perit Dial Int* 16:426-426
- Ifudu AH, Saddler MC, Murata GH, Malhotra D, Sena P, Simon D, Ellege L, Gibel LJ, 2002:** Symptomatic fluid retention in patients on continuous peritoneal dialysis. *J Am Soc Nephrol* 6:198-206

Ikizler JY, Depner T: Principles of hemodialysis, in Owen WF, Pereira BJG, Sayegh M, 2001: Dialysis and Transplantation (ed 1). Philadelphia, PA, Saunders

Johansson A-C, Attman P, Haraldsson B, 1999: Creatinine generation rate and lean body mass: A critical analysis in peritoneal dialysis patients. *Kidney Int* 51:855-859

Joki N, 2006: Relationship between serum albumin level before initiating hemodialysis and angiographic severity of coronary atherosclerosis in end-stage renal disease patients. *Nephrol Dial Transplant*; 21(6):1633 -1639.

Kaizu Y, 2003: Association between inflammatory mediators and muscle mass in long-term hemodialysis patients. *Am J Kidney Dis.*; 42:295-302.

Kalantar-Zadeh K, 2001: A malnutrition-inflammation score is correlated with morbidity and mortality in maintenance hemodialysis patients. *Am J Kidney Dis.*;38:1251-1263.

Kasama R, 2006: Vitamin B₆ and hemodialysis: the impact of high flux/high-efficiency dialysis and review of the literature. *Am J Kidney Dis.*; 8:680-686.

Kayatas M, 2006: Comparison of the non-invasive methods estimating dry weight in hemodialysis patients. *Ren. Fail.*; 28(3):217-222.

Kaysen GA, 2001: The microinflammatory state in uremia: causes and potential consequences. *J Am Soc Nephrol.*;12:1549-1557.

Keshaviah P, Emerson PF, Vonesh EF, Brandes JC, 2004: Relationship between body size, fill volume, and mass transfer area coefficient in peritoneal dialysis. *J Am Soc Nephrol* 4:1820-1826

- Keshaviah PR, Nolph KD, 1998:** Protein catabolic rate calculations in CAPD patients. *ASAIO Trans* 37:M400-M402, 1991
- Khan AH, Harford AM, Sena P, Eisenberg B, Wood B, Simon D, Goldman RS, Kanig SP, 1998:** Hand gangrene in diabetics on chronic dialysis. *ASAIO Trans* 37:638-643
- Khanna R, 2001:** The Toronto Western Hospital protocol. *Perit Dial Bull*; 1:101.
- Khanna R, Nolph KD, Oreopoulos DG, 2000:** Peritoneal dialysis in diabetics, in *The Essentials of Peritoneal Dialysis*. Boston, MA, Kluwer Academic Publishers; 99-100
- Kimmel PL, Bosch JP, 1998:** Effectiveness of renal fellowship training for subsequent clinical practice. *Am J Kidney Dis* 18:249-256
- Kjellstrand CM, Blagg CR, 2005:** Differences in dialysis practice are the main reasons for the high mortality rate in the United States compared to Japan. *Hemodial Int.*; 7: 67
- Kogirima M, 2006:** Low resting energy expenditure in middle-aged and elderly hemodialysis patients with poor nutritional status. *J Med Invest.*; 53(1-2):34-41.
- Kohaut EC, Waldo FB, Benfield MR, 2001:** The effect of changes in dialysate volume on glucose and urea equilibration. *Perit Dial Int* 14:236-239
- Konig PS, Lhotta K, Kronenberg F, Ioannidis M, Herold M, 1998:** CAPD: A successful treatment in patients suffering from therapy-resistant congestive heart failure. *Adv Perit Dial* 7:97-101
- Kopple JD, Jones MR, Keshaviah PR, Bergstrom J, Lindsay RM, Moran J, Nolph KD, Teehan BP, 2002:** A proposed glossary for dialysis kinetics. *Am J Kidney Dis* 26:963-981

- Korbet SM, Rodby RA, 2001:** Peritoneal membrane failure: Differential diagnosis, evaluation, and treatment. *Semin Dial* 7:128-137
- Kouidi E, 2007:** Exercise renal rehabilitation program: psychosocial effects. *Nephron.*; 77:152-158.
- Kramer HJ, 2006:** Increasing body mass index and obesity in the incident ESRD population. *J Am Soc Nephrol.*; 17(5):1453-1459.
- Krivitski NM, 2005:** Theory and validation of access flow measurement by dilution technique during hemodialysis. *Kidney Int.*; 48:244-250
- Kurella M, 2005:** Suicide in the end-stage renal disease program. *J Am Soc Nephrol.*; 16:774-781.
- Lameire NH, Vanholder R, Veyt D, Lambert M, Ringoir S, 1999:** A longitudinal, five year survey of urea kinetic parameters in CAPD patients. *Kidney Int* 42:426-432
- Le Meur Y, 2001:** Plasma levels and metabolism of AcSDKP in patients with chronic renal failure: relationship with erythropoietin requirements. *Am J Kidney Dis.*; 38:510-517.
- Ledebol, 2004:** Ultra pure dialysis fluid – direct and indirect benefits in dialysis therapy. *Blood Purify* ; 22: 20–25
- Leray-Moragues H, Kerkeni N, Bosc JY, Martin K, 2002:** Effective flow performances and dialysis doses delivered with permanent catheters: A 24-month comparative study of permanent catheters versus arterio-venous vascular accesses. *Nephrol Dial Transplant* 17:1286-1292
- Levey AS, Bosch J, Breyer J, Lewis J, Greene T, Rogers N, Roth D, 2004:** A more accurate method to estimate glomerular filtration

rate from serum creatinine: A new prediction equation. *Ann Intern Med* 130:461

Leyboldt JK, 2007: Hemodialyzer mass transfer-area coefficients for urea increase at high dialysate flow rates. The Hemodialysis (HEMO) study. *Kidney Int.*;51:2013-2017.

Lindsay RM, Spanner E, Heidenheim RP, LeFebvre JM, Hodsman A, Baird J, Allison MEM, 2007: Which comes first, Kt/V or PCR-chicken or egg? *Kidney Int Suppl* 42:S32-S36

Lo WK, Prowant BF, Gamboa SB, Moore HL, Nolph KD, 2001: How reproducible is daily creatinine recovery in chronic peritoneal dialysis. *Perit Dial Int* 14:286-288

Lopez-Gomez JM, 2004: Presence of a failed kidney transplant in patients who are on hemodialysis is associated with chronic inflammatory state and erythropoietin resistance. *J Am Soc Nephrol.*; 15(9):2494-2501.

Lorenzo V, Martn M, Rufino M, Hernandez D, Torres A, Ayus JC, 2004: Predialysis nephrologic care and a functioning arteriovenous fistula at entry are associated with better survival in incident hemodialysis patients: an observational cohort study. *Am J Kidney Dis.*; 43: 999–1007

Lowrie EG, 2005: The online measurement of peritoneal dialysis dose (Kt): clinical outcome as A function of body surface area. *Kidney Int.*;68(3):1344-1354.

Lutes R, Holley JL, Perlmutter J, Piraino B, 2000: Correlation of normalized protein catabolic rate to weekly creatinine clearance and KT/V in patients on peritoneal dialysis. *Adv Perit Dial* 9:97-100

- MacLeod AM, 2006:** Cellulose, modified cellulose and synthetic membranes in the haemodialysis of patients with end-stage renal disease. *Cochrane Database Syst Rev*
- MacRae JM, Ahmed A, Johnson N, Levin A, Kiaii M, 2005:** Central vein stenosis: A common problem in patients on hemodialysis. *ASAIO J* 51:77-81
- Mactier RA, 1998:** Investigation and management of ultrafiltration failure in CAPD. *Adv Perit Dial* 7:57-62
- Mactier RA, Perry M, Henderson IS, 2000:** Relationships between dialysis quantification and normalized protein catabolic rate in peritoneal dialysis. *Perit Dial Int* 13:508-511
- Maggiore Q, 2002:** The effects of control of thermal balance on vascular stability in hemodialysis patients: results of the European randomized clinical trial. *Am J Kidney Dis.*; 40(2):280-290.
- Maiorca R, Brunori G, Zubani R, Cancarini GC, Manili L, Camerini C, Movilli E, Pola A, d'Avolio G, Gelatti U, 2002:** Predictive value of dialysis adequacy and nutritional indices for mortality and morbidity in CAPD and HD patients. A longitudinal study. *Nephrol Dial Transplant* 10:2295-2305
- Maiorca R, Cancarini GC, Brunori G, Camerini C, Manili L, 2000:** Morbidity and mortality of CAPD and hemodialysis. *Kidney Int* 43:4-15
- Malhotra D, Tzamaloukas AH, Murata GH, Fox L, Goldman RS, Avasthi PS, 2003:** Serum albumin in continuous peritoneal dialysis: Its predictors and relationship to urea clearance. *Kidney Int* 50:1501-1507

- Marshall M, 2006:** Associations of hemodialysis dose and session length with mortality risk in Australian and New Zealand patients. *Kidney Int.*; 69:1229-1236.
- Mehrotra R, Blake P, Berman N, Nolph KD, 2002:** An analysis of dialysis training in the United States and Canada. *Am J Kidney Dis.*; 40: 152–160.
- Mendelssohn DC, Ethier J, Elder SJ, Saran R, Port FK, Pisoni RL, 2006:** Haemodialysis vascular Dialysis Outcomes and Practice Patterns Study (DOPPS II). *Nephrol Dial Transplant*; 21: 721–728.
- Mendley SR, Majkowski NL, 2003:** Protein intake and urea and nitrogen appearance in stable pediatric PD patients. *J Am Soc Nephrol* 7:1432-1432
- Mendoza FA, 2006:** Description of 12 cases of nephrogenic fibrosing dermopathy and review of the literature. *Semin Arthritis Rheum*; 35:238-249.
- Meyer KB, Espindle DM, DeGiacomo JM, Jenulson CS, Kurtin PS, Davies AR, 2001:** Monitoring dialysis patients' health status. *Am J Kidney Dis* 24:267-279
- Misra M, 2005:** Core curriculum. The basics of hemodialysis equipment. *Hemodial Int*; 9:30-36.
- Mitch WE and Walser M, 1998:** A proposed mechanism for reduced creatinine excretion in severe chronic renal failure. *Nephron* 21:248-254
- Murata GH, Tzamaloukas AH, Malhotra D, Saddler MC, Fox L, Wood B, Morgan K, Goldman RS, 2003:** Protein catabolic rate in patients on continuous peritoneal dialysis: A multivariate predictive model. *ASAIO J* 42:46-51

- Nakai S, Shinzato T, Sanaka T, Kikuchi K, Kitaoka T, Shinoda T, Yamazaki C, Sakai R, Omori H, Morita O, Iseki K, Kubo K, Tabei K, Masakane I, Fushimi K, Akiba T, 2004:** An overview of dialysis treatment in Japan. *J Jpn Soc Dial Ther.*; 34: 1121–1147.
- Nakamoto H, 2002:** Effect of optimization of hemodialysis dose on adequacy of hemodialysis. *Am J Kidney Dis.*; 40:1045-1054.
- National Kidney Foundation, 2006:** K/DOQI Clinical Practice Guidelines for hemodialysis Adequacy. *Am J Kidney Dis* 37:S65-S136
- Neal B, 2000:** Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Blood Pressure Lowering Treatment Trialists' Collaboration. Lancet*; 356:1955-1964.
- Nissenson AR, Agarwall R, Allon M, Cheung AK, Clark W, Depner T, Diaz-Buxo JA, Kjellstrand C, Klinger A, Martin KJ, Norris K, Ward R, Wish J, 2004:** Improving outcomes in CKD and ESRD patients: carrying the torch from training to practice. *Semin Dial.*; 17: 380–397.
- Nolph KD, Jensen RA, Khanna R, Twardowski ZJ, 2001:** Weight limitations for weekly urea clearances using various exchange volumes in continuous ambulatory peritoneal dialysis. *Perit Dial Int* 14:261-264
- Nolph KD, Twardowski ZJ, Khanna R, Moore HL, Prowant BF, 2002:** Predicted and measured daily creatinine production in CAPD: Identifying noncompliance.

Owen WF Jr, Lew NL, Yan Liu MS, Lowrie EG, Lazarus JM, 2000:

The urea reduction ratio and serum albumin concentration as predictors of mortality in patients undergoing hemodialysis. *N Engl J Med* 329:1001-1006

Owen WF, Chertow G, Lazarus JM, Lowrie EG, 2006: The dose of hemodialysis: Mortality responses by race and gender. *JAMA* 280:1-6

Perini S, LaBerge JM, Pearl JM, 2006: Tesio catheter: Radiologically guided placement, mechanical performance, and adequacy of delivered dialysis. *Radiology* 215:129-137

Powe NR, Klag MJ, Sadler JH, Anderson GF, Bass EB, Briggs WA, Fink NE, Levey AS, Levin NW, Meyer KB, Rubin HR, Wu AW, 2003: Choices for healthy outcomes in caring for end-stage renal disease. *Semin Dial* 9:9-11

Pritchard SS, 2003: Treatment modality selection in 150 consecutive patients starting ESRD therapy. *Perit Dial Int* 16:69-72, 1996

Pupim LB, Cuppari L, Ikizler TA, 2006: Nutrition and metabolism in kidney disease. *Semin Nephrol.*; 26(2):134-157.

Pupim RB, 2002: Intradialytic parenteral nutrition improves protein and energy homeostasis in chronic hemodialysis patients. *J Clin Invest.*; 110:483-492.

Rayan SS, Terramani TT, Weiss VJ, Chaikof EL, 2003: The LifeSite Hemodialysis machine *J Vasc Surg* 38:714-718

Rayner HC, Besarab A, Brown WW, Disney A, Saito A, Pisoni RL, 2004: Vascular access results from the Dialysis Outcomes and PracticePatterns Study (DOPPS): performance against Kidney

- Disease Outcomes Quality Initiative (K/DOQI) Clinical Practice Guidelines. *Am J Kidney Dis*; 44: 22–26.
- Renal Physicians Association, 2006:** Shared decision making (guideline regarding withdrawal from dialysis and palliative care). September 12
- Ritz E, 2006:** Lowering of blood pressure the lower, the better ? *J Am Soc Nephrol*; 17:2345-2352.
- Robertson BC, Juhasz NM, Walker PJ, Raymond KH, Taber TE, Adcock AA, 2002:** A prescription model for peritoneal dialysis. *ASAIO J* 41:116-126
- Robson D, Fox L, Goldman RS, Avasthi PS, 2003:** Creatinine clearance in continuous peritoneal dialysis: Dialysis dose required for a minimal acceptable level. *Perit Dial Int* 16:41-47
- Rocco MV, Jordan JR, Burkart JM, 1999:** Changes in peritoneal transport during the first month of peritoneal dialysis. *Perit Dial Int* 15:12-17
- Rodby RA, Firanek CA, Cheng YG, Korbet SM, 1999:** Reproducibility of studies of peritoneal dialysis adequacy. *Kidney Int* 50:267-271
- Ronco C, Bosch JP, Lew SQ, Feriani M, Chiaramonte S, Conz P, Brendolan A, La Greca G, 2001:** Adequacy of continuous ambulatory peritoneal analysis: Comparison with other dialysis techniques. *Kidney Int Suppl* 46:S18-S24
- Rudolph H, 2003:** Muscle and serum trace element profile in uremia. *Trans Am Soc Artif Intern Organs* ; 19:456-465.
- Sarkar JT, Agodoa LY, Jones CA, Port FK, 2006:** Body size, dose of hemodialysis, and mortality. *Am J Kidney Dis* 35:80-88.

- Salusky IB, 2002:** Nutritional management of pediatric patients on chronic dialysis, in Nissenson AR, Fine RN, Gentile DE (eds): Clinical Dialysis (ed 3). Norwalk, CT, Appleton and Lange, 1995, pp 535-548
- Sam R, Walker PJ, Raymond KH, 2006:** Composition and clinical use of hemodialysates. Hemodialysis Int.; 10:15-28.
- Saran R, Bragg-Gresham JL, Levin NW, Twardowski ZJ, Wizemann V, Saito A, Kimata N, Gillespie BW, Combe C, Bommer J, Akiba T, Mapes DL, Young EW, Port FK, 2006:** Longer treatment time and slower ultrafiltration in hemodialysis: Associations with reduced mortality in the DOPPS. Kidney Int ; 69: 1222–1228
- Schaefer F, Haraldsson B, Haas S, Simkova E, Feber J, Mehls O, 2005:** Estimation of peritoneal mass transport by three-pore model in children. Kidney Int 54:1372-1379
- Schaefer F, Klaus G, Mehls O, 2006:** The Mid-European Pediatric Peritoneal Dialysis Study Group: Peritoneal transplant properties and dialysis dose affect growth and nutritional status in children on chronic peritoneal dialysis. J Am Soc Nephrol 10:1786-1792
- Schmidt RJ, Cruz C, Dumler F, 2001:** Effective continuous ambulatory peritoneal dialysis following abdominal aortic aneurysm repair. Perit Dial Int 13:40-44
- Schoenfeld PY, Diaz-Buxo J, Keen M, Gotch FA, 2000:** The effect of body position (P), surface area (BSA), fill volume and intraperitoneal exchange volume (Vip) on the peritoneal transport constant. J Am Soc Nephrol 4:416

- Sehgal A, Snow RJ, Sinder ME, Amini SB, DeOreo PB, Silver MR., Cebul RD, 2004:** Barriers to adequate delivery of hemodialysis. *Am J Kidney Dis* 31:593-601
- Sherman RA, 2004:** Effect of variations in dialysis solution temperature on blood pressure during hemodialysis. *Am J Kidney Dis.*; 4:66-68.
- Smith L, Folden T, Youngblood B, Panlilio F, Gotch F, 2003:** Clinical evaluation of a peritoneal dialysis kinetic modeling set. *Adv Perit Dial* 12:46-48, 1996
- Sobh M, Sheeshaa H, Mahmoud K, Ghonim M, 1998:** The relationship between midarm circumference, laboratory parameters and by calculating the normalized protein catabolic rate (nPCR).. *Af. Neph. J.*; 20:10-19
- Sorof JM, Brewer ED, Portman RJ, 2004:** Ambulatory blood pressure monitoring and interdialytic weight gain in children receiving chronic hemodialysis. *Am J Kidney Dis* 33:667-674
- Stegmayr B, Banga R, Lundberg L, Wikdahl AM, Plum-Wirell M, 2003:** PD treatment for severe congestive heart failure. *Perit Dial Int* 16:231-235
- Stein A, Walls J, 2001:** The correlation between Kt/V and protein catabolic rate—A self-fulfilling prophecy. *Nephrol Dial Transplant* 9:743-745
- Stiller S, Bonnie-Schorn E, Grassmann A, Uhlenbusch-Korwer I, Mann H, 2001:** A critical review of sodium profiling for hemodialysis. *Semin Dial.*; 14: 337–347
- Struijk DG, Krediet RT, Koomen GCM, Boeschoten EW, Arisz L, 2001:** The effect of serum albumin at the start of continuous ambulatory peritoneal dialysis treatment on patient survival. *Perit Dial Int* 14:121-126

Szczech LA, Lowrie E, Li Z, Lazarus, JM, Lew NL, Owen WF, 2003:
Changing hemodialysis thresholds for optimal survival. Kidney
Int.; 59:738-45

Tattersall JE, Doyle S, Greenwood RN, Farrington K, 2001:
Maintaining adequacy in CAPD by individualizing the dialysis
prescription. Nephrol Dial Transplant 9:749-752

Tawney K, 2000: developing a dialysis rehabilitation program. Nephrol
Nurs J ; 27:524-539.

Teehan BP, Schleifer CR, Brown J, 1999: Urea kinetic modeling is an
appropriate assessment of adequacy. Semin Dial 5:189-192

Teehan BP, Schleifer CR, Brown J, 2001: Adequacy of continuous
ambulatory peritoneal dialysis: Morbidity and mortality in
chronic peritoneal dialysis. Am J Kidney Dis 24:990-1001

Tucker KL, 2005: High homocysteine and low B vitamin predict
cognitive decline in aging men: The Veterans Affairs
Normative Aging Study. Am J Clin Nutr.; 82:627-635.

**Twardowski ZJ, Nolph KD, Khanna R, Prowant BF, Ryan LP,
Moore HL, Nielsen MP, 2000:** Peritoneal equilibration tests.
Perit Dial Bull 7:138-147

Tzamaloukas AH, 2003: Pharmacokinetic analysis of creatinine generation
discrepancy as an index of noncompliance in CAPD. Adv Perit
Dial 12:61-65

Tzamaloukas AH, 2004: Serum tonicity, extracellular volume and
clinical manifestations in symptomatic dialysis-associated
hyperglycemia treated only with insulin. Int J Artif Organs;
27:751-758.

Tzamaloukas AH, 2005: Can an excess of estimated over-predicted
creatinine generation be a discriminating test for non-

compliance in continuous ambulatory peritoneal dialysis? J Am Soc Nephrol 6:1519-1520.

Tzamaloukas AH, Braun M, Malhotra D, Murata GH, 2001: Estimated versus predicted creatinine generation as an indicator of compliance with the prescribed dose of continuous peritoneal dialysis. Int J Artif Organs 19:151-155

Tzamaloukas AH, Murata GH, Malhotra D, Fox L, Goldman RS, 2004: An analysis of the determinants of urinary urea and creatinine clearance in patients on continuous peritoneal dialysis. Adv Perit Dial 13:38-41

Tzamaloukas AH, Murata GH, Malhotra D, Fox L, Goldman RS, Avasthi PS, 2002: The minimal dose of dialysis required for a target KT/V in continuous peritoneal dialysis. Clin Nephrol 44:316-321

Uribarri J, 2004: Acid production in chronic hemodialysis patients. J Am Soc Nephrol.;9:114-120.

US Renal Data System, 2005: Excerpts from the USRDS 2004 Annual Data Report: atlas of end-stage renal disease in the United States. Am J Kidney Dis.; 45: 1–280.

US Renal Data Systems, 2003: Hospitalization, in National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases (ed): US Renal Data Systems 1996 Annual Data Report, Bethesda, MD, p 111

Vonesh EF, Burkart J, McMurray SD, Williams PF, 2003: Peritoneal dialysis in kinetic modeling: Validation in a multicenter clinical study. Perit Dial Int 16:471-481

- Vonesh EF, Lysaght MJ, Moran J, Farrell P, 1998:** Kinetic modeling as a prescription aid in peritoneal dialysis. *Blood Purif* 9:246-270
- Vonesh EF, Story KO, O'Neill WT, 2006:** A multinational clinical validation study of PD Adequest 2.0. *Perit Dial Int* 19:556-571
- Wang J, 2001:** Anthropometry in body composition. An overview. *Ann N Y Acad Sci*; 904:317-326.
- Warady B, 2003:** Longitudinal evaluation of transport kinetics in children receiving peritoneal dialysis. *Pediatr Nephrol* 13:571-576
- Warady BA and Hossli S, 2000:** Peritoneal membrane transport function in children receiving long-term dialysis. *J Am Soc Nephrol* 7:2385-2391
- Ward RA, 2004:** Ultra pure dialysate.; 17:489-497
- Warren PJ, Brandes JC, 2003:** Compliances with the peritoneal dialysis prescription is poor. *J Am Soc Nephrol* 4:1627-1629
- Weijmer MC, Vervloet MG, Wee PM, 2004:** Compared to tunnelled cuffed haemodialysis catheters, temporary untunnelled catheters are associated with more complications already within 2 weeks of use. *Nephrol Dial Transplant*. 19:670-677
- Wolfe RA, Ashby VB, Daugirdas JT, Agodoa LY, Jones CA, Port FK, 2006:** Body size, dose of hemodialysis, and mortality. *Am J Kidney Dis* 35:80-88.
- Wu AW, Cagney KA, Fink NE, Meyer KB, Herbert RJ, Jenckes MW, Sadler JH, Powe NR, 2002:** Development of a quality life measure for renal dialysis patients. *Quality Life Res* 4:505-505

Wuerth D, 2003: Chronic peritoneal dialysis patients diagnosed with clinical depression: results of pharmacologic therapy. *Semin Dial*;16:424-427.

Zawada ET, 2001: Indications for dialysis, in Daugirdas JT, Ing TS (eds): *Handbook of Dialysis* (ed 2). Toronto, Canada, Little, Brown and Co, pp 6-7