

REFERENCES

<i>Abaci A, Ekici E, Oguzhan A (2004).</i> Cardiac troponins T and I in patients with end-stage renal disease/the relation with left ventricular mass and their prognostic value. Clin Cardiol 2004; 27: 704–9.
<i>Abaci A, Ekici E, Oguzhan A, Tokgoz B and Utas C</i> Cardiac troponins T and I in patients with end-stage renal disease: the relation with left ventricular mass and their prognostic value. Clin Cardiol. 2004 Dec;27(12):704-9.
<i>Adams J, Abendschein D, and Jaffe A (1993)</i> Biochemical markers of myocardial injury. Circulation 88: 750
<i>Adams J, Bodor GS, Davila-Roman VG, Delmez JA, et al (1993).</i> Cardiac troponin I: a marker with high specificity for cardiac injury. Circulation 88: 101-6
<i>Amann K, Wiest G, Zimmer G, Gretz N, Ritz E and Mall G (1992)</i> Reduced capillary density in myocardium of uremic rats. A stereological study. Kidney Int 42: 1079-1085
<i>Ammann P, Fehr T, Minder EI (2001).</i>

Elevation of troponin I in sepsis and septic shock. Intensive Care Med; 27: 965–9.
<i>Apple FS, Sharkey SV, Hoeft P (1997).</i> Prognostic value of serum cardiac troponin I and T in chronic dialysis patients: a 1 year outcome analysis. Am J Kidney Dis 1997; 29: 399–403.
<i>Apple FS, Murakami MM, Pearce LA. (2002)</i> Predictive value of cardiac troponin I and T for subsequent death in end-stage renal disease. Circulation; 106: 2941–5.
<i>Apple FS, Pearce LA, and Herzog CA (2004).</i> Multi-biomarker risk stratification of N-terminal pro-B- type natriuretic peptide, high sensitivity C- reactive protein, and cardiac troponin T and I in end stage renal disease for all-cause death. Clin chem. 50(12): 2279-2285.
<i>Atman PO, Samuelsson O, Johansson AC and Moberly JB (2003).</i> Dialysis modalities and dyslipidaemia. Kidney International, 63, supp 84: S110-S112.
<i>Aviles RJ, Askari AT, Lindahl B (2002).</i> Troponin T levels in patients with acute coronary syndromes, with or without renal dysfunction. N Engl J Med 2002; 346: 2047–52
<i>AY-M Wang, CW-K Lam, C-M Yu1, M Wang, IH-S (2006)</i> Chan, S-F Lui1 and JE Sanderson

Troponin T, left ventricular mass, and function are excellent predictors of cardiovascular congestion in peritoneal dialysis Kidney International 70, 444–452
<i>Boulier A, Jaussent I, Terrier N (2004).</i> Measurement of circulating troponin Ic enhances the prognostic value of C-reactive protein in haemodialysis patients. Nephrol Dial Transplant 2004; 19: 2313–8.
<i>Brunner F, Brynger H and Challah S et all., (1998).</i> Combined report on regular dialysis and transplantation in Europe IX. Proc. EDTA 16, 4.
<i>Chapelle JP, Dubois B, Aldnhoff MC, Gielen J and Rorive G. (2002)</i> Comparison of plasma cardiac troponins T and I in chronically haemodialyzed patients in relation to cardiac status and age. Clin. Chem.. lab. Med. 2002 Mar; 40 (3): 240-245.
<i>Choy, JB., Armastrong , PW., Ulan, RA., Campbell, PM. and Prosser, Cl.(2003).</i> Do cardiac troponins provide prognostic insight in heamdialysis patients? Can. J. Cardiol. 19(8): 907:911.
<i>Collinson P, Boa FG, Gaze DC (2001).</i> Measurement of cardiac troponins. Ann Clin Biochem 2001; 38: 423–49.
<i>Croitoru M, and Teagtmeyer H (1995).</i>

Spurious rises in troponin T in end-stage renal disease. The Lancet; 346. octobe 7.
<i>Danch J., Collins R., Appleby P. and Peto R. (1998)</i> Association of fibrinogen, CRP, albumin or leukocytic count with coronary heart disease J. Am. Coll. Cardiol. 279: 1477-1482.
<i>Danch J., Whincup P, Walker M., et al (2000)</i> Low grade inflammation and coronary heart disease : prospective study and meta analysis BMI 321: 199-204
<i>Danch J., ZocaliS. And Peto R. (1999)</i> Risk factor for coronary heart disease and acute phase protien. A population based study Eur. Heart J. 20: 954-959
<i>De Beer F.C., Dodd C and Lhotta K. (1982)</i> Isolation and charecterization of CRP and serum amyloid P component in the rat Immunology 45: 55-70
<i>De Beer F.C., Kaysen D and Bekar F. (1982)</i> Measurement of serum CRP concentration in myocardial ischemia and infarction Br. Heart J. 47: 239-243
<i>De Beer F.C., Lutas E and Hill M. N. (1982)</i> Low density and very low denisty lipoprotiens are selectively bound by aggregated CRP J. Exp. Med. 156: 230-242
<i>De Mutsert R and , Krediet RT</i>

Malnutrition, inflammation and atherosclerosis (MIA-syndrome) in dialysis patients Ned Tijdschr Geneeskd 2006 Sep 16;150(37):2023-7.
<i>Deegan PB, Lafferty ME, Blumsohn A, Henderson IS, McGregor E. (2001).</i> Prognostic value of troponin T in hemodialysis patients is independent of comorbidity. Kidney Int. Dec;60(6):2399-405
<i>deFilippi C, Wasserman S, Rosanio S, Tiblier E, Sperger H, Tocchi M, Christenson R, Uretsky B, Smiley M, Gold J, Muniz H, Badalamenti J, Herzog C and Henrich W. (2003)</i> Cardiac troponin T and C-reactive protein for predicting prognosis, coronary atherosclerosis, and cardiomyopathy in patients undergoing long-term hemodialysis JAMA. Jul 16; 290(3):353-9.
<i>Desjarlais F, Leblanc M (2004).</i> Signification clinique de l'augmentation de la troponine T cardiaque (cTnT) chez des patients insuffisants renaux sous dialyse. Annales de Biologie Clinique du Quebec 2004;41: 3–9.
<i>Diris JHC, Hackeng CM, Kooman J (2004).</i> Impaired renal clearance explains elevated troponin T fragments in haemodialysis patients. Circulation 2004; 109: 23–5.
<i>Diris JHC, Hackeng CM, Kooman J (2004).</i>

Impaired renal clearance explains elevated troponin T fragments in haemodialysis patients. Circulation 2004; 109: 23–5.
<i>Fatema K, Hirono O, Takeishi Y, Nitobe J, Kaneko K and Ito M.</i> Haemodialysis improves myocardial interstitial edema and left ventricular diastolic function in patients with end-stage renal disease: noninvasive assessment by ultrasonic tissue characterization. Heart Vessels. 2002 Sep;16(6):227-31.
<i>Fathi R, Isbel N, Haluska B, Case C, Johnson DW and Marwick TH</i> Correlates of subclinical left ventricular dysfunction in ESRD. Am J Kidney Dis. 2003 May;41(5):1016-25.
<i>Fehr T, Knoflach A, Ammann P, Pei P, Binswanger U. (2003).</i> Differential use of cardiac troponin T versus I in hemodialysis patients. Clin Nephrol. Jan; 59(1):35-9.
<i>Filho RP, Nordfors L, Lindholm B, and Hoff CM, et al., (2003).</i> Genetic approaches in the clinical investigation of complex disorders: Malnutrition, inflammation and atherosclerosis (MAI) as a prototype. Kidney international; 63: S162-S157.
<i>Foley RN, Parfrey PS, Sarnak MJ. (1998)</i> Clinical epidemiology of cardiovascular disease in

chronic renal disease. Am J Kidney Dis; 32 (Suppl. 3): S112–9.
<i>Frankel WL, Herold DA, Zeigler TW, et al., (1996).</i> Cardiac troponin T is elevated in asymptomatic patients with chronic renal failure Am J Clin Pathol; 106: 118-123
<i>Freda BJ, Tang WH, Van Lente F (2002).</i> Cardiac troponins in renal insufficiency: review and clinical implications J Am Coll Cardiol 2002; 40: 2065–71.
<i>Gabr AE, Ibrahim IA, Aloulou SM, Al-Alfi MA, Al-Abdlrahim KA. (2004).</i> Cardiac troponin T and end stage renal disease. Saudi Med J. Aug; 25(8):1015-9.
<i>Galle J, Quaschnig T, Seibold S, and Wanner C, et al., (2003)</i> Endothelial dysfunction and inflammation: What is the link? Kidney international; 63: S45-S49.
<i>Grieselli M., Sall N.D., and Parker R.A. (1999)</i> CRP and complement are important mediators of tissue damage in acute myocardial infarction J. Exp. Med. 190: 1733-1739
<i>Heeschen C, Hamm CW, Bruemmer J, Simoons ML (2000).</i> Predictive value of C-reactive protein and troponin T in patients with unstable angina: a comparative analysis. CAPTURE Investigators. Chimeric c7E3

AntiPlatelet Therapy in Unstable angina REfractory to standard treatment trial. J Am Coll Cardiol 2000; 35: 1535–42.
<i>Horwich TB, Patel J, MacLellan WR (2003).</i> Cardiac troponin I is associated with impaired hemodynamics, progressive left ventricular dysfunction, and increased mortality rates in advanced heart failure. Circulation; 108: 833–8.
<i>Hutchinson WL. (2000).</i> Immunoradiometric assay of circulating C- reactive protein: age-related values in adult general population. Clin. Chem.46: 934-938
<i>Iliou MC, Fumeron C, Benoit MO, Tuppin P, Courvoisier CL, Calonge VM, Moatti N, Buisson C, Jacquot C; Chronic Haemodialysis and New Cardiac Markers Evaluation (CHANCE) Study. (2001).</i> Factors associated with increased serum levels of cardiac troponins T and I in chronic haemodialysis patients: Chronic Haemodialysis And New Cardiac Markers Evaluation (CHANCE) study. Nephrol Dial Transplant. Jul; 16(7):1452-8.
<i>Iliou, M., Fumeron, C., Benoit, M., and Tuppin,P. (2001)</i> Factors associated with increased serum level of cardiac troponin T and I in chronic haemodialysis

patients : Chronic Heamodialysis And New Markers Evalution (CHANCE) study. Nephrol Dial Transplant 16:1452-1458.
<i>Imazo M, Demichelis B, Cecci E (2003).</i> Cardiac troponin I in acute pericarditis J Am Coll Cardiol; 42: 2144–8.
<i>Ishii J, Nomura M, Okuma T, Naruse T and Mori Y (2001).</i> Risk stratification using serum concentration of cardiac troponin T in patients with end stage renal disease on chronic maintenance dialysis. Clin. Chem.. Acta.312 (1-2): 69-79
<i>Jaffe A</i> Pandora's box is torn asunder. Am Heart J 1999; 138: 9–12.
<i>Jaffe A (2001).</i> Testing the wrong hypothesis: the failure to recognize the limitations of troponin assays. J Am Coll Cardiol 2001; 38: 999– 1001.
<i>Joki N, Nase H, Nakamura R (1997)</i> Onset of coronaryn artery disease prior to initiation of haemodialysis in patients with end-stage renal disease. Nephrol Dial Transplant; 12:718–23.
<i>Jung HH, Ma KR, Han H (2004).</i> Elevated concentrations of cardiac troponins are associated with severe coronary artery calcification in asymptomatic haemodialysis patients.

Nephrol Dial Transplant 2004; 19: 3117–23.
<i>Ketteler M, Wanner C, Matzger T, Bongartz P, and Floege J et al., (2003).</i> Deficiencies of calcium-regulatory proteins in dialysis patients: A novel concept of cardiovascular calcification in ureamia. Kidney international; 63: S84-S87.
<i>Kobori G, Moroi S, Yoshida H, Yamamoto S, Kamoto T, Okuno H and Ogawa O.</i> Marked improvement of cardiac function following renal transplantation: a case report Hinyokika Kiyo. 2003 Jan;49(1):17-9.
<i>Kronenberg F, Lingenhel A, Neyer U, Lhotta K and Konig P et al., (2003)</i> Prevalence of dyslipideamia risk factors in heamodialysis and CAPD patients Kidney international; 63: S113-S116.
<i>Kronenberg F, Lingenhel A, Neyer U, Lhotta K, Konig P and Auinger M</i> Prevalence of dyslipidemic risk factors in hemodialysis and CAPD patients. Kidney Int Suppl. 2003 May;(84):S113-6.
<i>Kuller L.H., Tracy R.P., Shaten J. and Meilhan E.N. (1996).</i> Relation of CRP and coronary heart disease in the MRFIT nested case control study Am J Epidemiol, 144:537-547
<i>Lagrand W. K., Niessen H.W.M., Wolbink G. J.,</i>

<i>Jaspars L.H. and Meur CJLM (1997)</i> CRP colocalizes with complement in human hearts during acute myocardial infarction. Circulation 95: 97-103
<i>Lamb EJ, Webb MC, Abbas NA (2004).</i> The significance of serum troponin T in patients with kidney disease: a review of the literature. Ann Clin Biochem 2004; 41: 1–9.
Lin YP, Chen CH, Yu WC, Hsu TL and Ding PY. Left ventricular mass and hemodynamic overload in normotensive hemodialysis patients. Kidney Int. 2002 Nov;62(5):1828-38
Lipshultz SE, Somers MJ, Lipsitz SR, Colan SD, Jabs K, Rifai N. (2003) Serum cardiac troponin and subclinical cardiac status in pediatric chronic renal failure. Pediatrics. Jul; 112(1 Pt 1):79-86
Lipshultz SE, Somers MJ, Lipsitz SR, Colan SD, Jabs K, Rifai N. (2003) Serum cardiac troponin and subclinical cardiac status in pediatric chronic renal failure. Pediatrics. Jul; 112(1 Pt 1):79-86
<i>Lipshultz, SE., Somers, MJ., Lipsitz, SR., Colan, SD., Jabs, K.and Rifai, N.(2003)</i> Serum cardiac troponin and subclinical cardiac status in pediatric chronic renal failure. Pediatrics 112(1 pt 1):79-86).
<i>Liuzzo G., Maury CPJ., and Nussel E. (1994).</i>

The prognostic value of C- reactive protien, and cerum amyloid A protienin sever unstable angina. N. ENG. J. Med. 331:417-424
<i>Mallamaci F, Zoccali C, Parlongo S, Tripepi G, Benedetto FA, Cutrupi S, Bonanno G, Fatuzzo P, Rapisarda F, Seminara G, Stancanelli B, Bellanuova I, Cataliotti A (2002)</i> Diagnostic value of troponin T for alterations in left ventricular mass and function in dialysis patients. <i>Kidney Int.</i> Nov; 62(5):1884-90.
<i>Mallamaci F, Zoccali C, Parlongo S, Tripepi G, Benedetto FA, Cutrupi S, Bonanno G, Fatuzzo P, Rapisarda F, Seminara G, Stancanelli B, Bellanuova I, Cataliotti A, Malatino LS. (2002).</i> Troponin is related to left ventricular mass and predicts all-cause and cardiovascular mortality in hemodialysis patients. <i>Am J Kidney Dis.</i> Jul; 40(1):68-75.
<i>Mark B. Pepys and Gideon M. Hircshfield (2003).</i> Center for analysis and acute phase protien J. Clin. Invest. 111:1805-1812
<i>Martin LC, Barretti P, Cornejo IV, Felipe MJ, Forti AH and Matsubara BB.</i> Influence of fluid volume variations on the calculated value of the left ventricular mass measured by echocardiogram in patients submitted to hemodialysis. <i>Ren Fail.</i> 2003 Jan;25(1):43-53.

<i>McLaurin MD, Apple FS, Voss EM, Herzog CA, Sharkey SW. (1997).</i> Cardiac troponin I, cardiac troponin T, and creatine kinase MB in dialysis patients without ischemic heart disease: evidence of cardiac troponin T expression in skeletal muscle. <i>Clin Chem.</i> Jun; 43(6 Pt 1):976-82.
<i>Musso P, Cox I, Vidane E, Zambon D, Panteghini M (1999).</i> Cardiac troponins elevation in chronic renal failure: prevalence and clinical significance. <i>Clin Biochem</i> 1999; 32: 125–30.
<i>Nuhbegovic S, Halilbasic A, Karamehic J, Halilovic E and Meric M</i> Effect of haemodialysis on functional parameters of the left ventricle. <i>Med Arh.</i> 1999;53(1):19-20.
<i>Ooi DS, House AA (1998).</i> Cardiac troponin T in haemodialysis patients. <i>Clin Chem</i> 1998; 44: 1410–6.
<i>Pepys M. B. (1999)</i> The Lumleian lecture. CRP and amyloidosis: from protien to drug? In Horizons in medicine. Volume 10. G. Williams, editor. Royal College Physicians of London London UK. 397-414
<i>Pepys M.B., Levely AS and Methe H. (2002)</i> Targeted pharmacological depletion of serum

amyloid P component for treatment of human amyloidosis Nature 417: 254-259
<i>Pepys MB, Dash AC and Ashley MJ (1977)</i> Clin Exp. Immunol. 30:32-37
<i>Pepys MB., Baltz ML. (1983).</i> Acute phase protein with special reference to C-reactive protein and related proteins (pentaxins) and serum amyloid A protein.
<i>Prinsen B, Koning EJ, Koomans HA and Berger R (2003).</i> Hypertriglyceridemia in patients with chronic renal failure: possible mechanism. Kidney International, 63, suppl 84: S121-S124.
<i>Raine A, Margreiter R and Brunner F et al., (1992).</i> Report on management of renal failure in Europe XXII. Nephrol. Dial. Transplant. 7 (suppl. 2): 7.
<i>Ricchiuti V, Voss EM, Ney A (1998).</i> Cardiac troponin T isoforms expressed in renal diseased skeletal muscle will not cause falsepositive results by the second generation cardiac troponin T assay by Boehringer Mannheim. Clin Chem 1998; 44: 1919–24.
<i>Ridker P.M. (2000)</i> Measurement of CRP for the targeting of statin therapy in the primary prevention of acute coronary

events N. Engl. J. Med., 344: 1959-1965
<i>Ridker P.M., Cushman M., Stamper M.J., Tracy R. P. and Hennekens C. H. (1997)</i> Inflammation, aspirin and the risk of cardiovascular disease in apparently healthy men N. Engl. J. Med. 336: 973-979.
<i>Roppolo LP, Fitzgerald R, Dillow J (1999)</i> Comparison of troponin T and I as predictors of cardiac events in patients undergoing chronic dialysis at a veteran's hospital: a pilot study. J Am Coll Cardiol; 34: 448–52.
<i>Sarnak MJ (2003).</i> Cardiovascular complications in chronic kidney disease. Am J Kidney Dis 2003; 41: 11–7.
<i>Sato Y, Yamada T, Taniguchi R</i> Persistently increased serum concentrations of cardiac troponin T in patients with idiopathic dilated cardiomyopathy are predictive of adverse outcomes. Circulation 2001; 103: 369–74.
<i>Shemin D, Bostom AG, and Dworkin LD (2001).</i> Residual renal function and mortality risk in hemodialysis patients. Am J Kidney Dis 38: 85-90.
<i>Shine B., and Pepys MB. (1981).</i> Solid phase radioimmunoassay for C-reactive

protien
<i>Shutov AM, Edigarova OM and Mostykov V.</i> Impact of a hemodialysis session on cardiac function in patients with chronic renal failure. <i>Ter Arkh.</i> 2004;76(9):43-7.
<i>Thompson SG, Kienast J, Pyke SDM, and Van De Loo (1995).</i> Hemostatic factorrs and the risk of MI or sudden death in patients with angina pectoris. N. ENG. J. Med. 332: 635- 641
<i>Townsend PJ, Barton PJR, Ycoub HH (1995).</i> Molecular cloning of human cardiac troponins isoforms. Expression in developing and failing hearts. J Mol Cell Cardiol; 27: 2223–36.
<i>Vigushin DM. Pepys MB. and Hawkins PN. (1993).</i> Metabolic and scintigraphic studies of radioiodinated humen C-reactive protien in health and disease. J. Clin. Invest. 91:1351-1357
<i>Wayand D, Baum H, Scha'tzle G (2000).</i> Cardiac troponin T and I in end-stage renal failure. Clin Chem 2000; 46: 1345–50.
<i>Wayand D, Baum H, Schatzle G (2000).</i> Cardiac troponin T and I in end-stage renal failure. Clin Chem 2000; 46: 1345–50.
<i>Willging S, Keller F, Steinbach G (1998).</i> Specificity of cardiac troponins I and T in renal disease.

Clin Chem Lab Med 1998; 36: 87–92.

Conway B, McLaughlin M, Sharpe P, and Harty J.

Use of cardiac troponin T in diagnosis and prognosis of cardiac events in patients on chronic haemodialysis.

[Nephrol Dial Transplant.](#) 2005 Dec;20(12):2759-64.