

REFERENCES

- Adachi M, Osawa Y, Uchinami H, Kitamura T, Accili and Brenner DA (2007):** The forkhead transcription factor FoxO1 regulates proliferation and transdifferentiation of hepatic stellate cells. *Gastroenterology*; 132: 1434-1446.
- Adachi T, Togashi H, Suzuki A, Kasai S, Ito J, Sugahara K and Kawata S (2005):** NAD(P)H oxidase plays a crucial role in PDGF-induced proliferation of hepatic stellate cell. *Hepatology*; 41: 1272-1281.
- Adrian JE, Poelstra K, Scherphof GL, Molema G, Meijer DK, Reker-Smit C, Morselt HW and Kamps JA (2006):** Interaction of targeted liposomes with primary cultured hepatic stellate cells: involvement of multiple receptor systems. *J Hepatol*; 44: 560-567.
- Anan A, Baskin-Bey ES, Bronk SF, Werneburg NW, Shah VH and Gores GJ (2006):** Proteasome inhibition induces hepatic stellate cell apoptosis. *Hepatology*; 43: 335-344.
- Ankoma-Sey V, Matli M, Chang KB, Lalazar A, Donner DB, Wong L, Warren RS and Friedman SL (1998):** Coordinated induction of VEGF receptors in mesenchymal cell types during rat hepatic wound healing. *Oncogene*; 17: 115-121.
- Apte MV and Wilson JS (2003):** Stellate cell activation in alcoholic pancreatitis. *Pancreas*; 27: 316-320.

- Arey LB (1974):** Developmental Anatomy. 7th ed. Philadelphia, London. WB Saunders Co; P. 354.
- Asahina K, Sato H, Yamasaki C, Kataoka M, Shiokawa M, Katayama S, Tatenno C and Yoshizato K (2002):** Pleiotrophin/heparin-binding growth-associated molecule as a mitogen of rat hepatocytes and its role in regeneration and development of liver. Am J Pathol; 160: 2191-2205.
- Athari A, Hanecke K and Jungermann K (1994):** Prostaglandin F2 alpha and D2 release from primary Ito cell cultures after stimulation with noradrenaline and ATP but not adenosine. Hepatology; 20: 142-148.
- AW TY (1985):** Hepatic drug metabolism and drug induced liver injury. In: Gitnick G (ed). Current hepatology. New York, Wily, Vol. 5; P.113-196.
- Baba S, Fujii H, Hirose T, Yasuchika K, Azuma H, Hoppo T, Naito M, Machimoto T and Ikai I (2004):** Commitment of bone marrow cells to hepatic stellate cells in mouse. J Hepatol; 40: 255-260.
- Bachem MG, Meyer D, Melchior R, Sell KM and Gressner AM (1992):** Activation of rat liver perisinusoidal lipocytes by transforming growth factors derived from myofibroblast like cells. A potential mechanism of self perpetuation in liver fibrogenesis. J Clin Invest; 89: 19-27.

- Bachem MG, Zhou Z, Zhou S and Siech M (2006):** Role of stellate cells in pancreatic fibrogenesis associated with acute and chronic pancreatitis. *J Gastroenterol Hepatol*; 21 Suppl 3: S92-S96.
- Balian G, Click EM and Bornstein R (1980):** Location of collagen –binding domain in fibronectin. *J Biol Chem*; 255: 3234.
- Ballardini G, Groff P, Badiali dGL, Schuppan D and Bianchi FB (1994):** Ito cell heterogeneity: desmin-negative Ito cells in normal rat liver. *Hepatology*; 19: 440-446.
- Bartok I, Toth J, Remenar E and Virach SZ (1979):** Ultrastructure of the hepatic perisinusoidal cells in man and other mammalian species. *Anat Rec*; 194: 571.
- Bataller R, Sancho-Bru P, Gines P and Brenner DA (2005):** Liver fibrogenesis: a new role for the renin-angiotensin system. *Antioxid Redox Signal*; 7: 1346-1355.
- Bataller R, Sancho-Bru P, Gines P, Lora JM, Al-Garawi A, Sole M, Colmenero J, Nicolas JM, Jimenez W, Weich N, Gutierrez-Ramos JC, Arroyo V and Rodes J (2003):** Activated human hepatic stellate cells express the renin-angiotensin system and synthesize angiotensin II. *Gastroenterology*; 125: 117-125.

- Bataller R, Schwabe RF, Choi YH, Yang L, Paik YH, Lindquist J, Qian T, Schoonhoven R, Hagedorn CH, Lemasters JJ and Brenner DA (2003):** NADPH oxidase signal transduces angiotensin II in hepatic stellate cells and is critical in hepatic fibrosis. *J Clin Invest*; 112: 1383-1394.
- Batres RO and Olsen JA (1987):** A marginal vitamin A status alters the distribution of vitamin A among parenchymal and stellate cells in rat liver. *J Nutr*; 117: 874-879.
- Bauer M, Zhang JX and Bauer I (1994):** ET-1 induced alteration of hepatic microcirculation: sinusoidal and extrasinusoidal sites of action. *Am J Physiol*; 267: s143-s149.
- Beaussier M, Wendum D, Schiffer E, Dumont S, Rey C, Lienhart A and Housset C (2007):** Prominent contribution of portal mesenchymal cells to liver fibrosis in ischemic and obstructive cholestatic injuries. *Lab Invest*; 87: 292-303.
- Belfort R, Harrison SA, Brown K, Darland C, Finch J, Hardies J, Balas B, Gastaldelli A, Tio F, Pulcini J, Berria R, Ma JZ, Dwivedi S, Havranek R, Fincke C, DeFronzo R, Bannayan GA, Schenker S and Cusi K (2006):** A placebo-controlled trial of pioglitazone in subjects with nonalcoholic steatohepatitis. *N Engl J Med*; 355: 2297-2307.
- Bennet G (1984):** Role of Golgi complex in the secretory process. In: *Cell biology of the secretory process*; P. 102.

- Benten D, Kumaran V, Joseph B, Schattenberg J, Popov Y, Schuppan D and Gupta S (2005):** Hepatocyte transplantation activates hepatic stellate cells with beneficial modulation of cell engraftment in the rat. *Hepatology*; 42: 1072-1081.
- Benyon RC and Arthur MJ (2001):** Extracellular matrix degradation and the role of hepatic stellate cells. *Semin Liver Dis*; 21: 373-384.
- Bicker JJ (2001):** Kruppel-like factors: three fingers in many pies. *J Biol Chem*; 276: 34355-34358.
- Bilzer M, Roggel F and Gerbes AL (2006):** Role of Kupffer cells in host defense and liver disease. *Liver Int*; 26: 1175-1186.
- Blomhoff R, Green MH, Berg T and Norum KR (1990):** Transport and storage of vitamin A. *science*; 250: 399-404.
- Blomhoff R, Helgerud P, Rasmussen M, Berg T and Norum KR (1982):** In vivo uptake of chylomicron [³H] retinyl ester by rat liver: evidence for retinol transfer from parenchymal to nonparenchymal cells. *Proc Natl Acad Sci USA*; 79: 7326-7330.
- Bonacchi A, Romagnani P, Romanelli RG, Efsen E, Annunziato F, Lasagni L, Francalanci M, Laffi G, Pinzani M, Gentilini P and Marra F (2001):** Signal transduction by the chemokine receptor CXCR3: activation of Ras/ERK, Src, phosphatidylinositol 3-kinase/Akt controls cell migration and proliferation in human vascular pericytes. *J Biol Chem*; 276: 9945-9954.

- Breitkopf K, Godoy P, Ciucan L, Singer MV and Dooley S (2006):** TGF-beta/Smad signaling in the injured liver. *Z Gastroenterol*; 44: 57-66.
- Brigstock DR (2003):** The CCN family: a new stimulus package. *J endocrinol* 178: 169-175.
- Bronfenmajer S, Schaffner F and Popper H (1966):** Fat-storing cells (lipocytes) in human liver. *Arch Path*; 82: 447-453.
- Brun P, Castagliuolo I, Pinzani M, Palu G and Martines D (2005):** Exposure to bacterial cell wall products triggers an inflammatory phenotype in hepatic stellate cells. *Am J Physiol Gastrointest Liver Physiol* 289: G571-G578.
- Bruni C and Porter KR (1965):** The fine structure of the parenchymal cell of the normal rat liver. General observation. *Am J Path*; 46 (5): 691-700.
- Buchholz M, Kestler HA, Holzmann K, Ellenrieder V, Schneiderhan W, Siech M, Adler G, Bachem MG and Gress TM (2005):** Transcriptome analysis of human hepatic and pancreatic stellate cells: organ-specific variations of a common transcriptional phenotype. *J Mol Med*; 83(10): 795-805.
- Buckley JK (1964):** Phase contrast observations on the endoplasmic reticulum of living cells in culture. *Am J Path*; 46: 19-24.

- Burkel WE (1970):** The fine structure of the terminal branches of the hepatic arterial system of the rat. *Anat Rec*; 167: 329-350.
- Burkel WE and Low FN (1966):** The fine structure of rat liver sinusoids, space of Disse and associated tissue space. *Am J Anat*; 118: 769-773.
- Burt AD, Le Bail B and Balabaud C (1993):** Morphological investigation of sinusoidal cells. *Semin Liver Dis*; 13: 21-38.
- Canbay A, Feldstein AE, Baskin-Bey E, Bronk SF and Gores GJ (2004):** The caspase inhibitor IDN-6556 attenuates hepatic injury and fibrosis in the bile duct ligated mouse. *J Pharmacol Exp Ther* 308: 1191-1196.
- Canbay A, Friedman S and Gores GJ (2004):** Apoptosis: the nexus of liver injury and fibrosis. *Hepatology* 39:273-278.
- Canbay A, Higuchi H, Bronk SF, Taniai M, Sebo TJ and Gores GJ (2002):** Fas enhances fibrogenesis in the bile duct ligated mouse: a link between apoptosis and fibrosis. *Gastroenterology* 123: 1323-1330.
- Canby A, Taimr P, Torok N, Higuchi H, Friedman S and Gores GJ (2003):** Apoptotic body engulfment by a human stellate cell line is profibrogenic. *Lab Invest* 83: 655-663.
- Cardell RR (1977):** Smooth endoplasmic reticulum in rat hepatocytes during glycogen deposition and depletion. *Int Rev Cytol*; 48: 221-225.

- Carloni V, Defranco RM, Caligiuri A, Gentilini A, Sciammetta SC, Baldi E, Lottini B, Gentilini P and Pinzani M (2002):** Cell adhesion regulates platelets-derived growth factor-induced MAP kinase and PI-3 kinase activation in stellate cells. *Hepatology*; 36: 582-591.
- Carruthers JS and Steiner JW (1962):** Fine structure of terminal branches of the biliary tree. *Arch of Pathology*; 74: 230-243.
- Casini A, Ceni E and Salzano R (1997):** Neutrophil-derived Superoxide anion induces lipid peroxidation and stimulates collagen synthesis in human hepatic stellate cells: role of nitric oxide. *Hepatology*; 25: 361-367.
- Cassiman D, Barlow A, Vender Borgh S, Libbrecht L and Pachnis V (2006):** Hepatic stellate cells do not derive from the neural crest. *J Hepatol*; 44(6): 1098-1104.
- Cassiman D, Denef C, Desmet VJ and Roskams T (2001):** Human and rat hepatic stellate cells express neurotrophins and neurotrophin receptors. *Hepatology*; 33: 148-158.
- Cederbaum AI (2006):** Cytochrome P450 2E1-dependent oxidant stress and upregulation of anti-oxidant defense in liver cells. *J Gastroenterol Hepatol* 21 Suppl; 3: S22-S25.
- Chan ES, Montesinos MC, Fernandez P, Desai A, Delano DL, Yee H, Reiss AB, Pillinger MH, Chen JF, Schwarzschild MA and Friedman SL (2006):** Adenosine A (2A) receptors play a role in the pathogenesis of hepatic cirrhosis. *Br J Pharmacol*; 148: 1144-1155.

- Charo IF and Ransohoff RM (2006):** The many roles of chemokines and chemokine receptors in inflammation. *N Engl J MED*; 354: 610-621.
- Choi SS, Sicklick JK, Ma Q, Yang L, Huang J, Qi Y, Chen W, Li yx, Goldschmidt-Clermont PJ and Diehl AM (2006):** Sustained activation of Rac1 in hepatic stellate cells promotes liver injury and fibrosis in mice. *Hepatology*; 44: 1267-1277.
- Clemens MG and Zhang JX (1999):** Regulation of sinusoidal perfusion: In vivo methodology and control by endothelins. *Semin Liver Dis*; 19: 383-396.
- Cormack DH (1987):** Ham's Histology. 9th ed. Philadelphia, London, Mexico City, New York: JB Lippincott Co; 525-539.
- Corpechot C, Barbu V, Wendum D, Kinnman N, Rey C, Poupon R, Housset C and Rosmorduc O (2002):** Hypoxia-induced VEGF and collagen I expression are associated with angiogenesis and fibrogenesis in experimental cirrhosis. *Hepatology*; 35: 1010-1021.
- Czochra P, Klopčič B, Meyer E, Herkel J, Garcia-Lazaro JF, Thieringer F, Schirmacher P, Biesterfeld S, Galle PR, Lohse AW and Kanzler S (2006):** Liver fibrosis induced by hepatic overexpression of PDGF-B in transgenic mice. *J Hepatol*; 45: 419-428.

- Davis BH, Kramer RT and Davidson NO (1990):** Retinoic acid modulates rat Ito cell proliferation, collagen, transforming growth factor beta production. *J Clin Invest*; 86: 2062-2070.
- De Bleser PJ, Xu G, Rombouts K, Rogiers V and Geerts A (1999):** Glutathione levels discriminate between oxidative stress and transforming growth factor-beta signaling in activated rat hepatic stellate cells. *J Biol Chem*; 274: 33881-33887.
- De Duve C and Wittianx R (1966):** Functions of lysosomes. *Ann Rev of Phys*; 28:435-440.
- DeLeve LD (1998):** Glutathione defense in non-parenchymal cells. *Semin Liver Dis*; 18: 403-413.
- Demuth DG and Molleman A (2006):** Cannabinoid signaling. *Life Sci*; 78: 549-563.
- Depierre JW and Danller G (1975):** structural aspects of the membrane of endoplasmic reticulum. *Biochem Biophys Act*; 411-415.
- Di Sario A, Bendia E, Taffetani S, Marzioni M, Candelaresi C, Pigni P, Schindler U, Kleemann HW, Trozzi L, Macarri G and Benedetti A (2003):** Selective Na⁺ /H⁺ exchange inhibition by cariporide reduces liver fibrosis in the rat. *Hepatology*; 37: 256-266.

- Ding X, Saxena NK, Lin S, Xu A, Srinivasan S and Anania FA (2005):** The roles of leptin and adiponectin: a novel paradigm in adipocytokine regulation of liver fibrosis and stellate cell biology. *Am J Pathol*; 166: 1655-1669.
- Dunphy WG and Rothman JE (1985):** Compartmental organization of the Golgi stalk cell. *J Cell Biol*; 42: 13-15.
- Ekataksin W and Kaneda K (1999):** Liver microvascular architecture: an insight into the pathophysiology of portal hypertension. *Semin Liver Dis*; 19: 359-382.
- Elias J and Sherrick JC (1969):** Morphology of the liver. Academic press, New York; 35-70.
- Elrick LJ, Leel V, Blaylock Mg, Duncan L, Drever MR, Strachan G, Charlton KA, Kourth M, Porter AJ and Wright MC (2005):** Generation of a monoclonal human single chain antibody fragment to hepatic stellate cells: a potential mechanism for targeting liver anti-fibrotic therapeutics. *J Hepatol*; 42: 888-896.
- Elsharkawy AM, Oakley F and Mann DA (2005):** The role and regulation of hepatic stellate cell apoptosis in reversal of liver fibrosis. *Apoptosis*; 10: 927-939.
- Eng FJ and Friedman SL (2001):** Transcriptional regulation in hepatic stellate cells. *Semin Liver Dis*; 21: 385-395.

- Enzan H, Hayashi Y, Miyazaki E, Naruse K, Tao R, Kuroda N, Nakayama H, Kiyoku H, Hiroi M and Saibara T (1999):** Morphological aspects of hepatic fibrosis and Ito cells (hepatic stellate cells), with special reference to their myofibroblastic transformation. In: liver Diseases and Hepatic Sinusoidal Cells, edited by Tanikawa K, Ueno T. Tokyo: Springer-Verlag; 219-231.
- Evans RM, Barish GD and Wang YX (2004):** PPARs and the complex journey to obesity. *Nat Med*; 10: 355-361.
- Faouzi S, Lepreux S, Bedin C, Dubuisson L, Balabaud C, Bioulac-Sage P, Desmouliere A and Rosenbaum J (1999):** Activation of cultured rat hepatic stellate cells by tumoral hepatocytes. *Lab Invest*; 79: 485-493.
- Fauci AS, Mavilio D and Kottlil S (2005):** NK cells in HIV infection: paradigm for protection or targets for ambush. *Nat Rev Immunol*; 5: 835-843.
- Fawcett DW (1994):** Bloom and Fawcett, Text of Histology. 12th ed. Philadelphia, London, Toronto: WB Saunders Co; P. 679.
- Fiorucci S, Antonelli E, Distrutti E, Severino B, Fiorentina R, Baldoni M, Caliendo G, Santagada V and Morelli A (2004):** PAR1 antagonism protects against experimental liver fibrosis. Role of proteinase receptors in stellate cell activation. *Hepatology*; 39: 365-375.

- Fiorucci S, Rizzo G, Antonelli E, Renga B, Mencarelli a, Riccardi L, Orlandi S, Pruzanski M, Morelli A and Pellicciari R (2005):** A farnesoid X receptor-small heterodimer partner regulatory cascade modulates tissue metalloproteinase inhibitor-1 and matrix metalloprotease expression in hepatic stellate cells and promotes resolution of liver fibrosis. *J Pharmacol Exp Ther*; 314: 584-595.
- Fischer R, Cariers A, Reinehr R and Haussinger D (2002):** Caspase 9-de-pendent killing of hepatic stellate cells by activated Kupffer cells. *Gastroenterology*; 123: 845-861.
- Flisiak R, Baraona E, Li J and Lieber CS (1993):** Effects of ethanol on prostanoid production by liver fat-storing cells. *Hepatology*; 18:153-159.
- Foucher J, Chanteloup E, Vergniol J, Castera L, Le Bail B, Adhoute X, Bertet J, Couzigou P and de Ledinghen V (2006):** Diagnosis of cirrhosis by transient elastography (FibroScan): a prospective study. *Gut*; 55: 403-408.
- Friedman G, Liu LM, Friedman SL and Boyles JK (1991):** Apolipoprotein E is secreted by cultured lipocytes of the rat liver. *J Lipid Res*; 32:107-114.
- Friedman SL (2008):** Hepatic Stellate Cells: Protean, Multifunctional, and Enigmatic Cells of the Liver. *Physiol Rev*; 88: 125-172.
- Freidman SL and Arthur MJ (2002):** Reversing hepatic fibrosis. *Science Medicine*; 8: 194-205.

Friedman SL, Rockey DC, McGuire RF, Maher JJ, Boyles JK and Yamasaki G (1992): Isolated hepatic lipocytes and Kupffer cells from normal human liver: morphological and functional characteristics in primary culture. *Hepatology*; 15: 234-243.

Friedman SL and Roll FJ (1987): Isolation and culture of hepatic lipocytes, Kupffer cells, sinusoidal endothelial cells by density gradient centrifugation with Stractan. *Anal Biochem*; 161: 207-218.

Friedman SL, Wei S and Blaner WS (1993): Retinol release by activated rat hepatic lipocytes: regulation by Kupffer cell-conditioned medium and PDGF. *Am J Physiol Gastrointest Liver Physiol*; 264: G947-G952.

Friedman SL, Yamasaki G and Wong L (1994): Modulation of transforming growth factor beta receptors of rat lipocytes during the hepatic wound healing response. Enhanced binding and reduced gene expression accompany cellular activation in culture and in vivo. *J Biol Chem*; 269: 10551-10558.

Friend DS and Gilula (1972): Variations in tight and gap junction in mammalian tissues. *J Cell Biol*; 53: 758-765.

Fujita T, Maesawa C, Oikawa K, Nitta H, Wakabayashi G and Masuda T (2006): Interferon-gamma down-regulates expression of tumor necrosis factor-alpha converting enzyme/a disintegrin and metalloproteinase 17 in activated hepatic stellate cells of rats. *Int J Mol Med*; 17: 605-616.

- Gaca MD, Zhou X, Issa R, Kiriella K, Iredale JP and Benyon RC (2003):** Basement membrane-like matrix inhibits proliferation and collagen synthesis by activated rat hepatic stellate cells: evidence for matrix-dependent deactivation of stellate cells. *Matrix Biol*; 22: 229-239.
- Galli A, Crabb D, Price D, Ceni E, Salzano R, Surrenti C and Casini A (2000):** Peroxisome proliferator-activated receptor gamma transcriptional regulation is involved in platelet-derived growth factor-induced proliferation of human hepatic stellate cells. *Hepatology*; 31: 101-108.
- Gao R and Brigstock DR (2003):** Low density lipoprotein receptor-related protein (LRP) is a heparin-dependent adhesion receptor for connective tissue growth factor (CTGF) in rat activated hepatic stellate cells. *Hepatol Res*; 27: 214-220.
- Gascon-Barre M, Demers C, Mirshahi A, Neron S, Zalzal S and Nanci A (2003):** The normal liver harbors the vitamin D nuclear receptor in nonparenchymal and biliary epithelial cells. *Hepatology*; 37: 1034-1042.
- Gasull X, Bataller R, Gines P, Sancho-Bru P, Nicolas JM, Gorbis MN, Ferrer E, Badia E, Gual A, Arroyo V and Rodes J (2001):** Human myofibroblastic hepatic stellate cells express Ca (+) - activated K (+) channels that modulate the effects of endothelin-1 and nitric oxide. *J Hepatol*; 35(6): 739-748.

- Geerts A (2001):** History, heterogeneity, developmental biology, functions of quiescent hepatic stellate cells. *Semin Liver Dis*; 21: 311-335.
- Geerts A (2004):** On the origin of stellate cells: mesodermal, endodermal or neuro-ectodermal? *J Hepatol*; 40: 331-334.
- Geerts A, Bouwens L and Wisse E (1990):** Ultrastructure and function of hepatic fat-storing and pit cells. *J Electr Micr Tech*; 14: 247-256.
- Gelfand EV and Cannon CP (2006):** Rimonabant: a selective blocker of the cannabinoid CB1 receptors for the management of obesity, smoking cessation and cardiometabolic risk factors. *Expert Opin Invest Drugs*; 15: 307-315.
- Gillibert-Duplantier J, Neaud V, Blanc JF, Bioulac-Sage P and Rosenbaum J (2007):** Thrombin inhibits migration of human hepatic myofibroblasts. *Am J Physiol Gastrointest Liver Physiol*; 293: G128-G136.
- Goddard CJ, Smith A, Hoyland JA, Baird P, McMahon RF, Freemont AJ, Shomaf M, Haboubi NY and Warnes TW (1998):** Localisation and semiquantitative assessment of hepatic procollagen mRNA in primary biliary cirrhosis. *Gut*; 43: 433-440.
- Goodenough DA and Revel JP (1970):** A fine structural analysis of the intercellular junction in the mouse liver. *J Cell Biol*; 45: 272-275.

- Greenwald RJ, Freeman GJ and Sharpe AH (2005):** The B7 family revisited. *Annu Rev Immunol*; 23: 515-548.
- Greenway CV and Stark RD (1971):** Hepatic vascular bed. *Physiol Rev*; 51: 23-30.
- Greenwel P, Rubin J, Schwartz M, Hertzberg EL and Rojkind M (1993):** Liver fat-storing cell clones obtained from a CCl₄-cirrhotic rat are heterogeneous with regard to proliferation, expression of extracellular matrix components, interleukin-6, and connexin 43. *Lab Invest*; 69: 210-216.
- Gressner AM (1995):** Cytokines and cellular crosstalk involved in the activation of fat-storing cells. *J Hepatol*; 22: 28-36.
- Gressner AM, Lahme B, Meurer SK, Gressner O and Weiskirchen R (2006):** Variable expression of cystatin C in cultured trans-differentiating rat hepatic stellate cells. *World J Gastroenterol*; 12: 731-738.
- Gressner AM, Weiskirchen R, Breitkopf K and Dooley S (2002):** Roles of TGF-beta in hepatic fibrosis. *Front Biosci*; 7: D793-807.
- Gressner O, Lahme B, Gressner AM, Demirci I and Weiskirchen W (2007):** Differential effects of TGF- α on connective tissue growth factor expression in hepatic stellate cells and hepatocytes. *J Hepatol*; 47: 699-710.
- Griffith LG and Swartz MA (2006):** Capturing complex 3D tissue physiology in vitro. *Nat Rev Mol Cell Biol*; 7: 211-224.

- Grompe M (2005):** The origin of hepatocytes. *Gastroenterology*; 128: 2158-2160.
- Gupta SC, Gupta CD and Gupta SB (1981):** Hepatovenous segments in the human liver. *J Anat*; 133: 1-6.
- Gutierrez-Ruiz MC and Gomez-Quiroz LE (2007):** Liver fibrosis: searching for cell model answers. *Liver Int*; 27: 434-439.
- Guyot C, Combe C, Balabaud C, Bioulac-Sage P and Desmouliere A (2007):** Fibrogenic cell fate during fibrotic tissue remodeling observed in rat and human cultured liver slices. *J Hepatol*; 46: 142-150.
- Han YP (2006):** Matrix metalloproteinases, the pros and cons, in liver fibrosis. *J Gastroenterol Hepatol* 21 Suppl; 3: S88-S91.
- Han YP, Zhou L, Wang J, Xiong S, Garner WL, French SW and Tsukamoto H (2004):** Essential role of matrix metalloproteinases in interleukin-1-induced myofibroblastic activation of hepatic stellate cell in collagen. *J Biol Chem*; 279: 4820-4828.
- Hashmi AZ, Hakim W, Kruglov EA, Watanabe A, Watkins W, Dranoff JA and Mehal WZ (2007):** Adenosine inhibits cytosolic calcium signals and chemotaxis in hepatic stellate cells. *Am J Physiol Gastrointest Liver Physiol*; 292: G395-G401.

- Hautekeete ML, Dodeman I, Azais-Braesco V, Van den Berg K, Seynaeve C and Geerts A (1998):** Hepatic stellate cells and liver retinoid content in alcoholic liver disease in humans. *Alcohol Clin Exp Res*; 22: 494-500.
- Hehlhans S, Haase M and Cordes N (2007):** Signaling via integrins: implications for cell survival and anticancer strategies. *Biochem Biophys Acta*; 1775: 163-180.
- Henderson NC, Mackinnon AC, Farnworth SL, Poirier F, Russo FP, Iredale JP, Haslett C, Simpson KJ and Sethi T (2006):** Galectin-3 regulates myofibroblast activation and hepatic fibrosis. *Proc Natl Acad Sci USA*; 103: 5060-5065.
- Hertzberg EL, Lawrence TS and Gilula NB (1981):** Gap junction communication. *Ann Rev Physiol*; 43: 479-483.
- Hines JE, Johnson SJ and Burt AD (1993):** In vivo responses of macrophages and perisinusoidal cells to cholestatic liver injury. *Am J Pathol*; 142: 511-518.
- Hudnall SD (1991)** Cyclosporin A renders target cells resistant to immune cytolysis. *Eur J Immunol*; 21: 221-226.
- Inagaki Y and Okazaki I (2007):** Emerging insights into transforming growth factor-beta Smad signal in hepatic fibrogenesis. *Gut*; 56: 284-292.
- Iredale JP (2001):** Hepatic stellate cell behavior during resolution of liver injury. *Semin Liver Dis*; 21: 427-436.

- Ito T (1951):** Cytological studies on stellate cells of Kupffer and fat storing cells in the capillary wall of human liver. *Acta Anat Nippon*; 26: 2.
- Iwamoto H, Sakai H, Kotoh K, Nakamuta M and Nawata H (1999):** Soluble Arg-Gly-Asp peptides reduce collagen accumulation in cultured rat hepatic stellate cells. *Dig Dis Sci*; 44: 1038-1045.
- Jarnagin WR, Rockey DC, Koteliensky VE, Wang SS and Bissell DM (1994):** Expression of variant fibronectins in wound healing: cellular source and biological activity of the EIIIA segment in rat hepatic fibrogenesis. *J Cell Biol*; 127: 2037-2048.
- Jezequel AM, Novelli G, Venturini C and Orlandi F (1984):** Quantitative analysis of the perisinusoidal cells in human liver; the lipocytes. *Front Gastrointestinal Res*; 8: 85-90.
- Jochum W, Passegue E and Wagner EF (2001):** AP-1 in mouse development and tumorigenesis. *Oncogene*; 20: 2401-2412.
- Jones AL and Mills ES (1984):** The liver and Gall Bladder. In: *Modern Concept of Gastrointestinal Histology*. Elsevier. New York. Amsterdam. Oxford; P. 708.
- Jones AL, Schmucker DL, Renston RL and Murakami T (1980):** The architecture of bile secretion. *Digest Dis Sci*; 25: 609-613.

- Junqueira LC, Carneiro J and Kelley RO (1992):** Basic Histology. 7th ed. Los Altos. California: Lange Medical Publications; P. 321-350.
- Kalinichenko VV, Bhattacharyya D, Zhou Y, Gusarova GA, Kim W, Shin B and Costa RH (2003):** Foxf1 +/- mice exhibit defective stellate cell activation and abnormal liver regeneration following CCl₄ injury. *Hepatology*; 37: 107-117.
- Kalluri R and Neilson EG (2003):** Epithelial-mesenchymal transition and its implications for fibrosis. *J Clin Invest*; 112: 1776-1784.
- Kaneda K, Ekataksin W and Sogawa M (1998):** Endothelin-1-induced vasoconstriction causes a significant increase in portal pressure of rat liver: localized constrictive effect on distal segment of preterminal portal venules as revealed by light and electron microscopy and serial reconstruction. *Hepatology*; 27: 735-747.
- Kaplowitz N (1986):** Drug induced hepatotoxicity. *Ann Int Med*; 104: 826-828.
- Kawada N, Kristensen DB, Asahina K, Nakatani K, Minamiyama Y, Seki S and Yoshizato K (2001):** Characterization of a stellate cell activation-associated protein (STAP) with peroxidase activity found in rat hepatic stellate cells. *J Biol Chem*; 276: 25318-25323.

- Kawada N, Seki S, Kuroki T and Kaneda K (1999):** ROCK inhibitor Y-27632 attenuates stellate cell contraction and portal pressure increase induced by endothelin-1. *Biochem Biophys Res Commun*; 266: 296-300.
- Kiassov AP, Van Eyken P, van Pelt JF, Depla E, Fevery J, Desmet VJ and Yap SH (1995):** Desmin expressing nonhematopoietic liver cells during rat liver development: an immunohistochemical and morphometric study. *Differentiation*; 59: 253-258.
- Kikuchi S, Griffin CT, Wang SS and Bissell DM (2005):** Role of CD44 in epithelial wound repair: migration of rat hepatic stellate cells utilizes hyaluronic acid and CD44v6. *J Biol Chem*; 280: 15398-15404.
- Kim KY, Rhim T, Choi I and Kim SS (2001):** N-acetylcysteine induces cell cycle arrest in hepatic stellate cells through its reducing activity. *J Biol Chem*; 276: 40591-40598.
- Kinoshita K, Iimuro Y, Fujimoto J, Inagaki Y, Namikawa K, Kiyama H, Nakajima Y, Otagawa K, Kawada N, Friedman SL and Ikeda K (2007):** Targeted and regulable expression of transgenes in hepatic stellate cells and myofibroblasts in culture and in vivo using an adenoviral Cre-LoxP system to antagonize hepatic fibrosis. *Gut*; 56: 396-404.

- Knittel T, Aurisch S, Neubauer K, Eichhorst S and Ramadori G (1996):** Cell-type-specific expression of neural cell adhesion molecule (N-CAM) in Ito cells of rat liver. Up-regulation during in vitro activation and in hepatic tissue repair. *J Am Pathol*; 149: 449-462.
- Knittel T, Dinter C, Kobold D, Neubauer K, Mehde M, Eichhorst S and Ramadori G (1999):** Expression and regulation of cell adhesion molecules by hepatic stellate cells (HSCs) of rat liver: involvement of HSC in recruitment of inflammatory cells during hepatic tissue repair. *Am J Pathol*; 154: 153-167.
- Kobayashi S, Seki S, Kawada N, Morikawa H, Nakatani K, Uyama N, Ikeda K, Nakajima Y, Arakawa T and Kaneda K (2003):** Apoptosis of T cells in the hepatic fibrotic tissue of the rat: a possible inducing role of hepatic myofibroblast-like cells. *Cell Tissue Res*; 311: 353-364.
- Koenig S, Probst I, Becker H and Krause P (2006)** Zonal hierarchy of differentiation markers and nestin expression during oval cell mediated rat liver regeneration. *Histochem Cell Biol*; 126(6): 723-734.
- Kordes C, Sawitza I, Muller-Marbach A, Ale-Agha N, Keitel V, Klonowski-Stumpe H and Haussinger D (2007):** CD133 (+) hepatic stellate cells are progenitor cells. *Biochem Biophys Res Commun*; 352:410-417.
- Krause WJ and Cutts JH (1994):** *Essentials of Histology*. Little, Brown and Company Boston, New York, Toronto, London; P. 472-476.

- Kruglov E, Correa P, Arora G, Yu J, Nathanson MH and Dranoff JA (2007):** Molecular basis for calcium signaling in hepatic stellate cells. *Am J Physiol Gastrointest Liver Physiol*; 292: G975-G982.
- Kruglov EA, Nathanson RA, Nguyen T and Dranoff JA (2006):** Secretion of MCP-1/CCL2 by bile duct epithelia induces myofibroblastic transdifferentiation of portal fibroblasts. *Am J Physiol Liver Gastrointest Liver Physiol Gastrointest Physiol*; 290: G765-771.
- Labrador JP, Azcoitia V, Tuckermann J, Lin C, Olasso E, Manes S, Bruckner K, Goergen JL, Lemke G, Yancopoulos G, Angel P, Martinez C and Klein R (2001):** The collagen receptor DDR2 regulates proliferation and its elimination leads to dwarfism. *EMBO Rep*; 2: 446-452.
- Lalazar A, Wong L, Yamasaki G and Friedman SL (1997):** Early genes induced in hepatic stellate cells during wound healing. *Gene*; 195: 235-243.
- Lee JS, Semela D, Iredale J and Shah VH (2007):** Sinusoidal remodeling and angiogenesis: a new function for the liver-specific pericyte? *Hepatology*; 45: 817-825.
- Leeson GR and Lesson TS (1981):** *Histology*. 4th ed. Philadelphia, London: WB Saunders Co; P. 383-398.
- Leeson TS, Lesson CR and Papro AA (1988):** *Text/Atlas of Histology*. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo: WB Saunders Co; P. 475-496.

- Lim YS, Lee HC and Lee HS (2007):** Switch of cadherin expression from E-to N-type during the activation of rat hepatic stellate cells. *Histochem Cell Biol*; 127: 149-160.
- Liu C, Gaca MD, Swenson ES, Vellucci VF, Reiss M and Wells RG (2003):** Smads 2 and 3 are differentially activated by transforming growth factor-beta (TGF-beta) in quiescent and activated hepatic stellate cells. Constitutive nuclear localization of Smads in activated cells is TGF-beta-independent. *J Biol Chem*; 278: 11721-11728.
- Liu X, Hu H and Yin JQ (2006):** Therapeutic strategies against TGF-beta signaling pathway in hepatic fibrosis. *Liver Int*; 26: 8-22.
- Loud AV (1968):** A quantitative stereological description of the ultrastructure of normal rat liver parenchymal cells. *J Cell Biol*; 37: 27-46.
- Maher JJ (1993):** Cell-specific expression of hepatocyte growth factor in liver. Upregulation in sinusoidal endothelial cells after carbon tetrachloride. *J Clin Invest*; 91: 2244-2252.
- Maher JJ (2001):** Interactions between hepatic stellate cells and the immune system. *Semin Liver Dis*; 21: 417-426.
- Maher JJ and Friedman SL (1993):** Parenchymal and nonparenchymal cell interactions in the liver. *Semin Liver Dis*; 13: 13-20.

- Maher JJ, Friedman SL, Roll FJ and Bissell DM (1988):** Immunolocalization of laminin in normal rat liver and biosynthesis of laminin by hepatic lipocytes in primary culture. *Gastroenterology*; 94: 1053-1062.
- Maher JJ, Lozier JS and Scott MK (1998):** Rat hepatic stellate cells produce cytokine-induced neutrophil chemoattractant in culture and in vivo. *Am J Physiol Gastrointest Liver Physiol*; 275: G847-G853.
- Mallat A, Preaux AM, Serradeil-Le Gal C, Raufaste D, Gallois C, Brenner DA, Bradham C, Maclouf J, Iourgenko V, Fouassier L, Dhumeaux D, Mavrier P and Lotersztajn S (1996):** Growth inhibitory properties of endothelin-1 in activated human hepatic stellate cells: a cyclic adenosine monophosphate-mediated pathway Inhibition of both extracellular signal-regulated kinase and c-Jun kinase and upregulation of endothelin B receptors. *J Clin Invest*; 98: 2771-2778.
- Marek CJ, Tucker SJ, Konstantinou DK, Elrick LJ, Haefner D, Sigalas C, Murray GI, Goodwin B and Wright MC (2005):** Pregnenolone-16 α - carbonitrile inhibits rodent liver fibrogenesis via PXR (pregnane X receptor)-dependent and PXR-independent mechanisms. *Biochem J*; 387: 601-608.
- Margolis RN, Cardel RR and Curnow RT (1979):** Association of glycogen synthetase phosphatase and phosphorylase phosphatase activities with membranes of hepatic smooth endoplasmic reticulum. *J Cell Biol*; 83: 148-150.

- Marra F (2002):** Chemokines in liver inflammation and fibrosis. *Fron Biosci*; 7: d1899-1914.
- Marra F, Aleffi S, Bertolani C, Petrai I and Vizzutti F (2005):** Adipokines and liver fibrosis. *Eur Rev Med Pharmacol Sci*; 9: 279-284.
- Marra F, Choudhury GG, Pinzani M and Abboud HE (1994):** Regulation of platelet-derived growth factor secretion and gene expression in human liver fat-storing cells. *Gastroenterology*; 107: 1110-1117.
- Marra F, Efsen E, Romanelli RG, Caligiuri A, Pastacaldi S, Batignani G, Bonacchi A, Caporale R, Laffi G, Pinzani M and Gentilini P (2000):** Ligands of peroxisome proliferator-activated receptor gamma modulate profibrogenic and proinflammatory actions in hepatic stellate cells. *Gastroenterology*; 119: 466-478.
- Marra F and Pinzani M (2002):** Role of hepatic stellate cells in the pathogenesis of portal hypertension. *Nefrologia* 22 Suppl; 5: 34-40.
- Marra F, Romanelli RG, Giannini C, Failli P, Pastacaldi S, Arrighi MC, Pinzani M, Laffi G, Montalto P and Gentilini P (1999):** Monocyte chemotactic protein-1 as a chemoattractant for human hepatic stellate cells. *Hepatology*; 29: 140-148.

- Marra F, Valente AJ, Pinzani M and Abboud HE (1993):** Cultured human liver fat-storing cells produce monocyte chemotactic protein-1. Regulation by proinflammatory cytokines. *J Clin Invest*; 92: 1674-1680.
- Marsden ER, Hu Z, Fujio K, Nakatsukasa H, Thorgeirsson SS and Evarts RP (1992):** Expression of acidic fibroblast growth factor in regenerating liver and during hepatic differentiation. *Lab Invest*; 67: 427-433.
- Mastumoto E and Hirosawa K (1983):** Some observations on the structure of liver sinus with special reference to the Vitamin A storing cell. *Am J Anat*; 167-193.
- Matsuoka M and Tsukamoto H (1990):** Stimulation of hepatic lipocyte collagen production by Kupffer cell-derived transforming growth factor beta: implication for a pathogenetic role in alcoholic liver fibrogenesis. *Hepatology*; 11: 599-605.
- Matsuura T, Nagamori S, Fujise K, Hasumura S, Homma S, Sujino H, Shimizu K, Niiya M, Kameda H and Hirosawa K (1989):** Retinol transport in cultured fat storing cells of rat liver. Quantitative analysis by anchored cell analysis and sorting system. *Lab Invest*; 61: 107-115.
- Maubach G, Lim MC, Zhang CY and Zhuo L (2006):** GFAP promoter directs lacZ expression specifically in a rat hepatic stellate cell line. *World J Gastroenterol*; 12: 723-730.
- McCuskey RS (2000):** Morphological mechanisms for regulating blood flow through hepatic sinusoids. *Liver*; 20: 3-7.

- McMinn RMH (1994):** Last's anatomy: Regional and Applied 9th ed. London: Churchill Livingstone; P. 343-351.
- Melhem A, Muhanna N, Bishara A, Alvarez CE, Ilan Y, Bishara T, Horani A, Nassar M, Friedman SL and Safadi R (2006):** Anti-fibrotic activity of NK cells in experimental liver injury through killing of activated HSC. *J Hepatol*; 45: 60-71.
- Melton AC, Datta A and Yee HF JR (2006):** [Ca²⁺] independent contractile force generation by rat hepatic stellate cells in response to endothelin-1. *Am J Physiol Gastrointest Liver Physiol*; 290: G7-G13.
- Melton AC and Yee HF (2007):** Hepatic stellate cell protrusions couple platelet-derived growth factor-BB to chemotaxis. *Hepatology*; 45: 1446-1453.
- Meyer DH, Bachem MG and Gressner AM (1990):** Modulation of hepatic lipocyte proteoglycan synthesis and proliferation by Kupffer cell-derived transforming growth factors type beta 1 and type alpha. *Biochem Biophys Res Commun*; 171: 1122-1129.
- Milani S, Herbst H, Schuppan D, Grappone C, Pellegrini G, Pinzani M, Casini A, Calabro A, Ciancio G and Stefanini F (1994):** Bur-roughs AK, Surrenti C. Differential expression of matrix-metalloproteinase-1 and -2 genes in normal and fibrotic human liver. *Am J Pathol*; 144: 528-537.

- Miller EJ and Matukas VT (1974):** Biosynthesis of collagen. The Biochemist's view. Fed Proc; 33: 1187-1190.
- Mitra SK (1966):** The terminal distribution of the hepatic artery with special reference to arterio-portal anastomosis. J Anat; 100: 651-654.
- Montesano R and Nicolescu P (1978):** Fenestrations in endothelium of rat liver sinusoids revisited by Freeze Fracture. Anat Rec; 100: 861-864.
- Moore KL and Persaud TVN (1993):** The developing clinically oriented embryology. 5th ed. Philadelphia, London, Toronto, Sydney: WB Saunders Co; P. 243.
- Moriwaki H, Blaner WS, Piantedosi R and Goodman DS (1988):** Effects of dietary retinoid and triglyceride on the lipid composition of rat liver stellate cells and stellate cell lipid droplets. J Lipid Res; 29: 1523-153.
- Murakami K, Abe T, Miyazawa M, Yamaguchi M, Masuda T, Matsuura T, Nagamori S, Takeuchi K, Abe K and Kyogoku M (1995):** Establishment of a new human cell line, LI90, exhibiting characteristics of hepatic Ito (fat-storing) cells. Lab Invest; 72: 731-739.

- Murphy FR, Issa R, Zhou X, Ratnarajah S, Nagase H, Arthur MJ, Benyon C and Iredale JP (2002):** Inhibition of apoptosis of activated hepatic stellate cells by tissue inhibitor of metalloproteinase-1 is mediated via effects on matrix metalloproteinase inhibition: implications for reversibility of liver fibrosis. *J Biol Chem*; 277: 11069-11076.
- Nagy NE, Holven KB, Roos N, Senoo H, Kojima N, Norum KR and Blomhoff R (1997):** Storage of vitamin A in extrahepatic stellate cells in normal rats. *J Lipid Res*; 38: 645-658.
- Nakane PK (1963):** Ito's fat storing cell of the mouse liver. *Anat Rec*; 145: 265-266.
- Nakata K, Leong GF and Brauer RW (1960):** Direct measurements of blood pressures in minute vessels of the liver. *Am J Physiol*; 199: 1181-1188.
- Nakatani K, Tanaka H, Ikeda K, Sakabe M, Kadoya H, Seki S, Kaneda K and Nakajima Y (2006):** Expression of NCAM in activated portal fibroblasts during regeneration of the rat liver after partial hepatectomy. *Arch Histol Cytol*; 69: 61-72.
- Natarajan SK, Thomas S, Ramachandran A, Pulimood AB and Balasubramanian KA (2005):** Retinoid metabolism during development of liver cirrhosis. *Arch Biochem Biophys*; 443: 93-100.

- Nieto N, Friedman SL and Cederbaum AI (2002):** Cytochrome P450 2E1-derived reactive oxygen species mediate paracrine stimulation of collagen I protein synthesis by hepatic stellate cells. *J Biol Chem*; 277: 9853-9864.
- Nieto N, Friedman SL and Cederbaum AI (2002):** Stimulation and proliferation of primary rat hepatic stellate cells by cytochrome P450 2E1-derived reactive oxygen species. *Hepatology*; 35: 62-73.
- Novikoff PM and Yam A (1978):** Site of lipoprotein particles in normal rat hepatocytes. *J Cell Biol*; 76: 1-5.
- Novo E, Marra F, Zamara E, Valfre di Bonzo L, Caligiuri A, Cannito S, Antonaci C, Colombatto S, Pinzani M and Parola M (2006):** Dose dependent and divergent effects of superoxide anion on cell death, proliferation, migration of activated human hepatic stellate cells. *Gut*; 55: 90-97.
- Oh SH, Cho KH, Yang BS and Roh YK (2006):** Natural compounds from Danshen suppress the activity of hepatic stellate cells. *Arch Pharm Res*; 29: 762-767.
- Ohata M, Lin M, Satre M and Tsukamoto H (1997):** Diminished retinoic acid signaling in hepatic stellate cells in cholestatic liver fibrosis. *Am J Physiol Gastrointest Liver Physiol*; 272: G589-G596.

- Okuno M, Sato T, Kitamoto T, Imai S, Kawada N, Suzuki Y, Yoshimura H, Moriwaki H, Onuki K, Masushige S, Muto Y, Friedman SL, Kato S and Kojima S (1999):** Increased 9, 13-di-cis-retinoic acid in rat hepatic fibrosis: implication for a potential link between retinoid loss and TGF-beta mediated fibrogenesis in vivo. *J Hepatol*; 30: 1073-1080.
- Olaso E, Salado C, Egilegor E, Gutierrez V, Santisteban A, Sancho-Bru P, Friedman SL and Vidal-Vanaclocha F (2003):** Proangiogenic role of tumor-activated hepatic stellate cells in experimental melanoma metastasis. *Hepatology*; 37: 674-685.
- Osada N, Mochida S, Inao M, Mashimo Y and Fujiwara K (2001):** Apoptosis in dissociation between DNA synthesis and cellular functions of activated hepatic stellate cells a study with carbon tetrachloride induced rat liver injury. *Biochem Biophys Res Commun*; 282(2): 524-528.
- Paik YH, Lee KS, Lee HJ, Yang KM, Lee SJ, Lee DK, Han KH, Chon CY, Lee SI, Moon YM and Brenner DA (2006):** Hepatic stellate cells primed with cytokines upregulate inflammation in response to peptidoglycan or lipoteichoic acid. *Lab Invest*; 86: 676-686.
- Paik YH, Schwabe RF, Bataller R, Russo MP, Jobin C and Brenner DA (2003):** Toll-like receptor 4 mediates inflammatory signaling by bacterial lipopolysaccharide in human hepatic stellate cells. *Hepatology*; 37: 1043-1055

- Pan Q, Li DG, Lu HM, Wang YQ, Zhang WZ and Xu QF (2005):** A new immortalized rat cell line, hepatic stellate cell-PQ, exhibiting characteristics of hepatic stellate cell. *Hepatobiliary Pancreat Dis Int*; 4: 281-284.
- Paradis V, Dargere D, Bonvoust F, Vidaud M, Segarini P and Bedossa P (2002):** Effects and regulation of connective tissue growth factor on hepatic stellate cells. *Lab Invest*; 82: 767-774.
- Paradis V, Perlemuter G, Bonvoust F, Dargere D, Parfait B, Vidaud M, Conti M, Huet S, Ba N, Buffet C and Bedossa P (2001):** High glucose and hyperinsulinemia stimulate connective tissue growth factor expression: a potential mechanism involved in progression to fibrosis in nonalcoholic steatohepatitis. *Hepatology*; 34: 738-744.
- Parola M, Leonarduzzi G, Robino G, Albano E, Poli G and Dianzani MU (1996):** On the role of lipid peroxidation in the pathogenesis of liver damage induced by long-standing cholestasis. *Free Radical Biol Med*; 20: 351-359.
- Parola M, Robino G, Bordone R, Leonarduzzi G, Casini A, Pinzani M, Neve E, Bellomo G, Dianzani MU, Ingelman-Sundberg M and Albano E (1997):** Detection of cytochrome P4503A (CYP3A) in human hepatic stellate cells. *Biochem Biophys Res Commun*; 238: 420-424.
- Passino MA, Adams RA, Sikorski SL and Akassoglou K (2007):** Regulation of hepatic stellate cell differentiation by the neurotrophin receptor p75NTR. *Science*; 315: 1853-1856.

- Patella S, Phillips DJ, Tchongue J, de Kretser DM and Sievert W (2006):** Follistatin attenuates early liver fibrosis: effects on hepatic stellate cell activation and hepatocyte apoptosis. *Am J Physiol Gastrointest Liver Physiol*; 290: G137-G144.
- Perri RE, Langer DA, Chatterjee S, Gibbons SJ, Gadgil J, Cao S, Farrugia G and Shah VH (2006):** Defects in cGMP-PKG pathway contribute to impaired NO-dependent responses in hepatic stellate cells upon activation. *Am J Physiol Gastrointest Liver Physiol*; 290: G535-G542.
- Pinzani M (2002):** PDGF and signal transduction in hepatic stellate cells. *Front Biosci*; 7: d1720-1726.
- Pinzani M, Carloni V, Marra F, Riccardi D, Laffi G and Gentilini P (1994):** Biosynthesis of platelet-activating factor and its 1O-acyl analogue by liver fat-storing cells. *Gastroenterology*; 106: 1301-1311.
- Pinzani M and Marra F (2001):** Cytokine receptors and signaling in hepatic stellate cells. *Semin Liver Dis*; 21: 397-416.
- Planaguma A, Claria J, Miquel R, Lopez-Parra M, Titos E, Masferrer JL, Arroyo V and Rodes J (2005):** The selective cyclooxygenase-2 inhibitor SC-236 reduces liver fibrosis by mechanisms involving non-parenchymal cell apoptosis and PPAR gamma activation. *FASEB J*; 19: 1120-1122.

- Polliack A, Gamliel H, Horshko C, Knook NL and Sleyster FC (1978):** Surface morphology and ultrastructure of isolated hepatic Kupffer and endothelial cells. *Biomedicine*; 29: 268-272.
- Poynard T, Mathurin P, Lai CL, Guyader D, Poupon R, Tainturier MH, Myers RP, Muntenau M, Ratziu V, Manns M, Vogel A, Capron F, Chedid A and Bedossa P (2003):** A comparison of fibrosis progression in chronic liver diseases. *J Hepatol*; 38: 257-265.
- Radaeva S, Sun R, Jaruga B, Nguyen VT, Tian Z and Gao B (2006):** Natural killer cells ameliorate liver fibrosis by killing activated stellate cells in NKG2D-dependent and tumor necrosis factor-related apoptosis-inducing ligand-dependent manners. *Gastroenterology* 130: 435-452.
- Ramm GA, Britton RS, O'Neill R and Bacon BR (1994):** Identification and characterization of a receptor for tissue ferritin on activated rat lipocytes. *J Clin Invest*; 94: 9-15.
- Rappaport AM (1973):** The microcirculatory hepatic unit. *Microvasc Res*; 6: 212-214.
- Reeves HL, Burt AD, Wood S and Day CP (1996):** Hepatic stellate cell activation occurs in the absence of hepatitis in alcoholic liver disease and correlates with the severity of steatosis. *J Hepatol*; 25: 677-683.

- Reimann T, Hempel U, Krautwald S, Axmann A, Scheibe R, Seidel D and Wenzel KW (1997):** Transforming growth factor-beta1 induces activation of Ras, Raf-1, MEK and MAPK in rat hepatic stellate cells. *FEBS Lett*; 403: 57-60.
- Reinehr RM, Kubitz R and Peters-Regehr T (1998):** Activation of rat hepatic stellate cells in culture is associated with increased sensitivity to endothelin1. *Hepatology*; 28: 1566-1577.
- Reynaert H, Thompson MG, Thomas T and Geerts A (2002):** Hepatic stellate cells: role in microcirculation and pathophysiology of portal hypertension. *Gut*; 50: 571-581.
- Rifaat and Scott LF (2002):** Hepatic fibrosis: Role of hepatic stellate cell activation. *J Biol Chem*; 275: 2247-2250.
- Rimola A, Londono MC, Guevara G, Bruguera M, Navasa M, Forns X, Garcia-Retortillo M, Garcia-Valdecasas JC and Rodes J (2004):** Beneficial effect of angiotensin-blocking agents on graft fibrosis in hepatitis C recurrence after liver transplantation. *Transplantation*; 78: 686-691.
- Rockey DC (2001):** Cellular pathophysiology of portal hypertension and prospects for management with gene therapy. *Clin Liver Dis*; 5: 851-865.
- Rockey DC (2001):** Hepatic blood flow regulation by stellate cells in normal and injured liver. *Semin Liver Dis*; 21: 337-350.
- Rockey DC (2003):** Vascular mediators in the injured liver. *Hepatology*; 37: 4-12.