

References

- Adams PC, Reboussin DM, Barton JC, et al (2005):** Hemochromatosis and iron-overload screening in a racially diverse population. *N Engl J Med*; 352(17): 76-78.
- Agarwal R, Peters TJ, Coombes RC, et al (2002):** Tamoxifen-related porphyria cutanea tarda. *Med Oncol*; 19(2):121-300.
- Alison MR, Poulson R, Forbes S, et al. (2002):** An introduction to stem cells. *J Pathol*; 134(19): 419–423.
- Alison MR, Poulson R, Jeffery R, et al. (2000):** Hepatocytes from non-hepatic adult stem cells. *Nature*; 406(93): 257-260.
- Alison MR, Vig P, Russo F, et al. (2004):** Hepatic stem cells: From inside and outside the liver?. *Cell*; 37(11): 1–21.
- Allen KJ (2003):** Organ donation and embryonic stem cell research are on the same ethical continuum. *N Engl J*; 148(54):383–392.
- Allen KJ, Cheah DY, Wright PFA, et al. (2004):** Liver cell transplantation leads to repopulation and functional correction in a mouse model of Wilson's disease. *J Gastroenterol Hepatol*; 19(11):1283– 90.
- Allen KJ, Lee XL, Cheah DY, et al. (2004):** The potential of bone marrow stem cells to correct liver dysfunction in a mouse model of Wilson's disease. *Cell*; 13(7–8):765– 73.

- Almeida-Porada G, Porada CD, Chamberlain J, et al (2004):** Formation of human hepatocytes by human hematopoietic stem cells in sheep. *Blood*; 104(112):2582–2590.
- Alvarez-Dolado M, Opardal R, Garcia-Verdugo JM, et al. (2003):** Fusion of bone-marrow-derived cells with purkinje neurons, cardiomyocytes and hepatocytes. *Nature*; 425 (61):968– 73.
- Ambrosino G, Varotto S, Strom SC, et al. (2005):** Isolated hepatocyte transplantation for Crigler–Najjar syndrome type 1. *Cell*; 14(34):151–157.
- Ambruso DR, McCabe ER, Anderson D, et al. (1985):** Infectious and bleeding complications in patients with glycogenosis Ib. *Science*; 139(7):691-7.
- Amit M and Itskovitz-Eldor J (2002):** Derivation and spontaneous differentiation of human embryonic stem cells. *Gut*; 15(3):225–232.
- Anderson DJ, Gage FH and Weissman IL (2001):** Can stem cells cross lineage boundaries?. *Nature*; 7(78): 393–395.
- Anjos-Afonso F, Siapati EK and Bonnet D (2004):** In vivo contribution of murine mesenchymal stem cells into multiple cell-types under minimal damage conditions. *J Cell Sci*; 117(23):5655–5664.
- Aoki T, Jin Z, Nishino N, et al. (2005):** Intrasplenic transplantation of encapsulated hepatocytes decreases mortality and improves liver functions in fulminant hepatic failure from 90% partial hepatectomy in rats. *Transplantation*; 79(44):783–790.

- Arahuetes RM, Sierra E, Codesal J, et al. (2001):** Optimization of the technique to isolate fetal hepatocytes, and assessment of their functionality by transplantation. *Life Sci*; 68(7): 763– 772.
- Arikura J, Inagaki M, Huiling X, et al. (2004):** Colonization of albumin-producing hepatocytes derived from transplanted F344 rat bone marrow cells in the liver of congenic Nagase’s analbuminemic rats. *J Hepatol*; 41(2):215– 221.
- Arroyo V, Gines P, Gerbes AL, et al. (2002):** Definition and diagnostic criteria of refractory ascites and hepatorenal syndrome in cirrhosis. *Hepatology*; 23(1): 164-76.
- Attaran M, Schneider A, Grote C, et al. (2004):** Regional and transient ischemia/reperfusion injury in the liver improves therapeutic efficacy of allogeneic intraportal hepatocyte transplantation in low-density lipoprotein receptor deficient Watanabe rabbits. *J Hepatol*; 41(2):837–844.
- Avital I, Feraresso C, Aoki T, et al. (2002):** Bone marrow-derived liver stem cell and mature hepatocyte engraftment in livers undergoing rejection. *Surgery*; 132(2):384– 390.
- Ballen KK (2005):** New trends in umbilical cord blood transplantation. *Blood*; 105(10): 86–92.
- Barberi T, Klivenyi P, Calingasan NY, et al. (2003):** Neural subtype specification of fertilization and nuclear transfer embryonic stem cells and application in parkinsonian mice. *Nat Biotechnol*; 21(10):1200 –1207.

Beerheide W, von Mach MA, Ringel M, et al. (2002): Downregulation of beta2-microglobulin in human cord blood somatic stem cells after transplantation into livers of SCID-mice: an escape mechanism of stem cells?. *Biochem Biophys Res Commun*; 294:1052–1063.

Bellentani S and Tiribelli C (2001): The spectrum of liver disease in the general population: Lessons from the Dionysos study. *J Hepatol*; 35: 531–537.

Bethesda N (2007): Stem cell up-to-date. *Hepatology*; 39(6):1477– 1487.

Beutler E, Felitti VJ, Koziol JA, et al. (2002): Penetrance of 845G--> A (C282Y) HFE hereditary haemochromatosis mutation in the USA. *Lancet I*; 359(92): 211-218.

Beutler E, Gelbart T and Miller W (2002): Severe jaundice in a patient with a previously undescribed glucose-6-phosphate dehydrogenase (G6PD) mutation and Gilbert syndrome. *Blood*; 28(2): 104-107.

Bianco P and Riminucci M (1998): The bone marrow stroma in vivo: Ontogeny, structure, cellular composition and changes in disease. In *Marrow Stromal Cell Culture. Handbooks in Practical Animal Cell Biology*, ed. J.N. Beresford and M.E. Owen, p. 1025. Cambridge, UK: Cambridge University Press. p. 244-289.

Bilir BM, Guinette D, Karrer F, et al. (2000): Hepatocyte transplantation in acute liver failure. *Liver Transplant*; 6:32–40.

Blanpain C, Lowry WE, Geohegan A, et al. (2004): Self renewal, multipotency, and the existence of two cell populations within an epithelial stem cell niche. *Cell* 118: 530–532.

Blei AT (2005): The pathophysiology of brain edema in acute liver failure. *Neurochem Int.*; 47(1-2):71-77.

Bodamer OA, Feillet F, Lane RE, et al. (2002): Utilization of cornstarch in glycogen storage disease type Ia. *Eur J Gastroenterol Hepatol.*; 14(11): 51-60.

Bongso A and Richards M (2004): History and perspective of stem cell research. In *Best Practice & Research Clinical Obstetrics & Gynaecology*, eds. N. Fisk & J. Itskovitz, London: Elsevier Ltd. p. 734-744.

Borlak J, Thum T, Landt O, et al. (2000): Molecular diagnosis of a familial nonhemolytic hyperbilirubinemia (Gilbert's syndrome) in healthy subjects. *Hepatology*; 32(4 Pt 1): 92-95.

BottaiD, Fiocco R, Gelain F, et al. (2003): Neural stem cells in the adult nervous system. *Stem Cell*; 12: 655–670.

Brewer GJ (2000): Recognition, diagnosis, and management of Wilson's disease. *Nature*; 223(1):39-46.

Bruix J, Sherman M, Llovet JM, et al. (2001): Clinical management of hepatocellular carcinoma. Conclusions of the Barcelona-2000 EASL conference. *J Hepatol*; 35: 421-430.

Burlina AB (2004): Hepatocyte transplantation for inborn errors of metabolism. *J Inherit Metab Dis.*; 27(1):373–383.

Cabrera-Abreu J, Crabtree NJ, Elias E, et al. (2004): Bone mineral density and markers of bone turnover in patients with glycogen storage disease types I, III and IX. *J Inherit Metab Dis.*; 27(1):1-9.

Cai J, Ito M, Nagata H, et al. (2002): Treatment of liver failure in rats with end-stage cirrhosis by transplantation of immortalized hepatocytes. *Hepatology*; 36:386–394.

Cai YF, Zhen ZJ, Min J, et al. (2004): Selection, proliferation and differentiation of bone marrow-derived liver stem cells with a culture system containing cholestatic serum in vitro. *J Gastroenterol*; 10(22): 8–12.

Cairo MS and Wagner JE (2000): Placental and/or umbilical cord blood: an alternative source of hematopoietic stem cells for transplantation. *Blood*; 90(12): 65–78.

Caldwell SH and Battle EH (1999): Ascites and spontaneous bacterial peritonitis. In: Schiff ER, Sorrell MF, Maddrey WC, eds. *Schiff's Diseases of the Liver*. 8th ed. Philadelphia, Pa: Lippincott-Raven. p.371-385.

Camargo FD, Finegold M and Goodell MA (2004): Hematopoietic myelomonocytic cells are the major source of hepatocyte fusion partners. *J Clin Invest.*; 113(9):1266– 1270.

- Campos MA, Wanner A and Zhang G (2005):** Trends in the diagnosis of symptomatic patients with alpha1-antitrypsin deficiency between 1968 and 2003. *Chest*; 128(3):1179-1186.
- Cantz T, Sharma AD, Jochheim-Richter A, et al. (2004):** Reevaluation of bone marrow-derived cells as a source for hepatocyte regeneration. *Cell*; 13(6): 659– 666.
- Cantz T, Zuckerman DM, Burda MR, et al. (2003):** Quantitative gene expression analysis reveals transition of fetal liver progenitor cells to mature hepatocytes after transplantation in uPA/RAG-2 mice. *Am J Pathol*; 162(1):37– 45.
- Carpenter MK, Rosler ES, Fisk GJ, et al. (2004):** Properties of four human embryonic stem cell lines maintained in a feeder-free culture system. *Science*; 229:243–258.
- Chan C, Berthiaume F, Nath BD, et al (2004):** Hepatic tissue engineering for adjunct and temporary liver support: critical technologies. *Liver Transplant*; 10:1331–1342.
- Chinzei R, Tanaka Y, Shimizu-Saito K, et al. (2002):** Embryoid-body cells derived from a mouse embryonic stem cell line show differentiation into functional hepatocytes. *Hepatology*; 36(1):22–29.
- Choi D, Oh HJ, Chang UJ, et al. (2002):** In vivo differentiation of mouse embryonic stem cells into hepatocytes. *Cell Transplant*; 11(4): 59–68.

- Chowdhury JR, Grossman M, Gupta S, et al (1991):** Long-term improvement of hypercholesterolemia after ex vivo gene therapy in LDLR-deficient rabbits. *Science*; 254:1802–1805.
- Clarke DL, Johansson CB, Wilbertz J, et al. (2000):** Generalized potential of adult stem cells. *Science* 288, 1660-1663.
- Costa E (2006):** Hematologically important mutations: bilirubin UDP-glucuronosyltransferase gene mutations in Gilbert and Crigler-Najjar syndromes. *Blood*; 36(1):77-80.
- Couchie D, Holic N, Chobert MN, et al. (2002):** In vitro differentiation of WB-F344 rat liver epithelial cells into the biliary lineage. *Nature*; 69:209–215.
- Cowan CA, Klimanskaya I, McMahon JP, et al. (2004):** Derivation of embryonic stem-cell lines from human blastocysts. *N Engl J Med* 350:1353–1356.
- Cutler C and Antin JH (2001):** Peripheral blood stem cells for allogeneic transplantation: a review. *Stem Cells*; 19(2):108-117.
- Cutler JA, Patel R, Mitchell MJ, et al. (2005):** The significance of published polymorphisms in 14 cases of mild factor VII deficiency. *Blood*; 16(2):91-95.
- Dabeva MD, Petkov PM, Sandhu J, et al. (2000):** Proliferation and differentiation of fetal liver epithelial progenitor cells after transplantation into adult rat liver. *Am J Pathol*; 156(6): 2017–

2031.

D'Amico G, Pagliaro L and Bosch J (1995): The treatment of portal hypertension: a meta-analytic review. *Hepatology*; 22(1): 332-354.

Danet GH, Luongo JL, Butler G, et al. (2002): C1qRp defines a new human stem cell population with hematopoietic and hepatic potential. *Natl Acad Sci*; 99(16): 441–445.

Demetriou AA, Reisner A, Sanchez J, et al. (1988): Transplantation of microcarrier-attached hepatocytes into 90% partially hepatectomized rats. *Hepatology*; 8:1006–1009.

De Vree JM, Ottenhoff R, Bosma PJ, et al. (2000): Correction of liver disease by hepatocyte transplantation in a mouse model of progressive familial intrahepatic cholestasis. *Gastroenterology*; 119:1720–1730.

Dhawan A, Mitry RR, Hughes RD, et al. (2004): Hepatocyte transplantation for inherited factor VII deficiency. *Transplantation*; 78:1812–1814.

Donovan PJ and de Miguel MP (2003): Turning germ cells into stem cells. *Nature*; 13: 463–471.

Eiges R, Schuldiner M, Drukker M, et al (2001): Establishment of human embryonic stem cell-transduced clones carrying a marker of undifferentiated cells. *Curr. Biol.*; 11: 514-518.

Eisenberg LM and Eisenberg CA (2003): Stem cell plasticity, cell fusion, and transdifferentiation. *Birth Defect Res. Part. C. Embryo*

Today; 69: 209-218.

Evans MJ and Kaufman MH (1981): Establishment in culture of pluripotential cells from mouse embryos. *Nature*; 292(5819):154– 156.

Farheen S, Sengupta S and Santra A (2006): Gilbert's syndrome: High frequency of the (TA) 7 TAA allele in India and its interaction with a novel CAT insertion in promoter of the gene for bilirubin UDP-glucuronosyltransferase 1 gene. *J Gastroenterol*; 12(14): 269-275.

Fausto N (2004): Liver regeneration and repair: hepatocytes, progenitor cells, and stem cells. *Hepatology*; 39(6):1477–1487.

Fiegel HC, Park JJ, Lioznov MV, et al. (2003): Characterization of cell types during rat liver development. *Hepatology* 37:148–154.

Filip S, English D and Mokry J (2004): Issues in stem cell plasticity. *J. Cell. Mol. Med.* 8, 572-577.

Fisher RA, Bu D, Thompson M, et al. (2000): Defining hepatocellular chimerism in a liver failure patient bridged with hepatocyte infusion. *Transplantation*; 69:303–307.

Fox IJ, Chowdhury JR, Kaufman SS, et al. (1998): Treatment of the Crigler–Najjar syndrome type I with hepatocyte transplantation. *N Engl J Med*; 338:1422–1426.

Francavilla A, Starzl TE, Barone M, et al. (1999): Studies on mechanisms of augmentation of liver regeneration by cyclosporine and FK 506. *Hepatology*; 14:140–143.

Friedman SL (1999): Hepatic fibrosis. In: Schiff ER, Sorrell MF, Maddrey WC, eds. Schiff's Diseases of the Liver. 8th ed. Philadelphia, Pa: Lippincott-Raven. p. 371-385.

Fuentealba IC and Aburto EM (2003): Animal models of copper-associated liver disease. *Comp Hepatol*; 112(9):1407– 1418.

Fujikawa T, Hirose T, Fujii H, et al. (2003): Purification of adult hepatic progenitor cells using green fluorescent protein (GFP)-transgenic mice and fluorescence-activated cell sorting. *J Hepatol* 39:162–170.

Fujikawa T, Oh SH, Pi L, et al. (2005): Teratoma formation leads to failure of treatment for type I diabetes using embryonic stem cell-derived insulin-producing cells. *Am J Pathol*; 166(6):1781– 1791.

Gisbert JP, García-Buey L, Alonso A, et al. (2004): Hepatocellular carcinoma risk in patients with porphyria cutanea tarda. *Eur J Gastroenterol Hepatol.*; 16(7):689-692.

Goodell MA (2003): Stem-cell „plasticity“: befuddled by the muddle. *Curr. Opin. Hematol.*; 10: 208-213.

Gordon GJ, Butz GM, Grisham JW, et al. (2002): Isolation, short-term culture, and transplantation of small hepatocyte-like progenitor cells from retrorsine exposed rats. *Transplantation*; 73:1236– 1243.

Groszmann RJ and Atterbury CE (2002): The pathophysiology of portal

hypertension: a basis for classification. *Semin Liver Dis.*; 2(3): 177-186.

Grove JE, Bruscia E and Krause DS (2004): Plasticity of bone marrow-derived stem cells. *Stem Cells*; 22: 487–500.

Gupta R and Parashar Y (2004): Crigler-najjar syndrome type II. *Indian J Pediat.*; 71(11):1043-1055.

Hamazaki T, Iiboshi Y, Oka M, et al. (2001): Hepatic maturation in differentiating embryonic stem cells in vitro. *Science*; 497(1):15– 19.

Hao QL, Zhu J, Price MA, et al. (2001): Identification of a novel, human multilymphoid progenitor in cord blood. *Blood*; 97:3683–3690.

Harada M, Kawaguchi T, Kumemura H, et al. (2005): The Wilson disease protein ATP7B resides in the late endosomes with Rab7 and the Niemann-Pick C1 protein. *Am J Pathol.*; 166:499–510.

Harrison DE and Astle CM (2001): Short- and long-term multilineage repopulating hematopoietic stem cells in late fetal and newborn mice: models for human umbilical cord blood. *Blood*; 90(1):174– 181.

Hatch HM, Zheng D, Jorgensen ML, et al. (2002): SDF-1a/CXCR4: a mechanism for hepatic oval cell activation and bone marrow stem cell recruitment to the injured liver of rats. *Stem Cells*; 4(4): 339– 351.

Hawley R G and Sobieski D A (2002): Somatic stem cell plasticity: to be or not to be. *Stem Cells*; 44: 195-197.

Heng BC, Yu H, Yin Y, et al. (2005): Factors influencing stem cell differentiation into the hepatic lineage in vitro. *J Gastroenterol Hepatol*; 20(7):975– 987.

Hengstler JG, Brulport M, Schormann W, et al. (2005): Generation of human hepatocytes by stem cell technology: definition of the hepatocyte. *Nature*; 54:61–74.

Herrmann FH, Wulff K, Auberger K, et al. (2000): Molecular biology and clinical manifestation of hereditary factor VII deficiency. *Semin Thromb Hemost.*; 26(4):393-400.

Herzenberg LA and De Rosa SC (2000): Monoclonal antibodies and the FACS: complementary tools for immunobiology and medicine. *Immunol. Today*; 21, 383- 390.

Herzog EL, Chai L and Krause DS (2003): Plasticity of marrow-derived stem cells. *Blood*; 102: 483–493.

Himmelfmann A and Fehr J (1999): Cloning of the hereditary hemochromatosis gene: implications for pathogenesis, diagnosis, and screening. *J Lab Clin Med*; 133(3): 229-236.

Hoppo T, Fujii H, Hirose T, et al. (2004): Thy1-positive mesenchymal cells promote the maturation of CD49f-positive hepatic progenitor cells in the mouse fetal liver. *Hepatology* 39:1362–1370.

Hovatta O, Mikkola M, Ye Z, et al. (2003): A culture system using human foreskin fibroblasts as feeder cells allows production of human

embryonic stem cells. Hum Reprod; 18:1404–1449.

Hwang WS, Lee BC, Lee CK, et al. (2005): Human embryonic stem cells and therapeutic cloning. J Vet Sci; 6(2):87– 96.

Hwang WS, Roh SI, Lee BC, et al. (2005): Patientspecific embryonic stem cells derived from human blastocysts. Science; 308(5729):1777– 1783.

Hwang WS, Ryu YJ, Park JH, et al. (2004): Evidence of a pluripotent human embryonic stem cell line derived from a cloned blastocyst. Science; 303(5664):1669–1674.

Ishikawa F, Drake CJ, Yang S, et al. (2003): Transplanted human cord blood cells give rise to hepatocytes in engrafted mice. Gut; 96:174– 185.

Iwasaki T (2004): Recent advances in the treatment of graft-versus-host disease. Clin Med Res.; 2(4):243-252.

Jackson K, Majka SM, Wang H, et al. (2001): Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells. J. Clin. Invest.; 107: 1-8.

Jalan R (2005): Acute liver failure: current management and future prospects. J Hepatol.; 42 Suppl(1): 115-123

Jang YY, Collector MI, Baylin SB, et al (2004): Hematopoietic stem cells convert into liver cells within days without fusion. Nat Cell Biol; 6(6):532 –539.

Jiang Y, Jahagirdar BN, Reinhardt RL, et al. (2002): Pluripotency of

mesenchymal stem cells derived from adult bone marrow. Nature; 418: 41–49.

Jones EA, Tosh D, Wilson DI, et al. (2002): Hepatic differentiation of murine embryonic stem cells. Exp Cell Res; 272 (1):15 –22.

Kakinuma S, Tanaka Y, Chinzei R, et al. (2003): Human umbilical cord blood as a source of transplantable hepatic progenitor cells. Stem Cells; 21(2):217–227.

Kamath PS, Wiesner RH, Malinchoc M, et al. (2001): A model to predict survival in patients with end stage liver disease. Hepatology; 33:464-470.

Kanazawa Y and Verma IM (2003): Little evidence of bone marrow-derived hepatocytes in the replacement of injured liver. Proc Natl Acad Sci USA. p. 556-570.

Kang XQ, Zang WJ, Song TS, et al. (2005): Rat bone marrow mesenchymal stem cells differentiate into hepatocytes in vitro. World J Gastroenterol; 11(22):379–384.

Kawahara T, Kasai S, Yagita H, et al. (2001): Critical role of Fas/Fas ligand interaction in CD28-independent pathway of allogeneic murine hepatocyte rejection. Hepatology; 26(4):944–948.

Keeffe EB (1998): Summary of guidelines on organ allocation and patient listing for liver transplantation. Liver Transpl Surg; 4(suppl 1): 108-114.

Kiessling AA and Anderson SC (2003): Human embryonic stem cells. Boston: Jones and Bartlett. Stem Cells; 20 (2):146– 154.

Kinoshita T, Sekiguchi T, Xu Mj, et al. (2000): Hepatic differentiation

induced by oncostatin M attenuates fetal liver hematopoiesis. Proc Natl Acad Sci; 96(13): 65– 70.

Koeberl DD, Kishnani PS and Chen YT (2007): Glycogen storage disease types I and II: treatment updates. J Inherit Metab Dis.; 30(2):159-164.

Koenig S, Stoesser C, Krause P, et al. (2005): Liver repopulation after hepatocellular transplantation: integration and interaction of transplanted hepatocytes in the host. Cell Transplant; 14:31–40.

Kogler G, Sensken S, Airey JA, et al. (2004): A new human somatic stem cell from placental cord blood with intrinsic pluripotent differentiation potential. J Exp Med; 200(2):123– 135.

Kollet O, Shivtiel S, Chen YQ, et al. (2003): HGF, SDF-1, and MMP-9 are involved in stress-induced human CD34+ stem cell recruitment to the liver. J Clin Invest; 112(2): 160– 169.

Kondo M, Weiss ML, Wagers AJ, et al. (2003): Biology of hematopoietic stem cells and progenitors: Implications for clinical application. Annu Rev Immunol 21:759–806.

Krause DS, Theise ND, Collector MI, et al. (2001): Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell. Cell; 105(3):369– 377.

Kuai XL, Cong XQ, Li XL, et al. (2003): Generation of hepatocytes from cultured mouse embryonic stem cells. Liver Transpl; 9(10):1094–1099.

Kumashiro Y, Asahina K, Ozeki R, et al. (2005): Enrichment of hepatocytes differentiated from mouse embryonic stem cells as

a transplantable source. *Transplantation*; 79(5):550– 557

Laconi E and Laconi S (2002): Principles of hepatocyte repopulation. *Semin Cell Dev Biol*; 13:433–438.

Laconi S, Pillai S, Porcu PP, et al. (2001): Massive liver replacement by transplanted hepatocytes in the absence of exogenous growth stimuli in rats treated with retrorsine. *Am J Pathol*; 158:771–777.

Lagasse E, Connors H, Al-Dhalimy M, et al. (2000): Purified hematopoietic stem cells can differentiate into hepatocytes in vivo. *Nat Med*; 6(11): 229– 234.

Lambrecht RW and Bonkovsky HL (2002): Hemochromatosis and porphyria. *Semin Gastrointest Dis.*; 13(2):109-119.

Laurson J, Selden C and Hodgson HJ (2005): Hepatocyte progenitors in man and in rodents—multiple pathways, multiple candidates. *Int J Exp Pathol*; 86(1):1– 18.

Le Blanc K and Ringden O (2005): Immunobiology of human mesenchymal stem cells and future use in hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant*; 11(5):321–334.

Lee KD, Kuo TK, Whang-Peng J, et al. (2004): In vitro hepatic differentiation of human mesenchymal stem cells. *Hepatology*; 40(6):1275– 1284.

Lee OK, Kuo TK, Chen WM, et al. (2004): Isolation of multipotent mesenchymal stem cells from umbilical cord blood. *Blood*;

103(5):1669–1675.

Lowes KN, Croager EJ, Olynyk JK, et al. (2003): Oval cell-mediated liver regeneration: role of cytokines and growth factors. *J Gastroenterol Hepatol*; 18(1):4– 12.

Lucey MR, Brown KA, Everson GT, et al. (1997): Minimal criteria for placement of adults on the liver transplant waiting list: A report of a national conference organized by the American Society of Transplant Physicians and the American Association for the Study of Liver Diseases. *Liver Transpl Surg*; 3:628-637.

Malhi H, Irani AN, Gagandeep S, et al. (2002): Isolation of human progenitor liver epithelial cells with extensive replication capacity and differentiation into mature hepatocytes. *J Cell Sci*; 115(13): 2679– 2678.

Malinchoc M, Kamath PS, Gordon FD, et al. (2000): A model to predict poor survival in patients undergoing transjugular intrahepatic portosystemic shunts. *Hepatology*; 31:864-871.

Mallet VO, Mitchell C, Mezey E, et al. (2002): Bone marrow transplantation in mice leads to a minor population of hepatocytes that can be selectively amplified in vivo. *Hepatology*; 35(4):799–811.

Mariani G, Herrmann FH, Dolce A, et al. (2005): Clinical phenotypes and factor VII genotype in congenital factor VII deficiency. *Nature*; 93(3):481-487.

Martin AP, Bartels M, Schreiber S, et al. (2006): Successful staged

kidney and liver transplantation for glycogen storage disease type Ib: A case report. *Transplant*; 38(10): 615-619.

Merion RM (2004): When is a patient too well and when is a patient too sick for a liver transplant?. *Liver Transplant*; 10: 69-73.

Mitchell KE, Weiss ML, Mitchell BM, et al. (2003): Matrix cells from Wharton's jelly form neurons and glia. *Stem Cells*; 21: 50–60.

Muraca M, Gerunda G, Neri D, et al. (2002): Hepatocyte transplantation as a treatment for glycogen storage disease type 1a. *Lancet II*; 359:317–318.

Nagata H, Ito M, Shirota C, et al. (2003): Route of hepatocyte delivery affects hepatocyte engraftment in the spleen. *Transplant*; 76:732–734.

Nakagami M, Morimoto T, Itoh K, et al. (2000): Patterns of restoration of remnant liver volume after graft harvesting in donors for living related liver transplantation. *Transplant*; 30 (1):195– 199.

National center for health statistics (2002): Classification of diseases and disability. Elsevier Ltd. p.343-355.

Neuberger J. (2000): Liver transplantation. *J Hepatol*; 32:198–207.

Newsome PN, Johannessen I, Boyle S, et al. (2003): Human cord blood-derived cells can differentiate into hepatocytes in the mouse liver with no evidence of cellular fusion. *Gastroenterology*; 124(7):1891– 990.

Nguyen AD, Pan CJ, Weinstein DA, et al. (2006): Increased scavenger receptor class B type I-mediated cellular cholesterol efflux and

antioxidant capacity in the sera of glycogen storage disease type Ia patients. *Mol Genet Metab*; 89(3):233-238.

Nowak G, Ericzon BG, Nava S, et al. (2005): Identification of expandable human hepatic progenitors which differentiate into mature hepatic cells in vivo. *Gut*; 54(7): 972– 979.

Oertel M, Rosencrantz R, Chen YQ, et al. (2003): Repopulation of rat liver by fetal hepatoblasts and adult hepatocytes transduced ex vivo with lentiviral vectors. *Hepatology*; 37(5):994– 1005.

Oh SH, Miyazaki M, Kouchi H, et al. (2000): Hepatocyte growth factor induces differentiation of adult rat bone marrow cells into a hepatocyte lineage in vitro. *Science*; 279(2):500– 504.

Okie S (2005): Stem-cell research—signposts and roadblocks. *N Engl J Med*; 353(1):1– 5.

Okumoto K, Saito T, Hattori E, et al. (2005): Differentiation of rat bone marrow cells cultured on artificial basement membrane containing extracellular matrix into a liver cell lineage. *J Hepatol*; 43(1):110– 116.

Overturf K, al-Dhalimy M, Ou C, et al. (2001): Serial transplantation reveals the stem-cell-like regenerative potential of adult mouse hepatocytes. *Am J Pathol*; 151(5):1273– 1280.

Parviz F, Matullo C, Garrison WD, et al. (2003): Hepatocyte nuclear factor 4alpha controls the development of a hepatic epithelium and liver morphogenesis. *Nat Genet*; 34(3): 292– 296.

Penvy L and Rao S (2003): The stem cell menagerie. *Trends Neurosci*; 26: 351–359.

- Pera MF (2001):** Scientific considerations relating to the ethics of the use of human embryonic stem cells in research and medicine. *Reprod Fertil Dev.*; 13(1):23– 29.
- Pera MF, Reubinoff B and Trounson A (2000):** Human embryonic stem cells. *J Cell Sci*; 113 (Pt 1): 5–10.
- Pera MF and Trounson AO (2004):** Human embryonic stem cells: Prospects for development. *Science*; 131: 5515–5525.
- Pereira SP, Langley PG and Williams R (1996):** The management of abnormalities of hemostasis in chronic liver failure. *Semin Liver Dis.*; 16(4):403-414
- Perri RE, Hahn SH and Ferber MJ (2005):** Wilson Disease-keeping the bar for diagnosis raised. *Hepatology*; 42(4):974.
- Petersen BE, Bowen WC, Patrene KD, et al. (1999):** Bone marrow as a potential source of hepatic oval cells. *Science*; 284(5417):1168– 1170.
- Petersen J, Ott M and von Weizsacker F (2001):** Current status of cell-based therapies in liver diseases. *Gastroenterol*; 39:975–980.
- Petit FM, Gajdos V and Francoual J (2004):** Allelic heterogeneity of Crigler-Najjar type I syndrome: a study of 24 cases. *Clin Genet.*; 66(6):571-572.
- Pinotti M, Toso R, Girelli D, et al. (2000):** Modulation of factor VII levels by intron 7 polymorphisms: population and in vitro studies. *Blood*; 95(11):343-348.

- Polson J and Lee WM (2005):** The management of acute liver failure. *Hepatology*; 41(5):1179-1197.
- Rambhatla L, Chiu CP, Kundu P, et al. (2003):** Generation of hepatocyte-like cells from human embryonic stem cells. *Cell Transplant*; 12(1):1– 11.
- Ren X, Hogaboam C, Carpenter A, et al. (2003):** Stem cell factor restores hepatocyte proliferation in IL-6 knockout mice following 70% hepatectomy. *J Clin Invest.*; 112(9):1407– 1418.
- Reubinoff BE, Pera MF, Fong CY, et al. (2000):** Embryonic stem cell lines from human blastocysts: Somatic differentiation in vitro. *Nat Biotechnol*; 18: 399–404.
- Roberts HR, Monroe DM and Hoffman M (2001):** Molecular biology and biochemistry of the coagulation factors and pathways of hemostasis. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn U, eds. *William's Hematology*. 2001. 6th ed. New York, NY: McGraw-Hill. p. 1409-1434.
- Rogers I and Casper RF (2003):** Stem cells: you can't tell a cell by its cover. *Hum Reprod Update*; 9(1):25– 33.
- Rogers I and Casper RF (2004):** Umbilical cord blood stem cells. In *Best Practice & Research Clinical Obstetrics & Gynaecology*, eds: N. Fisk & J. Itskovitz, London: Elsevier Ltd., 21: 383- 390.
- Roskams TA, Theise ND, Balabaud C, et al. (2004):** Nomenclature of the finer branches of the biliary tree: canals, ductules, and ductular

reactions in human livers. *Hepatology*; 39(6): 39– 45.

Rossant J (2001): Stem cells in the mammalian blastocyst. *Harvey Lect* 97: 17–40.

Ruhnke M, Ungefroren H, Zehle G, et al. (2003): Long-term culture and differentiation of rat embryonic stem cell-like cells into neuronal, glial, endothelial, and hepatic lineages. *Stem Cells*; 21(4):428– 436.

Ryan Edmond A, Lakey Jonathan RT, Rajotte Ray V, et al. (2001): Clinical outcomes and insulin secretion after islet transplantation with the Edmonton protocol. *Diabetes*; 50:710– 719.

Saji Y, Tamura S, Yoshida Y, et al. (2004): Basic fibroblast growth factor promotes the trans-differentiation of mouse bone marrow cells into hepatic lineage cells via multiple liver enriched transcription factors. *J Hepatol*; 41(4):545–550.

Sakaida I, Terai S, Yamamoto N, et al. (2004): Transplantation of bone marrow cells reduces CCl₄-induced liver fibrosis in mice. *Hepatology*; 40(6):1304– 1311.

Sandhu JS, Petkov PM, Dabevas MD, et al. (2001): Stem cell properties and repopulation of the rat liver by fetal liver epithelial progenitor cells. *Am J Pathol*; 159:1323–1334.

Sass DA and Shakil AO (2005): Fulminant hepatic failure. *Liver Transplant*; 11(6):594-605.

Sato Y, Araki H, Kato J, et al. (2005): Human mesenchymal stem cells

xenografted directly to rat liver differentiated into human hepatocytes without fusion. *Blood*; 106 (2):756– 763.

Schiodt FV and Lee WM (2003): Fulminant liver disease. *Clin Liver Dis.*; 7(2):331-349.

Schiodt FV, Rochling FA, Casey DL et al. (1997): Acetaminophen toxicity in an urban county hospital. *N Engl J.*; 337(16):1112-1117.

Schmidt LE and Dalhoff K (2005): Alpha-fetoprotein is a predictor of outcome in acetaminophen-induced liver injury. *Hepatology*; 41(1):26-31.

Schwartz RE, Reyes M, Koodie L, et al. (2002): Multipotent adult progenitor cells from bone marrow differentiate into functional hepatocyte-like cells. *J Clin Invest.*; 109(10):1291– 1302.

Seaberg RM and Van der Kooy D (2002): Adult rodent neurogenic regions: The ventricular subependyma contains neural stem cells, but the dentate gyrus contains restricted progenitors. *J Neurosci*; 22: 1784–1793.

Selden C and Hodgson H (2004): Cellular therapies for liver replacement. *Transpl Immunol.*; 12(3 – 4):273– 88.

Shafritz DA and Dabeva MD (2002): Liver stem cells and model systems for liver repopulation. *J Hepatol.*; 36:552–564.

Sharma AD, Cantz T, Richter R, et al. (2005): Human cord blood stem cells generate human cytokeratin 18-negative hepatocyte-like cells in injured mouse liver. *Am J Pathol*; 167:555–564.

- Shizuru JA, Negrin RS and Weissman IL (2005):** Hematopoietic stem and progenitor cells: clinical and preclinical regeneration of the hematolymphoid system. *Annu Rev Med.*; 56(1):509– 538.
- Short B, Brouard N, Occhiodoro-Scott T, et al. (2003):** Mesenchymal stem cells. *Arch Med Res.*; 34: 565–571.
- Singhal S and Mehta J (1999):** Reimmunization after blood or marrow stem cell transplantation. *Bone Marrow Transplant*; 23(7):637-646.
- Smith AG (2001):** Embryo-derived stem cells: Of mice and men. *Annu Rev Cell Dev Biol.*; 17: 435–462.
- Sobaniec-Lotowska ME and Lebensztejn DM (2006):** Ultrastructure of Kupffer cells and hepatocytes in the Dubin-Johnson syndrome: a case report. *World J Gastroenterol.*; 12(6):987-9.
- Socie G (2005):** Current issues in allogeneic stem cell transplantation. *Hematology*; 10 Suppl: 1:63.
- Socie G, Salooja N, Cohen A, et al. (2003):** Nonmalignant late effects after allogeneic stem cell transplantation. *Blood*; 101(9):373-385.
- Sokal EM, Smets F, Bourgois A, et al. (2006):** Hepatocyte transplantation in a 4-year-old girl with peroxisomal biogenesis disease: technique, safety, and metabolic follow-up. *Transplantation* 2003; 76:735–738. 158 A. Nussler et al. / *J Hepatol.*, 55:144–159.
- Sort P, Navasa M, Arroyo V, et al. (1999):** Effect of intravenous albumin on renal impairment and mortality in patients with cirrhosis and

spontaneous bacterial peritonitis. *N Engl J Med.*; 341(6): 403-409.

Stem Cell and Developmental Biology Writing Group's Report (2004):

Natl Inst Diabetes & Digestive & Kidney Diseases, NIH. 1–27.

Stockley RA (2000): Alpha-1-antitrypsin deficiency: what next?. *Thorax.*; 55(7):614-618.

Strick-Marchand H, Morosan S, Charneau P, et al. (2004): Bipotential mouse embryonic liver stem cell lines contribute to liver regeneration and differentiate as bile ducts and hepatocytes. *Proc Natl Acad Sci USA.*; 101(22): 60–65.

Strick-Marchand H and Weiss MC (2002): Inducible differentiation and morphogenesis of bipotential liver cell lines from wild-type mouse embryos. *Hepatology*; 36(4 Pt. 1):794– 804.

Strom S and Fisher R (2003): Hepatocyte transplantation: new possibilities for therapy. *Gastroenterology*; 124:568–571.

Strom SC, Chowdhury JR and Fox IJ (1999): Hepatocyte transplantation for the treatment of human disease. *Semin Liver Dis.*; 19:39–48.

Strom SC, Fisher RA, Thompson MT, et al. (1999): Hepatocyte transplantation as a bridge to orthotopic liver transplantation in terminal liver failure. *Transplantation*; 63(4):559– 569.

Suzuki A, Zheng Y, Kondo R, et al. (2000): Flow-cytometric separation and enrichment of hepatic progenitor cells in the developing mouse liver. *Hepatology*; 32:1230–1239.

- Takano H, Ohtsuka M, Akazawa H, et al. (2003):** Pleiotropic effects of cytokines on acute myocardial infarction: GCSF as a novel therapy for acute myocardial infarction. *Curr Pharm.*; 9(14): 21– 27.
- Tanabe Y, Tajima F, Nakamura Y, et al. (2004):** Analyses to clarify rich fractions in hepatic progenitor cells from human umbilical cord blood and cell fusion. *Biochem Biophys Res Commun.*; 324(2):711– 718.
- Tanimizu N, Nishikawa M, Saito H, et al. (2003):** Isolation of hepatoblasts based on the expression of Dlk/Pref-1. *J Cell Sci.*; 116:1775–1786.
- Tateno C and Yoshizato K (1996):** Long-term cultivation of adult rat hepatocytes that undergo multiple cell divisions and express normal parenchymal phenotypes. *Am J Pathol.*; 148:383–392.
- Temple S (2001):** The development of neural stem cells. *Nature*; 414: 112– 117.
- Terai S, Sakaida I, Yamamoto N, Ohata S, et al. (2003):** An in vivo model for monitoring trans-differentiation of bone marrow cells into functional hepatocytes. *J Biochem.*; 134(4): 551– 558.
- Theise ND, Nimmakayalu M, Gardner R, et al. (2000):** Liver from bone marrow in humans. *Hepatology*; 32(1):11 –16.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, et al. (1998):** Embryonic stem cell lines derived from human blastocysts. *Science*; 282(5391):1145– 1147.

- Turrini P, Monego G, Gonzalez J, et al. (2004):** Human hepatocytes in mice receiving pre-immune injection with human cord blood cells. *Nature*; 326(1): 66– 73.
- Vassilopoulos G, Wang PR and Russell DW (2003):** Transplanted bone marrow regenerates liver by cell fusion. *Nature*; 422(34):901– 904.
- Verfaillie CM (2004):** Adult stem cells: Tissue specific or not. *Handbook of Stem Cells*, Vol. 2, ed. R. Lanza. p. 13–20.
- von Mach MA, Hengstler JG, Brulport M, et al.(2004):** In vitro cultured islet-derived progenitor cells of human origin express human albumin in severe combined immunodeficiency mouse liver in vivo. *Stem Cells*; 22:1134–1141.
- Wagers AJ, Sherwood RI, Christensen JL, et al. (2002):** Little evidence for developmental plasticity of adult hematopoietic stem cells. *Science*; 297(50):256– 259.
- Wagers AJ and Weissman IL (2004):** Plasticity of adult stem cells. *Cell*; 116: 639–648.
- Wang X, Al-Dhalimy M, Lagasse E, et al. (2001):** Liver repopulation and correction of metabolic liver disease by transplanted adult mouse pancreatic cells. *Am J Pathol*; 158:571–579.
- Wang X, Foster M, Al-Dhalimy M, et al. (2003):** The origin and liver repopulating capacity of murine oval cells. *Proc Natl Acad Sci.*; 100(91): 81–88.
- Wang X, Ge S, McNamara G, et al. (2003):** Albuminexpressing

hepatocyte-like cells develop in the livers of immunodeficient mice that received transplants of highly purified human hematopoietic stem cells. *Blood*; 101(10):41– 48.

Wang X, Montini E, Al-Dhalimy M, et al. (2002): Kinetics of liver repopulation after bone marrow transplantation. *Am J Pathol*; 161(2):565–574.

Wang X, Willenbring H, Akkari Y, et al. (2003): Cell fusion is the principal source of bone-marrow-derived hepatocytes. *Nature*; 422(34):897– 901.

Wang Y, Nan X, Li Y, et al. (2005): Induction of umbilical cord blood-derived beta (2) m (–) c-Met (+) cells into hepatocytelike cells by coculture with CFSC/HGF cells. *Liver Transplant*; 11(6): 635– 543.

Weissman IL (2000): Stem cells: units of development, units of regeneration, and units in evolution. *Cell*; 100(1):157– 168.

Weissman IL (2000): Translating stem and progenitor cell biology to the clinic: Barriers and opportunities. *Science*; 287(57):142–146.

Wiesner RH, McDiarmid SV, Kamath PS, et al. (2001): MELD and PELD: Application of survival models to liver allocation. *Liver Transplant*; 7:567- 580.

Willenbring H, Bailey AS, Foster M, et al. (2004): Myelomonocytic cells are sufficient for therapeutic cell fusion in liver. *Nat Med*; 10(7):744– 748.

Wilpshaar J, Bhatia M, Kanhai HH, et al. (2002): Engraftment potential of human fetal hematopoietic cells in NOD/SCID mice is not

restricted to mitotically quiescent cells. *Blood*; 100(1):120–127.

Wilpshaar J, Noort WA, Kanhai HH, et al. (2001): Engraftment potential into NOD/SCID mice of CD34+ cells derived from human fetal liver as compared to fetal bone marrow. *Haematol*; 86(10):1021– 1028.

Wognum AW, Eaves AC and Thomas TE (2003): Identification and isolation of hematopoietic stem cells. *Nature*; 34(6):461–475.

Wolf B, Freehauf CL, Thomas JA, et al. (2003): Markedly elevated serum biotinidase activity may indicate glycogen storage disease type Ia. *J Inherit Metab Dis.*; 26(8):85-90.

Wright NA (2000): Epithelial stem cell repertoire: Clues to the origin of cell lineages, proliferative units and cancer. *Int J Exp Pathol.*, 81: 117–143.

Yamada T, Yoshikawa M, Kanda S, et al. (2002): In vitro differentiation of embryonic stem cells into hepatocyte-like cells identified by cellular uptake of indocyanine green. *Stem Cells*; 20 (2):146–154.

Yamamoto H, Quinn G, Asari A, et al. (2003): Differentiation of embryonic stem cells into hepatocytes: biological functions and therapeutic application. *Hepatology*; 37(5):983– 993.

Yin Y, Lim YK, Salto-Tellez M, et al. (2002): AFP+, ESC derived cells engraft and differentiate into hepatocytes in vivo. *Stem Cells*; 20(4):338– 346.

Zeng F, Chen M, Katsumata M, et al. (2005): Identification and characterization of engrafted human cells in human/goat xenogeneic transplantation chimerism. *DNA Cell Biol*; 24:403–409.

Zulewski H, Abraham EJ, Gerlach MJ, et al. (2001): Multipotential nestin positive stem cells isolated from adult pancreatic islets differentiate *ex vivo* into pancreatic endocrine, exocrine and hepatic phenotypes. *Diabetes*; 50: 521–525.