

REFERENCES

- Abulafi A.M and Williams N.S. (1994):** Local recurrence of colorectal cancer: the problem, mechanisms, management and adjuvant therapy. Br j Surg; 81:7.
- Adam R. (2003):** Chemotherapy and surgery: new perspectives on the treatment of unresectable liver metastases. Ann Oncol. 2:13.
- Ahlquist D, Wieand H, Moertal C. (1995):** Accuracy of fecal occult blood screening for colorectal neoplasia. J Am Med Assoc; 269: 1262.
- Albe X, Vassilakos P, Helfer-Guarnori K. (1990):** Independent prognostic value of ploidy in colorectal cancer. Cancer; 66:1168.
- Alektiar KM, Zelefsky MJ, Paty PB. (2000):** High-dose-rate intraoperative brachytherapy for recurrent colorectal cancer. Int J Radiat Oncol Biol Phys; 48: 219.
- Andre T, Boni C, Mounedji-Boudiaf L, et al, (2004):** oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for rectal cancer. N Engl J Med; 350:2343.
- Armitage N.C. and Hardcastle J.D. (1996):** Neoplasms: screening and diagnosis. Curr Opin Gastroenterol; 2:16.
- Arulampalam THA, Costa DC, Loizidou M, Visvikis D, (2001):** Positron emission tomography and colorectal cancer. Br J Surg; 88: 176.
- Barlow A.P, and Thompson M.H. (1993):** colonoscopic follow up after resection for colorectal cancer: a selective policy. Br J surg; 80:781.
- Beart R.W, Metzger P.P, O'Connell M.J. (1981):** Postoperative screening of patients with carcinoma of the colon. Dis Colon Rectum; 24:585

- Beart R.W, and O'Connell M.J. (1999):** Postoperative follow up of patients with carcinoma of the colon. Mayo Clin Proc; 58: 361.
- Bendavid Y, Latulippe JF, Younan R.J. (2002):** Phase I study on sentinel lymph node mapping in colon cancer: a preliminary report. J Surg Oncol; 79: 81.
- Ben-Eliyahu S, Page GG, Yirmiya R, Shakhar G. (1999):** Evidence that stress and surgical interventions promote tumor development by suppressing natural killer cell activity. Int J Cancer; 80: 880.
- Beynon J, Foy D.M, Roe A.M, Temple L.N, et al, (1996):** Endoluminal ultrasound assessment of local invasion in rectal cancer. Br J Surg; 73: 474.
- Bjorklund B. (1990):** The nature and clinical use of tissue polypeptide antigen (TPA). Tumor Diagnostic; 1:9. (abs).
- Broderick-Villa GA, O'Connell TX, DiFronzo LA. (2002):** Does tumor burden limit the success and accuracy of lymphatic mapping and sentinel lymph node biopsy in colorectal cancer? Ann Surg Oncol; 9: S71.
- Bruinvels DJ, Stiggelbout, AM, Kievit J. (1994):** Follow-up of patients with colorectal cancer: A meta-analysis. Ann Surg; 219:174.
- Buhler H, Seefeld U, Deyhle P, Buhler H, Seefeld U, Deyhle P. (1984):** Endoscopic follow-up after colorectal cancer surgery: Early detection of local recurrence? Cancer; 54:791.
- Burrow L, and Tartter P. (1998):** Effect of blood transfusion on colonic malignancy rate. Lancet; 2:662.
- Bussey H.J. (1988):** Metachronous carcinoma of the large intestine and intestinal polyps. Proc R Soc Med; 60:208.

- Camma C, Giunta M, Fiorica F, Pagliaro L, et al, (2000):** preoperative radiotherapy for resectable rectal cancer: a meta-analysis. JAMA; 284: 1008.
- Carne PW, Frye JN, Robertson GM, et al, (2004):** open operation for palliation of colorectal cancer. Dis Colon Rectum; 47:1455.
- Chapman AE, Levitt MD, Hewett P, Woods R, et al, (2001):** laparoscopic-assisted resection of colorectal malignancies: a systemic review. Ann Surg; 234: 590.
- Compton C.C, Fielding L.P, Burgart L.J. (2000):** Prognostic factors in colorectal cancer. Arch Pathol Lab Med; 124: 979.
- Cunningham D, Humblet Y, Siena M, et al, (2003):** Cetuximab (C225) alone or in combination with irinotecan (CPT-11) in patients with epidermal growth factor receptor (EGFR)-positive, irinotecan-refractory metastatic colorectal cancer (MCRC)[abstract]. Proc Am Soc Clin Oncol; 22:252
- Dahlberg M, Glimelius B, Pahlman L. (1999):** Changing strategy for rectal cancer is associated with improved outcome. Br J Surg; 86: 379.
- De Calan L, and Bourlier P. (1998):** Therapeutic choices for cancer of the rectum. J Chir; 135: 51.
- Delgado F, Bolufer JM, Grau E, Domingo C, et al, (1999):** Laparoscopic colorectal cancer resection: initial follow-up results. Surg Laparosc Endosc Perian Tech; 9:91.
- Delvillano B.C, and Zurawski V.R. (1993):** Carbohydrate antigenic determinant (CA 19-9) a monoclonal antibody defined tumor marker. Immunodiagnosics; 65:269.
- Devesa J.M, Morales V, Enriquez J.M. (1999):** The bases for a comprehensive follow up. Dis Colon Rectum; 31:636.

- Dong S, Traverse G, Johnson C. (2001):** Detecting colorectal cancer in stool with the use of multiple genetic targets. *Nat Cancer Inst*; 93: 858.
- Douillard JY, Cunningham D, Roth AD, et al, (2000):** Irinotecan combined with fluorouracil compared with fluorouracil alone as first line treatment for metastatic colorectal cancer: a multicenter randomized trial. *Lancet*; 355:1041.
- Dowdall JF, and Maguire OJ. (2002):** Experience of surgery for rectal cancer with total mesorectal excision in a general surgical practice. *Br J Surg*; 89: 1014.
- Durdey P, and Williams N.S. (1984):** The effect of malignancy and inflammatory fixation of rectal carcinoma as prognosis after rectal excision. *Br J Surg*; 71:787.
- Durdey P, Williams N.S, Brown D.A. (1999):** Serum carcinoembryonic antigen and acute phase reactant proteins in the preoperative detection of fixation of colorectal tumors. *Br J Surg*; 71: 881.
- Feil W, Wunderlich M, Kovats E. (1988):** Rectal cancer: Factors influencing the development of local recurrence after radical anterior resection. *Int J Colorectal.Dis*; 195:334
- Fong Y, Fortner H, Sun RL, Brennan MF, et al, (1999):** Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: Analysis of 1001 consecutive cases. *Ann Surg*; 21:230.
- Francois Y, Nemoz CJ, Baulieux J, Vigna IJ, et al, (1999):** Influence of the interval between preoperative radiation therapy and surgery on downstaging and on the rate of sphincter-sparing surgery for rectal cancer. *J Clin Oncol*; 17:2396.

- Fuchs CS. (1994):** A prospective study of family history and the risk factors of colorectal cancer. *N Engl J Med*; 331:254.
- Fucini C, Tommasi S.M, Rosi S, Malatantis G, et al, (1997):** follow up of colorectal cancer resected for cure: an experience with CEA, TPA, CA 19.9 analysis and second look surgery. *Dis Colon Rectum*; 30: 273.
- Fyrykolm GJ, Glimelius B, Pahlman L. (1993):** Preoperative or postoperative irradiation in adenocarcinoma of the rectum. *Dis Colon Rectum*; 36: 564.
- Galandiuk S, Wieand H.S, Moertel C.G. (1992):** Patterns of recurrence after curative resection of carcinoma of the colon and rectum. *Surg Gynecol Obstet.*; 174:27.
- Gasparini M, Buraggi G.L, Regalia E, Maffioli L. et al, (1994):** Comparison of radioimmunodetection with other imaging methods in evaluating local relapses of colorectal carcinoma. *Cancer*; 73: 846.
- Gerard A, Buyse M, Nordlinger. (1988):** Preoperative radiotherapy as adjuvant treatment in rectal cancer. *Ann Surg*; 208:606.
- Gervaz P, Rotholtz N, Wexner SD, You SY, et al, (2001):** Colonic J-pouch function in rectal cancer patients: impact of adjuvant chemotherapy. *Dis Colon Rectum*; 44:1667.
- Gill S, Loprinzi CL, Sargent DJ, et al, (2004):** pooled analysis of fluorouracil based adjuvant therapy for stage II and III rectal cancer: who benefits and by how much. *J Clin Oncol*; 22:1797.
- Glover, and Waugh. (1999):** Local staging of rectal cancer in comparison with histological finding. *Dis Colon and rectum*; 43: 338.
- Goldberg RM, Flemin TR, Rangen CM. (1998):** Surgery for recurrent colon cancer: strategies for identifying resectable

recurrence and success rates after resection. *Ann Intern Med*; 129: 27.

Goldberg RM, Sargent DJ, Morton RF. (2004): A randomized controlled trial of fluorouracil plus leucovorin, irinotecan and oxaliplatin combinations in patients with untreated metastatic rectal cancer. *J Clin Oncol*; 22:22.

Goligher J. Dnithie h, Nixon H. (1984). Surgical anatomy and physiology of the rectum. In surgery of the anus, rectum and colon. 5th ed, printed in Great Britain. pp 1.

Goligher J. Dnithie h, Nixon H. (1984): Rare tumor of the rectum. In surgery of the anus, rectum and colon. 5th ed., printed in Great Britain. pp1. Cited from **Raven (1948)**

Goligher J.C. (1998): What can be done to improve the results in colorectal cancer? *Scand J Gastroenterol*: 23:190.

Grunwald V, Hidalgo M. (2003): Devolving inhibitors of the epidermal growth factor receptor for cancer treatment. *J Natl Cancer Inst*; 95:851.

Gunderson L.L, and Martenson J.A. (1988): Irradiation of adenocarcinomas of the gastrointestinal tract malignancies. *Front Radiat Ther Oncol*; 22:127.

Gunderson L.L, O'Connell M.J, and Dozois R.R. (1992): The role of intra-operative irradiation in locally advanced primary and recurrent rectal adenocarcinoma. *World J Surg*; 16:495.

Gunderson L.L, Nelson H, Martenson J.A. (1996): Intraoperative electron and external beam irradiation with or without 5-fluorouracil and maximum surgical resection for previously unirradiated locally recurrent colorectal cancer. *Dis Colon Rectum*; 39:1379.

- Gunderson L.L, Haddock M.G, Nelson H. (1997):** Locally recurrent colorectal cancer: IOERT and EBRT $\pm$ 5-FU and maximal resection. Intraoperative Radiation Therapy in the Treatment of Cancer. Basal Karger; pp.31.
- Gunderson L.L, Haddock M.G, Nelson H. (1999):** Indications for and results of combined modality treatment of colorectal cancer. Acta Oncol; 38:7.
- Habib N.A, Peck M.A, Sawyer C.N, (1983):** Does fixity affect prognosis in colorectal tumors ? Br J Surg; 70:423.
- Hall N.R, Finan P.J, Stephenson B.M. (1994):** The role of CA-19-9 and CEA in surveillance following curative resection of colorectal cancer. Br J Cancer; 70:549.
- Hallbook O, Pahlman L, Krog M, Wexner SD, et al, (1996):** Randomized comparison of straight and colonic J pouch anastomosis after low anterior resection. Ann Surg; 224:58. Cited from **Paterson (1912)**
- Hamelin R, Laurent-Puig P, Olschwang S. (1994):** Association of p53 mutations with short survival in colorectal cancer. Gastroenterology; 106:42.
- Harper S.G. (1995):** Use of Intrarectal Ultrasound in the staging of rectal cancer. CurrentSurgery; 52:469.
- Hartley JE, Mehigan BJ, Qureshi AE, Duthie GS, et al, (2001):** Total mesorectal excision: assessment of the laparoscopic approach. Dis Colon Rectum; 44: 315.
- Hartley JE, Monson JR. (2002):** The role of laparoscopy in the multimodality treatment of colorectal cancer. Surg Clin North Am; 82:1019.

Hartly JE, Guillem J, Lopez R, (2003): Resection of locally recurrent rectal cancer in the presence of distant metastases. *Ann surg oncol*; 3:227.

Heah SM, Seow-Choen F, Eu KW, Ho YH, et al, (2002): Prospective randomized trial comparing sigmoid versus descending colonic J-pouch after total rectal excision. *Dis Colon Rectum*; 45:322.

Heald R.J, Husband E.M, and Ryall R.D.H. (1982): The mesorectum in rectal cancer surgery—the clue to pelvic recurrence? *Br J Surg*; 69:613.

Heald R.J, and Ryall R.D. (1986): Recurrence and survival after total mesorectal excision for rectal cancer. *Lancet*; 1:1479.

Hodgman C.G, MacCarty R.L, Wollf B.G. (1996): Preoperative staging of rectal carcinoma by computed tomography and magnetic resonance imaging. *Dis Colon Rectum*; 29: 446.

Hofiman GC, Baker JW, Doxey JB, Hubbard GW, et al, (2001): Minimal invasive surgery for colorectal cancer. Initial follow-up. *Ann Surg*; 223:790.

Hojo K, Sawada T, Moriya Y. (1989): An analysis of survival and voiding, sexual function after wide ileopelvic lymphadenectomy in patients with carcinoma of the rectum compared to conventional abdominosacral lymphadenectomy. *Dis Colon Rectum*; 32:128.

Hurwitz H, Fehrenbacher L, Novotny W, et al, (2004): Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl Med*; 350:2335.

- Jain RK. (2001):** Normalizing tumor vasculature with anti-angiogenic therapy: a new paradigm for combination therapy. *Nat Med*; 7: 748.
- Jimenz R, Shoup M, Cohen A. (2003):** Contemporary outcomes of total pelvic exenteration in the treatment of colorectal cancer. *Dis colon rectum*; 46: 1619.
- Johnson RJ, Jenkins J.P, Isherwood James R.D, Schoeield P.F. (1997):** Quantitative magnetic resonance imaging in rectal cancer. *Br J Radiol*; 60: 461.
- Juan Rosai. (1981):** Gastrointestinal tract. In Ackerman's surgical pathology. 6th ed., printed in united states of America; pp 502.
- Kanellos I. (2003):** Surgical management of colorectal cancer. *Medimond int proc*; 4:275.
- Kapiteijn E, Marijnen CAM, Nagtegaal ID, Putter H, et al, (2000):** Preoperative radiotherapy combined with total mesorectal excision for respectable rectal cancer. *N Engl J Med*; 345:638.
- Kemeny N, Huang Y, Cohen A. (1999):** Hepatic arterial infusion of chemotherapy after resection of hepatic metastases from colorectal cancer. *N Engl J Med*; 341: 2039.
- Keynes W.M. (1999):** Implantation from the bowel lumen in cancer of the large intestine. *Ann Urg*; 153: 357.
- Kikuchi R, and Takano M. (1995):** Management of early invasive colorectal cancer. Risk of recurrence and clinical guidelines. *Dis Colon and Rectum*; 38: 1286.

- Kim N.K, Yasmineh W.G, Freier E.F. (1998):** Value of alkaline phosphatase, 5-nucleotidase, gamma glutamyltransferase, and glutamate dehydrogenase activity measurements in serums in diagnosis of metastasis to the liver. Clin Chem; 23:2034.
- Kramer T, Share R, Kiel K. (1991):** Intraoperative radiation therapy of colorectal cancer. New York Pergamon Press; pp. 308.
- Krestin G.P, Steinbrich E, Friedmann G. (1998):** Recurrent rectal cancer: diagnosis with MR imaging versus CT. Radiology; 168: 307.
- Krook J.E, Moertel C.G, Gunderson L.L. (1991):** Effective surgical adjuvant therapy for high-risk rectal carcinoma. N Engl J Med; 324:709.
- Kuullier E, Rullier A, Laurent C, Vendrely V, et al, (2002):** Guidelines for conservative surgery after preoperative radio chemotherapy for rectal carcinoma: pathologic assessment of distal intramural spread. Meeting of the European Association of coloproctology. Eriangen; 56.
- Lee J.K, Stanley R.J, Sagel S.S, Levitt R.G, Mclennan B.L, (1991):** CT appearance of the pelvis after abdominoperineal resection for rectal carcinoma. Radiology; 141:737.
- Lee J.K, Chow NH, Wang ST, et al, (200)** Prognostic value of vascular endothelial growth factor expression in colorectal patients. Eur J Cancer; 36:748.
- Lezoche F, Paganini AM, Feliciotti F, (2002):** Transanal endoscopic microsurgery excision. World J surgery; 305:317.

- Longo EW. (2002):** The preoperative assessment and postoperative surveillance of patient with colon and rectal cancer. Surg Clin N Am; 82:1091.
- Lopez-Kostner F, Lavery I.C, Hool G.R. (1997):** Total mesorectal excision is not necessary for cancers of the upper rectum. Surgery; 124:612.
- Mackay SG, Pager CK, Joseph D, Solomon M. (2003):** Assessment of accuracy of trans-rectal U/S in anorectal neoplasia. Dis Colon Rectum; 7:134.
- Madsen P.M, and Christiansen J. (1986):** Distal intramural spread of rectal carcinoma. Dis Colon Rectum; 29:279.
- Magrini S, Nelson H, Gunderson L.L. (1996):** Sacropelvic resection and intraoperative electron irradiation in the management of recurrent anorectal cancer. Dis Colon Rectum; 39:1.
- Manson P.N, Corman M.L, Collier J.A. (2002):** Anterior resection for adenocarcinoma. Am J Surg; 131:431.
- Marijnen CA, Nagtegaal ID, Klein Kranenbarg E, Hermans J, et al, (2001):** Downstaging after short-term preoperative radiotherapy in rectal cancer patients. J Clin Oncol; 19: 1976.
- Martin E.W, Minton J.P, and Carey L.C. (1985):** CEA-directed second-look surgery in the asymptomatic patient after primary resection of colorectal carcinoma. Ann Surg; 202:310.
- Martin E.W, and Carey L.C. (2002):** Second-look surgery for colorectal cancer. Ann Surg; 214:321.
- Martling A, Cedemiark B, Johansson H, Rutqvist LE, et al, (2002):** The surgeon as a prognostic factor after the

introduction of total mesorectal excision in the treatment of rectal cancer. Br J Surg; 89;1008.

Mason AY. (2002): Surgical access to the rectum, a transsphincteric exposure. Proc R Soc Med; 63:91.

Matsouka H, Masaki T, Iakahara T. (2002): Preoperative staging by multidetector computed tomography in rectal cancer. cancer; 2: 82.

McAnena OJ, Heald RJ, Lockhart-Mummery HE. (1998):
Curative and functional results of total mesorectal excision with ultra-low anterior resection in the management of carcinoma of the lower one-third of the rectum. Surg Gynecol Obstet; 170:517.

McDermott F.T, Hughes E.S, Pihl E. (1985): Local recurrence after potentially curative resection for rectal cancer in a series of 1008 patients. Br J Surg; 72:34.

Mellgren A, Sirivongs P, Rothenherger DA. (2002): Is local excision adequate therapy for early rectal cancer? Dis Colon Rectum; 43: 1064.

Merrie AE, van Rij AM, Phillips LV, Rossaak JI. et al, (2001): Diagnostic use of the sentinel node in colon cancer. Dis Colon Rectum; 44: 410.

Minton J.P, Hoehn I.L, Gerber D.M. (2000): Results of a 400 – patient carcinoembryonic antigen second look colorectal cancer study. Cancer; 55: 1284.

Moertel C.G, Fleming T.R, Macdonald J.S. (1990).
Levamisole and fluorouracil for adjuvant therapy of resected rectal carcinoma. N Engl J Med; 322:352.

- Moreira L.F, Hizuta A, Iwagaki H, Tanaka N, Orita K. (1994):** Lateral lymph node dissection for rectal carcinoma below the peritoneal reflection. Br J Surg; 81: 293.
- Morikawa E, Yasutomi M, Shindou K. (1994):** Distribution of metastatic lymph nodes in colorectal cancer by the modified clearing method. Dis Colon Rectum; 37:219. Cited from **Jamieson (1909)**
- Morrow M, and Enker W.E. (1984):** Late ovarian metastases in carcinoma of the colon and rectum. Arch Surg; 119:1385.
- Naegle C.E, Schaffer J.L, and Heppenstall R.B. (1984):** Compartment syndrome complicating prolonged use of lithotomy position. Surgery; 110:566.
- Nag S, Martinez-Monge R, Martin EW. (1999):** Intraoperative electron beam radiotherapy in recurrent colorectal carcinoma. J Surg Oncol; 72: 66.
- Nagtegaal ID, Marijnen CA, Kranenbarg EK. (2002):** Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimeter but two millimeters is the limit. Am J Surg Pathol; 26: 350.
- Nava H.R. and Pagana T.J. (1992):** Postoperative surveillance of colorectal carcinoma. Cancer; 49:1043.
- Nelson RL, Persky V, Turyk M. (1999):** Determination of factors responsible for the declining incidence of colorectal cancer. Dis Colon Rectum; 42:741.
- Nissan A, Guillem JG, Paty PB. (2001):** Abdominoperineal resection for rectal cancer at a specialty center. Dis Colon Rectum; 44: 27.

- Northover J.M.A. (1985):** Carcinoembryonic antigen and recurrent colorectal cancer. *Br J Surg*; 72:S44.
- Nyam D, and Nelson H. (1997):** Recurrent colorectal cancer, *Surgery of the Colon & Rectum*. New York. Churchill Livingstone; 22:58.
- Olsen H.W, Lawrence W.A, Snook C.W. (1988):** Review of recurrent polyps and cancer in 500 patients with initial colonoscopy for polyps. *Dis Colon Rectum*; 31:222.
- Orefice S, Gennari L, Mor C. (1981):** The value of the CEA test in the diagnosis of metastases of adenocarcinoma of the gastroenteric tract. *Tumori*; 67:109.
- Oshima M, Nesb Akken A, Nygaard K. (2001):** Bladder and sexual dysfunction after mesorectal excision for rectal cancer. *Br J Surg*; 87: 2107.
- Ota DM. (2000):** Is intraoperative lymph node mapping and sentinel lymph node biopsy for colorectal carcinoma necessary? *Ann Surg Oncol*; 7: 82. Cited from Miles (1926)
- Parkin D.M, Pisani P, Ferlay J. (1997):** Global cancer statistics. *CA Cancer J Clin*; 49:35.
- Parks AG, and Percy IP. (2000):** Resection and sutured colo-anal anastomosis for rectal carcinoma. *Br J Surg*; 69:301.
- Peter L, Lawrence H, Martin M. (1995):** Anatomy of Alimentary system. In *Gray's Anatomy*. 38th ed., printed in Great Britain. pp1683.
- Pezim M.E, and Nicholls R.J. (1994):** Survival after high or low ligation of inferior mesenteric artery during curative surgery for rectal cancer. *Ann Surg*; 200:729.

- Phillips R.K, Hittinger R, Blesovsky L. (1984):** Local recurrence following curative surgery for large bowel cancer, the rectum and rectosigmoid. Br J Surg; 71:17.
- Phillips R.K, Hittinger R, Blesovsky L. et al, (1999):** Local recurrence following curative surgery for large bowel cancer. Br J Surg; 71:12.
- Pilipshen S.J, Heilweil M, Quan S.H. (1999):** Patterns of pelvic recurrence following definitive resections of rectal cancer. Cancer; 53:1354.
- Plebani M, Basso D, Roveroni G, Depaoll. (2002):** N-Terminal peptide of type 3 procollagen. A possible predictor of colorectal carcinoma recurrence. Cancer; 79: 1299.
- Pollet W.G, and Nicholls R.J. (1984):** The relationship between the extent of distal clearance and survival and local recurrence rates after curative anterior resection for carcinoma of the rectum. Ann Surg; 198:159.
- Quirke P, Durdey P, Dixon M.F. (1986);** Local recurrence of rectal adenocarcinoma due to inadequate surgical resection: Histological study of lateral tumor spread and surgical excision. Lancet; 2:996.
- Radice E, Nelson H, Mercill S. (1999):** Primary myocutaneous flap closure following resection of locally advanced pelvic malignancies. Br J Surg; 86:349.
- Ranbarger K.R, Johnston W.D, and Chang J.C. (1982):** Prognostic significance of surgical perforation of the rectum during abdomino perineal resection for rectal carcinoma. Am J Surg; 143:186.
- Rantala A, and Ovaska J. (1995):** Palliative laser treatment of rectal cancer. Scand J Gastroenterol; 30:177.

- Read D.R, Hambrick E, Abcarian H, Levine H. (1997):** The preoperative diagnosis of hepatic metastases in cases of colorectal carcinoma. *Dis Colon Rectum*; 20:101.
- Rex D, Lrhman G, Hawes R, Ulbright Smith J. (2003):** Screening colonoscopy in asymptomatic average-risk persons with negative fecal occult blood tests. *Gastroenterology*; 100:64.
- Rhomberg W, Eiter H, Hergan K. (1994):** Inoperable recurrent rectal cancer: Results of a prospective trial with radiation therapy. *Int J Radiat Oncol Biol Phys*; 30:419.
- Rich T, Gunderson L.L, Lew R. (1983):** Patterns of recurrence of rectal cancer after potentially curative surgery. *Cancer*; 52:1317.
- Romano G, De Rosa P, Vallone G, Rotondo A. (1995):** Intrarectal ultrasound and computed tomography in the pre and postoperative assessment of patients with rectal cancer. *Br J Surg*; 72:5117.
- Rullier E, Zerbib F, Laurent C, Bonnel C. et al, (2002):** Intersphincteric resection with excision of internal anal sphincter for conservative treatment of very low rectal cancer. *Dis Colon Rectum*; 42:1168.
- Safi F, Link K.H, Beger H.G. (1993):** Is follow up of colorectal cancer patients worthwhile ? *Dis Colon Rectum*; 36: 635.
- Sagar P.M, and Pemberton J.H. (1996):** Surgical management of locally recurrent rectal cancer. *Br J Surg.*; 83: 293.
- Saha S, Wiese D, Badin J. (2000):** Technical details of sentinel lymph node mapping in colorectal cancer and its impact on staging. *Ann Surg Oncol*; 7: 120.

- Sailer M, Fuchs KH, Fein AI, Thiede A. (2002):** Randomized clinical trial comparing quality of life after straight and pouch coloanal reconstruction. Br J Surg; 89: 1108.
- Saltz BL. (2002):** adjuvant therapy of the cancer of the colon and rectum. Surg Clin N Am; 82:1035.
- Sardi A. (1991):** Multiple operations for recurrent colorectal cancer. Semin Surg Oncol; 7:146.
- Sauer, and Bacon. (2000):** Use of ultrasonography in the staging of rectal cancer. Current surgery; 52: 469.
- Sauter J, Stangl R, Altendor-Hofmann A. (1990):** Hepatic metastases from colorectal carcinoma: Impact of surgical resection on the natural history. Br J Surg; 77:1241.
- Schiessel R, Wunderlich M, Herbst F. (2002):** Local recurrence of colorectal cancer, effect of early detection and aggressive surgery. Br J Surg; 73:342.
- Schneebaum S, Arnold M.W, Young D. (1999):** Role of carcinoembryonic antigen in predicting respectability of recurrent colorectal cancer. Dis Colon Rectum; 36: 810.
- Simunovic M, Sexton R, Rempel E, Moran Bj, Healed Rj. (2003):** Optimal preoperative assessment and surgery for cancer rectum. Br J Surgery; 7:83-91.
- Suzuki K, Gunderson L.L, Devine R.M. (1995):** Intraoperative irradiation after palliative surgery of locally recurrent rectal cancer. Cancer; 75:939.
- Suzuki K, Dozois R.R, Devine R.M. (1996):** Curative reoperations for locally recurrent rectal cancer. Dis Colon Rectum; 39:730.

- Takahara T. (2002):** preoperative staging by computed tomography in patient with rectal carcinoma. *Am J Surg*; 184:131.
- Temple JW. (2000):** Locally recurrent rectal cancer: role of composite resection of extensive pelvic tumors with strategies for minimizing risk of recurrence. *J Surg oncol*; 73:47.
- Tornquist A, Ekelund G, Leandoer L. (2000):** The value of intensive follows up after curative resection for colorectal carcinoma. *Br J Surg*; 69: 725.
- Umpleby H.C, Fermor B, Symes M.O, Williamson R.C.N. (1999):** Viability of exfoliated carcinoma cells. *Br J Surg*; 71: 659.
- Van Custem S, Kapiteijn E, Nagegaal ID, Putter H. (2001):** Preoperative radiotherapy combined with total mesorectal excision for respectable rectal cancer. *N Engl J Med*; 345:646.
- Vernava A.M, Longo W.E, Virgo K.S. (1994):** Current follow-up strategies after resection of colon cancer. *Dis Colon Rectum*; 37:573.
- Wang J.Y, Tang R, and Chiang J.M. (1994):** Value of carcinoembryonic antigen in the management of colorectal cancer. *Dis Colon Rectum*; 37:272.
- Wassif S.B, Langenhorst B.L, and Hop C.J. (1994):** The contribution of preoperative radiotherapy in the management of borderline operability rectal cancer. *New York Grune & Stratton*; pp 621.
- Wibe A, Rendedal PR, Svensson E. (2002):** Prognostic significance of the circumferential resection margin

following total mesorectal excision for rectal cancer. Br J Surg; 89: 327.

Willett CG, Tepper JE, Cohen AM, Orlow E, et al, (2002): Failure patterns following curative resection of colonic carcinoma. Ann Surg; 200: 685.

Willett CG, boucher Y, di-tomaso, et al, (2004): Direct evidence that the VEGF-specific antibody bevacizumab has antivasculr effects in human rectal cancer. Nat Med; 10:145.

Williams N.S, Dixon M.E, Johnston D. (1983): Reappraisal of the 5 cm rule of distal excision for carcinoma of the rectum: A study of distal intramural spread and of patients' survival. Br. J. Surg; 70:150.Cited from **Beirne (1933)**

Williams N.S, Durdey P, Johnston D. (2000): The outcomes following sphincter saving resection and abdominoperineal resection for low rectal cancer. Br J Surg; 72:595.

Witzig T.E, Loprinzi C.L, Gonchoroff N.J. (1994): DNA ploidy and cell kinetic measurements as predictors of recurrence and survival in stages B2 and C colorectal adenocarcinoma. Cancer; 68:879.

Wolmark N. (1983): The prognostic significance of tumor location and bowel perforation in Dukes' B and C colorectal cancer. Ann Surg; 198:743.

Wolmark N, and Fisher B. (1986): An analysis of survival and treatment failure following abdominoperineal resection and sphincter saving resection in Dukes' B and C rectal carcinoma. Ann Surg; 204:480.

Wong JH, Steineman S, Calderia C, Bowles J, Namiki T. (2001): Ex vivo sentinel node mapping in carcinoma of the colon and rectum. Ann Surg; 233: 515.

Wood TF, Spirt M, Rangel D. (2001): Lymphatic mapping improves staging during laparoscopic colectomy for cancer. Surg Endosc; 15: 715

Yong L, Deam-m, Monson JRT, Darzi A. (2001): Systematic review of laparoscopic surgery for colorectal malignancy. Siirg Endosr 2WI; 15: 1431.

Young-Fadok, T.M, Wolff B.G, Nivatvongs S. (1998): Prophylactic oophorectomy in colorectal carcinoma: Preliminary results of a randomized, prospective trial. Dis Colon Rectum; 41:277.

Zaheer S, Pemberton J.H, Farouk R. (1998): Surgical treatment of adenocarcinoma of the rectum. Ann Surg; 227:800.