

References

Abarquez RF (2001).

Management options for hypertension syndrome.

Int J Clin Pract ; **55(8)**: 537-545.

Abassi ZA, Tate JE, Golomb E and Keiser HR (1992).

Role of neutral endopeptidase in the metabolism of endothelin.

Hypertension ; **20**: 81-95.

Abboud FM (1982).

The sympathetic system in hypertension.

Hypertension ; **4**: 208-225.

Agewall S, Persson B and Fageberg B (1993).

Microalbuminuria in treated hypertensive men at high risk of coronary disease.

J Hypertens ; **11**: 461-469.

Agrawal B, Berger A, Walk K and Luft FC (1996).

Microalbuminuria screening by reagent strip predicts cardiovascular risk in hypertension.

J Hypertens ; **14**: 223-228.

Alberts GF, Peifley KA, Johns A, *et al.* (1994).

Constitutive endothelin-1 overexpression promotes smooth muscle cell proliferation via an external autocrine loop.

J Biol Chem ; **269**: 10112-10118.

Allen K, Shykoff BE and Izzo JL (1998).

Cognitive appraisal of threat or challenge predicts hemodynamic responses to mental arithmetic and speech tasks.

Am J Hypertens ; 11: 134A.

Anderson EA, Hoffman RP, Balon TW and Mark AL (1991).

Hyperinsulinemia produce both sympathetic neural activation and vasodilatation in normal humans.

J Clin Invest ; 87: 2246-2252.

Arnesen E, Refsum H, Bonna KH *et al.* (1995).

Plasma Hcy and coronary heart disease.

Int J Epidemiol ; 24 : 704 - 709.

Aronow WS, Ahn C, Kronzon I and Gutstein H (1997).

Association of plasma renin activity and echocardiographic left ventricular hypertrophy with frequency of new coronary events and new atherothrombotic brain infarction in older person with systemic hypertension.

Am J cardiol ; 79 (11): 1543-1545.

Banskota NK, Taub R and Zellner K (1989).

Characterization of induction of protooncogene c-myc and cellular growth in human vascular smooth muscle cells by insulin and IGF-I.

Diabetes ; 38: 123-129.

Barham O and Trinder P (1972).

Quantitative enzymatic colorimetric determination of uric acid in serum, plasma and urine.

Analyst ; 97 : 142.

Baron AD (1996).

Insulin and the vasculature.

J Invest Med ; 44: 406-412.

Beckman JS and Koppenol WH (1996).

Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and the ugly.

Am J Physiol ; 271:C1424-C1437.

Beever G , Lip GYH and O'Brien E. (2001)

Pathophysiology of hypertension .

BMJ ; 322 : 912 - 916.

Benigni A, Perico N, Gaspari F, Zoja C, Bellizi L and Remuzzi G (1990).

Increased renal endothelin production in rats with renal mass reduction.

J Am Soc Nephrol ; 1: 411.

Benigni A, Zola C, Corna D, et al. (1996).

Blocking both type A and B endothelin receptor in the kidney attenuates renal injury and prolongs survival in rats with remnant kidney.

Am J Kidney Dis ; 27: 416-423.

Berkin KE and Ball SG. (2001)

Essential hypertension : The heart and hypertension.

Heart ; 86 : 467 - 475.

Bjorntorp P, Krotkiewski M and Sjostrom L (1983).

Impact of obesity on metabolism in men and women: Importance of regional adipose tissue distribution.

J Clin Invest ; 72: 72-1150.

Bianchi S, Biggazzi R, Baldari G, Sgherri GP and Campese VM (1994).

Diurnal variation of blood pressure and microalbuminuria in essential hypertension.

Am J Hypertens ; 7: 23-29.

Bianchi S, Bigazzi R, Muscelli E, Baldari G, Pecori N and Natali A (1995).

Insulin resistance in microalbuminuric hypertensives.

Hypertension ; 26: 789-795.

Bigazzi R, Bianchi S, Campese VM and Baldari G (1992).

Prevalence of microalbuminuria in a large population of patients with mild to moderate essential hypertension.

Nephron ; 61: 94-97.

Bigazzi R, Bianchi S, Baldari G and Campese VM (1998).

Microalbuminuria predicts cardiovascular and renal insufficiency in hypertension.

J Hypertens ; 16: 1325-1333.

Binggeli C, Corti R, Sudano I, Lusher T and Noll G (2002).

Effects of chronic calcium channel blockade on sympathetic nerve activity in hypertension.

Hypertension ; 39(4): 892-896.

Bland JM and Altman DG (1983).

Measurement in medicine: The analysis of method comparison studies.

J Hypertens ; 12(3): 315-331.

Blankenhorn DH and krausch DM (1989).

Reversal of atherosclerosis and sclerosis: The two components of arteriosclerosis.

Circulation ; 79 : 1-7.

Blummenfeld JD, Alderman MH, Sealey JE, *et al.* (1998).

Plasma renin activity and acute myocardial infarction: More evidence for a positive association.

J Am Coll Cardiol ; 31: 310A.

Bogaty P , Mure P and dumesnil JG. (2002).

New insight into diastolic dysfunction as the cause of left sided heart failure associated with systemic hypertension.

Am J Cardiol ; 89 : 341 - 345.

Boger RH, Bode-Boger SM, Brandes RP, Phivthong-ngam L, Bohme M,

Nafe R, Mugge A and Frolich JC (1997).

Dietary L-arginine reduces the progression of atherosclerosis in cholesterol-fed rabbits: comparison with lovastatin.

Circulation ; 96:1282-1290.

Boger RH, Bode-Boger SM, Szuba A, Tsao PS, Chan JR, Tangphao O,

Blaschke TF and Cooke JP (1998).

Asymmetric dimethylarginine (ADMA): a novel risk factor for endothelial dysfunction: its role in hypercholesterolemia.

Circulation ; 98:1842-1847.

Borgeson DD, Grantham JA, Williamson EE, *et al.* (1998).

Chronic oral endothelin type A receptor antagonism in experimental heart failure.

Hypertension ; **31**: 755-770.

Bostom AG, Shemin D, Lapane KL, *et al.* (1995).

Hyperhomocysteinaemia and traditional cardiac, vascular disease risk factors in end-stage renal disease patients on dialysis: A case-control study.

Atherosclerosis ; **114**: 93-103.

Boysen G , Brander T , Christensen H *et al.*(2003).

Homocysteine and risk of recurrent stroke .

Stroke ; **34(5)** : 1258-1261.

Brenner BM and Rector SM (eds.) (1987).

Structure and function of the renal circulation. In: Renal Physiology in Health and Disease. *Chapter 1, WB Saunders Company, Philadelphia, London* p: 1-26.

Brenner BM and Rector SM (eds.) (2000).

Essential hypertension. In: The Kidney. 6th edition, Chapter 46, WB Saunders Company, Philadelphia, London p: 1967-2006.

Bruner CA, Mangiapane ML, Fink GD and Webb RC (1988).

Area postrema ablation and vascular reactivity in deoxycorticosterone-salt-treated rats.

Hypertension ; **11**: 668-673.

Burns PN (1987).

The physical principles of doppler and spectral analysis.

J Clin Ultrasound ; **154**: 567-590.

Burtis CA and Ashwood ER (1994).

Textbook of clinical chemistry. 2nd Edition: WB. Saunders co: Philadelphia.
page 562.

Cacoub P, Dorent R, Maistre G, *et al.* (1993).

Endothelin-1 in primary pulmonary hypertension and the Eisenmenger syndrome.

Am J Cardiol ; **72**: 448-450.

Cain A, Khalil E and Raouf A (2002).

Pathophysiology of essential hypertension: role of pump, the vessel, and the kidney.

Semin Nephrol ; **22(1)**: 3-16.

Camilleri JP, Berry CI, Fiessinger JN, Bariety J and Springer V (1989).

Disease of the arterial system.

Berlin Heidelberg ; **2**: 8-16.

Campese VM, Bianchi S and Bigazzi R (1999).

Association between hyperlipidemia and microalbuminuria in essential hypertension.

Kid Int ; **56(71)**: S10-S13.

Capron L, Jarnet J and Houssete E (1988).

Growth promoting effects of diabetes and insulin on arteries.

Diabetes ; **35**: 973-978.

Caro JF (1991).

Insulin resistance in obese and non obese men.

J Clin Endocrinol Metab **73**: 691-695.

Casale PN, Milner A and Devereux RB (1985).

Value of left ventricle mass in predicting cardiovascular morbid events in hypertensive men.

Circulation **72**: 130.

Casale PN, Devereux RB and Milner M (1986).

Value of echocardiographic measurement of left ventricular mass in predicting cardiovascular morbid events in hypertensive men.

Ann Intern Med ; **105**: 173-178.

Celermajer DS, Sorensen KE, Bull C, Robinson Jand Deanfield JE (1994).

Endothelium-dependent dilation in the systemic arteries of asymptomatic subjects related to coronary risk factors and their interaction.

J Am Coll Cardiol ; **24**:1468–1474.

Cerasola G, Cottone S, Mule G, Nardi E and Riccobene R (1996).

Microalbuminuria, renal dysfunction and cardiovascular complication in essential hypertension.

J Hypertens ; **14**: 915-920.

Chen LH, Liu ML, Hwang HY, Chen LS, Korenberg J and Shane B (1997).

Human methionine synthase. cDNA cloning gene localization and expression.

J Biol Chem; **272**: 3268-3634.

Chico A, Perez A, Cardoba A, Arcelus R, Carreras G and Blanco (1998).

Plasma homocysteine as a risk factor for cardiovascular disease.

Diabetologia ; **41**: 684-693.

Chobanian AV, Bakris GL , Black HR et al., 2003.

The seventh report of the joint national committee on prevention , detection , evaluation and treatment of high blood pressure.

JAMA ; **289**(19) : 2560

Christen AI, Sanchez RA, Baolivo HP, Armentano RL, Risk MR and Fischer ET (1997).

Non-invasive assessment of systemic elastic behaviour in hypertensive patients.

Med Prog Technol ; **21**: 5-11.

Clavell A, Stingo AJ, Margulies KB, et al. (1995).

Role of endothelin receptor subtypes in the in vivo regulation of renal function.

Am J Physiol **268**: 455-460.

Clavell A, Mattingly M, Nir A, et al. (1996).

Angiotensin-converting enzyme inhibition modulates circulating tissue endothelin in chronic canine thoracic inferior vena cava constriction.

J Clin Invest 97: 1286-1292.

Clozel JP, Kuhn H and Hefti F (1989).

Decreases of vascular hypertrophy in four different types of arteries in spontaneously hypertensive rats.

Am J Med 87: 925-955.

Cobe SM (1998).

Lipids in hypertension patients.

Am J Hypertens 11(7) : 887-889.

Cook BB , Treiber FA , Mensah G , Jindal M and Kapuku GK. (2001).

Family history of hypertension. and left ventricular mass in youth.

Am J Hypertens ; 14(4 pt1) : 351 - 356.

Cooke J P (2000).

Does ADMA Cause Endothelial Dysfunction?

Arteriosclerosis, Thrombosis, and Vascular Biology;20:2032.

Cooke J P (2003).

Flow, NO, and atherogenesis

PNAS ; 100(3): 768 - 770.

Cooke JP, Rossitch E Jr, Andon NA, Loscalzo J and Dzau VJ (1991).

Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator.

J Clin Invest ; **88**:1663–1671.

Cooke JP and Dzau VJ (1997) .

Nitric oxide synthase: role in the genesis of vascular disease.

Annu Rev Med ; **48**:489–509.

Corea L, Bentivoglio M, Verdechia P and Motolese M (1984).

Plasma norepinephrine and left ventricular hypertrophy in systemic hypertension.

Am J cardiol ; **53**: 1299-1303.

Cowley AW and Roman RJ (1996).

The role of the kidney in hypertension.

JAMA ; **275**: 1581-1589.

Crimm RH, Flak JM, Byington R, *et al.* (1990).

A comparison of antihypertensive drugs effects on the progression of extracranial carotid atherosclerosis.

Drug ; **40(2)**: 38-43.

Cuzzalo F , Mallamaci F , Tripepi G *et al.* (2001).

Urinary adrenomedullin is related to ET - 1 and salt intake in patients with essential hypertension.

Am J Hypertens ; **14(3)** : 224 - 230.

Davenport AP, Ashby MJ, Easton P, Ell AA, Bedford J, Dickenson C, Capper SJ and Brown MJ (1990).

Radioimmunoassay measuring endothelin-like immunoreactivity in human plasma: Comparison of levels in patients with essential hypertension and normotensive control subjects.

Clin Sci ; **78**: 261-264.

Decsi T and Molnar D (2003)

Insulin resistance syndrome in children : pathophysiology and potential management strategies .

Paediatr Drugs ; **5(5)** : 291-299

Desouza C , Keabler MC, Namara D *et al.* (2002).

Vitamins decrease CVD associated with medication.

Drugs ; **62 (4)** : 605 - 616.

Despres JP, Moorjani S and Lupien PJ (1990).

Regional distribution of body fat, plasma lipoproteins, and cardiovascular disease.

Atherosclerosis ; **10**: 497-511.

Devereux RB , Alonso DR and Lutas GM. (1986).

Echocardiographic assessment of left ventricular hypertrophy

Am J Cardiol ; **57**: 480 - 498.

Devereux RB , Lutas EM and Casale PN. (1984).

Standarization of M - mode echocardiographic left ventricular anatomic measurement.

J AM Coll Caediol ; **4** : 1222 - 1230.

Dibona GF (1991).

Role of the renal nerves in hypertension.

Semin Nephrol ; 11: 503-511.

Doria A, Fioetto P, Avogaro A, Carraro A and Nosadini R (1991).

Insulin resistance is associated with high sodium-lithium counter transport in essential hypertension.

Am J Physiol ; 261: 684-691.

Drayer JM, Weber MA and Laragh JH (1981).

Echocardiography in the evaluation of patients with mild or borderline hypertension. Symposia Specialists; Miami; 21-23.

Dunn FG and Pfeffer MA. (1999).

Left ventricular hypertrophy and hypertension .

N Engl J Med ; 340 : 1279 - 1280.

Durand P ,prost M , Loreau N , Lussier S and Blache D (2001).

Impaired Hcy metabolism and atherothrombotic disease.

Labortory investigation ; 81: 645 - 672.

Durrett LR and Ziegler MG (1990).

A sensitive radioenzymatic assay for catechol drugs.

J neurosci Res; 5: 587-598.

Eisenhofer G, Esler MD, Goldstein Ds and Kopin IJ (1991).

Neuronal uptake, metabolism, and release of tritium-labeled norepinephrine during assessment of its plasma kinetics.

Am J Physiol ; 261: 505-515.

Esler M, Julius S and Zweifler A (1977).

Mild high-renin essential hypertension: Neurogenic human hypertension.

N Engl J Med ; 296: 405-411.

Facchini FS, Stoohs RA and Reaven GM (1996).

Enhanced sympathetic nervous activity. The linchpin between insulin resistance, hyperinsulinemia, and heart rate.

Am J Hypertens ; 9: 31013-1017.

Falkner B, Hulman S, Tannenbaum J and Kushner H (1990).

Insulin resistance and blood pressure in young black men.

Hypertension ; 16: 706-711.

Fareh J, Touyz RM, Schiffrin EL and Thibault G (1996).

Endothelin-1 and angiotensin II receptors in cells from rat hypertrophied heart, receptor regulation and intracellular Ca²⁺ modulation.

Circ Res ; 78: 302-311.

Fawcett JK and Scott JE (1960).

Determination of blood urea by Betheldl's reaction.

J Clin Path ; 13: 156-159

Ferrar DJ, Green HD, Wagner WP and Bond GM (1980).

Reduction in pulse wave velocity and improvement of aortic distensibility accompanying regression of atherosclerosis in the Rhesus monkey.

Circ Res; 47: 425-432.

Ferrannini E, Buzzigoli G and Bonaddona R (1987).

Insulin resistance in essential hypertension.

N Engl J Med ; **317**: 350-357.

Ferrari P, Weidmann P, Shaw S and Heynen G (1991).

Altered insulin sensitivity, hyperinsulinemia and dyslipidemia in individuals with hypertensive parents.

Am J Med ; **91**: 589-596.

Ferrario CM, Barnes KL and Bohonek S (1991).

Neurogenic hypertension produced by lesions of the nucleus tractus solitarius alone or with sinoaortic denervation in the dog.

Hypertension ; **3**: 112-118.

Ferro CJ , Haynes WG , Hand MF and Webb DJ. (2002).

Forearm vasoconstriction to endothelin-1 is impaired , but constriction to sarafotoxin 6 c and vasodilatation to B Q - 123 unaltered in patient with essential hypertension

Clinical Science ; **103** : 533 - 585.

Fisher DL, Hurwitz B, Jeunemaitre X, Kopkins PN and Williams GH (2002).

Familial aggregation of low-renin hypertension.

Hypertension ; **39(4)**: 914-918.

Fontbonne A, Tchobroutsky G and Eschwege E (1990).

Coronary heart disease mortality risk: Plasma insulin level is a more sensitive marker than hypertension or abnormal glucose tolerance in overweight males.

Int J Obes ; **12**: 557-565.

Fontbonne AM and Eschwege EM (1991).

Insulin and Cardiovascular disease. Paris prespective study.

Diab Care; **14**: 461-469.

Frantze F, Faaren AL, Alfheim I and Nordhei AK (1998).

An enzyme conversion immunoassay for determining total homocysteine in plasma or serum.

Clin Chem ; **44**: 311-316.

Friedwald WT, Levy RI and Frederickson DS (1978).

Estimation of LDL-C without the use of preparative ultracentrifugation.

Clin Chem ; **18**: 499-502.

Fries ED (1990).

Origins and development of antihypertensive treatment. In: Hypertension: Pathophysiology, diagnosis and Management. Laragh JH, Brenner BM (eds). New York Reven , p: 2093-2105.

Fujisaki H, Ito H, Hirata Y, et al. (1995).

Natriuretic peptides inhibit angiotensin II-induced proliferation of rat cardiac fibroblasts by blocking endothelin-I gene expression.

J Clin Invest ; **96**: 1059-1065.

Fujita K, Matsumura Y, Kita S, *et al.* (1995).

Role of endothelin-1 and the ETA receptor in the maintenance of deoxycorticosterone acetate-salt induced hypertension.

Br J Pharmacol ; 114: 925-930.

Fujiwara N, Osanai T, Kamada T, Katoh T, Takahashi K, Okumura K (2000).

Study on the relationship between plasma nitrite and nitrate level and salt sensitivity in human hypertension: modulation of nitric oxide synthesis by salt intake.

Circulation;101: 856-861.

Fukuroda T, Fujikawa T, Ozaki S, *et al.* (1994).

Clearance of circulating endothelin-1 by ER-B receptors in rats.

Biochem Biophys Res Commun ; 119: 1461-1465.

Fukuroda T, Ozaki S, Ihara M, *et al.* (1994).

Synergistic inhibition by BQ-123 and BQ-788 of endothelin-1-induced contractions of the rabbit pulmonary artery.

Br J Pharmacol ; 113: 336-338.

Garg UC, Hassid A (1989).

Nitric oxide-generating vasodilators and 8-bromocyclic guanosine monophosphate inhibit mitogenesis and proliferation of cultured rat vascular smooth muscle cells.

J Clin Invest ; 83:1774-1777.

Garrow TA (1996).

Purification, kinetics properties, and cDNA cloning of mammalian betaine-homocysteine methyltransferase.

J Biol Chem ; **271**: 22831-22838.

Glasser SP (2001).

Hypertension syndrome and cardiovascular events.

Post Grad Med ; **110(5)**: 29-36.

Goddard J and Webb DJ (2000).

Plasma endothelin concentration in hypertension.

J Cardiovasc Pharmacol ; **35(4)**: 525-531.

Goldstein DS (1983).

Plasma catecholamines and hypertension: An analytic review.

Hypertension ; **5**: 86-99.

Goldstein DS, Levinson PD and Zimlichman R (1985).

Clonidine suppression testing in essential hypertension.

Ann Intern Med ; **102**: 42-48.

Goonasekera CD , Shah V , Rees DD and Dillon MJ (2000).

Vascular endothelial cell activation associated with increased plasma asymmetric dimethyl arginine in children and young adults with hypertension : a basis for atheroma.

Blood press ; **9 (1)** : 2-3.

Gosse P, Jullien V, Jarnier P and Clementy J (1999).

Echocardiographic definition of left ventricular hypertrophy in hypertension.

J Hum Hypertens ; 13(8): 503-509.

Goyette P, Summer JS, Milos R, et al. (1994).

Human methylenetetrahydrofolate reductase: Isolation of cDNA, mapping and mutation identification.

Nature Genet ; 7: 195-200.

Graettinger WF, Cheung DG and Weber MA (1990).

Wave configuration as can indicator of the relationship between blood pressure and the heart in normotensive adloscents.

Chest ; 97: 896-900.

Graettinger WF, Neutel JM, Smith DH and Weber MA (1991).

Left ventricular diastolic filling alterations in normotensive young adults with a family history of systemic hypertension.

Am J Cardiol ; 68: 51-56.

Graham IM, Slott JH, Ulane RE et al. (1997).

Plasma Hcy as a risk factor for vascular disease.

JAMA; 277(22): 1775 - 1781.

Granata AR, Ruggiero DA and Park DH (1985).

Brain stem area with C1 epinephrine neurons mediates baroreflex vasodepressor reponses.

Am J Physiol ; 248: 547-567.

Grantham JA, Schirger JA, Williamson EE, *et al.* (1998).

Enhanced endothelin-converting enzyme immunoreactivity in early atherosclerosis.

J Cardiovasc Pharmacol ; 31: 22-26.

Grubbs AL and Ergul A. (2001).

A review of endothelin and hypertension in African American individuals.

Ethn Dis ; 11(4): 741 - 748.

Guimaraes CL , Trentin PG and Rae GA. (2002).

Endothelin ETB receptor mediated mechanisms.

Clinical science ; 103 : 3405 - 3445.

Guo GB and Abboud FM (1984).

Angiotensin II attenuates baroreflex control of heart rate and sympathetic activity.

Am J Physiol ; 246: 80-89.

Gupta AK, Clark RV and Kirchner KA (1992).

Effects of insulin on renal sodium excretion.

Hypertension ; 19: 178-182.

Guyton AC, Hall JE, Coleman TG, *et al.* (1995).

The dominant role of the kidneys in long-term arterial pressure regulation in normal and hypertensive state. In: Hypertension: Pathophysiology, Diagnosis, and Management. Laragh JH, Brenner BM (eds.). 2nd Raven Press, New York, p: 1311-1326.

Hand MF, Haynes WG and Webb DJ(1998).

Hemodialysis and L-arginine, but not D-arginine, correct renal failure-associated endothelial dysfunction.

Kidney Int ; **53**:1068–1077.

Handler CE, Child A and Light ND (1985).

Mitral valve prolapse, aortic compliance and skin collagen in joint hypermobility syndrome.

Br Heart J ; **54**: 501-508.

Haynes WG and Webb DJ (1993).

Endothelium-dependent modulation of responses to endothelin-1 in human veins.

Clin Sci ; **84**: 427.

Haynes WG, Davenport AP and Webb DJ (1993).

Endothelium: Progress in pharmacology and physiology.

Trends Pharmacol Sci ; **14**: 225-228.

Haynes WF, Strachan FE and Webb DJ (1995).

Endothelin ET-A and ET-B receptors mediate vasoconstriction of human resistance and capacitance vessels in vivo.

Circulation ; **92**: 357-363.

Hingorani AD, Liang CF, Fatibene J, Lyon A, Monteith S, Parsons A, Haydock S, Hopper RV, Stephens NG, O'Shaughnessy KM and Brown MJ (1999).

A common variant of the endothelial nitric oxide synthase (Glu298" src="adma does_files/rarr.gif" border=0Asp) is a major risk factor for coronary artery disease in the UK.

Circulation ; **100**:1515–1520.

Hocher B, Rohmeiss P, Thone-Reineke C, et al. (1997).

Endothelin-1 transgenic mice develop glomerulosclerosis, interstitial fibrosis and renal cysts, but not hypertension.

J Clin Invest ; **99**: 1380-1389.

Hoffman A, Boomsma F and Schalekamp MADH (1979).

Raised blood pressure and plasma noradrenaline from an open population.

BMJ ; **1**: 1536-1979.

Hoffman A, Grossman E, Goldstein DS, Gill JR and Keiser MR (1997).

Urinary excretion rate of endothelin-1 in patients with essential hypertension.

Kid Inter ; **45**: 556-560.

Holzapfel GA , Gasser TC and stadler M. (2002).

A structural model of the viscoelastic behavior of arterial walls.

Euro J of Mechanics ; **21** : 441 - 463 .

Hoogeveen EK, Kostense PJ, Beks BJ et al.(1998).

Hyperhomocystienemia is associated with increased risk of CVD.Arterioscler Thromb Vasc Biol ; **18**:133-138.

Horky K, Kopecha J, Greogorova I and Dvorakova J (1976).

Relationship between renin activity and urinary catecholamine in various types of hypertension.

Endokrinologie ; 67:331-342.

Hua L, Li C, Xia D, Qu P, Zhang W and Feng X (2000).

Relationship between hypertensive LVH and levels of endothelin and nitric oxide.

Hypertens Res ; 23(4): 377-380.

Huang A, Vita JA, Venema RC and Keaney J (2000).

Ascorbic acid enhances endothelial nitric oxide synthase activity by increasing intracellular tetrahydrobiopterin.

J Biol Chem; 275:17399–17406.

Ichihara A, Inscho EW and Imig JD (1997).

Role of the nerves in afferent arteriolar reactivity in angiotensin-induced hypertension.

Hypertension ; 29: 442-449.

Iglarz M and Schiffrin EL (2003).

Role of endothelin - 1 in hypertension .

Curr Hypertens Rep ; 5(2) : 144-148.

Imazu M. (2002) .

.Hypertension and insulin resistance.

Current hypertension reports ; 4 : 477 - 482.

Nephron Exp Nephrol ; 93(3) : 106-111

Iwasaki S, Homma T, Matsuda Y and Kon V (1995).

Endothelin receptor subtype-B mediates autoinduction of endothelin-1 in rat mesangial cells.

J Biol Chem ; 270: 6997-7003.

Izzo IR (1989).

Sympathoadrenal activity, catecholamines, and the pathogenesis of vasculopathic hypertensive target organ damage.

Am J Hypertens ; 2: 3055-3125.

Izzo JL (1996).

Biochemical assessment of sympathetic activity. Primer on the autonomic nervous system.

San Diego Academic ; 116-122.

Izzo JL (2000).

Systolic hypertension, arterial stiffness, and vascular damage: role of the renin angiotensin system.

Blood Press Monit ; 5: 257-261.

Izzo JL, Licht MR and Smith RJ (1987).

Chronic effects of vasodilator (pinacidil), alpha adrenergic blockage (Prazocin) and angiotensin converting enzyme inhibition (captopril) in systemic hypertension.

Am J Cardiol ; 60: 303-308.

Jackson RV, Williamson MM and Gordon RD (1983).

Long term minoxidil therapy and renal function, cardiac function, hypertrichosis and blood pressure.

Aust NZ J Med ; **3**: 39-44.

Jain S , Ram H , Kumari S and Khullar M (2003).

Plasma homocysteine levels in patients with essential hypertension and their siblings .

Ren Fail ; **25(2)** : 195-201 .

Jamerson K, Smith SD and Amerena J (1994).

Vasoconstriction norepinephrine causes less forearm insulin resistance than a reflex sympathetic vasoconstriction.

Hypertension ; **23**: 1006-1011.

Jong SC, Stehouwer CD, Mackaay AJ, Van-den-Berg M and Rauwerda JA (1997).

High prevalence of hyperhomocysteinemia and a symptomatic vascular disease in siblings of young patients with vascular disease and hyperhomocysteinemia.

Arterioscler Thromb Vasc Biol ; **17(11)**: 2655-2662.

Joshi S , Gupta S , Tank S *et al.* (2003)

Essential hypertension : antecedents in children.

Indian Pediatr ; **40(1)** : 24-29

Julius S, Brant D and Buda AJ (1989).

Neurogenic pressor episodes fail to cause hypertension, but to induce cardiac hypertrophy.

Hypertension ; 13: 422-429.

Julius S, Krause L and Schork NJ (1991).

Hyperkinetic borderline hypertension in Tecumsh. Michigan

J Hypertens ; 9: 77-84.

Jullien V, Gosse P, Ansoborlo P, Lemetyer P and Clementyl J (1998).

Elevated serum insulin levels in patients with essential hypertension and microalbuminuria.

Hypertension ; 23: 681-687.

Kahleova R , Palyzova D , Zvara K *et al.* (2002).

Essential hypertension in adolescents association with insulin resistance and with metabolism of homocysteine and vitamins.

Am J Hypertens ; 15 (10) : 857 - 864.

Kakimoto Y, Akazawa S (1970).

Isolation and identification of N^G , N^G -, and N^G , N^G -dimethyl-arginine, N -mono-, di-, and trimethyllysine, and glucosylgalactosyl-, and galactosyl-{PRIVATE "TYPE=PICT;ALT={delta}"}-hydroxylysine from human urine.

J Biol Chem ; 245:5751-5758.

Kang SS , Wang PWK and Malinow. (1992).

Hyperhomocysteinemia as a risk factor for occlusive vascular disease.

Ann Rev Nutr ; 12 : 279 - 298.

Kang SS and Wang PWK (1996).

Genetic and nongentic factors for moderate hyperhomocyst(e)inemia.

Atherosclerosis ;119: 135-138.

Kannel WB, Brand N and Skinner JJ (1987).

The relation of adiposity to blood pressure and development of hypertension.

Ann Intern Med ; 67: 48.

Kapellrud H, Bangstad HJ, Berg K and Hanssen KF (1991).

Serum lipoprotein (a) concentrations in IDDM patients with microalbuminuria.

Brit Med J ; 303: 675-678.

Kawarada A, Shimazu H, Yamokoshi K and Kamiyoo A (1986).

Non invasive automatic measurement of arterial elasticity in human fingers and rabbit forelegs using photoelectric plethysmography.

Med Biol Eng Comput ; 24: 591-596.

Keane WF (1994).

Lipids and the kidney.

Kidney Int ; 46: 910-920.

Keane WF, Mulcahy WS, Kasiske BL, Kim Y and O'Donnell MP(1991)

Hyperlipidemia and progressive renal disease.

Kidney Int ; 39: 541-548.

Kelm M, Schafer S, Minger S, Heydthausen M and Strauer BE (1996).

Left ventricular mass is linked to cardiac noradrenaline in normtensive and hypertensive patients.

J Hypertens ; 14(11): 1357-1364.

Kielstein J, Böger R, Bode-Böger S, Schäffer J, Barbey M, Koch K and Frölich J(1999).

Asymmetric dimethylarginine plasma concentrations differ in patients with end-stage renal disease: relationship to treatment method and atherosclerotic disease.

J Am Soc Nephrol; 10:594–600.

Kielstein J T, Bode-Boger S M, Frolich J C, Ritz E, Haller H, and Fliser D (2003).

Asymmetric Dimethylarginine, Blood Pressure, and Renal Perfusion in Elderly Subjects

Circulation ; 107(14): 1891 - 1895.

Kirby DA, pinto JMB and Lown B (1990).

Changes in ventricular fibrillation threshold during the development of perinephritic hypertension.

Am J Hypertens ; 3: 741-748.

Kishore J and Harjai MBBS. (1999).

Potential new cardiovascular risk factors :Left ventricular hypertrophy, homocysteine , triglycerides , oxidative stress and fibrinogen.

Ann Intern Med ; 131(5) : 376 - 386.

Kissebah AH, Vydelingum N and Murraray R (1982).

Relation of body fat distribution to metabolic complication to obesity.

J Clin Endocrinal Metab ; 54: 254.

- Kitamura K, Tanaka T, Kato J, Ogawa T, Eto T and Tanaka K (1989).**
Immunoreactive endothelin in rat kidney inner medulla: markedly decreased
in spontaneously hypertensive rats.
Biochem Biophys Res Commun ; **162**: 38-44.
- Klahr SD, Schreiner G and Ichikawa I (1988).**
The progression of renal disease.
N Engl J Med ; **318**: 1657-1666.
- Koda M, Sasaguri M, Miura S, Tashiro E and Arakawa K (1998).**
Plasma renin activity could be a useful predictor of LVH in patients with
essential hypertension.
J Hum Hypertens ; **12(7)**: 455-461.
- Kohno M, Yasunari K and Murakawa K (1990).**
Plasma immunoreactive endothelin-1 in essential hypertension.
Am J Med ; **88**: 614-618.
- Kontis SB (1985).**
Application of microprocessors and doppler ultrasound (Ph. D thesis),
London, University of London.
- Kool MJF, Struijker-Boudier HA and Wijnen JA (1992).**
Effects of diurnal variability and exercise training on properties of large
arteries.
J Hypertens ; **10(6)**: 549-552.

Kosako T, Suzuki N, Matsumoto H, Itoh I, Yasuhara T and Fujino M (1989).

Synthesis of the vasoconstrictor peptide endothelin in kidney cells.

FEBS Lett ; **249**: 42-46.

Krieger DR and Landsberg L (1995).

Obesity and hypertension. New York Raven 2367.

Krum H, Viskoper RJ, Lacourciere Y, *et al.* (1998).

The effect of an endothelin receptor antagonist, bosentan, on blood pressure in patients with essential hypertension.

N Engl J Med ; **338**: 784-790.

Kullo IJ and Burnett JC (1996).

C-type natriuretic peptide: The vascular component of the natriuretic peptide system. In: Sowers JR (ed). Contemporary Endocrinology of the Vasculature.

Humana ; p: 79-93.

Lambert GW. (2000).

Central nervous system norepinephrine metabolism in hypertension..

Current hypertension. reports ; **2** : 302 - 308.

Landsberg L (1994).

Pathophysiology of obesity-related hypertension: Role of insulin and the sympathetic nervous system.

J Cardiovasc Pharmacol ; **23**: 51.

Landsberg L and Young JB (2000).

Physiology and pharmacology of the autonomic nervous system. In: Essential of Internal Medicine. Harison J (eds.), Vol 1. WB Saunders, London, p: 438-450.

Laogun AA (1977).

In vivo arterial compliance in health and disease. Ph.D thesis, London, University of London.

Largh JH (1987).

Role of the renin-angiotensin-aldosterone axis in human hypertensive disorders. In: Prespective in Hypertension, vol 1.

The Kidney in Hypertension. Kaplan NM, Brenner BM, Laragh JH (eds.), Raven Press, New York, p: 3552.

Largh JH and Sealey JE (1991).

Renin-angiotensin-aldosterone system and the renal regulation of sodium, potassium and blood pressure homeostasis. In: Handbook of Physiology, Sect. 8, Renal Physiology, Windhager EE (ed.), Oxford University, New York, p: 1409-1541.

Largh JH and Sealey JE (1995).

Renin system understanding for analysis and treatment of hypertensive patients: A means to quantify the vasoconstrictor elements, diagnose curable renal and adrenal causes, assess risk of cardiovascular morbidity, and find the best-fit drug regimen. In: Hypertension: Pathophysiology, Diagnosis and Management. Laragh JH, Brenner BM (eds.), 2nd edition, Raven Press, New York, p: 1813-1838.

Larsen K (1972).

Determination of serum creatinine by Jaffe's reaction.

Clin Chem Acta ; 41: 209.

Larsson B, Svardsudd K and Welin L (1984).

Abdominal adipose tissue and risk of cardiovascular disease and death.

BJM ; 289: 1257.

Laskey WK, Parker HG, Ferrari GA and Noordergraaf A (1990).

Estimation of total systemic arterial compliance in humans.

J Appl Physiol ; 69: 112-119.

Leadly RJ, Zhu JL and Goetz KL (1991).

Effects of endothelin-1 and sarafotoxin S6b on regional hemodynamics in the conscious dog.

Am J Physiol ; 260: 1210-1217.

Lebovitz HE. (2001).

Insulin resistance definition and consequences.

Exp Clin Endocrinal Diabetes ; 109 : 5135 - 5148.

Leiper JM, Santa Maria J, Chubb A, MacAllister RJ, Charles IG, Whitley GS and Vallance P (1999).

Identification of two human dimethylarginine dimethylaminohydrolases with distinct tissue distributions and homology with microbial arginine deiminases.

Biochem J ; 343(pt 1):209-214.

Leonardis CZD, Mallamaci SPF and Postorino M (1995).

Urinary and plasma endothelin 1 in essential hypertension and in hypertension secondary to renoparenchymal disease.

Nephrol Dial Transplant ; 10: 1320-1323.

Leoncin G , Sacchi G and Manera M . (2002).

Microalbuminuria is an integrated marker of subclinical organ damage in primary hypertension.

J of Hypertension ; 16(6) : 399 - 404.

Lerman A, Click RL, Narr BJ, *et al.* (1991).

Elevation of plasma endothelin associated with systemic hypertension in humans following orthotopic liver transplantation.

Transplantation ; 51: 646-650.

Lerman A, Sandok EK, Hidebrand FL and Burnett JC, (1992).

Inhibition of endothelium-derived relaxing factor enhances endothelin-mediated vasoconstriction.

Circulation ; 85: 1894-1898.

Lever M, Sizeland PCB, Boson LM, Haymer CM, Robson RA and Chambers ST (1994).).

Abnormal glycine betaine content of the blood and urine of diabetic and renal patients.

Clin Chim Acta ; 230: 69-79.

Levy D, Garrison RJ and Savage DD (1990).

Prognostic implication of echocardiograph determined left ventricular mass in the Framingham heart study.

N Engl J Med ; 332: 1561-1566.

Levy D , Savage DD , Garrison RJ *et al.* (1987).

Echo - criteria for left ventricular hypertrophy : The Framingham heart study.

Am J Cardiol ; 59 : 556 - 560.

Le Tourneau T, Van Belle E, Corseaux D, Vallet B, Lebuffe G, Dupuis B, Lablanche JM, McFadden E, Bauters C and Bertrand ME (1999).

Role of nitric oxide in restenosis after experimental balloon angioplasty in the hypercholesterolemic rabbit: effects on neointimal hyperplasia and vascular remodeling.

J Am Coll Cardiol ; 33: 876-882.

Licht MR and Izzo JL (1989).

Humoral effect of norepinephrine on renin release in humans.

Am J Hypertens ; 2: 788-791.

Lim U and Cassano PA . (2002).

Hcy and blood pressure.

Am J of Epidemiology ; 156 : 1105 - 1113.

Louis WJ, Doyle AE and Anavekar S (1973).

Plasma norepinephrine levels in essential hypertension.

N Engl J Med ; 288: 599-601.

Luft FC and Agrawal B (1999).

Microalbuminuria as a predictor factor for cardiovascular disease.

J Cardiol Pharamcol ; **33**(1): S11-S15.

Makenzie IS , Wilkinson IB , Cockcroft JR. (2002).

Assesment of arterial stiffness in clinical practice

J Med ; **95** : 67 - 74.

Mangiapane ML, Skoog KM and Rittenhouse P (1989).

Lesion of the area protrema region attenuates hypertension in spontaneously hypertensive rats.

Citc Res ; **64**: 129-135.

Mansoor MA, Bergmark C, Svardal AM, Lonning PE, and Ueland PM (1995).

Redox status and protein binding of plasma homocysteine and other aminothiols in patients with early-onset peripheral vascular disease: Homocysteine and peripheral vascular disease.

Arterioscler Thromb :**15**: 232–240.

Marsden PA, Brenner Bm and Ballerman BJ (1990).

Mechanism of angiotensin action on vascular smooth muscle, the adrenal, and the kidney.

New York Raven, p: 1247-1272.

Masuda H and Azuma H (2002).

Biological and pathophysiological roles of endogenous methylarginines as inhibitors of nitric oxide synthase .

Nippon Yakurigaku Zasshi . ; **119** (1) : 29-35.

Masuda H, Goto M, Tamaoki S and Azuma H (1999).

Accelerated intimal hyperplasia and increased endogenous inhibitors for NO synthesis in rabbits with alloxan-induced hyperglycaemia.

Br J Pharmacol ;126: 211–218.

Masuo K, Mikami M, Ogihara T and Tuck ML (1998).

Familial hypertension, insulin, sympathetic activity and blood pressure elevation.

Hypertension ; 32(1): 96-100.

Matsuoka H, Itoh S, Kimoto M, Kohno K, Tamai O, Wada Y, Yasukawa H, Iwami G, Okuda S and Imaizumi T (1997).

Asymmetrical dimethylarginine, an endogenous nitric oxide synthase inhibitor, in experimental hypertension.

Hypertension ; 29(pt 2):242–247.

Mclenachan JM and Dargie HJ (1990).

Ventricular arrhythmias in hypertensive left ventricular hypertrophy in relationship to coronary artery disease, left ventricular dysfunction and myocardial fibrosis.

Am J Hypertens ; 3: 735-740.

Meehan WP, Buchanan TA and Hsueh W (1994).

Chronic insulin administration elevates blood pressure in rats.

Hypertension ; 23: 1012-1018.

Mendis S, Athauda SB, Naser M and Takahashi K (1999).

Association between hyperhomocysteinemia and hypertension in Sri-Lanakans.

J Ent Med Res ; 27(1): 38-44.

Messent JW, Elliot TG, Hill RG, Jarret RJ and Keen H (1992).

Prognostic significance of microalbuminuria in IDDM: A twenty-three-year follow up study.

Kidney Intern ; 41: 836-839.

Mettei P , Virdis A , Ghiadoni L *et al.* (1997).

Endothelin function in hypertension..

J of Nephrology ; 10 (4) : 192 - 197.

Miller JW, Nadeau MR, Smit D and Selhub J (1994).

Vitamin B6 deficiency vs folate deficiency: Comparison of responses to methionine loading in rats.

Am J Clin Nutr ; 59: 1033-1039.

Miyazaki H, Matsuoka H, Cooke JP, Usui M, Ueda S, Okuda S and Imaizumi T (1999).

Endogenous nitric oxide synthase inhibitor: a novel marker of atherosclerosis.

Circulation ; 99:1141-1146.

Mogensen CE (1984).

Microalbuminuria predicts clinical proteinuria.

New England Journal of medicine; 310: 356-360.

Moncia G, Carugo S, Grassi G, Lantarotti A, Schiavina R and Sega R (2002).

Prevalence of left ventricular hypertrophy in hypertensive patients without and with blood pressure control. Data from PAMELA.

Hypertension ; **39(3)**: 744-749.

Moreau P and Dao HH (2001).

An update on the status of endothelin receptor antagonists for hypertension.

Expert Opin Investig Drugs ; **10(11)**: 1937-1946.

Moreland S, McMullen DM, Delaney CL, et al. (1992).

Venous smooth muscle contains vasoconstrictor ET-B-like receptors.

Biophys Res Commun ; **184**: 100-106.

Mudd SH, Levy HL, and Scovby F. (1995).

The metabolic basis of inherited diseases.

New York Mac Graw Hill ; 693-734.

Mulvaney MJ (1995).

Structural changes in the resistance vessels in human hypertension. In: Hypertension: Pathophysiology, Diagnosis, and Management. Laragh JH, Brenner BM (eds.). 2nd Raven Press, New York, p: 503-513.

Murad F(1996).

The 1996 Albert Lasker Medical Research Awards: signal transduction using nitric oxide and cyclic guanosine monophosphate.

JAMA ; **276**: 1189-1192.

Nabel EG, Selwyn AP and Ganz P(1990).

Large coronary arteries in humans are responsive to changing blood flow: an endothelium-dependent mechanism that fails in patients with atherosclerosis.
J Am Coll Cardiol ; **16**:349–356.

Najbauer J, Johnson BA, Young AL and Aswad DW(1993).

Peptides with sequences similar to glycine, arginine-rich motifs in proteins interacting with RNA are efficiently recognized by methyltransferase(s) modifying arginine in numerous proteins.

J Biol Chem ; **268**:10501–10509.

Natali A, Taddei S, Galvan AG, Baldi S, Virdis A, Salvetti A and Ferrannini E (1997).

Insulin sensitivity, vascular reactivity, and clamp-induced vasodilatation in essential hypertension.

Circulation ; **96**: 849-855.

Neutel JM, Smith DH and Graettinger WF (1992).

Heredity and hypertension: Impact on metabolic characteristics.
Am Heart J ; **124**: 435-440.

Neutel JM, Smith DH and Graettinger WF (1993).

Metabolic characteristics of hypertension.

Am Heart J ; **126**: 924-929.

Neutel JM and Smith DH (1995).

Metabolic and cardiovascular characteristics of hypertension.
Cardiol Clin ; **13**(4): 539-547.

Neutel JM, Smith DHG and Weber MA (1999).

Is high blood pressure a late manifestation of the hypertension syndrome.

Am J Hypertens ; 12: 2155-2235.

Neutel JM (2000).

Why lowering blood pressure is not enough: The hypertension syndrome and the clinical context of cardiovascular risk reduction.

Heart Dis ; 2(5): 370-374.

Neutel JM. (2001).

Beyond the sphygmomanometric numbers :Hypertension as a syndrome.

Am J Hypertens ; 14 : 250 S - 257S.

Nicholson JP, Resinck LM and Laragh JH (1989).

Hydrochlorothiazide is not additive to verapamil in treating essential hypertension.

Arch Intern Med ; 149: 125.

Nichols WW and O'Rourke MF (1998).

McDonalds blood flow in arteries. London, Arnold, 4th edition. p: 134-148.

Niranjan V, Telemaque S, deWit D, *et al.* (1996).

Systemic hypertension induced by hepatic overexpression of human preproendothelin-1 in rats.

J clin Invest ; 98: 2364-2372.

Numaguchi K, Egashira K, Takemoto M, Kadokami T, Shimokawa H, Sueishi K and Takeshita A (1995).

Chronic inhibition of nitric oxide synthesis causes coronary microvascular remodeling in rats. *Hypertension* ; 26(pt 1):957–962.

Nygard O , Vollset SE , Refsum H , *et al.*(1995).

Plasma homocysteine and cardiovascular risk profile. The Hardaland Homocysteine Study.

JAMA ; 274 :1526 - 1533.

Oemar BS, Tschudi MR, Godoy N, Brovkovich V, Malinski T and Luscher TF (1998).

Reduced endothelial nitric oxide synthase expression and production in human atherosclerosis.

Circulation ; 97:2494–2498.

Ogawa T, Kimoto M and Sasaoka K (1987).

Occurrence of a new enzyme catalyzing the direct conversion of $N^G.N^G$ -dimethyl-L-arginine to L-citrulline in rats.

Biochem Biophys Res Commun ; 148:671–677.

Ogihara T, Rakugi H, Mikami H and Masuo K (1995).

Enhancement of insulin sensitivity by troglitazone lowers blood pressure in diabetic hypertensives

Am J Hypertens ; 8: 316-320.

Ohara Y, Petersen TE and Harrison DG (1993).

Hypercholesterolemia increases endothelial superoxide anion production.
J Clin Invest ; 91:2546–2551.

Opsahl JA, Abraham PA, Halestenson CE and Keane WF (1988).

Correlation of office and ambulatory blood pressure measurements with urinary albumin and N-acetyl-beta-D-glucosaminidase excretions in essential hypertension.

Am J Hypertens ; 1: 5117-5120.

O'rourke MF (1985).

Basic concepts for the understanding of large arteries in hypertension.

J Cardiovasc Pharmacol ; 7: 14-21.

O'rourke MF (1992).

Arterial function in health and disease. Edinberg-London and New-York:
Churchill Livingstone, 23-66; 94-169; 185-232.

O'rourke MF. (2002).

From theory into practice : Arterial Hemodynamics in clinical hypertension.

J Hypertension ; 20 : 1901 - 1915.

O'rourke Mf , staessen JA , Vlachopoulou C *et al.* (2002).

Clinical application of arterial stiffness.

AM J Hypertens ; 15 : 426 - 444.

Orsio S, Benigni A, Bruzzi I, *et al.* (1993).

Renal endothelin gene expression is increased in remnant kidney and correlates with disease progression.

Kid Int ; 43: 354-358.

Osei K (1999).

Insulin resistance and systemic hypertension.

Am J Cardiol ; **84**: 33-36.

Osei K and Schuster DP (1994).

Ethnic differences in secretion, sensitivity, and hepatic extraction of insulin in black and white Americans.

Diabetes Med ; **11**: 755-762.

Osei K and Schuster DP (1996).

Effects of race and ethnicity on insulin sensitivity, blood pressure and heart rate in 3 ethnic populations.

Am J Hypertens ; **9**: 1157-1164.

Parums DV (1996).

Vascular disease.

Essential Clinical Pathol ; **2**: 272-274.

Parving HH, Jensen HE, Mogenson CE and Evrine PE (1974).

Increased urinary albumin excretion rate in benign essential hypertension.

Lancet ; **1**: 1190-1192.

Pedrinelli R, Bello VD, Catapano G and Ferrari M (1993).

Microalbuminuria is a marker of left ventricular hypertrophy.

Arteriosclerosis Thrombosis ; **13**: 900-906.

Pedrinelli R, Dell'Omo G, Di-Bello V, Pontremoli R and Mariani M (2002).

Microalbuminuria, an integrated marker of cardiovascular risk in essential hypertension.

J Hum Hypertens ;16(2): 79-89.

Pereira MA , Jacobs DR , Horn LV *et al.* (2002).

Dairy consumption , obesity and insulin resistance in young adults.

JAMA ; 287 : 2081 - 2089.

Pernow J, Kaijser J, Lannart L, *et al.* (1996).

Comparable potent coronary constrictor effects of endothelin-1 and big endothelin-1 in humans.

Circulation ; 94: 2077-2082.

Peterson GH, Jenser RE and Parnell MI (1960).

Mechanical properties in vivo.

Circ Res ; 8: 622-639.

Petrie A and Sabin C.(2000).

Medical statistics at a glance. Blackwell Science Limited , oxford.

Petrie JR, Ueda S and Elliott HL (1996).

Endothelial nitric oxide production and insulin sensitivity: A physiological link with implications for pathogenesis of cardiovascular disease.

Circulation ; 93: 1331-1333.

Petrova TV , Stryuk RI , Babrovnisky IP *et al.* (2001).

Relation between excess body weight , hypertension , hyperinsulinemia and impaired glucose tolerance.

Egypt Heart J ; **16** : 72 - 76

Pontremoli R, Revera M, Bezante GP, Viazzi F and Deferrari G (1999).

Left ventricular geometry and function in patients with essential hypertension and microalbuminuria.

J Hypertens ; **17(7)**: 993-1000.

Pratt JH (2000).

Low renin hypertension: more common than we think.

Cardiol Rev ; **8(4)**: 202-206.

Pyorala K, Savolainen E and Kaukola S (1985).

Plasma insulin as coronary heart disease risk factor.

Acta Med Scand ; **701**: 38-52.

Rabelink TJ, Kaasssjager KAH, Boer P, *et al.* (1995).

Effects of endothelin-1 on renal function in humans: Implications for physiology and pathophysiology.

Kidney Int ; **46**: 376-378.

Raji A , William GH , Jeunemaitre X *et al.*(2001).

Insulin resistace in hypertension.

J Hypertens ; **19 (1)** : 99 - 105.

Ramasamy S, Drummond GR, Ahn J, Storek M, Pohl J, Parthasarathy S and Harrison DG (1999).

Modulation of expression of endothelial nitric oxide synthase by nordihydroguaiaretic acid, a phenolic antioxidant in cultured endothelial cells.

Mol Pharmacol ; 56:116–123.

Randall OS (1984).

Effect of arterial compliance on systolic blood pressure and cardiac function.

Clin Exp Hypertens ; 4: 1045.

Rasmussen TE, Jougasaki M, Supaporn T, *et al.* (1998).

Cardiovascular actions of ET-B activation in vivo and modulation by receptor antagonism. hypertension.

Am J Hypertens ; 12(3): 315-331.

Reaven GM and Laws A (1994).

Insulin resistance, compensatory hyperinsulinemia, and coronary heart disease.

Diabetologia ; 37: 948-952.

Reaven GM, Lithell H and Landsberg L (1996).

Hypertension and associated metabolic abnormalities. The role of insulin resistance and the sympathoadrenal system.

N Engl J Med ; 334: 374-381.

Refsum H , Ueland P , Nygard O and Vollset SE.(1998).

Homocysteine and CVD.

Annu Rev Med ; 49 : 31 - 62.

Reisnick LM, Nicholson JP, Laragh JH (1985).

Alterations in calcium metabolism mediate dietary salt sensitivity in essential hypertension.

Trans Assoc Am Physicians ; 98: 313.

Remuzzi G , Perico N and Benigni A. (2002).

New therapeutics that antagonize endothelin.

Net Rev Drug Discov ; I(12) : 986 - 1001.

Robertson WB and Strong JP (1968).

Atherosclerosis in persons with hypertension and diabetes mellitus.

Lab Invest ; 18: 538-551.

Robinson K , Arheart K , Refsum H *et al.*(1998).

Low circulating folate and vitamin B6 concentrations : Risk factor for stroke , peripheral vascular disease and coronary disease.

Circulation ; 97 : 437 - 443.

Rocchini AP (2000).

Obesity hypertension, salt sensitivity and insulin resistance.

Nutr Metab Cardiovasc Dis ; 10(5): 287-299.

Rocchini AP, Mao HZ, Bobu K and Rocchini AJ (1998).

Clonidine prevents insulin resistance and hypertension in obese dogs.

Hypertension ; 32: 592.

- Rodico JL, Campo C and Ruilope LM (1998).**
Microalbuminuria in essential hypertension.
Kid Int ; 68: 551-554.
- Rosa EC , Moises VA , Sesso RC *et al.* (2001).**
Cardiac geometry in essential hypertension.
Arq Bress Cardiol ; 76(5) : 362 - 368.
- Rosa TT and Palatini P (2000).**
Clinical value of microalbuminuria in hypertension.
J Hypertens ; 18(6): 645-654.
- Ruige JB, Assendelft WJ and Dekker JM (1998).**
Insulin and risk of cardiovascular disease.
Circulation ; 97:996-1001.
- Ruilope LM, Campo C, Rodriguez-Artalejo F and Rodicio JL (1996).**
Blood pressure and renal function.
J Hypertens ; 14: 1259-1263.
- Ruiz-openza N, Hirayama K, Akimoto K and Herrera VLM (1998).**
Molecular characterization of a dual endothelin-1/angiotensin II receptor.
Mol Med ; 4: 96-108.
- Saad MF, Lillioja S, Nyomba BL, Castillo C and Bennett PH (1991).**
Racial differences in the relation between blood pressure and insulin resistance.
N Engl J Med ; 324: 733-739.

Safar ME, Laurent S and London GM (1987).

Structural and functional modifications of peripheral large arteries in hypertensive patients.

J Clin Hypertens ; **3**: 360-367.

Safar ME and O'Rourke MF (1994).

The arterial system in hypertension . Dordrecht, Kluwer. p: 134-148.

Sakai S, Miyauchi T, Kobayashi M, *et al.* (1996).

Inhibition of myocardial endothelin pathway improves long-term survival in heart failure.

Nature ; **384**: 353-355.

Sakai S, Miyauchi T, Saruta T, *et al.* (1996).

Endogenous endothelin-1 participates in the maintenance of cardiac function in rats with congestive heart failure.

Circulation ; **93**: 1214- 1222.

Sakamoto H, Sasaki S, Hirata Y, Imai T, Ando K, Ida T, Sakurai T and Marumo F (1990).

Production of endothelin-1 by rat cultured mesangial cells.

Biochem Biophys Res Commun ; **169**: 462-468.

Salomaa V, Riley W, Kark JD, Nardo C and Folsom AR (1995).

NIDDM and fasting glucose and insulin concentrations are associated with arterial stiffness indexes.

Circulation ; **91**: 1432-1443.

Samuelsson OG, Wilhelmsen L, Anderson OK, *et al.* (1987).

Cardiovascular morbidity in relation to change in blood pressure and serum cholesterol levels in treated hypertension.

JAMA ; 258: 1768-1776..

Sander GE and Giles TD (2002).

Hypertension and lipids : Lipids factors in hypertension syndrome. *Current hypertension reports* ; 4 : 458 - 463.

Savage DD, Drayer JIM, Hernery WL, Mathews EC, Ware JH, Gardin JM and Laragh JH (1979).

Echocardiographic assessment of cardiac anatomy and function in hypertensive subjects.

Circulation ; 59: 623-632.

Scannapieco G, Pauletto P and Pagnan A (1988).

Lipoprotein binding to cultured aortic smooth muscle cells from normotensive and hypertensive rats.

J Hypertens ; 6 (4): S269-S271.

Schächinger V, Britten MB and Zeiher AM (2000).

Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease.

Circulation ; 101:1899-1906.

Schiffrin EL (1998).

Endothelin and endothelin antagonists in hypertension.

J Hypertens ; 16: 1891-1895.

Schiffrin EL (2001).

Role of endothelin in hypertension and vascular disease.

Am J Hypertens ; 14 (6 pt2) : 833 - 895.

Schiffrin E and Thibault G (1991).

Plasma endothelin in human hypertension.

Am J Hypertens ; 4: 303-308.

Schilleci G , Vaudo G , Pasqualini L *et al.* (2002).

Left ventricular mass and systolic dysfunction in essential hypertension.

J of Human Hypertens ; 16 (2) :117 - 122.

Schaefer RL and Anderson RB. (1989).

Student edition of minitab , statistical softwar copy write wesley publishing
company Inc. London New York Paris

Schmieder RE, Veelken R, Gatzka CD and Ruddel H (1994).

Predictors for hypertensive nephropathy.

J Hypertens ;13: 357-365.

Schmieder RE, Veelken R and Schobel H (1997).

Glomerular hyperfiltration during sympathetic nervous system activation in
early essential hypertension.

J Am Soc Nephrol ; 8: 893-900.

Schwartz SM, deBlois D and O'Brein ERM (1995).

The intima: Soil for atherosclerosis and restenosis.

Cir Res ; 77: 445-465.

Sealey JA. (1991).

Plasma renin activity and plasma prorenin assays.

Clin. Chem ; 37/10 (8) : 1811 - 1819 .

Sealey JE and Largh JH (1995).

The renin-angiotensin-aldosterone system for the normal regulation of blood pressure and sodium and potassium homeostasis. In: Hypertension: Pathophysiology, Diagnosis, and Mangement. Laragh JH, Brenner BM (eds.). 2nd Raven Press, New York, p: 1763-1797.

Sen S (1983).

Regression of cardiac hypertrophy: Experimental animal model.

Am J Med ; 75: 87-93.

Sharabi Y, Boolman R, Rosenthal T, Grossman E and Sela B (1999).

Homocysteine levels in hypertensive patients with history of cardiovascular events.

Am J Hypertens ; 12(8): 766-771.

Silvestre JS, Robert V, Heymes C, *et al.* (1998).

Myocardial production of aldosterone and corticosterone in the rat.

J Biol Chem ; 273: 4883-4891.

Simonson MS and Herman WH (1993).

Protein kinase C and protein tyrosine kinase activity contribute to mitogenic signaling by endothelin-1: Cross-talk between G protein-coupled receptors and pp60c-src.

J Biol Chem ; 268: 9347-9357.

Skvorak JP, Nazian SJ and Dietz JR (1995).

Endothelin acts as a paracrine regulator of stretch-induced arterial natriuretic peptide release.

Am J Physiol ; 269: 1093-1098.

Stamler R, Stamler J and Riedlinger WF (1978).

Weight and blood pressure findings in hypertension screening of 1 million Americans.

JAMA ; 240: 1607-1610.

Stampfer MJ, Malinow MR, Willet WC *et al.*(1992).

A prospective study of plasma homocysteine and risk of myocardial infarction in physicians

JAMA ; 268: 877 - 881.

Stampfer MJ and Malinow MR (1995).

Can lowering homocystein levels reduces cardiovascular risk?

N Engl J Med ; 332: 328-329.

Stella ML, Failla M, Mangoni AA and Mancia G (1998).

Effects of isolated systolic hypertension and essential hypertension on large and middle-sized artery compliance.

Blood Press ; 7: 96-102.

Stem J and Lewis WHP (1957).

Determination of calcuim in blood by colorimetric method.

Clin Chem Acta ; 2: 576

Stergiopoulos N, Meister JJ and Westerhof N (1994).

Simple and accurate way for estimating total and segmental arterial compliance and pulse pressure method.

Ann Biomed Eng ; 22: 392-397.

Stien EA. (2002).

Managing dyslipidemia in the high risk patients.

Am J cardiol ; 89 (supp 1) : 50c - 57c.

Stout RW (1990).

Insulin and atheroma.

Diabetes Care ; 13: 631-654.

Summer MJ, Cannon TR, Munding JW, *et al.* (1992).

Endothelin ETA and ETB receptors mediate vascular smooth muscle cell contraction.

Br J Pharmacol ; 107: 858.

Suwaidi JA, Hamasaki S, Higano ST, Nishimura RA, Holmes DR Jr and Lerman A (2000).

Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction.

Circulation ; 101:948-954.

Suzuki N, Miyauchi J, Tomobe Y *et al.* (1990).

Plasma concentration of endothelin -1 in spontaneously hypertensive rats and in DOCA - salt hypertensive rats .

Biochem Biophys Res Commun ; 167: 941-947.

Taddei S, Virdis A and Mattei P (1992).

Endothelium-dependent forearm vasodilatation is reduced in normotensive subjects with familial history of hypertension.

J Cardiovasc Pharmacol ; **20**: 5193-5195.

Takkyuddin MA, Cervenka JH and Hsaio RJ (1990).

Chromogranin A storage and release in hypertension.

Hypertension ;**15**: 237-246.

Tang J, Frankel A, Cook RJ, Kim S, Paik WK, Williams KR, Clarke S and Herschman HR (2000).

PRMT1 is the predominant type I protein arginine methyltransferase in mammalian cells.

J Biol Chem ; **275**:7723–7730.

Thakur V, Richards R and Reisin E (2001).

Obesity, hypertension, and the heart.

Am J Med Sci ; **321**(4): 242-248.

Tietz NW (1976).

Fundamentals of clinical chemistry , WB. Saunders co., Philadelphia, PA, page 243.

Toblli JE , Bellid CA , Iavicoli OR *et al.* (2002).

Pulse wave velocity and urinary albumin excretion in hypertensive patients.

Medicina ; **62**(6) : 544-550

Tomura S, Kawada K, Saito K, Lin-Lin Y, Endou K, Hirano C, Yanagi H, Tsuchiya S and Kiyoko S (1999).

Prevalence of microalbuminuria and relationship to the risk of cardiovascular disease in the Japanese population.

Am J Nephrol ; 19: 13-20.

Tonolo G, Ciccarese M, Brizzi P, Puddu L, Secchi G and Maioli M (1997)

Reduction of albumin excretion rate in normotensive term simvastatin treatment.

Diabetes Care ; 20: 1891-1895.

Toto-Moukono J, Achimastos A, Asmar RG, Hugues CJ and Safar ME (1986).

Pulse wave velocity in patients with obesity and hypertension.

Am Heart J ; 12: 136-140.

Tracy RE, Newman WP, Strong JP and Bernson GS (1995).

Histologic features of atherosclerosis and hypertension from autopsies of young individuals in a defined geographic population.

Atherosclerosis ; 116: 163-179.

Tsao P, McEvoy LM, Drexler H, Butcher EC and Cooke JP (1994).

Enhanced endothelial adhesiveness in hypercholesterolemia is attenuated by L-arginine.

Circulation ; 89:2176-2182.

Tsao P, Theilmeier G, Singer AH, Leung LL and Cooke JP (1994).

L-Arginine attenuates platelet reactivity in hypercholesterolemic rabbits.
Arterioscler Thromb ; 14:1529–1533.

Tsioufis C , Stefanadis C , Toutouza M *et al.* (2002).

Microalbuminuria is associated with unfavourable cardiac geometric adaptation in essential hypertension..
J of Hypertens ; 26 (4) : 249 - 254.

Tsioufis C , Tzioumis C , Marinakis N *et al.*(2003).

Microalbuminuria is closely related to impaired arterial elasticity in untreated patients with essential hypertension .

Ueland BM, Refsum H, Stabler SP and Malinow R (1993).

Total homocysteine in plasma or serum : Methods and clinical application.
Clin Chem; 39: 1764-1779.

Uusitupa MI, Niskanen LK and Siitonen O (1990).

Five year incidence of atherosclerotic vascular disease in relation to general risk factors, insulin levels, and abnormalities in lipoprotein composition in NIDD and nondiabetic subjects.
Circulation ; 82: 27-36.

Vague J (1956).

The degree of musculin differentiation of obesities: A factor determinig predisposition to diabetes, atherosclerosis, gout, and uric calculous disease.
Am J Clin Nutr ; 4: 20.

Vallance P, Leone A, Calver A, Collier J and Moncada S (1992).

Accumulation of an endogenous inhibitor of nitric oxide synthesis in chronic renal failure.

Lancet ; **339**:572–575.

Van Zweiten PA, van Meel JC and Timmermans PB (1983).

Pharmacology of calcium entry blockers' interaction with vascular alpha-adrenoceptors.

Hypertension ; **5**: 118.

Vasquez-Vivar J, Kalyanaraman B, Martasek P, Hogg N, Masters BS, Karoui H, Tordo P and Pritchard KA Jr (1998).

Superoxide generation by endothelial nitric oxide synthase: the influence of cofactors.

Proc Natl Acad Sci U S A ; **95**:9220–9225.

Verhoef P, Hennekens CH, Malinow MR *et al.* (1994).

A prespective study of plasma Hcy and risk of ischemic stroke. *Stroke* ; **25** : 1924 - 1930.

Verma S, Yao L, Dumont AS and Meneill JH (2000).

Metaformin treatment corrects insulin vascular resistance in hypertension.

J Hypertens ; **18**(10): 1445-1450.

Vlachakis ND and Aledort L (1980).

Hypertension and propranolol therapy: Effect on blood pressure, plasma catecholamines, and platelet aggregation.

Am J Cardiol ; **45**: 321-325.

Wachtell K , Palmieri V , Olsen MH *et al.* (2002).

Urine albumin / creatinine ratio and echocardiographic left ventricular structure and function in hypertensive patients.

Am Heart J ; 143 (2) : 319 - 326.

Wagner OF, Christ G, Wojta J, *et al.* (1992).

Polar secretion of endothelin-1 by cultured endothelial cells.

J Biol Chem ; 267: 16066-16068.

Wang X, Feuerstein GZ and Ohlstein EH (1995).

Temporal expression of ECE-1, ET-1, ET-3, ETA, ETB receptor mRNAs after balloon angioplasty in the rat.

J Cardiovasc Pharmacol ; 26: 22-25.

Ward KD, Sparrow D and Vokonas PS (1994).

The relationships of abdominal obesity, hyperinsulinemia and saturated fat intake to serum lipid levels.

Int J Obese ; 18: 137-144.

Warram JH, Martin BC and Krolewski AS (1990).

Slow glucose removal rate and hyperinsulinemia precede the development of type II diabetes in the offspring of diabetic parents.

Ann Intern Med ; 113: 909-915.

Watanabe K, Sekiya M, Tsuruoka T, Funada J and Kameoka H (1999).

Effect of insulin resistance on left ventricular hypertrophy and dysfunction in essential hypertension.

J Hypertens ; 17(8): 1153-1160.

Weber MA, Drayer JM and Baird WM (1986).

Echocardiographic evaluation of left ventricular hypertrophy.

J Cardiovasc Pharmacol; 8: 561-569.

Weber MA, Smith DHG, Neutel JM and Greattinger WF (1991).

Cardiovascular and metabolic characteristic of hypertension.

Am J Med ; 21(1A): 45-105.

Weber MA, Smith DM and Neutel JM (2000).

Differing roles of BMI and the renin angiotensin system in mediating the hypertension syndrome.

Am J Nephrol ; 20(3): 169-174.

Wei CM, Lerman A, Rodeheffer R, *et al.* (1994).

Endothelin in human congestive heart failure.

Circulation ; 89: 1580-1586.

Wilcken DEL and Wilcken B (1976).

The pathogenesis of coronary artery disease. A possible role for methionine metabolism .

J Clin Invest ; 57 : 1079-1082.

Williams PR, Hunt SC, Hasstedt SJ, *et al.* (1990).

Multigenic human hypertension: Evidence for subtypes and hope for haplotypes

Am J Hypertens; 8: 539-546.

Wold DS , Law M , Morris G *et al.* (2003).

Falate and risk of cardiovascular disease.

BMJ ; 326(7397) : 1035-1039.

Wright JS, Cruickshank JK, Kontis S, Caroline D and Gosling RG (1990).

Aortic compliance measured by noninvasive doppler ultrasound of a method and its reproducibility.

Clin Science, **78**: 463-468.

Yanagisawa M, Kunhara H, Kimura S, et al. (1988).

A novel potent vasoconstrictor peptide produced by vascular endothelial cells.

Nature ; **332**: 411-415.

Yang Z, Richard V, von Segesser L, et al. (1990).

Threshold concentrations of endothelin-1 potentiate contractions to norepinephrine and serotonin in human arteries. A new mechanism for vasospasm?

Circulation ; **82**: 188-195.

Yin FCP, Liu Z and Brin KP (1987).

Estimation of arterial compliance. In: Ventricular/Vascular Coupling. Yin FCP (ed.), New York, Springer, Verlage.

Zimmerman BG (1981).

Adrenegic facilitation by angiotensin: Does it serve a physiological function?

Clin Sci ; **60**: 343-348.

Zizek B and Poredos P (2001).

Insulin resistance to endothelial dysfunction in hypertensive patients and in normotensive offspring of subjects with essential hypertension.

J Intern Med ; **249**(2): 189-197.