



REFERENCES

- Abbasi, F.; Brown, B.W.; Lamendola, C. et al. (2002):** Relationship between obesity, insulin resistance, and coronary heart disease risk. *J Am Coll Cardiol* 40:937-943
- Ahima, R.S.; Saper, C.B.; Flier, J.S. and Elmquist, J.K. (2000):** Leptin regulation of neuroendocrine systems. *Front Neuroendocrinology*, 21, 263– 307.
- Ailhaud, G. (1996):** Early adipocyte differentiation. *Biochem. Soc. Trans.* 24: 400-402.
- Alpert, M.A.(2001):** Obesity cardiomyopathy; pathophysiology and evolution of the clinical syndrome. *Am J Med Sci.* 321: 225–236.
- Altioik, S.; M. xu, and Spiegelman, B. M. (1998):** PPARgamma induces cell cycle withdrawal: inhibition of E2F/DP DNA-binding activity via down-regulation of PP2A. *Genes Dev.* 11: 1987-1998
- Aoki, K.; Homma, M.; Hirano, T.; Oka, K.; Satoh, S.; Mukasa, K.; Ito, S. and Sekihara, H. (2001):** mRNA and enzyme activity of hepatic 11 β -hydroxysteroid dehydrogenase type 1 are elevated in C57BL/KsJ-db/db mice. *Life Science*, 69, 2543– 2549.
- Aruna, B.; Ghosh, S.; Singh, A.K.; Mande, S.C.; Srinivas, V.; Chauhan, R.; et al.(2003):** Human recombinant resistin protein displays a tendency to aggregate by forming intermolecular disulfide linkages. *Biochemistry* 42:10554-10559.
- Attar, E. and Bulun, S.E. (2006):** Aromatase and other steroidogenic genes in endometriosis: translational aspects *oxford j* 12:49- 56
- Aziz, N.M. (2004)** Visceral Obesity and Breast Cancer Risk, *Cancer* 74, 632–639.



- Bajaj, M.; Suraamornkul, S.; Hardies, L.J.; Pratipanawatr, T. and DeFronzo, R.A. (2004):** Plasma resistin concentration, hepatic fat content, and hepatic and peripheral insulin resistance in pioglitazone-treated type II diabetic patients. *Int J Obes Relat Metab Disord* 28:783–789.
- Bakker, A.H.; Van Dielen, F.M.; Greve, J.W.; Adam, J.A.; and Buurman, W.A. (2004):** Preadipocyte number in omental and subcutaneous adipose tissue of obese individuals. *Obesity Research*, 12, 488– 498.
- Ballard-Barbash, R. (2004)** Anthropometry and Breast Cancer. Body Size—A Moving Target, *Cancer* 74, 1090–1100.
- Ballatori, N. and Truong, A.T. (2002):** Glutathione as a primary osmotic driving force in hepatic bile formation. *Am J Physiol* 263:G617–24.
- Bandyopadhyay, B.; Fan, J.; Guan, S.; Li, Y.; Chen, M.; Woodley, D.T.; and Li, W. (2006):** A “traffic control” role for TGFbeta3: orchestrating dermal and epidermal cell motility during wound healing. *J. Cell.Biol.* 172: 1093–1105.
- Barak, N.; Ehrenpreis, E.D.; Harrison, J.R.; Sitrin, M.D.(2002):** Gastroesophageal reflux disease in obesity: pathophysiological and therapeutic considerations. *Obesity Rev* 3:9–15.
- Baratta, M. (2002):** Leptin—from a signal of adiposity to a hormone mediator in peripheral tissues. *Med Sci Monit* , 8:RA282-RA292.



- Bastard, J.P.; Vidal, H.; Jardel, C.; Bruckert, E.; Robin, D.; Vallier, P.; Blondy, P.; Turpin, G.; Forest, C. and Hainque, B. (2007):** Subcutaneous adipose tissue expression of plasminogen activator inhibitor-1 gene during very low calorie diet in obese subjects. *Int J Obes Relat Metab Disord* 24,70–74.
- Bastelica, D.; Morange, P.; Berthet, B.; Borghi, H.; Lacroix, O.; Grino, M.; Juhan-Vague, I. & Alessi, M.C. (2005):** Stromal cells are the main plasminogen activator inhibitor-1-producing cells in human fat: evidence of differences between visceral and subcutaneous deposits. *Arteriosclerosis Thrombosis and Vascular Biology*, 22, 173– 178.
- Begum, N.; Sandu, O.A.; Ito, M.; et al. (2006):** Active Rho kinase (ROK-B) associates with insulin receptor substrate-1 and inhibits insulin signaling in vascular smooth muscle cells. *J Biol Chem* 277:6214-6222
- Berg, A.H.; Combs, T.P.; Du, X.; Brownlee, M. and Scherer, P.E.(2005):** The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med* 7:947 –953
- Berg, G.; Delaive, K.; Manfreda, J.; Walld, R. and Kryger, M.H.(2002):** The use of health-care resources in obesity–hypoventilation syndrome. *Chest* 120:377–383.
- Berger, K.I.; Ayappa, I.; Chatr-Amontri, B.; Marfatia, A.; Sorkin, I.B.; Rapoport, D.M. and Goldring, R.M.(2001):** Obesity hypoventilation syndrome as a spectrum of respiratory disturbances during sleep. *Chest* 120:1231–1238.



- Berner, H. S., Lyngstadaas, S. P.; Spahr, A.; Monjo, M.; Thommesen, L.; Drevon, C.A.; Syversen, U.; Reseland, J. E. (2004).** Adiponectin and its receptors are expressed in bone-forming cells. *Bone* 35: 842-849.
- Blobel, G. A.; Nakajima, T.; Eckner, R.; Montminy, M. and Orkin, S. H. (1998):** CREB-binding protein cooperates with transcription factor GATA-1 and is required for erythroid differentiation. *Proc. Natl. Acad. Sci. USA* **95**:2061–2066.
- Bodary, P.F.; Westrick, R.J.; Wickenheiser, K.J.; Shen, Y. and Eitzman, D.T. (2002)** Effect of leptin on arterial thrombosis following vascular injury in mice. *Journal of the American Medical Association*, 287, 1706– 1709.
- Boden, G. and Hoeldtke, R.D.(2003):** Nerves, fat, and insulin resistance. *N Eng! J Med* 349:1966-1967
- Boden, G.(2006):** Interaction between free fatty acids and glucose metabolism. *Curr Opin Gin Nutr Metab Care* 5:545-549
- Bogacka, I.; Xie, H.; Bray, G.A.; Smith, S.R. (2005):** Pioglitazone induces mitochondrial biogenesis in human subcutaneous adipose tissue in vivo. *Diabetes* 54:1392–1399.
- Bouchard C.A.; Tremblay, A.; Nadeau, J.; Dussault, J.P.; Despres,G.; Theriault, P.J.; Lupien, O.; Serresse, M.R.; Boulay, and Fournier, G. (1990):** Long-term exercise training with conestant energy intake. 1: Effect on body composition and selected metabolic variables. *Int. J. obesity* 14:57-73.
- Bray, G.A.(1998):** Classification and evaluation of the obesities. *Med Clin North Am* 73:161-184



- Brun R.P.; Kim, J.B.; Hv, E.; Altioek, S. and Spiegelman, B.M. (1996):** Adipocyte differentiation: a transcriptional regulatory cascade. *Curr. Opin. Cell. Boil.* 8: 826-832.
- Brun, R.P.P.; Tontonoz, B.M.; Forman, R.; Ellis, J.; Chen, R.; Evans, M. and Spiegelman, B.M. (1996):** Differential activation of adipogenesis by multiple PPAR isoforms. *Genes Dev.* 10: 974-984.
- Bruun, J.M.; Pedersen, S.B.; Richelsen, B. (2003):** Regulation of interleukin 8 production and gene expression in human adipose tissue *in vitro*. *J Clin Endocrinol Metab* 86:1267–1273
- Buerger, C.N.; Plock, P.; Dehghanyar, Joukhadar, C. and Kloft, C. (2006):** Pharmacokinetics of unbound linezolid in plasma and tissue interstitium of critically ill patients after multiple dosing using microdialysis. *Antimicrob. Agents Chemother.* 50:2455-2463.
- Bugianesi, E.; Gastaldelli, A.; Vanni, E.; Gambino, R.; Cassader, M.; Baldi, S.; Ponti, V.; Pagano, G.; Ferrannini, E.; Rizzetto, M. (2005):** Insulin resistance in non-diabetic patients with non-alcoholic fatty liver disease: sites and mechanisms. *Diabetologia* 48:634–642.
- Bugianesi, E.; Gastaldelli, A.; Vanni, E.; Gambino, R.; Cassader, M.; and Baldi, S. (2005):** Insulin resistance in non-diabetic patients with non-alcoholic fatty liver disease: sites and mechanisms. *Diabetologia* 48: 634-642. Links.
- Calvo, J.C.; Rodbard, D.; Katki, A.; Chernick, S. and Yanagishita, M. (1991):** Differentiation of 3T3-L1 preadipocytes with 3-isobutyl-1-methylxanthine and dexamethasone stimulates cell-associated



and soluble chondroitin 4-sulfate proteoglycans. *J. Biol. Chem.* 266: 11237-11244.

Camp, H.S.; Ren, D.; and Leff, T. (2002): Adipogenesis and fat-cell function in obesity and diabetes. *Trends Mol Med* 8:442–447.

Campbell, D.J. (1997): Tissue rennin-angiotensin system: sites of angiotensin formation. *J. Cardio. Vasc. Pharmacol.* 10 (suppl.4): SI-88.

Cannon, B. and Nedergaard J.(2004): Brown adipose tissue: function and physiological significance. *Physiol Rev.* 84:277-359

Casimir, D.A.; Miller, C.W. and Ntambi, J.M. (1996): Preadipocyte differentiation blocked by prostaglandin stimulation of prostanoid FP2 receptor in murine 3T3-L1 cells. *Differentiation* 60: 203-210.


Cassader, M.; Gambino, R.; and Musso, G. (2006): Postprandial triglyceride-rich lipoprotein metabolism and insulin sensitivity in nonalcoholic steatohepatitis patients. *Lipids* 36:1117-1124.

Castellon, R. and Hamdi, H.K. (2007). "Demystifying the ACE polymorphism: from genetics to biology.". *Curr. Pharm. Des.* **13** (12): 1191-8.


Cederberg, A.; Gronning, L.M.; Ahren, B.; Tasken, K.; Carlsson. P.; and Enerback, S. (2001) FOXC2 is a winged helix gene that counteracts obesity, hypertriglyceridemia, and diet-induced insulin resistance. *Cell* 106: 563–573.



- Cesari, M.; Penninx, B.W.; Newman, A.B.; Kritchevsky, S.B.; Nicklas, B.J.; Sutton-Tyrrell, K.; Rubin, S.M.; Ding, J.; Simonsick, E.M.; Harris, T.B. and Pahor, M. (2003):** Inflammatory markers and onset of cardiovascular events: results from the Health ABC study. *Circulation*, 108, 2317– 2322.
- Chagnac, M.; Hernandez, E., and Morales, E. (2001):** Clinical features and long – term outcome of obesity-associated focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 9: 1790-8.
- Chan, J.L.; Heist, K.; DePaoli, A.M.; Veldhuis, J.D. and Mantzoros, C.S. (2003)** The role of falling leptin levels in the neuroendocrine and metabolic adaptation to short-term starvation in healthy men. *Journal of Clinical Investigations*, 111, 1409–1421.
- Chang, H.; Brown, C.W.; and Matzuk, M.M. (2002).** Genetic analysis of the mammalian transforming growth factor- β superfamily. 23: 787–823.
- Chavey, C.; Mari, B.; Monthouel, M.N.; Bonnafous, S.; Anglard, P.; Van Obberghen, E. and Tartare-Deckert S.(2006):** Matrix metalloproteinases are differentially expressed in adipose tissue during obesity and modulate adipocyte differentiation. *J Biol Chem* 278: 11888–11896.
- Chawla, A.; and Lazar, M.A.(1994):** Peroxisome proliferator and retinoid signaling pathways co-regulate preadipocyte phenotype and survival. *Proc. Natl. Acad. Sci. USA* 91: 1786-1790.
- Chehab, F.F.; Qiu J.; Mounzih. K.; Ewart-Toland, A.; Ogus S.(2002)** Leptin and reproduction. *Nutr Rev* 60:S39-S46discussion S68–84,85–7.



- Chen, G.; Paka, L.; Kako, Y.; Singhal, P.; Duan, W. and Pillarisetti, S. (2004):** *J. Biol.Chem.* 276, 49142-49147
- Chen, N.X.; Hausman, G.J. and Wright J.T. (1996):** Influence of thyroxine in vivo on preadipocyte development and insulin-like growth factor-I and IGF binding protein secretion in fetal stromal vascular cell cultures. *Obesity Res.* 4: 357-366,
- Chen, Z.; Torrens, J.I.; Anand, A.; Spiegelman, B.M.; and Friedman, J.M. (2005):** Krox20 stimulates adipogenesis via C/EBP β -dependent and -independent mechanisms. *Cell Metab* 1:93–106.
- Cherry, C.L.; Lal, L.; Thompson, K.A.; McLean, C.A.; Ross, L.L.; and Hernandez J. (2005)** Increased adipocyte apoptosis in lipoatrophy improves within 48 weeks of switching patient therapy from Stavudine to abacavir or zidovudine. *J Acquir Immune Defic Syndr.* Mar 1 38(3):263-7.
- Choy, L.; Rosen, B. and Spiegelman, B.M. (2002):** Adipsin and an endogenous pathway of complement from adipose cells. *J Biol Chem* 267: 12736-12741.
- Christiansen, T.; Richelsen, and Bruun, J.M. (2005):** Monocyte chemoattractant protein-1 is produced in isolated adipocytes, associated with adiposity and reduced after weight loss in morbid obese subjects. *Int J Obes Relat Metab Disord* 29:146–150.
- Christos, S.; and Mantzoros, DSc M.D.(2005):** The Role of Leptin in Human Obesity and Disease: A Review of Current Evidence
- Cianflone, K. and Xia, Z. (2003):** Critical review of acylation-stimulating protein physiology in humans and rodents. *Biochimica et Biophysica Acta*, 1609, 127– 143.



- Cigolini M. and Smith U.(2003):** Human adipose tissue in culture. VIII. Studies on the insulin-antagonistic effects of gluco-corticoids. *Metabolism* 28:502-510
- Clement, K.; Vaisse, C.; Lahlou, N.; Cabrol, S.; Pelloux, V. and Cassuto, D. (1998)**A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature*, 392, 398– 401.
- Cnop, M.; Landchild, M.J.; Vidal, J.; Havel, P.J.; Knowles, N.G.; Carr, D.R.; Wang, F. and Hull, R.L. (2002):** The concurrent accumulation of intra-abdominal and subcutaneous fat explains the association between insulin resistance and plasma leptin concentrations: distinct metabolic effects of two fat compartments. *Diabetes* 51:1005–1015
- Combs, T.P.; Nagajyothi, Mukherjee S.; de Almeida, C.J.; Jelicks, L.A.; Schubert, W.; Lin, Y. and Jayabalan, D.S.(2005)** The adipocyte as an important target cell for *Trypanosoma cruzi* infection. *J Biol Chem* 280:24085–24094.
- Coppack, S.W. (2001)** Pro-inflammatory cytokines and adipose tissue. *Proceedings of the Nutrition Society*, 60, 349– 356.
- Corry, D.B. and Tuck, M.L.(2005):** Selective aspects of the insulin resistance syndrome. *Curr Opin Nephrol Hypertens* 10:507-514
- Crandall D.L.; Herzlinger, H.G., Saunders, D.D. and Kral, J.G. (1994):** Developmental aspects of adipose tissue rennin-angiotensin system: therapeutic implications. *Drug Dev. Res.* 32: 117-125.
- Cserjesi, P.D.; Brown, K.L.; Ligon, G.E.; Lyons, N.G.; Copeland, D.J.; Gilbert, N.A.; Jenkins, and Olson E.N. (1995):** Scleraxis: a basic helix-loop-helix protein that prefigures skeletal formation during mouse embryogenesis. *Development* 121: 1099-1110, 1995.



Czech, M. P. (2002): *Mol. Cell* 9

Dagenais, G.R.; Yi, Q.; Mann, J.F.; Bosch, J.; Pogue, J. and Yusuf, S.(2005): Prognostic impact of body weight and abdominal obesity in women and men with cardiovascular disease. *Am Heart J.* 149: 54–60.

Dani, C. A.G.; Smith, S.; Dessolin, P.; Leroy, L.; Staccini, P.; Pvillegeois, C.; Darimont, and. Ailhaud. G. (1997):Differentiation of embryonic stem cells into adipocytes in vitro. *J. Cell Sci.* 110: 1279-1285.

Darimont, C.; Vassaux, G.; Ailhaud, G. and Negrel, R. (1994): Differentiation of preadipose cells: paracrine role of prostacyclin upon stimulation of adipose cells by angiotensin-II. *Endocrinology* 135: 2030-2036.

Diamond, J.; Guns Germs, and Steel(2005):the fates of human societies. New York, Norton.

Díez, J.J. and Iglesias. P. (2003). The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur. J. Endocrinol.* 148: 293-300.

Dixon, J.B.; Bhathal, P.S.; and O'Brien, P.E. (2003): Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese. *Gastroenterology* 121:91–100.

Dötsch, J.; Meissner, U. and Rascher, W.(2003): Leptin-induced weight loss is not solely mediated by anorexia. *Eur J Endocrinol* 148:11-12.

Drenick, E.J.; Bale, G.S.; Seltzer, F. and Johnson, D.G.(2000): Excessive mortality and causes of death in morbidly obese men. *J Am Med Assoc.* 243: 443–445.




- Dulloo, A.G.; Stock, M.J. and Solinas, G. (2002):** Leptin directly stimulates thermogenesis in skeletal muscle. *FEBS Lett* 515:109-113
- Dvorak, H.F.; Brown, L.F.; Detmar, M. and Dvorak, A.M.(2005):** Vascular permeability factor/vascular endothelial growth factor, microvascular hyperpermeability, and angiogenesis. *Am J Pathol* 146: 1029–1039.
- El- Atat, F.; Aneja, A.; Macferlane, S., and Sowers, J. (2003):** Obesity and hypertension. *Endocrinol Metab Clin N Am*; 32: 823-54
- El-Serag, H.B.(2005):** Obesity and disease of the esophagus and colon. *Gastroenterol Clin North Am* 234:63–82.
- Engelhardt, M.J.; Geerlings, M.I. and Meijer, J. (2004),** Inflammatory proteins in plasma and the risk of dementia: the Rotterdam Study. *Arch Neurol* 61:668-672.
- Engeli, S.; Schling, P. and Gorzelnia, K.(2005):** The adipose - tissue renin-angiotensin-aldosterone system: role in the metabolic syndrome? *Int J Biochem Cell Biol* 35:807-825
- Entenmann, G. and Hauner, H.(1996):** Relationship between replication and differentiation in cultured human adipocyte precursor cells. *Am. J. Physiol.* 270(*Cell Physiol.* 39): C1011-C1016.
- Eric, S. and Daar, M.D.(2007):** Physical and Biochemical Changes in HIV Disease MedicineNet, Accessed 22 September 2007
- Esposito, K.; Pontillo, A.; Di Palo, C.; Giugliano, G.; Masella, M.; Marfella, R. and Giugliano, D. (2003):** Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *Journal of the American Medical Association*, 289, 1799– 1804.



- Evans, J.L.; Goldfine, I.D.; Maddux, B.A. and Grodsky, G.M.(2003):** Oxidative stress and stress-activated signaling pathways a unifying hypothesis of type 2 diabetes. *Endocrinol Rev*, 23:599-622
- Fain, J.N.; Madan, A.K.; Hiler, M.L.; Cheema, P. and Bahouth, S.W. (2004):** comparison of the release of adipokines by adipose tissue, adipose tissue matrix, and adipocytes from visceral and subcutaneous abdominal adipose tissues of obese humans. *Endocrinology* 145:2273-2282.
- Fan, W.; Boston, B.A.; Kesterson, R.A.; Hruby, V.J. and Cone, R. D. (2003) :**Roleof melanocortinerbic neurons in feeding and the agouti obesity syndrome. *Nature (Lond.)* 385: 165–168.
- Farahat, M. N.; Yanni, G.; Poston, R.;and Panayi, G. S. (2003).** Cytokine expression in synovial membranes of patients with rheumatoid arthritis and osteoarthritis. *Ann. Rheum. Dis.* 52: 870-875.
- Faraj, M.; Havel, P.J.; Phelis, S.; Blank, D.; Sniderman, A.D.and Cianflone, K.(2003):** Plasma acylation-stimulating protein, adiponectin, leptin, and ghrelin before and after weight loss induced by gastric bypass surgery in morbidly obese subjects. *J Clin Endocrinol Metab*, 88 :1594 –1602.
- Farer, L.S.; Lowell, A.M. and Meador, M.P. (2000)** Extrapulmonary tuberculosis in the United States. *Am J Epidemiol* 109: 205–217.
- Farooqi, I.S. (2003):**clinical spectrum of obesity and mutations of the melanocortin 4 receptor gene. *N Engl J Med* 348:1160.




- Fasshauer, M.; Klein, J.; Kralisch, S.; Klier, M.; Lossner, U.; Bluher, M. and Paschke, R. (2004):** Monocyte chemoattractant protein 1 expression is stimulated by growth hormone and interleukin-6 in 3T3-L1 adipocytes. *Biochem Biophys Res Commun* 317:598–604
- Federico, G.; Street, M.E.; and Maghnie, M. (2006).** "Assessment of serum IGF-I concentrations in the diagnosis of isolated childhood-onset GH deficiency: a proposal of the Italian Society for Pediatric Endocrinology and Diabetes (SIEDP/ISPED)". *J. Endocrinol. Invest.* 29 (8): 732-7.
- Festa, A.; D'Agostino, R.; Jr, Tracy, R.P. and Haffner, S.M. (2002 April)** Insulin resistance atherosclerosis study. Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of type 2 diabetes: the insulin resistance atherosclerosis study. *Diabetes*, 51, 1131– 1137.
- Firth, S.M. and Baxter, R.C. (2004):** Cellular actions of the insulin-like growth factor binding proteins. *Endocr Rev* 23:824–854.
- Fisher, J.S.; Van Pelt, R.E.; Zinder, O.; Landt, M. and Kohrt, W.M.(2001):** Acute exercise effect on postabsorptive serum leptin. *J Appl Physiol* 91: 680–686.
- Flier, J.S. (1998)** Clinical review 94: what's in a name? In search of leptin's physiologic role. *Journal of Clinical Endocrinology and Metabolism*, 83, 1407– 1413.
- Flier, J.S. and Maratos-Flier, E. (1998):** Obesity and hypothalamus novel peptides for new pathways. *Cell* 92: 437-440.
- Fogo, A.B. (2003):** The potential for regression of renal scarring. *Curr Opin Nephrol Hypertens* 12: 223–225.



- Fonseca-Alaniz, M.H.; Takada, j.; Alonso-Vale, M.I.C.; and Lima, F.B. (2007):** Adipose tissue as an endocrine organ: from theory to practice. *J. Pediatr.* (Rio J.) vol.83 no.5 suppl.0 Porto Alegre.
- Ford, E.S.; Giles, W.H. and Dietz, W.H. (2002).** Prevalence of metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. *JAMA* **287**(3):356-359.
- Frayn, K.N.; Karpe, F.; Fielding, B.A.; Macdonald, I.A.; and Coppack, S.W.(2003):** Integrative physiology of human adipose tissue. *Int J Obes Relat Metab Disord* 27:875-888.
- Fried, S.K.; Bunkin, D.A.; and Greenberg, A.S.(1998):** Omental and subcutaneous adipose tissues of obese subjects release interleukin-6: depot difference and regulation by glucocorticoid. *J Clin Endocrinol Metab* 83: 847-850.
- Friedman, J.M.(2003):** A war on obesity, not the obese. *Science* 299:856-858
- Fruebis, J.; Tsao, T.S.; and Javorschi, S. (2001):** Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci USA* , 98:2005-2010.
- Funahashi, H.; Yada, T.; Suzuki, R.; and Shioda,S. (2003):** Distribution, function, and properties of leptin receptors in the brain. *Int Rev Cytol* 224:1-27.
- Gaillard, D.; Wabitsch, M.; Pipy, B.and Negrel, R.(1991):** Control of terminal differentiation of adipose precursor cells by glucocorticoids. *J. Lipid Res.* 32: 569-579.



- Gale, S.M.; Castracane, V.D. and Mantzoros, C.S.(2004):** Energy homeostasis, obesity and eating disorders: recent advances in endocrinology. *J Nutr* 134:295-298.
- Garg, A.; Peshock, R. M. and Fleckenstein, J. L.(2001) :** Adipose tissue distribution pattern in patients with familial partial lipodystrophy (Dunnigan variety). *J. Clin. Endocr. Metab.* 84: 170-174.
- Ge, H.F.; Huang, L.; Pourbahram, T. and Li, C.(2002):** Generation of soluble leptin receptor by ectodomain shedding of membrane-spanning receptors in vitro and in vivo. *J Biol Chem* 277:45898-45903.
- Gielen, S.; and Hambrecht, R.(2004):** The childhood obesity epidemic: impact on endothelial function. *Circulation* 109:1911-1913
- Giovannucci, E.(2005):** Insulin and colon cancer. *Cancer Causes Control*, 6: 164-179.
- Glackin, c. A.; Winters, K. A.; Murray, E.J.B. and Murray, S. S.(1994):** Transcripts encoding the basic-helix-loop-helix factor twist are expressed in mouse embryos, cell lines, and adult tissues. *Mol. Cell. Differ.* 2: 309-328.
- Goldberg, I. J.(1996):** Lipoprotein lipase and lipolysis: central roles in lipoprotein metabolism and atherogenesis. *J. Lipid Res.* 37: 693-707.
- Goldstein, B.J.(2002):** Insulin resistance as the core defect in type 2 diabetes mellitus. *Am J Cardiol* 90 suppl 3G-10G
- Gonzalez, .F. (2005):** Getting fat: two new players in molecular adipogenesis. *Cell Metab* 1:85–86.



- Goodfriend, T.L.; Kelley, D.E.; and Goodpaster, D.H (2004):** Visceral obesity and insulin resistance are associated with plasma aldosterone levels in women. *Obes Res* 7:355-362
- Green, H., and Kehinde, O. (1997):** Formation of normally differentiated subcutaneous fat pads by an established preadipose cell line. *J. Cell. Physiol.* 101: 169-171.
- Greenwood, M.R.C. and Johnson, P.R.(1993):** Genetic differences in adipose tissue metabolism and regulation. *Ann. N.Y. Acad. Sci.* 676:253-269.
- Gregoire, F. M.; Johnson, P. R.; and Greenwood, M. R.(1995):** Comparison of the adipoconversion of preadipocytes derived from lean and obese Zucker rats in serum-free cultures. *Int. J. Obes. Relat. Metab. Disorders* 19: 664-670.
- Gregoire, Francine, M.; Cynthia, M.; Smas, and Hei Sook Sul. (2004):** Understanding Adipocyte Differentiation *Physiological Reviews* Vol. 78 , pp. 783-809
- Grundy, S.M.; Brewer, H.B., and Jr Cleeman, J.I.(2004):** Definition of metabolic syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Arterioscler Thromb Vasc Biol* 24:e l3-e l8
- Grundy, S.M.; Hansen, B.; Smith, S.C.; and Jr.(2004):** Clinical management of metabolic syndrome: report of the American Heart Association/National Heart, Lung, and Blood Institute/American Diabetes Association conference on scientific issues related to management. *Arterioscler Thromb Vasc Biol* 24:e l9-e24



- Guzick, D.S.; Wing, R.; Smith, D.; Berga, S.L.; and Winters, S.J. (2007)** Endocrine consequences of weight loss in obese, hyperandrogenic, anovulatory women. *Fertil Steril* 61,598–604.
- Guzik, T.J.; Marvar, P.J.; Czesnikiewicz-guzik, M. and Korbust, R.(2007):** Perivascular Adipose Tissue As A Messenger Of The Brain-Vessel Axis: Role In Vascular Inflammation And Dysfunction *Journal Of Physiology And Pharmacology*, 58, 4, 591.610
- Hall, J.E.; Crook, E.D., and Jones, D.W. (2002):** Mechanisms of obesity associated cardiovascular and renal disease. *Am J Med Sci* 324: 127-37.
- Hanley, A.J.; Karter, A.J.;and Festa, A.(2006):** Factor analysis of metabolic syndrome using directly measured insulin sensitivity: the Insulin Resistance Atherosclerosis Study. *Diabetes* 51:2642-2647
- Harris, F. M.; Tesseur, I.; Brecht, W. J.; Xu, Q.; Mullendorff, K.; Chang, S.; Wyse-Coray, T.; Mahley, R. W.; and Huang, Y. (2004)** *J. Biol. Chem.* **279**, 3862-3868
- Harrison, T.R.; Jeffrey, S.; Flier, Eleftheria and Maratos-Flier.(2005):**obesity 64:422-429.
- Hauner, H.; Rorhig, K., and Petruschke, T.(1995):** Effects of epidermal growth factor (EGF), platelet-derived growth factor (PDGF) and fibroblast growth factor (FGF) on human adipocyte development and function. *Eur. J. Clin. Invest.* 25: 90-96.
- Hausman, G. J.; Wright, J. T., and Richardson, R. L.(1993)** The influence of extracellular matrix substrata on preadipocyte development in serum-free cultures of stromal-vascular cells. *J. Anim. Sci.* 74: 2117-2128.



- Havel, P.J. (2001):** Peripheral signals conveying metabolic information to the brain: short-term and long-term regulation of food intake and energy homeostasis. *Exp Biol Med (Maywood)* 226 :963 –977.
- Heilbronn, L.K.; Rood, J.; Janderova, L.; Albu, J.B.; Kelley, D.E.; Ravussin, E., and Smith, S.R. (2004)** Relationship between serum resistin concentrations and insulin resistance in nonobese, obese, and obese diabetic subjects. *J Clin Endocrinol Metab* 89,1844–1848.
- Hershcopt, R.J., and Bradlow, H.L. (1997)** Obesity, Diet, Endogenous Estrogens, and the Risk of Hormone-Sensitive Cancer, *Am. J. Clin. Nutr.* 45, 283–289.
- Hileman, S.M.; Pierroz, D.D.; Masuzaki, H.; Bjorback, C.; El-Haschimi, K.; and Banks, W.A. (2002):** Characterization of short isoforms of the leptin receptor in rat cerebral microvessels and of brain uptake of leptin in mouse models of obesity. *Endocrinology* 143:775-783.
- Hill, J.M.; Bhattacharjee, P.S. and Neumann, D.M.(2007):** "Apolipoprotein E alleles can contribute to the pathogenesis of numerous clinical conditions including HSV-1 corneal disease". *Exp Eye Res* , 84(5):801-11.
- Hilton, L.K. and Loucks, A.B.(2000)** Low energy availability, not exercise stress, suppresses the diurnal rhythm of leptin in healthy young women. *Am J Physiol Endocrinol Metab* 278: E43–E49.
- Hino, M.; Nakao, T.; Yamane, T.; Ohta, K.; Takubo, T. and Tatsumi, N. (2006):** Leptin receptor and leukemia. *Leuk Lymphoma* 36:457–461.



- Hiratsuka, A.; Adachi, H.; Fujiura, Y.; Yamagishi, S.; Hirai, Y.; Enomoto, M.; Satoh, A.; Hino, A.; Furuki, K., and Imaizumi, T.(2005):** Strong association between serum hepatocyte growth factor and metabolic syndrome. *J Clin Endocrinol Metab* 90: 2927–2931.
- Ho, L.; Qin, W. And Pomp, P.N. (2004):** Diet-induced insulin resistance promotes amyloidosis in a transgenic mouse model of Alzheimer's disease. *FASEB J* 18:902-904.
- Hofman, P.L.; Regan, F. and Jackson, W.E. (2004):** Premature birth and later insulin resistance. *N Engl J Med* 351: 2179-2186
- Hoggard, N.; Johnstone, A.M.; and Faber, P. (2004):** Plasma concentrations of alpha-MSH, AgRP and leptin in lean and obese men and their relationship to differing states of energy balance perturbation. *Clin Endocrinol (Oxf)* 61:31-9
- Holmes, R.C. and Meara, R.H.(2003):** Morphea, sclerotic panatrophly and disseminated granuloma annulare. *Clin Exp Dermatol.* 2003 Mar;8(2):201-3.
- Hong, J.H.; Hwang, E.S.; McManus, M.T.; Amsterdam, A.; Tian, Y.; Kalmukova, R. and Mueller, E. (2005):** TAZ, a transcriptional modulator of mesenchymal stem cell differentiation. *Science* 309:1074–1078
- Hotamisligil, G.S.; Peraldi, P.; Budavari, A.; Ellis, R.; White, M.F. and Spiegelman, B.M.(2006)** IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF- α - and obesity-induced insulin resistance. *Science* 271: 665-668.



- Hotamisligil, G.S.; Shargill, N.S. and Spiegelman, B.M. (2000):** Adipose expression at tumour necrosis factor- α direct role in obesity-linked insulin resistance science (Washington. D.C.) 254: 87-91.
- House, A.A. and Walley, V.M.(2005):** Right heart failure due to ventricular adiposity: ‘adipositas cordis’—an old diagnosis revisited. *Can J Cardiol.* 12: 485–489.
- Hube, F.; Birgel, M.; Lee, Y.M. and Hauner, H.(1999):** Expression pattern of tumour necrosis factor receptors in subcutaneous and omental human adipose tissue: role of obesity and non-insulin-dependent diabetes mellitus. *Eur J Clin Invest* 29: 672-678.
- Hug, C., and Lodish, H.F. (2005):** Medicine. Visfatin: a new adipokine. *Science* 307:366–367.
- Hunter, H.; Cobban, H.J. and Vandenabeele, P. (2003):** Tumor necrosis factor- α -induced activation of RhoA in airway smooth muscle cells: role in the Ca^{2+} sensitization of myosin light chain20 phosphorylation *Mol Pharmacol* 63:714-721
- Hyman, B.T.; Stoll, L.L. and Specter, A.A. (1992):** Pfostaglandin production by 3T3-L1 cells in culture. *Biochim, Biophys. Acta.* 713: 375-385.
- Inese Čakstiņa; Una Riekstiņa; Ruta Muceniece; and Jānis Ancāns(2007):** Isolation, characterization and analysis of *TGF β* gene expression in adipose-derived precursor cells cultured in different growth media *Acta Universitatis Latviensis, Vol. 723, Biology, pp. 181–190.*
- Jemal, A.; Siegel, R.; Ward, E.; Murray, T.; Xu, J. and Thun, M. J. (2007).** Cancer Statistics, *CA Cancer J. Clin.*, 57, 43-66



- Jeon, E.S.; Moon, H.J.; Lee, M.J.; Song, H.Y. and Kim, Y.M. (2006).** Sphingosyl phosphorylcholine induces differentiation of human mesenchymal stem cells into smooth-muscle like cells through a TGF-beta-dependent mechanism.. 1: 4994–5005.
- Jequier, E. (2002):** Pathways to obesity. *Int J Obes Relat Metab Disord* 26(suppl):S12-S17
- Johnson, B.H.; Standndge, M.K. and Moustaid, N. (2005):** Angiotensinogen gene expression in adipose tissue: analysis of obese models and hormonal and nutritional control. *Am. J. Physiol.* 273: 236-242.
- Johnson, P.R. and Greenwood, M.R.C. (1998):** The adipose tissue. In: "Cell and Tissue Biology: A Textbook of histology." (ed. By L. Weiss), 6th edition, urban and Schwarzenberg, Baltimore, MD. Pp. 191-209.
- Johnson, P.R.; Stern, J.S.; Greenwood, M.R.C.; Zucker, L.M. and Hirsch, J.(1973):** Effect of early nutrition on adipose cellularity and pancreatic insulin release in the Zuker rat.*J. Nutrition.* 103:738-743.
- Jones, S.A.; Horiuchi, S.; Topley, N.; Yamamoto, N. and Fuller, G.M.(2001):** The soluble interleukin 6 receptor: mechanisms of production and implications in disease. *FASEB J* 15: 43-58.
- Julie Janeway; Karen Sparks, M.B.E., and Steven, R. (2005):** .GERD and the Obese Patient, obesity action caoition j, Modesitt SC, van Nagell JR. Impact of obesity on the incidence and treatment of gynecologic cancers: a review. *Obstet Gynecol Survey.* 60: 683-692. Links



- Kadowaki, T. (2003):** Insights into insulin resistance and type 2 diabetes from knockout mouse models. *J Clin Invest* 106:459–465
- Kadowaki, T.; Kubota, N.; Terauchi, Y. and Yamauchi, T. (2002):** Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 277:25863–25866
- Kadowaki, T.; Kubota, N.; Terauchi, Y.; Yamauchi, T., and Kubota, T. (2002):** Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 277:25863–25866
- Kadowaki, T.; Yamauchi, T.; Kamon, J.; Ito, Y. and Tsuchida, A. (2003):** Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 423:762–769
- Kahn, C.R.(2002):** Banting lecture insulin action, diabetogenes, and the cause of type II diabetes. *Diabetes* 43: 1066-1084
- Kalant, D.; Cain, S.A.; Maslowska, M.; Sniderman, A.D., and Cianflone, K. (2005):** The chemoattractant receptor-like protein C5L2 binds the C3a des-Arg77/acylation-stimulating protein. *J Biol Chem* 278 :11123 –11129.
- Kamai, Y.; Mikawa, S.; Endo, K.; Sakai, H.; and Komando, T. (1996):** Regulation of insulin-like growth factor-I expression in mouse preadipocyte Ob1771 cells. *J. Biol. Chem.* 271: 9883-9886.
- Kambham, N.; Markowitz, G.S.; Valery, A.M.; Lin, J., and Agati, V.D.(2001)** Obesity–related glomerulopathy: An emerging epidemic. *Kidney Int*; 59: 1498-1509.
- Kang, S.; Bajnok, L.; Longo, K.A.; Petersen, R.K.; Hansen, J.B.; Kristiansen, K. and MacDougald, O.A. (2005):** Effects of Wnt signaling on brown adipocyte differentiation and metabolism mediated by PGC-1 α . *Mol Cell Biol* 25:1272–1282.



- Kaser, A. S. Hfer, A.; Ebenbiehler, C.F.; Tilg, H., and Patsch, J.R. (2003):** Resistm messenger-RNA expression is increased by promflammatory cytokines in vitro. *Biochem. Biophys. Res. Comman* 309: 286-290.
- Katagiri, H.; Yamada, T., and Oka, Y.(2007):** Adiposity and cardiovascular disorders: disturbance of the regulatory system consisting of humoral and neuronal signals. *Circ Res.* **101**:27–39.
- Kern, P.A.; Di Gregorio, G.B.; Lu, T.; Rassouli, N., and Ranganathan, G. (2003):** Adiponectin expression from human adipose tissue: Relation to obesity, insulin resistance, and tumor necrosis factor-alpha expression. *Diabetes* 52: 1779–1785
- Kershaw, E.E. and Flier, J.S. (2004):** Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab* 89: 2548–2556.
- Khovidhunkit, W.; Kim, M.S., and Memon, R.A. (2004):** Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host *J. Lipid Res* 45: 1169-1196
- Kim, B.; van Golen, C.M., and Feldman, E.L. (2004):** Insulin-like growth factor-I signaling in human neuroblastoma cells. *Oncogene* 23:130–141.
- Kim, S.Y.; Park, S.M., and Lee, S.T. (2006).** "Apolipoprotein C-II is a novel substrate for matrix metalloproteinases". *Biochem. Biophys. Res. Commun.* 339 (1): 47–54.



- Kissebah, A. H., and Krakower, G. R.(1994):** Regional adiposity and morbidity. *Physiol. Rev.* 74: 761-811.
- Klassen, H.J.; Imfeld, K.L.; Kirov, I.I.; Tai, L.; Gage, F.H., and Young, M.J. (2003).** Expression of cytokines by multipotent neural progenitor cells. *Cytokine* 22: 101–106.
- Klein, S.; Fontana, L.; Young, V.L.; Coggan, A.R.; Kilo, C.; Patterson, B.W., and Mohammed, B.S.(2004).** Absence of an effect of liposuction on insulin action and risk factors for coronary heart disease. *N Engl J Med* 350: 2549–2557.
- Koerner, A.; Kratzsch, J., and Kiess, W. (2005):** Adipocytokines: leptin–the classical, resistin–the controversial, adiponectin–the promising, and more to come. *Best Pract Res Clin Endocrinol Metab* 19:525–546.
- Kohn, A. D.; Summers, S. A.; Birnbaum, M. J., and Roth, R. A. (1996):** Expression of a constitutively active Akt Ser/Thr kinase in 3T3-L1 adipocytes stimulates glucose uptake and glucose transporter 4 translocation.*J.Biol.Chem.*271:31372-31378.
- Komazawa, N. (2004)** Enhanced insulin sensitivity, energy expenditure and thermogenesis in adipose-specific Pten suppression in mice. *Nat Med* 10: 1208–1215
- Kopelman, P.G. (1998):** Effects of obesity on fat topography: metabolic and endocrine *determinants*. In: *Clinical Obesity* (1st ed.), edited by Kopelman PG, and Stock MJ.. Oxford, UK: Blackwell Science, p. 158-175.
- Korpilahti, K.; Syvanne, M.; Engblom, E.; Hamalainen, H.; Puukka, P., and Ronnema, T. (2005):** Components of the insulin resistance syndrome are associated with progression of



atherosclerosis in non-grafted arteries 5 years after coronary artery bypass surgery. *Eur Heart J.* 19: 711–719.

Koutkia, P.; Canavan, B.; Johnson, M.L.; DePaoli, A., and Grinspoon, S. (2003): Characterization of leptin pulse dynamics and relationship to fat mass, growth hormone, cortisol, and insulin. *Am J Physiol Endocrinol Metab* 285:E372-E379.

Kovacs, K.; Lloyd, R.V.; Jin, L.; Tsumanuma, I.; Vidal, S.; Horvath, E.; Schweithauer, B.W.; Couce, M.E., and Burguera, B. (2006): Leptin and leptin receptor in anterior pituitary function. *Pituitary* 4: 33-47.

Krizova, J.; Sulkova, S.; Bednarova, V.; Kotrilikova, E., and Haluzik, M. (2002): Soluble leptin receptor levels in patients with anorexia nervosa. *Endocr Res* 28:199-205.

Kubo, A., and Corley, D.A. (2006): Body mass index and adenocarcinomas of the esophagus or gastric cardia: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 15: 872–8.

Kucio, C.; Besser, P., and Jonderko, K.(1988): Gallbladder motor function in obese versus lean females. *Eur J Clin Nutr.* 42:121-4.

Kugelmas, M.; Hill, D.B.; Vivian, B.; Marsano, L., and McClain, C.J. (2003): Cytokines and NASH: a pilot study of the effects of lifestyle modification and vitamin E. *Hepatology* 38:413–419.

Kuniyasu, A.; Ohgami, N.; Hayashi, S.; Miyazaki, A., and Horiuchi, S. (2003) CD36-mediated endocytic uptake of advanced glycation end products (AGE) in mouse 3T3-L1 and human subcutaneous adipocytes. *FEBS Lett* 537: 85–90.



- Kurrimbux, D.; Gaffen, Z.; Farrell, C.L.; Martin, D., and Thomas, S.A. (2004):** The involvement of the blood-brain and the blood-cerebrospinal fluid barriers in the distribution of leptin into and out of the rat brain. *Neuroscience* 123:527-536.
- Lacasa, D.; Garcia, E.; Henrion, D.; Agli, B., and Giudicelli, Y. (1997):** Site-related specificities of the control by androgenic status of adipogenesis and mitogen-activated protein kinase cascade/c-fos signaling pathways in rat preadipocytes. *Endocrinology* 138: 3181-3186.
- Lafontan, M., and Berlan, M.(2003):** Do regional differences in adipocyte biology provide new pathophysiological insights? *Trends Pharmacol Sci.* 24:276-83.
- Lammert, A.; Kiess, W.; Bottner, A.; Glasow, A., and Kratzsch, J.(2001):** Soluble leptin receptor represents the main leptin binding activity in human blood. *Biochem Biophys Res Commun* 283:982-988.
- Lane, M. D.; Lin, F. T.; Macdouglass, O. A. and Vasseur-Cognet. M. (1996):** Control of adipocyte differentiation by CCAAT/enhancer binding protein alpha (C/EBP alpha). *Int. J. Obesity Related Metab. Disorders* 20, Suppl.: S91-S96, 255.
- Lane, M.D. and Tang, Q.Q. (2005):** From multipotent stem cell to adipocyte. *Birth Defects Res A Clin Mol Teratol* 73:476–477
- Lazartigues, E.; Feng, Y., and Lavoie, J.L. (2007).** "The two fACES of the tissue renin-angiotensin systems: implication in cardiovascular diseases.". *Curr. Pharm. Des.* 13 (12): 1231-45



- Le Lay, S.L.; Boucher, J.; Rey, A.; Castan-Laurell, I.; Krief, S., and Ferre, P. (2005):** Decreased resistin expression in mice with different sensitivities to a high-fat diet. *Biochem Biophys Res Commun* 289:564-567.
- Lee, W.Y., and Mokhlesi, B. (2003):** Diagnosis and management of obesity hypoventilation syndrome in the intensive care unit. *Crit Care Clin* (In press.)
- Leonardsson, G. (2004)** Nuclear receptor corepressor RIP140 regulates fat accumulation. *Proc Natl Acad Sci USA* 101: 8437–8442
- Lepak, N. M., and Serron, G.(1996):** Prostaglandin F₂ alpha stimulates transforming growth factor-alpha expression in adipocyte precursors. *Endocrinology* 136: 3222-3229.
- Levels, J. H.; Marquart, J. A.; Abraham, P. R.; van den Ende, A. E.; Molhuizen, H. O.; van Deventer, S. J., and Meijers, J. C. (2005).** "Lipopolysaccharide is transferred from high-density to low-density lipoproteins by lipopolysaccharide-binding protein and phospholipid transfer protein." *Infect Immun* 73(4): 2321-6.
- Levels, J. H.; Marquart, J. A.; Abraham, P. R.; van den Ende, A. E.; Molhuizen, H. O.; van Deventer, S. J., and Meijers, J. C. (2005).** "Lipopolysaccharide is transferred from high-density to low-density lipoproteins by lipopolysaccharide-binding protein and phospholipid transfer protein." *Infect Immun* 73(4): 2321-6.
- Levine, J.A.; Eberhardt, N.L., and Jensen, M.D.(2001):** Role of nonexercise activity thermogenesis in resistance to fat gain in humans. *Science* 283:212-214




- Lim, H.S.; MacFayen, R.J., and Lip, G.Y.H. (2004):** Diabetes mellitus, the renin-angiotensin-aldosterone system, and the heart. *Arch Intern Med* 164:1737-1748
- Lin, Y.; Rajala, M. W.; Berger, J. P.; Moller, D. E.; Barzilai, N., and Scherer, P. E. (2003).** Hyperglycemia-induced production of acute phase reactants in adipose tissue. *J. Biol. Chem.* 276:42077-42083.
- Linscheid, P.; Seboek, D.; Zulewski, H.; Keller, U., and Muller, B. (2005).** Autocrine/paracrine role of inflammation-mediated calcitonin gene-related peptide and adrenomedullin expression in human adipose tissue. *Endocrinology* 146:2699-2708.
- List, J.F., and Habener, J.F.(2003):** Defective melanocortin 4 receptors in hyperphagia and morbid obesity. *N Engl J Med* 348:1160-1163
- Litherland, G.J.; Hajduch, E., and Hundai, H.S. (2005):** Intracellular signaling mechanisms regulating glucose transport in insulin-sensitive tissues (review). *Mol Membr Biol* 84. 18:195-204
- Lou, S.Q.;Mercado-Pimentel, M.E., and Runyan, R.B.(2008).** Multiple transforming growth factor- isoforms and receptors function during epithelial - mesenchymal cell transformation in the embryonic heart. *Cells Tissues Organs* 185: 146–156.
- Luther, J., and Glesby, M.J. (2007):** Dermatologic adverse effects of antiretroviral therapy: recognition and management. *Am J Clin Dermatol.* 8(4):221-33.
- Lyon, C.J., and Hsueh, W.A. (2003):** Effect of plasminogen activator inhibitor-1 in diabetes mellitus and cardiovascular disease. *Am J Med* 115, Suppl 8A: 62S–68S.



- Ma, L.J.; Mao, S.L.; Taylor, K.L.; Kanjanabuch, T.; Guan, Y.; Zhang, Y.; Brown, N.J.; Swift, L.L., and McGuinness, O.P. (2004):** Prevention of obesity and insulin resistance in mice lacking plasminogen activator inhibitor 1. *Diabetes* 53: 336–346.
- Macdouglad, O. A., and Lane, M. D. (1995):** Transcriptional regulation of gene expression during adipocyte differentiation. *Annu. Rev. Biochem.* 64: 345-373.
- Magni, P.(2003):** Hormonal control of the neuropeptide Y system. *Curr Protein Pept Sci* 4:45-57.
- Mahalia, S.; Desruisseaux; Nagajyothi; Maria, E.; Trujillo; Herbert, B.; Tanowitz, and Philipp, E.(2007):** Scherer Adipocyte, Adipose Tissue, and Infectious Disease. *Infect Immun.* March; 75(3): 1066–1078.
- Majka, M.; Janowska-Wieczorek, A.; Ratajczak, J.; Ehrenman, K.; Pietrzkowski, Z.; Kowalska, M.A.; Gewirtz, A.M.; Emerson, S.G., and Ratajczak, M.Z. (2001).** Numerous growth factors, cytokines, and chemokines are secreted by human CD34(+) cells, myeloblasts, erythroblasts, and megakaryoblasts and regulate normal hematopoiesis in an autocrine/paracrine manner. *Blood* 15: 3075–3085.
- Mandrup, S., and Lane, M. D. (1997):** Regulating adipogenesis. *J. Biol. Chem.* 272: 5367-5370.
- Mann, C.J.; Yen, F.T.; Grant, A.M., and Bihain, B.E. (2001):** Mechanism of cholesteryl ester transfer in hypertriglyceridemia. *J Clin Invest.* 88:2059–2066.



- Marette, A. (2004):** Mediators of cytokine-induced insulin resistance in obesity and other inflammatory settings. *Curr Opin Clin Nutr Metab Care* 5:377-383
- Margetic, S.; Gazzola, C.; Pegg, G.G., and Hill, R.A. (2002):** Leptin: a review of its peripheral actions and interactions. *Int J Obes Relat Metab Disord* 26:1407-1433.
- Maria, E.; Trujillo, and Philipp, E. (2006):** Adipose Tissue-Derived Factors: Impact on Health and Disease. *J endocrine reviews*.
- Marques, B.G.; Hausman, D.B.; Latimer, A.M.; Kras, K.M.; Grossman, B.M., and Martin, R.J. (2005):** Insulin-like growth factor I mediates high-fat diet-induced adipogenesis in Osborne-Mendel rats. *Am J Physiol Regulatory Integrative Comp Physiol* 278: R654-R662
- Marugame, T.; Lee, K., and Eguchi, H. (2002).** Relation of impaired glucose tolerance and diabetes to colorectal adenomas in Japan. *Cancer Causes Control*, 13, 917-21.
- Marx, J. (2003):** Cellular warriors at the battle of the bulge. *Science* 299:846-9
- Massiera, F.; Bloch-Faure, M.; Ceiler, D.; Murakami, K.; Fukamizu, A., and Gasc, J.M. (2001)** Adipose angiotensinogen is involved in adipose tissue growth and blood pressure regulation. *FASEB Journal*, 15, 2727– 2729.
- Matsuzawa, Y.; Shimomura, I., and Nakamura, T. (2005):** Pathophysiology and pathogenesis of visceral fat obesity. *Ann N Y Acad Sci* 676:270-278 -



- Matteoni, C.A.; Younossi, Z.M.; Gramlich, T.; Boparai, N.; Liu, Y.C., and McCullough, A.J. (1999):** Nonalcoholic fatty liver disease: a spectrum of clinical and pathological severity. *Gastroenterology* 116:1413–1419.
- Matthews, D.R.; Hosker, J.P., and Rudensky. (2005):** Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412-9.
- Mauriege, P.; Prud'homme, D., and Lemieux, S. (1999):** Regional differences in adipose tissue lipolysis from lean and obese women: existence of postreceptor alterations. *Am J Physiol* 269:E341-E350
- Maurin, T.; Saillan-Barreau, C.; Cousin, B.; Casteilla, L.; Doglio, A., and Penicaud, L. (2005):** Tumor necrosis factor- α stimulates HIV-1 production in primary culture of human adipocytes. *Exp Cell Res* 304:544–551.
- Mehra, M.R.; Uber, P.A., and Park, M.H. (2004):** Obesity and suppressed B- type natriuretic peptide levels in heart failure. *J Am Coll Cardiol* 43:1590-1595.
- Merkel, M.; Eckel, R. H., and Goldberg, I. J. (2006):** *J. Lipid Res.* 43, 1997–2006.
- Miettinen, T.A.; Olkkonen, V.M.; Jauhiainen, M., and Ehnholm, C. (2005).** Altered hepatic lipid and apolipoprotein A-I metabolism in mice lacking phospholipid transfer protein.
- Minokoshi, Y., and Kim, Y.B. (2002)** Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature*, 415, 339– 343.



- Misra, A.; Peethambaram, A., and Garg, A. (2004):** Clinical features and metabolic and autoimmune derangements in acquired partial lipodystrophy: report of 35 cases and review of the literature. *Medicine (Baltimore)*. 2004 Jan;83(1):18-34.
- Miyazawa-Hoshimoto, S.; Takahashi, K.; Bujo, H.; Hashimoto, N., and Saito, Y.(2003):** Elevated serum vascular endothelial growth factor is associated with visceral fat accumulation in human obese subjects. *Diabetologia* 46: 1483–1488.
- Mizuno, T.M.; Kleopoulos, S.P.; Bergen, H.T.; Roberts, J.L.; Priest, C.A., and Mobbs, C.V. (1999):** Hypothalamic pro-opiomelanocortin mRNA is reduced by fasting in ob/ob and db/db mice, but is stimulated by leptin. *Diabetes*. 47:294-7.
- Mizuno, T.M.; Kleopoulos, S.P.; Bergen, H.T.; Roberts, J.L.; Priest, C.A., and Mobbs, C.V. (1999):** Hypothalamic pro-opiomelanocortin mRNA is reduced by fasting in ob/ob and db/db mice, but is stimulated by leptin. *Diabetes*. 47:294-7.
- Moan, A.; Hide, I.K., and Kjeldsen, S.E. (2006):** Metabolic and adrenergic characteristics of young men with insulin resistance. *Blood Pressure* 1996;5(Suppl 1):30-37
- Moeller, G.; and Adamski, J. (2006):** Multifunctionality of human 17 β -hydroxysteroid dehydrogenases. *Mol. Cell. Endocrinol.* 248, 47-55
- Moerman, E.J.; Teng, K.; Lipschitz, D.A., and Lecka-Czernik, B. (2004):** Aging activates adipogenic and suppresses osteogenic programs in mesenchymal marrow stroma/stem cells: the role of PPAR- γ 2 transcription factor and TGF- β /BMP signaling pathways. *Aging Cell* 3:379–389.



- Mohamed-Ali, V.; Pinkney, J.H., and Coppack, S.W. (2004):** Adipose tissue as an endocrine and paracrine organ. *Int J Obes* 22: 1145-1158.
- Mohanty, P.; Aljada, A.; Ghanim, H.; Hofmeyer, D.; Tripathy, D.; Syed, T.; Al-Haddad, W.; Dhindsa, S., and Dandona, P. (2004):** Evidence for a potent antiinflammatory effect of rosiglitazone. *J Clin Endocrinol Metab* 89:2728–2735
- Mokdad, A.H.; Bowman, B.A., and Ford, E.S. (2005):** The continuing epidemic of obesity and diabetes in the United States. *JAMA* 286:1195-1200
- Montagnani, M.; Golovchenko, I., and Kirn, I. (2004):** Inhibition of phosphatidylinositol 3-kinase enhances mitogenic actions of insulin in endothelial cells. *J Biol Chem* 277:1794-1799
- Montez, J.M.; Soukas, A.; Asilmaz, E.; Fayzikhodjaeva, G.; Fantuzzi, G. and Friedman, J.M. (2005)** Acute leptin deficiency, leptin resistance, and the physiologic response to leptin withdrawal. *Proceedings of the National Academy of Sciences, USA*, 102, 2537– 2542.
- Moore, L.L.; Bradlee, M.L.; and Singer, M.R. (2004).** BMI and waist
- Morin, C.L.; Eckel, R.H.; Marcel, T. and Pagliassotti, M.J. (1999):** High fat diets elevate tissue-derived tumor necrosis factor- α activity. *Endocrinology* 138: 4665-4671.
- Moustaid, N., and SUL, H. S. (1991):** Regulation of expression of the fatty acid synthase gene in 3T3-L1 cells by differentiation and triiodothyronine. *J.Biol.Chem.*266:18550-18554.



- Mullarkey, C. J.; Edelstein, D., and Brownlee, M.(2000):** Free radical generation by early glycation products: a mechanism for accelerated atherogenesis in diabetes. *Biochem. Biophys. Res. Commun.*, 173: 932-939.
- Nakagawa, T.; Hu, H.; Zharikov, S.; Tuttle, K.R.; Short, R.A., and Glushakova, O. (2006).** "A causal role for uric acid in fructose-induced metabolic syndrome". *Am J Phys Renal Phys* **290** (3): F625–F631.
- Nakajima, H.; Morishita, R., and Yamamoto, K. (2000):** Mitogenic and antiapoptotic actions of hepatocyte growth factor through ERK, STAT3, and AKT in endothelial cells. *Hypertension*. 37:581–586
- Nakajima, T.; Fujioka, S.; Tokunaga, K.; Matsuzawa, Y., and Tarui, S.(2000):** Correlation of intraabdominal fat accumulation and left ventricular performance in obesity. *Am J Cardiol*. 64: 369–373.
- Napslitanol. (1993):** The differentiation of white adipose cells an electron microscope study. *J Cell BioL.C*: 663.
- National Institutes of Health, National Heart, Lung, and Blood Institute.(1998)** Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults (National Institutes of Health Publication No. 98-4083). Rockville, MD, U.S. Department of Health and Human Services, Public Health Service
- Neumeier, M.; Weigert, J.; Schäffler, A.; Wehrwein, G., and Müller-Ladner, U. (2006):** Different effects of adiponectin isoforms in human monocytic cells. *J Leukoc Biol* 79:803–808.




- Nilsson, M.; Johnsen, R.; Ye, W.; Hveem, K., and Lagergren, J.(2003):** Obesity and estrogen as risk factors for gastroesophageal reflux symptoms. *JAMA*. 290:66-72.
- Nilsson, M.; Lundegårdh, G.; Carling, L.; Ye, W., and Lagergren, J.(2002):** Body mass and reflux oesophagitis: an oestrogen-dependent association? *Scand J Gastroenterol*. 37:626-30.
- Nindl, B.C.; Kraemer, W.J.; Arciero, P.J., and Samatallee, N. (2002)** Leptin concentrations experience a delayed reduction after resistance exercise in men. *Med Sci Sports Exerc* 34: 608–613.
- Ninomiya-Tsuji, J.; Torti, F. M., and Ringold, G. M.(1993)** Tumor necrosis factor-induced c-myc expression in the absence of mitogenesis is associated with inhibition of adipocyte differentiation. *Proc.Natl. Acad. Sci. USA* 90: 9611-9615.
- Ntambi, J.M., and Young-Cheul, K. (2000):** Adipocyte differentiation and gene expression. *J Nutr* 130:3122S–3126S
- Ogden, C.L.; Carroll, M.D. and Flegal, K.M. (2008):** High Body Mass Index for Age Among US Children and Adolescents, 2003—2008. *JAMA*. 299(20):2401—2405.
- Oishi, Y.; Manabe, I.; Tobe, K.; Tsushima, K.; Shindo, T.; Fujiu, K, Nishimura, G.; Maemura, K., and Yamauchi, T. (2005):** Krüppel-like transcription factor KLF5 is a key regulator of adipocyte differentiation. *Cell Metab* 1:27–39.
- Okubo, M.; Horinishi, A., and Saito, M. (2007).** "A novel complex deletion-insertion mutation mediated by Alu repetitive elements leads to lipoprotein lipase deficiency". *Mol. Genet. Metab*. 92 (3): 229–33.



- Ortega Gonzalez, A.; Peces-Barba, G.; Fernandez, I., and Chumbi, R. (2002):** Evolution of patients with chronic obstructive pulmonary disease, obesity hypoventilation syndrome or congestive heart failure in a respiratory monitoring unit. *Arch Bronconeumol* 42:423–429
- Ortiga-Carvalho, T.M.; Oliveira, K.J.; Soares, B.A.; and Pazos-Moura, C.C. (2002):** The role of leptin in the regulation of TSH secretion in the fed state: in vivo and in vitro studies. *J Endocrinol* 174: 121-125.
- Otto, T.C.; Lane, M.D., and Cox, M.M. (2005):** Adipose development: from stem cell to adipocyte. *Crit Rev Biochem Mol Biol* 40:229–242
- Ozata, M.; Oktenli, C.; Gulec, M.; Ozgurtas, T.; Bulucu, F., and Caglar, K. (2002):** Increased fasting plasma acylation-stimulating protein concentrations in nephrotic syndrome. *J Clin Endocrinol Metab* 87 :853 –858.
- P.P.(1997):** Tumor necrosis factor increases serum leptin levels in humans. *J Clin Endocrinol Metab.* 82:4080-2.
- Pai, J.K.; Pischon, T.; Ma, J.; Manson, J.E.; Hankinson, S.E.; Joshipura, K., and Curhan, G.C. (2004)** Inflammatory markers and the risk of coronary heart **disease** in men and women. *New England Journal of Medicine*, 16 (351), 2599– 2610.
- Pairault, J., and Green, H.(1979):** A study of the adipose conversion of suspended 3T3 cells by using glycerophosphate dehydrogenase as differentiation marker. *Proc. Natl. Acad. Sci. USA* 76: 5138-5142.




- Pajvani, U. B.; Trujillo, M. E.; Combs, T. P.; Iyengar, P.; Jelicks, L.; Roth, K. A.; Kitsis, R. N., and Scherer, P. E. (2005).** Fat apoptosis through targeted activation of caspase 8: a new mouse model of inducible and reversible lipoatrophy. *Nat. Med.* 11:797-803.
- Park, H.S.; and Lee, K.U. (2003):** Postmenopausal women lose less visceral adipose tissue during a weight reduction program. *Menopause*, 10:222-7.
- Parpal, S.; Karlsson, M.; Thorn, H.; and Stralfors, P.(2006)**sterol depletion disrupts caveolae and insulin receptor signaling for metabolic control via insulin receptor substrate 1, but not for mitogen-activated protein kinase control. *J Biol Chem* ;276:9670-9678
- Patel, N.; Tong, P.; Khayat, Z.A.; Huang, C.; Ueyama, A.; and Klip, A. (2004).**Insulin-induced cortical actin remodeling promotes GLUT4 insertion at cell membrane ruffles. *J. Clin. Invest.* 108, 371–381.
- Pausova, Z.; Deslauriers, B., and Gaudet, D. (2005)** Role of tumor necrosis factor-alpha gene locus in obesity and obesity-associated hypertension in French Canadians *Hypertension* , 36:14-19
- Perfetto, F.; Tarquini, R.; Simonini, G.; Bindi, G.; Mancuso, F.; Guiducci, S.; Matucci-Cerinic, M., and Falcini, F. (2005):** Circulating leptin levels in juvenile idiopathic arthritis: a marker of nutritional status? *Ann Rheum Dis* 64:149–152.




- PetrusChke, T., and Huner, H.(1993)** Tumor necrosis factor-alpha prevents the differentiation of human adipocyte precursor cells and causes delipidation of newly developed fat cells. *J. Clin. Endocrinol. Metab.* 76: 742-747.
- PetrusChke, T.; ROHRIG, K., and Huner, H.(1994)** Transforming growth factor beta (TGF-beta) inhibits the differentiation of human adipocyte precursor cells in primary culture. *Int. J. Obesity Related Metab. Disorders* 18: 532-536.
- Pickup, J.C.; Chusney, G.D.; Thomas, S.M., and Burt, D.(2000)** Plasma interleukin-6, tumour necrosis factor α and blood cytokine production in type 2 diabetes. *Life Sci* 67: 291-300.
- Poirier, P., and Despres, J.P.(2003):** Exercise in weight management of obesity. *Cardiol Clin.* 19: 459–470.
- Porras, A., and Santos, E. (1996).**The insulin/Ras pathway of adipocytic differentiation of 3T3 L1 cells: dissociation between Raf-1 kinase and the MAPK/RSK cascade. *Int. J. Obesity Related Metab. Disorders* 20, Suppl. 3: S43-S51.
- Porte, D. J.; Baskin, D.G.; and Schwartz, M.W.(2002):** Leptin and insulin action in the central nervous system. *Nutr Rev* 60 :S20 – S29; discussion S68–S87.
- Preston, T.; Brownell, H. and Raptis, L.(1997)** The timing of insulin/c-Ras signal is critical for its effect upon the differentiation of C3H10T1/2-derived preadipocytes. *Cancer Lett.* 115: 165-171.
- Prins, J. B., andO'Rahilly, S.(1997)** Regulation of adipose cell number in man. *Clin. Sci.* 92: 3-11
- Puranen, T.; Poutanen, M.; Ghosh, D.; Vihko, R., and Vihko, P.(2003):** *Endocrinology*.138:3532–3539.



- Radic, R.; Nikolic, V.; Karner, I.; Kosovic, P.; Kurbel, S., and Selthofer, R.(2003)** Circadian rhythm of blood leptin in obese and non-obese people. *Coll Antropol* ;27:555-561.
- Rajala, M. W., and Scherer, P. E. (2003).** The adipocyte: at the crossroads of energy homeostasis, inflammation, and atherosclerosis. *Endocrinology* 144: 3765-3773.
- Rakugi, H.; Kamide, K., and Ogihara, T.(2005)** Vascular signaling pathways in the metabolic syndrome. *Curr Hypertens Rep*;4;105-111
- Rampino, T.; Libetta, C., and De Simone, W. (2003):** Hemodialysis stimulates hepatocyte growth factor release. *Kidney Int.* 53:1382–1388
- Ramsay, T.G.(2000).** Fat cells. *Endocrinol Metab Clin North Am* 25: 847-879
- Rasgon, N., and Jarvik, L. (2004):** Insulin resistance, affective disorders, and Alzheimer's disease: review and hypothesis. *J Gerontol A Biol Sci Med Sci* 59:178-183.
- Rask, E.; Olsson, T., and Soderberg, S. (2001)** Tissue-specific dysregulation of cortisol metabolism in human obesity. *J Clin Endocrinol Metab*;86:1418-1421
- Reaven, G., and Banting Lecture. (2001).** Role of insulin resistance in human disease. *Diabetes* , 37:1595-1607.
- Reaven, G.M.; Lithell, H., and Landsberg, L.(2003)** Mechanisms of disease: hypertension and associated metabolic abnormalities. The role of insulin resistance and the sympathoadrenal system. *N Engl J Med* , 334:374-381




- Rebuffe-Scrive, M.; Enk, L. ; Crona, N.; Loannroth, P.; Abrahamsson, L.; Smith, U., and Bjorntorp, P.(2005):** Fat cell metabolism in different regions in women: effect of menstrual cycle, pregnancy and lactation. *J. Clin. Invest.* 75: 1973–1976.
- Reilley, M.P., and Racier, D.J.(2003)** The metabolic syndrome. More than the sum of its parts? *Circulation*; 108:1546-1551
- Reisin, E.; Abel, R., and Modan, M. (2002):** Effect of weight loss without salt restriction on the reduction of blood pressure in overweight hypertensive patients. *N Engl J Med* 298(1):1-6
- Renahan, A.G.; Zwahlen, M., and Minder, C. (2004):** Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis. *Lancet* **363**: 1346–53.
- Richelson, B. (2002):** Release and effects of prostaglandins in adipose tissue. *Prostaglandins leukotrienes Essential Fatty Acids*; 47: 171-182.
- Richon, V. M.; Lyle, R. E., and Mcgehee, J,R. (1997)** Regulation and expression of retinoblastoma proteins p107 and p130 during 3T3-L1 adipocyte differentiation. *J. Biol. Chem.* 272: 10117-10124,
- Rincon ,M.; Muzumdar, R.; Atzmon, G., and Barzilai, N. (2005).** "The paradox of the insulin/IGF-1 signaling pathway in longevity.". *Mech. Ageing Dev.* 125 (6): 397-403.
- Rittershaus, C.; Miller, D.P.; Thomas, L.J.; Picard, M.D., and Honan, C.M. (2000):** Vaccine-induced antibodies inhibit CETP activity in vivo and reduce aortic lesions in a rabbit model of atherosclerosis. *Arterioscler Thromb Vasc Biol.* 20:2106–2112.



- Robenek, H.; Hofnagel, O.; Buers, I.; Robenek, M. J.; Troyer, D., and Severs, N. J. (2006):** Adipophilin-enriched domains in the ER membrane are sites of lipid droplet biogenesis. *J. Cell Sci.* 119: 4215–4224
- Roberts, A.B.; Glick, A.B., and Sporn, M.B. (1998):** Interrelationships between two families of multifunctional effectors: retinoids and TGF- β . In: Morris-Kay G. (ed.) Oxford Science Publications, pp. 137-148.
- Romijn, J.A.; and Fliers, E.(2005)** Sympathetic and parasympathetic innervation of adipose tissue: metabolic implications. *Curr Opin Clin Nutr Metab Care.* 8(4):440-4.
- Rosen, E. D.; Walkey, C. J.; Puigserver, P., and Spiegelman, B. M. (2006).** Transcriptional regulation of adipogenesis. *Genes Dev.* 14:1293–1307.
- Rosen, E.D. (2006):** The transcriptional basis of adipocyte development. *Prostaglandins Leukot Essent Fatty Acids* 73:31–34.
- Rosicka, M.; Krsek, M.; Matoulek, M.; Jarkovska, Z.; Marek, J., and Justova, V. (2003)** Serum ghrelin levels in obese patients: the relationship to serum leptin levels and soluble leptin receptor levels. *Physiol Res* ;52:61-66.
- Rotter Sopasakis, V.; Larsson, B.M.; Johansson, A.; Holmang, A. and Smith, U. (2004)** Short-term infusion of interleukin-6 does not induce insulin resistance *in vivo* or impair insulin signalling in rats. *Diabetologia*, 47, 1879– 1887.




- Ruan, H.; Hachohen, N., and Golub, T.R. (2005).** Tumor necrosis factor- α suppresses adipocyte-specific genes and activates expression of preadipocyte genes in 3T3-L1-adipocytes. *Diabetes*, 51:1319-1336
- Rubins, H.B.; Robins, S.J., and Collins, D.(2000):** Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. *N Eng J Med* 341(6):410-8
- Ruotolo, G., and Howard, B.V. (2002)** Dyslipidemia of the metabolic syndrome. *Curr Cardiol Rep*; 4:494-500
- Rye, K.A.; Clay, M.A., and Barter, P.J.(1999):** Remodelling of high density lipoproteins by plasma factors. *Atherosclerosis*. 145:227–238.
- Sabbagh, A.S.; Otrock, Z.K., and Mahfoud, Z.R. (2007).** "Angiotensin-converting enzyme gene polymorphism and allele frequencies in the Lebanese population: prevalence and review of the literature.". *Mol. Biol. Rep.* 34 (1): 47-52.
- Saghizadeh, M.; Ong, J.M., and Barvey, W.T. (2003).** The expression of TNF- α by human muscle: relationship to insulin resistance. *J Clin Invest* , 97:1111-1116
- Saleh, J.; Summers, L.K.; Cianflone, K.; Fielding, B.A.; Sniderman, A.D., Frayn, K.N. (2005):** Coordinated release of acylation stimulating protein (ASP) and triacylglycerol clearance by human adipose tissue in vivo in the postprandial period. *J Lipid Res* 39:884 –891.




- Salmenniemi, U.; Ruotsalainen, E., and Pihlajamaki, J. (2004).** Multiple abnormalities in glucose and energy metabolism and coordinated changes in levels of adiponectin, cytokines, and adhesion molecules in subjects with metabolic syndrome. *Circulation*;110:3842-3848
- Saltiel, A.R.(2007):** You are what you secrete. *Nat Med* ;7:887-888
- Samuel, V.T.; Liu, Z.X.; Qu, X.; Elder, B.D.; Bilz, S.; Befroy, D.; Romanelli, A.J., and Shulman, G.I. (2004):** Mechanism of hepatic insulin resistance in non-alcoholic fatty liver disease. *J Biol Chem* 279:32345–32353
- Sarafidis, P.A., and Bakris, G.L. (2006):** Insulin and endothelin: an interplay contributing to hypertension development? *J Clin Endocrinol Metab.* 92:379–385.
- Sartipy, P., and Loskutoff, D.J. (2003):** Monocyte chemoattractant protein 1 in obesity and insulin resistance. *Proc Natl Acad Sci USA* 100:7265–7270.
- Sasso, F.C.; Carbonara, O., and Nasti, R. (2003):** glucose metabolism and coronary heart disease in patients with normal glucose tolerance. *JAMA* ;291:1857-1863
- Schäffler, A.; Ehling, A.; Neumann, E.; Herfarth, H.; Tarner, I.; Gay, S.; Schölmerich, J., and Müller-Ladner, U. (2003).** Adipocytokines in synovial fluid. *J. Am. Med. Assoc.* 290: 1709-1710.
- Scherer, P. E.; Williams, S.; Fogliano, M.; Baldini, G. , Andlodish, H. F.(1995)** A novel serum protein similar to C1q, produced exclusively in adipocytes. *J. Biol. Chem.* 270: 26746-26749,.




- Schmidt, W.; Poll-Jordan, G., and Loffler, G.(1990)** Adipose conversion of 3T3-L1 cells in a serum-free culture system depends on epidermal growth factor, insulin-like growth factor I, corticosterone, and cyclic AMP. *J. Biol. Chem.* 265: 15489-15495.
- Schwartz, M.W.; Woods, S.C.; Porte, D.; Seeley, R.J., and Baskin, D.G. (2007):** Central nervous system control of food intake. *Nature* 404:661-71.
- Schwarz, E. J.; Regiato, M. J., and Shao, D. S.(1997)** Retinoic acid blocks adipogenesis by inhibiting C/EBPbeta-mediated transcription. *Mol. Cell. Biol.* 17: 1552-1561,.
- Seckl, J.R. (2004):** 11beta-hydroxysteroid dehydrogenases: changing glucocorticoid action. *Current Opinions in Pharmacology*, 4, 597– 602.
- Serrero, G., and Millis, D.(1996).** Physiological role of epidermal growth factor on adipose tissue development in vivo. *Proc. Natl. Acad. Sci. USA* 88: 3912-3916.
- Sfeir, Z.; Ibrahimi, A.; Amri, E.; Grimaldi, P.; and Abumrad, N.(1997)** Regulation of FAT/CD36 gene expression: further evidence in support of a role of the protein in fatty acid binding/transport. *Prostaglandins Leukotrienes Essential Fatty Acids* 57: 17-21.
- Shen, W.J.; Patel, S.; Hong, R., and Kraemer, F.B. (2000):** Hormone-sensitive lipase functions as an oligomer. *Biochemistry*, 39, 2392-8.
- Shi, Y. and Massague, J. (2003).** Mechanisms of TGF- β signaling from cell membrane to the nucleus. *Cell* 113: 685–700.



- Shillabeer, G.; Kumar, V.; Tibbo, E. and Lay, D.C. (1998):** Arachidonic acid metabolites at the lipo-oxygenase as well as the cyclooxygenase pathway may be involved in regulating preadipocyte differentiation. *Metabolism* 47: 461-466.
- Shimizu, H.; Shimomura, Y.; Nakanishi, Y.; Futawatari, T.; Ohtani, K.; Sato**
- Shirasaka, T.; Takasaki, M., and Kannan, H. (2003).** Cardiovascular effects of leptin and orexins. *Am J Physiol Regul Integr Comp Physiol*; 284: R639-R651.
- Shugart, E. C.; Levenson, A. S., and Constance C. M. (1995)** Differential expression of *gas* and *gadd* genes at distinct growth arrest points during adipocyte development. *Cell Growth Differ.* 6: 1541-1547.
- Siggins, S.; Bykov, I.; Hermansson, M.; Somerharju, P.; Lindros, K., Singh, P.P.; Singh, M., and Mastana, S.S. (2006).** "APOE distribution in world populations with new data from India and the UK.". *Ann. Hum. Biol.* 33 (3): 279-308.
- Sinha, R.; Dufour, S.; Petersen, K.F.; LeBon, V.; Enoksson, S.; Ma, Y.Z.; Savoye, M.; Rothman, D.L.; Shulman, G.I., and Caprio, S. (2005):** Assessment of skeletal muscle triglyceride content by (1)H nuclear magnetic resonance spectroscopy in lean and obese adolescents: relationships to insulin sensitivity, total body fat, and central adiposity. *Diabetes*, 51:1022-
- Smas, C. M.; Chen, L. and h. S. Sul. (1997):** Cleavage of membrane-associated pref-1 generates a soluble inhibitor of adipocyte differentiation. *Mol. Cell. Biol.* 17: 977-988.




- Sowers, J.R.; and Lester, M.(2004):** Diabetes and cardiovascular disease. *Diabetes Care* 22 (suppl 3):c14-c20.
- Spate, U., and Schulze, P.C.(2004):** Proinflammatory cytokines and skeletal muscle. *Curr Opin Clin Nutr Metab Care* 7:265-269.
- Spiegelman, B.M., and Farmer, S. R. (1982)**Decreases in tubulin and actin gene expression prior to morphological differentiation of 3T3 adipocytes. *Cell* 29: 53-60.
- Spiegelman, B.M.; Choy, L.; Hotamisligil, G.S.; Graves, R.A., and Tontonoz, P. (2003):** Regulation of adipocyte gene expression in differentiation and syndromes of obesity/diabetes. *J Biol Chem* 268: 6823-6826.
- Spuler, S.; Kalbhenn, T.; Zabojszcza, J.; van Landeghem, F. K. H.; Ludtke, A.; Wenzel, K.; Koehnlein, M.; Schuelke, M.; Ludemann, L.; Schmidt, H. H.(2007) :** Muscle and nerve pathology in Dunnigan familial partial lipodystrophy. *Neurology* 68: 677-683.
- Sriram, K.; Benkovic, S.A.; Miller, D.B.; and O’Callaghan, J.P.(2002):** Obesity exacerbates chemically induced neurodegeneration. *Neuroscience*. 115: 1335–1346.
- St. Jeor, S.T.(1993)** The role of weight manegment in the health of women. *J. Amer. Dietetic Assoc.* 93:1007-1012.
- Stamets, K.; Taylor, D.S.; Kunselman, A.; Demers, L.M.; Pelkman, C.L. and Legro, R.S. (2004)** A randomized trial of the effects of two types of short-term hypocaloric diets on weight loss in women with polycystic ovary syndrome. *Fertil Steril* 81,630–637.



- Strazzullo, P.; Barbato, A.; Galletti, F.; Barba, G.; Siani, A.; Iacone, R.; D'Elia, L.; Russo, O.; Versiero, M.; Farinaro, E. and Cappuccio, F.P.(2006):** Abnormalities of renal sodium handling in the metabolic syndrome. Results of the Olivetti Heart Study. *J Hypertens.* 24:1633–1639.
- Suryawan, A., and Hu, C.Y.(1997)** Effect of retinoic acid on differentiation of cultured pig preadipocytes. *J. Anim. Sci.* 75: 112-117.
- Suryawan, A.; Swanson, L. V. and Hu, C.Y.(1997):** Insulin and hydrocortisone, but not triiodothyronine, are required for the differentiation of pig preadipocytes in primary culture. *J. Anim. Sci.* 75: 105-111.
- Susen, S.; Sautiere, K.; Mouquet, F.; Cuilleret, F.; Chmait, A.; McFadden, E.P.; Hennache, B.; Richard, F. and de Groote, P.(2007):** Serum hepatocyte growth factor levels predict long-term clinical outcome after percutaneous coronary revascularization. *Eur Heart J* 26: 2387–2395.
- Swierczynski, J.; Korczynska, J.; Goyke, E.; Adrych, K.; Raczynska, S.; and Sledzinski, Z.(2007):** Serum hepatocyte growth factor concentration in obese women decreases after vertical banded gastroplasty. *Obes Surg* 15: 803–808.
- Tabe, Y.; Konopleva, M.; Igari, J. and Andreeff, M. (2004):** Spontaneous migration of acute promyelocytic leukemia cells beneath cultured bone marrow adipocytes with matched expression of the major histocompatibility complex. *Rinsho Byori* 52:642–648.



- Taegtmeyer, H.; McNulty, P. and Young, M.E. (2002):** Adaptation and maladaptation of the heart in diabetes, part I: general concepts. *Circulation* 105:1727-1733.
- Targher, G.; Bertolini, L.; Poli, F.; Rodella, S.; Scala, L.; Tessari, R.; Zenari, L. and Falezza, G. (2005):** Nonalcoholic fatty liver disease and risk of future cardiovascular events among type 2 diabetic patients. *Diabetes* 54:3541–3546
- Tartaglia, L.A. (1997):** The leptin receptor. *J. Biol. Chem.* 272: 6093-6096.
- Tezuka, M.; Irahara, M.; Ogura, K.; Kiyokawa, M. and Tamura, T. (2002):** Effects of leptin on gonadotropin secretion in juvenile female rat pituitary cells. *Eur J Endocrinol* 146: 261- 266.
- The Nordic Myeloma Study Group Seidel C, Borset, M.; Turesson, I.; Abildgaard, N.; Sundan, A. and Waage, A. (2004):** Elevated serum concentrations of hepatocyte growth factor in patients with multiple myeloma. *Blood*. 91:806–812
- Thorne, A.; Lonnqvist, F.; Nordfors, L.; Jansson, M.; Schalling, M. and Arner, P.(2003):** Leptin secretion from adipose tissue in women. Relationship to plasma levels and gene expression. *J Clin Invest.* 99:2398-404.
- Tilg, H., and Moschen, A. R. (2006):** Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nat. Rev. Immunol.* 6: 772–783.



- Timchenko, N. A.; Wilde, M.; Nakanishi, M.; Smith, J.R. and Darlington, G. J.(1996):** CCAAT/enhancer-binding protein alpha (C/EBP alpha) inhibits cell proliferation through the p21 (WAF-1/CIP-1/SDI-1) protein. *Genes Dev.* 10: 804-815, 276.
- Tiziana Ronti, Graziana Lupattelli, and Elmo Mannarino (2006)** The endocrine function of adipose tissue: an update *Clinical Endocrinology* 64 (4) , 355–365
- Tomas, E.; Tsao, T.S.; Saha, A.K.; Murrey, H.E.; Zhang, C.C. and Itani, S.L.(2002):** Enhanced muscle fat oxidation and glucose transport by ACRP 30 globular domain: acetyl CoA carboxylase inhibition and AMP-activated protein kinase activation. *Proc Natl Acad Sci U S A* 99:16309-16313.
- Tong, Q.; Dalgin, G.; Xu, H.; Ting, C.N.; Leiden, J.M. and Hotamisligil, G.S. (2000)** Function of GATA transcription factors in preadipocyte–adipocyte transition. *Science* 290: 134–138
- Tong, Q.; Tsai, J.; Tan, G.; Dalgin, G. and Hotamisligil, G.S. (2005)** Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Mol Cell Biol* 25: 706–715
- Trauner, M . and Boyer, J.L.(2003):** Bile salt transporters: molecular characterization, function, and regulation. *Physiol Rev* ;83:633–71.
- Trujillo, M.E.; Lee, M.J.; Sullivan, S.; Feng, J.; Schneider, S.H.; Greenberg, A.S. and Fried, S.K. (2006):** Tumor necrosis factor α and glucocorticoid synergistically increase leptin production in



human adipose tissue: role for p38 mitogen-activated protein kinase. *J Clin Endocrinol Metab* 91:1484–1490.

Tsai, J.; Tong, Q.; Tan, G.; Dalgin, G. and Hotamisligil, G.S. (2005)

Interaction between GATA and the C/EBP family of transcription factors is critical in GATA-mediated suppression of adipocyte differentiation. *Mol Cell Biol* 25: 706–715

Tsigos, C.; Papanicolaou, D.A.; Mitsiadis, C.S.; Kyrou, I. and

Chrousos, G.P.(2003). Dose effects of recombinant interleukin-6 on pituitary hormone secretion and energy expenditure. *Neuroendocrinology* 66: 54-62.

Turner, A.J. and Hooper, N.M. (2002): The angiotensin-converting

enzyme gene family: genomics and pharmacology. *Trends Pharmacol Sci* 23:177–183.

Tutuian, R. and Castell, D.O.(2007): Pressure details from the weight-

challenged gastroesophageal junction: more than the usual suspects. *Gastroenterology* 130:988–9.

Ukkola, O. and Santaniemi, M.(2002) Adiponectin: a link between

excess adiposity and associated comorbidities? *J Mol Med*;80:696-702

Umek, R.M.; Friedman, A. D. and Mcknight, L.(1991) CCAAT-

enhancer binding protein: a component of a differentiation switch. *Science* 251: 288-292.

Unger, R.H.; Zhou, Y.T. and Orci, L.(2003): Regulation of fatty acid

homeostasis in cells: novel role of leptin. *Proc Natl Acad Sci USA*. 96:2327-32.




- Urs, S.; Smith, C.; Campbell, B.; Saxton, A.M.; Taylor, J.; Zhang, B.; Snoddy, J.; Jones Voy, B. and Moustaid-Moussa, N. (2004):** Gene expression profiling in human preadipocytes and adipocytes by microarray analysis. *J Nutr* 134:762–770.
- Valsamakis, G.; Anwar, A.; Tomlinson, J.W. and Shackleton, C.H. (2004):** 11 β -hydroxysteroid dehydrogenase type 1 activity in lean and obese males with type 2 diabetes mellitus. *Journal of Clinical Endocrinology and Metabolism*, 89, 4755– 4761.
- Van der Kooy, K.; Leenen, R. and Seidell, J.C. (1995).** Waist-hip ratio is a poor predictor of changes in visceral fat. *Am J Clin Nutr* , 57:327-333
- Van der Vusse, G.J.; Van Bilsen, M. and Glatz, J.F. (2000):** Cardiac fatty acid uptake and transport in health and disease. *Cardiovasc res* 45:279-293.
- Van Snick, J.(1996).** Interleukin-6: an overview. *Annu Rev Immunol* 8: 253-278.
- Vanderstraeten-Gregoire, F. (1989)** *La Differentiation In Vitro des Precurseurs des Cellules Adipeuses de Rat* (PhD thesis). Louvain: Universite Catholique de Louvain.
- Vassaux, G.R.; Negrel, G.; Ailhaud, and Gaillard, D. (2003):** Proliferation and differentiation of rat adipose precursor cells in chemically defined medium: differential action of anti-adipogenic agents. *J. Cell. Physiol.* 161: 249-256, 1994.
- Veniant, M.M. and LeBel, C.P.(2003)** Leptin: from animals to humans. *Curr Pharm Des*;9:811-818.



- Vidal, H. (2001)** Gene expression in visceral and subcutaneous adipose tissues. *Ann Med*;33:547-555
- Vihko, P.; Isomaa, V. and Ghosh, D. (2003):** Structure and function of 17 β -hydroxysteroid dehydrogenase type 1 and type 2. *Molec. Cell. Endocrin.*, 171, 71-76
- Vojarova, B.; Weyer, C. and Lindsay, R.S. (2005).** High white blood cell count is associated with a worsening of insulin sensitivity and predicts the development of type 2 diabetes. *Diabetes*, 51:455 - 461
- Wabitsch, M.E.; Heinze, H.; Hauner, R.M.; Shymko, W. M.; Teller, P.; De meyts, and Ilondo, M.M. (1996):** Biological effects of human growth hormone in rat adipocyte precursor cells and newly differentiated adipocytes in primary culture. *Metabolism* 45: 34-42.
- Wajchenberg, B.L. (2004):** Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev* 21:697–738.
- Wake, D.J.; Rask, E.; Livingstone, D.E.; Soderberg, S.; Olsson, T. and Walker, B.R. (2003):** Local and systemic impact of transcriptional up-regulation of 11 β -hydroxysteroid dehydrogenase type 1 in adipose tissue in human obesity. *Journal of Clinical Endocrinology and Metabolism*, 88, 3983– 3988.
- Wallace, A.M. (2000)** Measurement of leptin and leptin binding in the human circulation. *Ann Clin Biochem* ;37:244-252.



- Wallace, A.M.; McMahon, A.D.; Packard, C.J.; Kelly, A.; Shepherd, J.; Gaw, A. and Sattar, N. (2001)** Plasma leptin and the risk of cardiovascular disease in the West of Scotland Coronary Prevention Study (WOSCOPS). *Circulation*, 104, 3052– 3056
- Wang, H.; Johnson, G. L.; Liu, X. and Malbon, C. C. (1996):** Repression of adipogenesis by adenylyl cyclase stimulatory G-protein alpha subunit is expressed within region 146-220. *J. Biol. Chem.* 271: 22022-22029.
- Wang, H.M.; Goligorsky, S. and Malbon, C. C. (1997):** Temporal activation of Ca^{2+} -calmodulin-sensitive protein kinase type II is obligate for adipogenesis. *J. Biol. Chem.* 272: 1817-1821.
- Wang, M. (2005):** The role of glucocorticoid action in the pathophysiology of the metabolic syndrome. *Nutrition and Metabolism*, 2 (2), 3.
- Wang, T.J.; Farise, H. and Levy, D. (2004):** Obesity and the risk of new-onset atrial fibrillation. *JAMA* ;292: 2471-2477
- Webber, J. (2003)** Energy balance in obesity. *Proc Nutr Soc*, 62:539-543.
- Wei, E.K.; Giovannucci, E.; Fuchs, C.S.; Willett, W.C.; Mantzoros, C.S. and Weigert, J. (2005):** The adiponectin paralog CORS-26 has anti-inflammatory properties and is produced by human monocytic cells. *FEBS Lett* 579:5565–5570.
- Weiner, F. R.; Smith, P. J.; Wertheimer, S. and Rubin. C. S. (1991)** Regulation of gene expression by insulin and tumor necrosis factor alpha in 3T3-L1 cells. Modulation of the transcription of genes encoding acyl-CoA synthetase and stearyl-CoA desaturase-1. *J. Biol. Chem.* 266: 23525-23528.



- Weisberg, S. P.; McCann, D.; Desai, M.; Rosenbaum, M.; Leibel, R. L. and Ferrante, Jr. A. W. (2005).** Obesity is associated with macrophage accumulation in adipose tissue. *J. Clin. Investig.* 112:1796-1808.
- Weyer, C.; P'unahashi, T.; Tamaka, S.; Hotta, K.; Matsuzawa, Y.; Pratley, E. and Tataranni, P.A. (2004):** Hypoadiponectinemia in obesity and type II diabetes and association with insulin resistance and hyperinsulinemia. *J Clin. Endocrinol. Metab.*; 86: 1930-1935.
- Wilhelmsen, L.; Rosengren, A.; Eriksson, H. and Lappas, G.(2005):** Heart failure in the general population of men—morbidity, risk factors and prognosis. *J Intern Med.* 249: 253–261.
- Winder, W.W.(2004).** AMP-activated protein kinase: possible target for treatment of type 2 diabetes. *Diabetes Technol*, 2:441-448
- Wolfsdorf, J.; Sadeghi-Nejad, A. and Senior, B. (2008)** Leptin-replacement therapy in lipodystrophy. *New England Journal of Medicine*, 346.
- Wolk, A.; Lindblad, P. and Adami, H. O. (2006)** Nutrition and Renal Cell Cancer, *Cancer Causes & Control* 7, 5–18.
- Wong, M.L.; Licinio, J.; Yildiz, B.O.; Mantzoros, C.S.; Prolo, P. and Kling, M. (2004).** Simultaneous and continuous 24-hour plasma and cerebrospinal fluid leptin measurements: dissociation of concentrations in central and peripheral compartments. *J Clin Endocrinol Metab*, 89:258-265.
- Wootan, M.G.; Bernlohr, D.A. and Storch, J.(2003):** Mechanism of fluorescent fatty acid transfer from adipocyte fatty acid binding protein to membranes. *Biochemistry* 32: 8622-8627



- Wu, J.; Srinivasan, S.V.; Neumann, J.C. and Lingrel, J.B. (2005):** The KLF2 transcription factor does not affect the formation of preadipocytes but inhibits their differentiation into adipocytes. *Biochemistry* 44:11098–11105.
- Wu, Z. (1999a)** Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell* 98: 115–124
- Wu, Z.; Bucher, N. L. and Farmer, S. R.(1996)** Induction of peroxisome proliferator-activated receptor gamma during the conversion of 3T3 fibroblasts into adipocytes is mediated by C/EBPbeta, C/EBPdelta, and glucocorticoids. *Mol. Cell. Biol.* 16: 4128-4136.
- Xing, H.; Northrop, J. P. and Grove, J. R.(1997)** TNF alpha-mediated inhibition and reversal of adipocyte differentiation is accompanied by suppressed expression of PPARgamma without effects on Pref-1 expression. *Endocrinology* 138: 2776-2783.
- Xu, A.; Wang, Y.; Keshaw, H.; Xu, L.Y.; Lam, K.S. and Cooper, G.J.(2005):** The fatderived hormone adiponectin alleviates alcoholic and nonalcoholic fatty liver disease in mice. *J Clin Invest* 112: 91-100.
- Xu, A.; Wang, Y.; Xu, J. Y.; Stejskal, D.; Tam, S.; Zhang, J.; Wat, N. M.; Wong, W. K. and Lam , K. S. (2006).** Adipocyte fatty acid-binding protein is a plasma biomarker closely associated with obesity and metabolic syndrome. *Clin. Chem.* 52: 405–413.
- Xu, F.P.; Chen, M.S.; Wang, Y.Z.; Yi, Q.; Lin, S.B.; Chen, A.F. and Luo, J.D. (2004)** Leptin induces hypertrophy via endothelin-1-



reactive oxygen species pathway in cultured neonatal rat cardiomyocytes. *Circulation*, 110, 1269– 1275.

Xue, J. C.; Shwartz, E. J.; Chawla, A. and Lazar, M. A(1996). Distinct stages in adipogenesis revealed by retinoid inhibition of differentiation after induction of PPARgamma. *Mol. Cell. Biol.* 16: 1567-1575.

Yaffe, K.; Kanaya, A. and Lindquist, K. (2004):The metabolic syndrome, inflammation, and risk of cognitive decline. *JAMA* 292:2237-2242.


Yamauchi, J.; Hirasawa, A. and Miyamoto, Y. (2000) Role of Db1's big sister in the anti-mitogenic pathway from alpha1B-adrenergic receptor to c-Jun N-terminal kinase. *Biochem Biophys Res Commun*;296:85-92

Yamauchi, T., Oike, Y.; Kamon, J.; Waki, H.; Komeda, K. and Tsuchida, A. (2002) Increased insulin sensitivity despite lipodystrophy in Crebbp heterozygous mice. *Nat. Genet.* 30:221–226.

Yatagai, T.; Nishida, Y.; Nagasaka, S.; Nakamura, T.; Tokuyama, K.; Shindo, M.; Tanaka, H. and Ishibashi, S.(2003): Relationship between exercise training-induced increase in insulin sensitivity and adiponectinemia in healthy men. *Endocr J* 50: 233–238

Yeh, W. C.; Bierer, B. E. and Mcknight, S. L.(1995) Rapamycin inhibits clonal expansion and adipogenic differentiation of 3T3-L1 cells. *Proc. Natl. Acad. Sci. USA* 92: 11086-11090.

Yokota, T.; Meka, C.S.; Kouro, T.; Medina, K.L.; Igarashi, H. and Takahashi, M. (2003): Adiponectin, a fat cell product, influences the earliest lymphocyte precursors in bone marrow cultures by



activation of the cyclooxygenase-prostaglandin pathway in stromal cells. *J Immunol* 171:5091–5099.

Youngstrom, T.G. and Bartness, T.J.(1995): Catecholaminergic innervation of white adipose tissue in Siberian hamsters. *Am. J. Physiol.* 268(*Regulatory Integrative Comp. Physiol.* 37): R744-R751

Yu, H. and Rohan, T. (2000). Role of the insulin-like growth factor family on cancer development and progression. *J Natl Cancer Inst*, 92, 1472-89.

Yudkin, J.S.; Eringa, E. and Stehouwer, C.D. (2005) ‘Vasocrine’ signalling from perivascular fat: a mechanism linking insulin resistance to **vascular** disease. *Lancet*, 365, 1817– 1820.

Yudkin, J.S.; Kumari, M.; Humphries, S.E. and Mohamed-Ali,V.(2000): Inflammation, obesity, stress and coronary heart disease: is interleukin-6 the link? *Atherosclerosis* 148: 209-214.

Yusuf, S.; Dagenais, G. and Pogue, J. (2004): vitamin E risk patients: the heart outcomes prevention evaluation study investigators. *N Engl J Med* 342:154-160.

Zabeau, L.; Lavens, D.; Peelman, F.; Eyckerman, S. and Vandekerckhove, J. (2003): The ins and outs of leptin receptor activation. *FEBS Lett* 546:45-50.

Zaccaria, M.; Ermolao, A.; Roi, G.S.; Englaro, P.; Tegov, G.; and Varnier, M.(2002): Leptin reduction after endurance races differing in duration and energy expenditure.*Eur J Appl Physiol*87:108–111.



- Zackova, M.; Skobisova, E.; Urbankova, E.; Jezek, P. (2005).** Activating omega-6 polyunsaturated fatty acids and inhibitory purine nucleotides are high affinity ligands for novel mitochondrial uncoupling proteins UCP2 and UCP3. *J Biol Chem* 278(23): 20761-20769.
- Zhou, M.; Lin, B.Z.; Coughlin, S.; Vallega, G.; Pilch, P.F. (2005).** UCP-3 expression in skeletal muscle: Effects of exercise, hypoxia, and AMP-activated protein kinase. *Am J Physiol Endocrinol Metab* 279(3): E622-E629.
- Zierath, J.R.; Livingston, J.N.; Thorne, A.(1998):** Regional difference in insulin inhibition of non-esterified fatty acid release from human adipocytes: relation to insulin receptor phosphorylation and intracellular signaling through the insulin receptor substrate-1 pathway. *Diabetologia* 41:1343-1354
- Zieske, A.W.; Malcom, G.T. and Strong, J.P.(2002):** Natural history and risk factors of atherosclerosis in children and youth: the PDAY study. *Pediatr Pathol Mol Med.* 21: 213–237.
- Zumbach, M.S.; Boehme, M.W.J.; Wahl, P.; Stremmel, W.; Ziegler, R. and Nawroth, P.P. (1997):** Tumor necrosis factor increases serum leptin levels in humans. *J Clin Endocrinol Metab* 82: 4080-4082