
REFERENCES

- Abu-Ghareeb, A. (1968) :Serodiagnosis methods in patients with typhoid fever Gaz, Egypt pediat. Assoc. 16 : 232.
- Ahvonon, P., Elli, J. and Hao, K. (1969) : Marked cross agglutination between Brucella and a subtype of Yersinia enterocolitica. Acta. Path. Microbiol. Scandinav. 75 : 291-95.
- Alder, B.B. and Splitter G.A. (1988) : Determination of bovine lymphocyte responses to extracted proteins of Brucella abortus by using protein immunoblotting. Infection and Immunity. 56 (10) : 2581-86.
- Aleixo, A.G. and Balasurbramanian, S. (1988) : A fluorescent enzyme immunoassay for Salmonella detection. J. of immunoassay. 9 (1) : 83-95.
- Allardyce, R.A. and Bienestock, J. (1984) : The mucosal immune system in health and disease with an emphasis on parasitic infection. Bull W.H.O. 62 : 7-25.
- Appassakij, H.N., Napatown, B., Suttipant, S. and Tssnee, S. (1987) : Enzyme-linked immunosorbent assay for detection of Salmonella typhi protein antigen. J. Clin. Microbiol. 25 : 273 : 277.
- Ashida, T.J. (1949) :Studies on virulence antigen of salmonella and its isolation J. Exper. Med. 20 : 181.
- Ashkenazi, S., Thomas, G., Barbara, E. and Larry, K. (1988) : Quantitative analysis and partial characterization of cytotoxin produced by Salmonella strains. Infection nad

-
- Immunology. 56 (12) : 3089-94.
- Ashkanazi, S., Cleary, K.P., Pickering, L.K. and Cleary, T.G. (1990) : The association of shigella toxin and other cytotoxins with the neurologic manifestation of shigellosis. J. Infect. Dis. 161 (5) : 961.
- Baker, E., Roberta, E., Richard, B. and Matthew, A. (1954) : The Vi antigens of Enterobacteriaceae. J. Immunol. 83 : 680-86.
- Bang, B. (1897) : Cited from Topley and Wilson (1984) : Principles of bacteriology virology and immunity 7th Edition. Vol.3 p. 411.
- Barrett, T.J., Synder, J.D., Blake, P.A. and Feeley, J.C. (1982) : Enzyme-linked immuno sorbent assay for detection of S. typhi Vi antigen in urine from typhoid patients. J. Clin. Microbiol. 15 : 235-37.
- Bastuji, B.G., Raml, A.B., Gorad, D. and Joseph, N.L. (1990) : Sodium dodecyl sulfate immunoblotting analysis of smooth lipopolysaccharide heterogeneity among Brucella biovars related to A and B specificities. J. Clin. Microbiol. 28 (10) : 2169-74.
- Bayer and Raehl (1904) : J. Med. Res. 12 : 313. Cited from Topley and Wilson (1983). Principles of Bacteriology & Virology and Immunology. Ch. 37: 332 - 51.
- Beal, C., Viens, P., Grant, R. and Hughes, J. (1970) : A new technique for sampling duodenal contents demonstration of upper small bowel pathogen. Am. J. Trop. Med. and Hyg. 19 : 349-52.
- Benavente, L., Gotuzzo, E., Gurra, J. and Brauo, N. (1984) : Diagnosis of typhoid fever using a string capsule device, Trans.

- Roy.Soc. Trop. Med and Hyg.78 : 404-406.
- Bergey, E. (1984) : Mannual of systematic Bacter. Vol. 1 : 381-86.
- Bevan, L. E. W. (1921-1922) : Trans. R. Soc. Trop. Med. Hyg. 15, 215 Cited from Topley and Wilson (1984) : Principle of bacteriology, virology and immunity. 7th Edition. Vol.3 p.142.
- Black, R.E., Myron, M., Mary, L.C., Sandford, B., and Sammel, B.F. (1987) :Prevention of shigellosis by a Salmonella typhi-Shigella sonnei bivalent vaccine. J. of Infections. Diseases. 155 (6) : 1260-65.
- Bokkenheuser, V., Smitty, P., and Richardson, N.(1964) : A challenge to the validity of the Vi test for the detection of chronic typhoid carriers.Am. J. public Hlth.54 : 1501-1503.
- Boyd, J.S.K. (1931) : Identification of a new group of shigella in infections dysentery. J. R. Army. Med. Corp. 57 (151).
- Breadstreet C.M.P. , Agnes, J. and Hilary, E. (1970): Intradermal test and serological tests in suspected cases of brucella in man . Lancet 2 :653.
- Bruce, D. (1887) : Practitioner 39, 161. Cited from Topley and Wilson (1984) : Principles of Bacteriology, Virology and Immunonology. 7th Edition Vol.3. p. 141.
- Buchanan, M. and Faber, L. (1980) : 2-mercapto-ethanol Brucella agglutination test. J. Clin. Microb. 11 : 691.
- Bundle, D.R., Perry, M.P., and Dubrry, G. (1988) : Characterization of Brucella polysaceharide B. Infection and Immunity. 56 (5) : 1101-1106.
- Burrows, J.W.M. and Lewert, R.M. (1963) : Text book of Microbiology 21 : 555.

-
- Butler, T. Speelman, P., Kabir, I and Barmuall, J. (1986) : Colonic dysfunction during shigelosis. *J. infect. Ois. Children.* 154 : 817.
- Christie, A.B. (1987) : *Infections diseases : Epidemiology and clinical practice.* p. 1136.
- Cleary, T.G., Mathewson, J.J., Fairs, E., and Pickering, L.K. (1985) : Shiga-like cytotoxin production by enteropathogenic *E. coli* serogroups. *Infect Immun.* 47 : 335-37.
- Collins, F.M. (1974) : Vaccines and cell mediated immunity. *Bact. Reviews.* 38 : 371-402.
- Corbel, M.J. (1975) : The serological relationship between *Brucella* species, *Yersinia enterocolitica* serogroup IX and salmonella serotypes of kauffmann white group N. *J. Hyg. Camb.* 75 : 151-171.
- Corbel, M.J. and Morgan, W.J. (1976) : *J. Bact.* 11:221.
- Coovadia, Y.M., Singh, V., Bhana, R.H., and Moodely, N. (1986) : Comparson of passive haemagglutination test with Widal agglutination test for serological diagnosis of typhoid fever in an endemic area. *J.Clin. Path.* 39 : 680-83..
- Cruickshank, J.G. (1939) : *J. Hyg.* 39 : 224.
- Cruickshank, Duguid and Collier. (1975) : *Practical Microbiology.* B. 13 th. edition. p. 350.
- Cruickshank, R., Duguid, J.P. and Swain, R.A. (1981): *Medical Microbiology*, 13 th ed. London and Newyork, Livingstone.
- Cupta, A.K., and Reo, K.M. (1981) : Simultaneous detection of *S. typhi* antigen and antibody in serum by counter-immunoelectrophoresis for an early and rapid

- diagnosis. J. Immunol. Meth.30 :349-53.
- Daniels, E., Schnerson, R., Egans, W. Robbins, J.B.(1989) : Characterization of *S. paratyphi* C Vi polysaccharide. Infection and Immunity. 57 : 3159-64.
- DE-Klerk, and Anderson (1985) : Comparative evaluation of ELISA in laboratory diagnosis of Brucellosis. J. Clinic. Micro. 21 : 381.
- Dham, S.K., and Thompson, R.A. (1982) : Humoral and cell mediated immune responses in chronic typhoid carriers. Clinical and Experimental Immunology.50: 34-40.
- Diaze, R., Marau, C.E. and Rivero, A.(1976) : Comparison of counter-immunoelectrophoresis with other serological tests in the diagnosis of human brucellosis.Bulleten of the W.H.O.53 :305.
- Diaze, R., Maravi, C.E. and Rivero, A.(1978) : Rose Bengal plate agglutination and counter immunoelectrophoresis test on spinal fluid in diagnosis of *Brucella* meningitis. J. Clin. Microbiol. 7 : 236.
- Dodds, K.L., Malcolm, B. and McDonald, I.J. (1987) : Electrophoretic and immunochemical study of the liposaccharide produced by chemostat grown *E-Coli* : 0 157. J. of General Microbial. 133 : 2679-87.
- Douglass, J.T. and Palmer, D.A. (1988) : Use of monoclonal antibodies to identify the distribution of A and M epitopes on smooth *Brucella* species. J. of Clinical Microbiol. 26 (7) : 1353-56.
- Duguid, J.P., Anderson, E.S; and Campbell, I. (1966) : Morphological and biological characters of salmonella species J. of pathology and bacteriology. 92 : 107.
- Duguid, J.P. and Gillies, R.R. (1958) : J. Path. Bact. 75 : 519.

- Duguid, J.P., Marmion, B.P., and Swain, R.H.A. (1981) : Medical Microbiology. Vol. I : Microbial infection. 13 th edition by Mackie and McCartney, Churchill livingstone. Inc. Great Britain. Ch. 26.
- Duguid, J.P., Marssion, B.P., and Swain, R.H.A. (1982) : Medical Microbiology Vol. 11. Twelfth edition. p. 420.
- Eberth, C.J. (1880) : Virchow Arch. 81 : 58. Cited from Topley and Wilson (1983) . Principles of Bacteriology & Virology and Immunology. Ch. 37: 332 - 51.
- Edelman, R., and Levine, M.M. (1986) : Summary of an interational workshop on typhoid fever. Review of infect. Dis. 8 (3) : May-June, 329-49.
- Edelman, R. and Levine, M.M (1983) : Summary of an international workshop on typhoid fever J. Infect. Dis. 147 : 1108-18.
- Edwards, P.R. (1951) : Public Health. Report. 66 : 637-39.
- Edwards, P.R. and Fife, M.A. (1961) : Appl. Microbiol. 9 : 478.
- Edwards, P.R. and Ewing, W.H. (1962) : Identification of Enterobacteriaceae. Burgass. publish Co., Minneopolis, Minnesota, U.S.A. Ch.2.
- Edwards, M.B.E., Agens, J. and Pateicia, C.M. (1970) : Comparsion of the indirect flourescence antibody test agglutination, complement fixation test and coombs test for Brucella antibody. J. Clin. Patho. 23 : 161.
- Edwards, P.R. and Ewing, W.H. (1972) : Identification of Enterobacteriaceae. 3 rd edition, Burgen, Minneapdis, Ch.3, 4 and 5.

- Edward, E.A. and Hilderbrand, R.L. (1976) : New method of grouping β hemolytic streptococci by coagglutination. Appl. Microbiol. 28 : 972.
- Ekpo, P., Suttipant, S., Napatwan, B. and Stitayo, S. (1990) : Monoclonal antibodies to 52-Kilodalton protein of salmonella typhi. J. of Clin. Microbiol. 28 (8) : 1818-21.
- El-Behairy, F., Hassanien, Nafie, M., El-Rehaivy, M. and Mandour, A. (1972) : A clinico microbiological study of infantile diarrhea in Upper Egypt. Pediat. Assoc. 20 : 195.
- El-Diwani, M., Gaber, M., Eissa, A., and Amin. Y. (1960) : A clinico bacteriological study of gastroenteritis in Egyptian infants. Egypt. Pediat. Assoc., 8 : 364.
- Elsinghorst, E.A., Baro, L.S. and Kopecko, D.J. (1989) : Penetration of human intestinal epithelial cells by salmonella : molecular cloning and expression of salmonella typhi invasion determinants in E. Coli. Proc. Nalt. Sci.. U.S.A. 86 : 5173-77.
- Engvall P. and Perlmann, P. (1971) : Enzyme-linked immunosorbent assay (ELISA) Quantitative assay of immunoglobulin G. Immun chemistry, 8 : 871-74.
- Evan, A.C. (1918) : Further on Bacterium Abortus and related bacteria. J. Infect. Dis. 22 : 280.
- Evan, A.S. and Feldman, H.A. (1984) : Bacterial infection of human-epidemiology and control. Plenum Medical Book. Co. New York and London. Ch-36 : Typhoid fever by Richard B Hornick.
- Felix, A. and Pitt, M. (1934b) : Virulence of B. typhosus and

- resistance to O antibody. J. Pathol. Bacteriol. 38 : 409-20.
- Felix, A., and Pitt, R.M. (1951) : The pathogenic and immunogenic activities of S. typhi in relation to its antigenic constituents. J. Hyg. 49 : 92-110.
- Felix, M. and Thomas, D. (1982) : Medical Microbiol. p. 297-303.
- Fingold, S.M., and Martin, W.J. (1982) : Diagnostic Microbiology, 6th ed, C.V. Mosby st-lowis Missouri, U.S.A.
- Finlay, B.B. and Falkow, S. (1989) : Salmonella as intracellular parasite. Mol. Microbiol. 3 : 1833-41.
- Finlay, B.B., Heffron, F. and Falkow, S. (1989) : Penetration of salmonella through a polarized Median -Darby canine kidney epithelial cell monolayer. Science, 243 : 940-43.
- Flexner, S. (1900) : Cited from wilson and miles, 1975.
- Francoi, J.V., Bradera, U.A. and Leary, A.B. (1992) : Characterization of shigella serotypes D (S. sonnei) O polysaccharide and enterbacterial R. Lipopolysaccharide core by use of mouse monoclonal antibodies. Infect. and Immun. 60 (7) : 2741-47.
- Gaffkey, (1884) : Cited from Toply and wilson, 1975.
- Galan, J.E. and Curtiss III, R. (1989) : Cloning and molecular characterization of genes whose products allow salmonella typhimurium to penertrate tissue culture cell. Proc. Natl. Acad. Sc. U.S.A. 86 : 6383-87.
- Galian, J.E., and Curtiss, III. R. (1991) : Disterbution of the Inv A, - B, - C and - D genes of salmonella typhimurrium among other salmonella serovars : Inv A mutants of

salmonella typhi are deficient for entry into Mammalian cells. Infection and Immunity. 59 (9) : 2001-8.

Georgala, D.L, and Boothroyd, M. (1965) : Isolation of S. typhi from different specimens Can. J. Res. 22 : 48.

George, A. and Kamat, R.S. (1987) : Regulation of cell-mediated immunity in mice immunized with *S. enteritidis*. J. Med. Microbiol. 23 : 239-46.

George, A., Rath, S. and Kamat, R.S. (1986) : Differential regulation of effector responses of cell mediated immunity in experimental salmonellosis. Clinical and Exper. Immunology. 63 : 327-33.

Gianella, R.A., Broitman, S.A., and Zamecheck, N. (1973) : Pathogenesis of acute bacterial diarrhoeal disorders. J. Infect. Dis. 128 : 69-75.

Gianella, R.A., Broitman, S.A. and Zamecheck, N. (1970) : Salmonellosis; Relevance of reduced gastric acid secretion to severity of diarrhea and effect on small bowel. Gastroenterology. 60 : 666.

Glenn, J. and Thomas, E. (1981) : Manual of allergy and immunology diagnosis. and therapy of Brucella. First Edition. p. 378.

Gilman, R.H., Terminel, M., Levine, M.M., Hernandez, M., and Hornick, R.B. (1975) : Relative efficacy of blood, urine, rectal swab, bone-marrow, and rose-sport cultures for recovery of S. typhi in typhoid fever. Lancet 1 : 1211-1213.

Graigie, J. and Brandon, K.F. (1936) : The identification of the V and W forms of *B. typhosus*. and the occurrence of V form in

- cases of typhoid fever and in carriers. J. Pathol. Bacterial. 43 : 249-60.
- Greisman, S.E., Hornick, R.B., and Woodward, t.E. (1964) Role of endotoxin in pathogenesis of typhoid fever . J. Clin. Invest. 43 : 1747.
- Guerra-Caceras, J.G., Gotuzzo, H.E., Grosby, D.E., and Carillio, P. (1979) : Diagnostic value of bone marrow culture in typhiod fever. Trans. Roy. Soc. Trop. Med. and Hyg. 73 (6) : 680.
- Guo, Y.N., Hsu, H.S., Mumaw, V.R., and Nakoneczna, I. (1986a) : Electron microscopy studyies on the bactericidal action of inflummatory leucocytes in murine Salmonellosis. J. of Medical. Microbiol. 21 : 151-59.
- Hardy, A.V., Mackel, D.C., and Hamerick, D. (1953) : Cultural methods in diagnosis of salmonella .Armed. Forces. Med. J. 4 : 393.
- Hartman, A.B., Maria, M.R., and Clydel, S. (1991) : Moleuclar analysis of variant plasmid forms of a bivalent salmonella typhi-shigella sonnei vaccine strain. J. of Clin. Microbiol. 29 (1) : 27-32.
- Hefferman, E., Ferreccio, C. and Robbins, J.B. (1987) :J. Infect. Dis. 155 : 1254 - 59.
- Hernandez-Velarde, R., Diazm G. and Munoz, H. (1980) : Use of the ELISA technique for diagnosis of typhoid fever. I. Standarization of technique. Arch. Invest. Med (Mex). 11 : 137-267.
- Herrington, D.A., Van-de-Verg, L., Formal, S.B., and Leuine, M.M. (1990) : Studies in volunteers to evaluate candidate Shigella vaccines. Vaccine. 8 (4) : 353-7.

- Hirschl, A., Stanck, G., Rotter, Chary, M.L., and Niemetz, A.H. (1985) : Antibody response to somatic antigen of S. typhi in areas endemic and non endemic for S. typhi in areas endemic and non endemic for typhoid fever. Infect. Control. 6 (3) : 110.
- Hoeprich, P.O. (1983) : Infectious Diseases. A modern treatment of infections processes. 3rd edition by Haper and Row, Pub, philadelphia. Ch. 64. by Richard B. Hornick.
- Hoffman, R.E., and Shillam, P.J. (1990) : The use of hygiene and antimicrobial therapy to control out break of shigellosis. Am. J. Dis. Child. 144 (2) : 219.
- Hoffman, S.L., Pujabi, N.H. Rockhill, R.C., and Puluagish, P.C. (1984) : Doudenol string capsule culture compared with bone marrow, blood, rectal swab cultures for diagnosis of typhoid fever. J. Infect. Dis. 149 (2) : 157.
- Hornick, R.B. (1981) : Typhoid fever and salmonella infection. In : Tropical and Geographic Medicine. p. 718.
- Hornick, R.B., Greisman, S.E., Woodward, T.E. and Dupont, H.L. (1970) : Typhoid fever : pathogenesis and immunologic control. N. Engl. J. Med. 283 : 686-91.
- Huckstep, R.L. (1962) : Typhoid fever and salmonella infection. Text book. E & S. Living stone, Ltd, ch. 394.
- Hunter, A. and Colbert, B. (1956) : Flocculation tests for Brucella. J. Immunity. 77 : 232-41.
- Hurvell, B. Ahvonen, P. and Thal, E. (1971) : Serological cross relations between different Brucella species and Yersinia

- enterocolitica. Act. Vet. Scand. 12 : 86-94.
- Hynes, M., (1942) : The isolation of intestinal pathology by selective media . J. Path Bact. 54: 193.
- Ibrahim, I. Brande. (1983) : Basic science and Medical applied. 374-76.
- Jawetz, E., Melnick, J.L., and Adelberg, E.A. (1980) : Review of Medical Microbiology. 14th. ed. Lange. Med. Publ. California, U.S.A.
- Jelliffe, D.B., and Stourfield, J.B. (1978) : Diseases of children in the subtropics and Tropics, 3rd, edition, Edward Arnold, Ltd, London.
- Jesudason, M., Sridharam, G. and John, T.J. (1991) : Antigenic sharing of salmonella typhi, non flagellar proteins with other salmonella and other shigella and E. Coli. Indian. J. Exp. Biol. 29 (4) : 338-41.
- Jimenez, V.L. and John, F. (1990) : Heterogeneity of Lipopolysaccharide phenotype among *S. typhi* strains. J. of Infectious Dis. 162 : 763-64.
- Jimenez, V.L., Adelberg, E.A., and Melnick, J.L. (1987) : C₃b generation is affected by the structure of the O-antigen polysaccharide from salmonella. J. Immunol. 139 : 1253-60.
- John, T.J., Sivadasan K., and Kurien, B. (1984) : Evaluation of passive bacterial agglutination for the diagnosis of typhoid fever. J. Clin. Microbiol. 20 : 751-53.
- John, R.L., and Roy, T.S. (1986) : Role of the Vi antigen of *S. typhi* in resistance to host defense in vitro. J. Lab. Clin. Med. 108 : 506-516.

- John , E.S., Ayad. E. and Zohair, F. (1982) : Diagnosis of human brucellosis with ELISA. *Lancet*. July, 3 :19.
- Johnston, W.M. and Lior, H.(1986) : Cytotoxic and cytotoxic factor produced by campylobacter jejuni, campylobacter coli and campylobacter laridis .*J. Clin Microbiol.* 24: 275 - 81.
- Johnston, E.M., Knanskop, F., and Baran, L.S. (1965) : Genetic mapping of Vi and somatic antigenic determinants in salmonellosis. *J. Bact.* 90 : 302-308.
- Jones, M.A., Bruins, S.C. and Kurien, B. (1980) : Evaluation of radial immunodiffusion test with polysaccharide B antigen for diagnosis of bovine brucellosis. *J. Clin. Microbiol.* 12 : 753-60.
- Joklik, W.K., Willett, H.P. and Amos, D.B. (1992): Cited from Zinnser Microbiology. 20th edition Ch. 35. P. 559-61.
- Jongh, L., McCabe, W.R. and Sivadasan, K. (1985) : Detection of antibodies against lipopolysaccharide of E-Coli and salmonella R. and S strains by immunoblotting. *Infect. and Immun.* 50 (3) : 716-20.
- Kauffmann, F. (1935) : *Z. F. Hyg.* 117-62.
- Kauffmann, F. (1936) : *Z. F. Bact.* , 116-17.
- Kauffmann, F. (1937) : Cited from Edwards, P.R. and Ewing, W.H. (1969). *Identification of Enterobacteriaceae*.
- Kauffmann, F. and Petersen, A. (1955) : *Acta. path. Microbiol. Scand.* 36 : 173. Cited from Topley and Wilson (1983) .*Principles of Microbiology & Virology and Immunology*. Ch. 37:323 - 51.
- Kauffmann, P.R. (1951) : *Public Health, Report.* 66 : 637-39.

- Kay, D., Palmeri, M., Eykman, L. Rocha, H. and Hook, E.W. (1966) : Comparison of bile and trypticase soy broth for isolation of salmonella from blood. *Amer. J. Clin. Path.* 36 (8) : 408-410.
- Kerr, W.R., Joyce, D., Poyne, D. and Robertson, L. (1966) : The laboratory diagnosis of chronic brucellosis *lancet*, 20 : 1181.
- Ketyi, I., Pacsa, L. and Kuch . B. (1979) : Shigella dysenteriae 1-like cytotoxic enterotoxins produced by salmonella strains . *Act. Microbiol. Acad . Sci.* 26: 217 - 23.
- Keusch, G.T. and Jacewicz, M. (1977) : The pathogenesis of shigella diarrhea. and relationship of shiga toxin, neurotoxin and cytotoxin. *J. Infect. Dis.* 131 : 533.
- Keusch, G.T. (1986) : Infectious Diarrhea. Shigella. In Gorbach, S.L. (editor.). Boston, Blucknell Scientific Publication, 31.
- Koster, F.T., Boonpucknavig, V. and Sujaho, S. (1984) : Renal histopathology in the hemolytic uremic syndrom. following shigellosis. *Clin. Nephrology.* 21 : 126.
- Laemmli, U.K. (1970) : Cleavage of structural protein during assembly of the head of bacteriophage T4. *Nature*, 227 : 680-685.
- Leading, A. (1978) : Typhoid and its serology. *Brit. Med. J.* 1 : 389.
- Lee, D.R., Schamitman, C.A. and Pugsley, A.P. (1979) : Major proteins of outer protein antigen of salmonella *J. Bact.* 138 : 861-70.

-
- Lee, C.A. and Falkow, S. (1990) : The ability of salmonella to enter mamalian cells is affected by bacterial growth state. Proc. Natl. Acad. U.S.A. 87, 1304-8.
- Leifson, E. (1935) : Cultural characters of shigella and its biochemical activites. J. Path. Bact. 40 : 581.
- Leive, L.L., and Jimenez, L.V. (1986) : Lipopolysaccharide O-antigen structure controls alternative pathway activation of complement effect on phagocytosis and virulence of salmonella. Amer. Society for Microbiol. 14-18.
- Le Minor, L. and Ben Hamida, F. (1962) : Studies of biochemical activies of some Enterobacteriaceae Ann. Inst. Pasteur. 102 : 267.
- Leong, D., Diaz, R., Rudabach, J. and Wilson, J.B. (1970) : Some structural and biological properties of brucella endotoxin. Infection and Immunity. 1 (2) : 174-82.
- Levine, M.M., Gardos, O., Gilman, R.H., Woodward, W.,E., and Waldman, W. (1978) : Diagnostic value of the widal test in areas endemic for typhoid fever. Am. J. Trop. Med. Hyg. 27 (4) : 795-800.
- Levine, M.M. and Edelman, R. (1990) : Future vaccines against enteric pathogens. Infect. Dis. Clin. North. Am., 4 (1) : 105-121.
- Libasi, R., Armando, F. and Theseil, B. (1988) : Protection against salmonella typhi infection in mice after immunization with outer membrane protein isolated from salmonella typhi, 9, 12, d, Vi. Infect. Immun. 56 : 2953-59.
- Lindberg, A.A. and Holme, T. (1972) : Evaluation of some extraction methods for the preparation of bacterial lipopolysaccharide for structural analysis. Acta. Path.

Microbiol. Scand. Section B. 80 : 751-59.

Lindberg, A.A. (1980) : Bacterial virulence factors with particular reference to salmonella bacteria. Scand. J. Infect. Dis. Suppl. 24 : 86-99.

Lowery, B.O., Nira, J.R., Levis, F. and Rose, J. (1951) : Protein measurement with the folin phenol reagent. J. Biol. Chem. 193 : 265-75.

Machei, A and McCantney, C. (1989) : Practical Medical Microbiology. B. 13 th edition. p. 529.

Madanagoplan, N., Supramaniam, S. and Sunitis C. (1975) : J. Trop. Med and Hyg. 2 : 27.

Mandle, A.D., Baron, L.S. and Backler, C.E. (1959) : Role of Vi in Salmonella paratyphi C infections. Proc. Soc. Exp. Biol. Med. 100 : 653-56.

Martini, E. and Lenz., O. (1902) : Cited from Wilson and Miles, 1975 5th Edition.

Martin, W.Z., Mock, W.E. and Ewing, W.H. (1968) : Antigenic analysis of shigella sonnei by gel diffusion. J. Microbiol. 14 : 737.

Mattsby, B.I., Genski, P. and Alving, C.R. (1984) : Heterogeneity of lipid A : comparison of lipid A types from different gram negative bacteria. J. Bact. 159 : 900-904.

Mckendrick, G.D.W. (1973) : Prices Textbook of the practice of Medicine Ed. Scott. R.B. Oxford University. Press, London, p. 51.

Merril, R.M., Grabus, J. and Hornick R.B. (1981) : Science. 211 : 1437-38.

- Miller, R.M., Grabus, J. and Hornick, R.B. (1972) : Lack of enhanced oxygen consumption by polymorphonuclear leucocytes on phagocytosis of virulent *S. typhi*. *Science*. 175 : 1011-1014.
- Mittal, K.R. and Tizard, I. (1981) : Serological cross-reaction between *brucella abortus* and *yersinia enterocolitica* serotype O9. *The Veterinary Bull.* Vol. 51 (7) : 501 -507.
- Mortimer, P.S., John, L.I. and Albert, B. (1980) : *Annual Rev. of Microbiol.* 34 : 370.
- Munfprd, R.S., Wear, R.E., Patton, C. and Feldman, R.A. (1979) : Human disease caused by *brucella ovis*. *J. Amer. Med. Ass.* 231 : 1267-69.
- Murray, and Thompson, K. (1990) : *Medical Microbiol . The C.V. Mosby Company . U.S.A.* P.106.
- Nadler, H., Dolan, C., Forgaces, P., and George, H. (1982) : *Brucella suis*; an unusual casuse of suppurative lymphadenitis in an outpatient. *J. Clin. Microbiol.* (3) : 575-76.
- Nakae, T. (1976) : Properties and function of pore proteins of genus *salmonella* .*J. Biol. Chem.* 251 : 2176-78.
- Nardillo, S., Pizzella, T., Russo, M. and Galanti, B. (1983) : ELISA determination of IgM anti LPS in the early phase of typhoid fever. *Boll. 1st. Sieroter. milan* 62 (4) : 372.
- Nardiello, S., Pizzella, T. Russo, M. and Galanti, B. (1984) : Serodiagnosis of Typhoid fever by ELISA determination of annti. *S. typhi* Lipopolysaccharide antibodies. *J. Clin. Microb.* 20 (4) : 718-21.
- Nartikova, V.F., Egorova, T.P. Belkin, P.Z., Fedosova, V.G., and Levenson, V.I. (1991) : *Zh-Microbiol-Epidemiol. Immunobiol.* 4 : 14-17.

- Neill, R.J., Gensk, P., Formal, S.B., and Newland, J.W. (1988) : Deletion of shiga toxin gene in a cholerae-resistant derivative of shigella dysenteriae type I that retains virulence. J. Infect. Dis. 158 : 737.
- Neter, E., Westphal, O., Luderitz, O. and Gerznski, A.E. (1962) : The bacterial haemagglutination test for demonstration of antibodies to Enterobacteria. Ann. N.Y. Acad. Sci. 66 : 141.
- Nolan, C.M., Feely, J.C., White, J.P. and Wong, K.H. (1980) : Evaluation of a new assay for Vi antibody in chronic carriers of S. typhi J. Clin. Microbiol. 12 : 22-26.
- O'Brein, A.D., Laveck E.D. and Formal, S.B. (1982): Production of shigella dysenteriae type 1 like cytotoxin by E. coli. J. Infect. Dis. 146: 763 - 69.
- O'Brein, A.D., and Holmes, R.K. (1987) : Shiga and shiga-like toxins. Microbiol. Rev. 51 : 206-220.
- Ortiz, V., Isibasi, A. and Kumate, J. (1989) : Immunoblot detection of class-specific humoral immune response to outer membrane proteins isolated from salmonella typhi in humans with typhoid fever. J. of Clin. Microbiol. 27 (7) : 1640-45.
- Osborn, M.Z. and Wutt, H.C. (1980) : proteins of the outer memberane of gram negative bacteria. Ann. Rev. Microbial. 34 : 364-422.
- Parrett, D., Nidosonk. H. and White, R.G. (1977) : Radioimmunoassay of Ig M, Ig E and Ig A brucella antibodies. lancet, 21 : 1075-77.
- Perry, M.B., Bundle, D.R., Perry, J.A. and Griffith, D.W. (1986) : The

- structure of the antigenic lipopolysaccharide O-chain produced by salmonella Urbana and salmonella godesberg. Carbohydr-Res. 156 : 107-22.
- Perry, M.B., Maclean, L. and Griffith, D.W. (1986) : Structure of the O-chain polysacchride of the phenol-phase soluble lipopolysaccharide of E. Coli O : 157 Biochem. Cell-Biol. 64 (1) : 21-28.
- Persson, M.A., Erik, E., Lennart, H. and Edward, S.C.I. (1986) : Immunoglobulin G (Ig G) and Ig A subclass pattern of human antibodies to shigella flexneri and salmonella serogroup B and D lipopolysaccharide O antigens. Infection and Immunity, 52 (3) : 834-39.
- Pordo, D.T, Cleary, L.K. and Johnson, P.C. (1986) : The relation between production of cytotoxin and clinical features in shigellosis. J. Infect. Dis. 154 : 149-155.
- Qadri, A., Gosh, S. Prsash, K., Kumar, r. and Moudgil, K.D. (1990) : Sandwich enzyme imunoassays for detection of salmonella typhi. J. Immunoassay, 11 (2) : 251-69.
- Qadri, A., Satish, K., Gupta, K. and Gursaran, T. (1988) : Monoclonal antibodies delinate multiple epitopes on the O antigens of S. typhi lipopolysaccharide. J. of Clin. Microbiol. 26 (11) : 2292-96.
- Rajagopalan, P., Kumar, R. and Malaviya, A. (1981) : Immunological studies in typhoid fever : Immunoglobulins, C3, antibodies, rheumatoid factor and circulating immune complexes in patients with typhoid fever. Clin. Exp. Immunol. 47 : 68-73.
- Rajagopalan, P., Kumar, R. and Malaviya, A. (1982) : Immunological studies in typhoid fever cell mediated immune response and lymphocyte subpopulation in patients with typhoid

- fever. Clin. Exp. Immunol. 47 : 269-74.
- Reddin, J.L., Anderson, R.K., Jewnness, R. and Spink, W.W. (1965) : Significance of 7S and macroglobulin agglutining in human brucellosis. N. Eng. Med. 272 : 1263.
- Reitmeyer, J.C., Peterson, J.W., and Wilson, K.J. (1986) : Salmonella cytotoxin : a component of bacterial outer membrane. Microbiol. Patholog. 1 : 503-510.
- Rennels, M.B. and Levine, M.M. (1986) : Studies on diarrheal diseases in children. Pediat. Infect. Dis. 5 : 519-25.
- Report, (1958) : Int. Bull. Bact. Nomencl. 8:25.
- Ribi, E., Haskins, W.T., Landy, M. and Milner, K.C. (1961) : Immunological response and pathogenesis of salmonella Bact. Rev. 25 : 427-36.
- Robinson, D.T. (1967) : Diagnostic value of widal test in typhoid fever .New. Engl. J. Med. 277 : 171.
- Robbins, J.D. and Robbins, J.B. (1984) : Reexamination of the protective role of the capsular polysaccharide (Vi antigen). J. Infectious. Dis. 150 : 436-449.
- Robert, H., Ronica, M. and Reizn, J.I. (1987) : Prevention of shigellosis by a salmonella typhi-shigella sonnei bivalent vaccine. J. of Infectious. Dis. 155 (6) : 1260-65.
- Rockhill, R.C., Rurnams, L.W., Lesmana, M. and Dennis, D.T. (1980) : Detection of S. typhi, D, Vi and d antigens, by a slide coagglutination in urine from patients with Typhoid fever. J. Clin. Microbiol. 11 : 213-16.
- Rodhe, R. (1967) : J. Appl. Bact. 205 : 404.
- Rowe, J., Gianella, R.A. and Formal, S.B. (1976) : Antigenic structure

of shigella species J. Syst. Bact. 26. : 76.

- Rubin, F.A., Tcapecko, D.J., Noon, R.F. and Baronl, L.S. (1985) : Development of a DNA probe to detect salmonella typhi. J. Clin. Micriobiol. 22 : 600-605.
- Rubin, F.A., Mcwhirter, P.D., Burr, D., and Lane, E. (1990) : Rapid diagnosis of Typhoid fever through. identification of S. typhi within 18 hours of specimen aquisition by culture of mononuclear cell platelet fraction of blood. J. of Clin. Microbiol. 28 (4) : 825-27.
- Ryan, W.J. (1972) :Comparsion between different methods for diagnosis of typhoid fever .J. Med. Microbiol. 5 : 533.
- Salvatore, N., Tereasa, P., Michele, R. and Bruno, G. (1984) : Serodiagnosis of typhoid fever by ELISA determination of antisalmonella typhi. lipopolysaccharide antibodies. J. of Clinic. Microbiol. 20 : 718-21.
- Sandiford, J.P., Luby, J.P. and Dietshy, J.M. (1982) : The science and Practice of Clinical Medicine Vol. 8 Infectious Disease.
- Sarasombath, S., Banchin, N., Sukosol, T., and Manasatit, S. (1987) : J. of Clinc. Microbiol. 25 (6) : 1088-93.
- Sarasombath, S. Lertmemongkolchai, G., and Banchuin, N. (1988) : Characterization of monoclonal antibodies to protein antigen of S. typhi. J. Clin. Microbiol. 26 (3) : 508-12.
- Sanpectchsong, V. and Tharavanij, S. (1980) : Immunological study of typhoid fever in man. Southeast Asian. J. Trop. Med. Pub. HLth, 11 (4) : 510-515.
- Sarma, V.N.B., Malaviya, A.N., Kumar, R. and Bakhtary, M.K. (1977) : Development of immune response during

- typhoid fever in man. Clin. Exp. Immunol. 28 : 35-93.
- Sato, K. Inaba, Y. and Mtumoto, M. (1981): Isolation of human rota virus in cell culture . Arch. Virol. 69: 155 - 160.
- Schmidt, G.I., Fromme, H, and Mayer, H. (1970) : Immunochemical studies on core lipopolysaccharide of Enterobacteriaceae of different genera. Eur. J. Biochem. 14 : 357-66.
- Schroeder, S.A. (1968) : Interpretation of serologic test for typhoid fever. J. Am. Med. Assco. 206 : 839-40.
- Schubert, H.H., Edward, P.R. and Ramsy, C.H. (1959) : Changes of serum complement in experimental typhoid fever .J. Bact. 77 : 648.
- Schultz, C. (1989) : Isoenzyme analysis of Typhoid-Shigella and Escherichia-shigella hybrid vaccines and their parental strains. J. Clin. Microbiol. 27 (12): 2838-41.
- Seid, R.C., Denni, J.K., Jerald, C., Louis, S. and Samuel, F. (1984) : Unusual Lipopolysaccharide antigens of a salmonella typhi oral vaccine strain expressing the shigella sonnei form I antigen. J. of Biological Chem. 259 (14) : 9028-34.
- Sharma, U.K. Jayasheela, M. and Sarxena, S.N. (1987) : Enterotoxigenicity and antibiotic resistance in salmonella strains isolated from patients of gastroenteritis. Indian. J. Med. Res. 85 : 608-11.
- Shehabi, A.A. (1981) : The value of a single widal test in the diagnosis of acute typhoid fever. Trop. Geogr. Med. 33 (1) : 113-116.
- Shiga, K. (1898) : Cited from Wilson and Miles, 1975.
- Smith, T.H, and Raegh, A. (1903) : J. Med. Res. 10 : 89 Cited from

Topley and Wilson, 1983. 7th edition.

Sonne, C. (1915) : Cited from Wilson and Miles, (1975).

Spaun, J. (1952) : Acta. Path. Microbiol. Scand. 31 : 462.

Spink, W. (1956) : The nature of Brucellosis. p. 207

Sprintz, H., Gangarosa, E.J., Willams, M and Woodward, T.E.(1966) :
Histopathology of the upper small intestines in typhoid
fever. Am. J. Dig. Dis. New Series. 11 : 615.

Stableforth, A.W. (1954) : Bull. World Health. Organization-cited
from Hunter and Colbert (1956). J. Immunology. 77 :
232-41, 1956.

Stokes, J.L. and Bayne, H.G. (1957) :Identification and cultural
characters of salmonellae. J. Bact. 74 : 200.

Susan, J., George, C., Frederick, T., and Wallis, L. (1981) :
Serological diagnosis of brucellosis. J. Clin. Microbiol.
13 : 398.

Susan, B., Willam, F., Chum, N., Arnold, F. and Roger, M. (1986) :
ELISA with major outer membrane protein of *Brucella*
melitensis to measure immune response to *Brucella*
species. J. Clin. Microbiol. 24 : 566.

Takeuchi, A. and Sprinz, H. (1967) : Pathogenesis of *Salmonella typhi*
in typhoid fever. Amer. J. Path. 51 : 137-161.

Tansg, R.W., Chan, K.H., Chan, P.Y., Wan, K.C. and Schlect, S.
(1987) : A murine monoclonal antibody specific for the
outer core oligosaccharide of salmonella
lipopolysaccharide. Infection and Immunity. 55 (1) :
211-16.

- Tantivanich, S., Chongsanguan, M., and Tharawanij, S. (1984) : A simple and rapid diagnostic test for Typhoid fever. Southeast, Asian, J. Trop. Med. Pub., 15 (3) :317-22.
- Taylor, W.L. and Schelhart, D. (1968) : Isolation of shigella, comparsion of enrichment broths with stools. Applied Microbiol. 16 : 1383.
- Taylor, D.N., Harris, J.R., Barrett, J.J., and Blacke, P.A. (1983) : Detection of urinary Vi antigen as a diagnosis test for typhoid fever. J. Clin. Microbiol. 18 (4) : 872-78.
- Thevanesan, V., Arseculerante, S. N. and Athauda, P.K. (1982) : Cell mediated and humoral immune responses in human typhoid fever. Trop. Geogr. Med. 34 : 13-17.
- Thomas, J.C., Watson, K.C., and Hevston, A.S. (1954) : Use of streptokinase bile salt broth for clot cultures in diagnosis of Typhoid fever. J. Clin. Path. 7 : 50.
- Thomason, B.M., Worter, M.A. and Sanders, E. (1964) : Bact. Proc. p. 56.
- Thomason, B.M., Cowart, G.S., and Cherry, W.B. (1965) : Current status of immunofluoresence techniques for rapid detection of shigella in faecal specimens. Applied. Microbiol. 13 : 605.
- Timothy, J. and Barret, T. (1985) : Improvment of the indirect haemagglutination assay for Salmonella typhi Vi antibodies by use of glutaraldehyde-fixed erythrocytes. J. of Clin. Microbiol. 22 (4) : 662-63.
- Tiwari, H. and Kamat, R.S. (1986) : Cross reactions in cell mediated immunity of enteric fever. J. of Medical Microbia. 21 : 233-37.

-
- Topley, W.W. and Wilsons, G.S. (1975) : Principles of Bacteriology, Virology and Immunology. 6th Editions London, Arnold, Ch. 30, 68.
- Topley and Wilsons (1983) : Principles of Bacteriology, Virology and Immunology Vol. 2 systematic bact. Ch. 40 : 406-21.
- Topley and Wilsons (1983) : Principles of Bacteriology, Virology and Immunology Vol. 2 systematic bact. Ch. 37 : 338-40.
- Topley and Wilsons (1985) : Principles of Bacteriology, Virology and Immunology 7th edition. Great Britian. Ch.68.
- Topley, W.W. and Ayrton, J. (1924) : J. Hyg. 23 : 198.
- Towbin, H., Theophil, S. and Julian, G. (1979) : Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets. procedure and some applications. Proc. Natl. Acad. Sci. u.S.A., 26 (9) : 4350-54.
- Traum, J.E. (1914) : Rep. chief. Bur. Animal. Industry. P. 30. Cited from Topley and Wilson (1984) : Principle, of Bacteriology, Virology and Immunology. 7th Vol.3 : p.142.
- Tsai, C.M. and Carl. F.E. (1982) : A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. Anal. biochem. 119 : 115-119.
- Tsang, R.S.W. and Chau, P.Y. (1981) : Serological diagnosis of Typhoid fever by counter immunoelectrophoresis. Br. Med. J. 282-1501.
- Tsang, R.S.W., Chau, P.Y., La Boorg, J.T. and Rowley, D. (1981) : Antibody resonse to the lipopolysaccharide and protein antigens of S. typhi during typhoid infection. Clin. Exp. Immunol. 46 : 508-514.

- Turk, D.C., Porter, I.A., Duerden, B.I., and Reid, T.M. (1983) : A short text Book of Medical Microbiology 5th Edition by Hodder and stoughton, London, Toronto, Ch.7.
- Van-de-verg, L., Herrington, D.A., Murphy, J.R. and Levire, M.M. (1990) : Specific immunoglobulin A secreting cells in peripheral blood of humans following oial immunization. with a bivalent salmonella typhi-shigella sonnei vaccine of infection by pathogenic shigella sonnei. Infect. Immun. 58 (6) : 2002-4.
- Versilova, P.A., Gerryseva, M.J., Aslanja, R.G. and Knjazena , E.N. (1974) : Diagnosis of human and animal Brucella by indirect haemagglutination test. Bull. W.H.O. 51 : 191-197.
- Watson, K.C. (1955) : Isolation of S. typhi from blood stream. J. Lab. Clin. Med. 46 : 128-134.
- Weber, K. (1968): Nature, 218: 1116 .
- Weber, K. and Osborn, M. (1969) : The reliability of molecular weight determination by dodecyl sulfate- polyacrylamide gel electrophoresis . J.Biol . Chem 244 : 4406 - 12.
- Webster, M.E.L. and Freeman, M.E. (1952) : Studies on Vi antigen II. Purification of Vi antigen from E. coli : 5396/38. J. Immunol. 69 : 135-142.
- Weil, E. and Folix, A. (1920) : Laboratory diagnosis and epidemiology of salmonella and shigella. Charles. C. Thomas, springfield, III, p. 1-24.
- Wheat, r.w. (1964) : In Bacterial endotoxin 76-80.
- Wheeler, K.M. and Mickle, F.L. (1945) : Antigens of shigella sonnei. J. Immunol. 51 : 257.
- White, P.B. (1926) : Med. Res. Council, Gr. Brit. Spec. Rep. Ser. No. :

103.

- Whiteside, R.E. and Baker, E.E. (1959) : The Vi antigens of enterobacteriaceae. II imunologic and biologic properties. J. Immunol. 83 : 687-96.
- Wilke, G.I. and Moenning, V. (1987) : Monoclonal antibodies nad characterization of epitopes of smooth brucella lipopolysaccharide Anne. Inst. Pastenr. Microbiol. 138 : 549-60.
- Wilson, G.S., and Miles, A.A. (1964) : Enteric infection : Cited from Topley and Wilson. Principles of bacteriology , virology and immunology. 5th. Ed. Vol.2, p. 1863-75.
- Wong, K.H. and John, C.F. (1972) : Isolation of Vi antigen and simple method for its measurment. Applied. Microbial. 24 (4) : 628-33.
- World health organisation (1964) : Expert commites on Brucella. Cited from Spink, 1956. The nature of Brucella. p. 207.
- Zagmunt, M.S., Dubray, G., Bundle, D.R. and Perry, M.P. (1988) : Purified native haptens of Brucella abortus B14 nd Brucella melitensis 16 M reveal the lipopolysaccharide origin of the antigens. Ann. Inst. Pasteur. Microbiol. 139 : 421-33.
- Zammit, T. (1905) : Rep. Comm. Medit. Fev. part. III. P. 83. Cited from Topley and Wilson (1984) . Principles of Bacteriology, Virology and Immunology. 7th. Edition. Vol.3 : p. 141.
- Zuulunov. A., Lerman, M., Ash Kenazi, S. and Dinari, G. (1990) : The prognosis of convulsions during childhood shigellosis. Eur. j. Pedia. 149 (1) : 293.