

References

- Alicia, J.M.; Leach, C.S.; Teresa, C.H. and William, R.J. (1999):**
CDC guideline for prevention of surgical site infection.
*Am.J.Infect.Control.*27 (2):97-132.

- Alrawi, S.; Houshan, L, L.; Satheesan, R.; Raju, R.; Cunningham, J.
and Acinapura, A. (2001):**
Glove reinforcement: an alternative to double gloving.
Infect.Control.Hosp.Epidemiol 22(8): 526-527.

- Ako, N.A.I.; Adejuyigbe, O.; Adewami, T.O. and Lawal, O.O. (1992):**
Sources of intraoperative bacterial colonization of clean surgical
Wounds and subsequent post-operative wound infection in
Nigerian Hospitals.
*East.Afr.Med.J.*69 (9): 500-507.

- Amira, M.A. (1996):**
Study of post-operative wound infection incidence and possible
Etiological factors in great Cairo. M.Sc.Thesis.Microbiology.
Zagazig University.

- Baird, D. (1996):**
Staphylococcus: cluster forming Gram positive cocci.
In: Mackie&McCartney, *Practical Medical Microbiology*. Collee, J.
G.; Marmion, B.P.; Fraser, A.G. and Simmons, A. (eds).fourteenth
edition, Churchill, Livingstone.P.245-261.

- Bakr, A. A. (1988):**
Bacteriological study of post-operative wound sepsis.
M.Sc. Thesis. Microbiology. Benha University.

- Bennett, E.; Welsby, I.; Dunn, T.J.; Young, L. R.; Wahl, T.A.;
Diers, T. L.; Phillips-Bute. B.G.; Newman, M. F. and
Mythen, M. G. (1999):
The use of postoperative morbidity survey to evaluate
Patients with prolonged hospitalization after routine,
Moderate-risk, elective surgery.
Aneth. Analg. 89:514.

- Bisno, A.L. and Stevens, D.L. (1996):
Streptococcal infection of skin and soft tissues.
N. Engl. J. Med. 334:240.

- Blumberg, N. (1997):
Allogenic transfusion and infection: Economic and
Clinical implication.
Seminars in Hematology.34, Suppl2: 34-40.

- Brooks, G. F.; Butel, J. S. and Morse, S. A. (1998):
The *Staphylococci* in: Jawetz, Melnick & Adelberg's
Medical Microbiology. 21st edition, Appleton &
Lange P. 197-202.

- Brooks, G. F.; Janet, S. and Stephens, A. (1998):
Legionellae in: Jawetz, Melnick & Adelberg's
Medical Microbiology. 21st edition, Appleton
And Lang: 273-277.

- Brown, R.; Collee, J. G. and Pexton, R. L. (1996):
Bacteroid, *Fusobacterium* and other Gram negative
Anaerobic rods: anaerobic cocci: identification of anaerobes.
In: Mackie & McCartney, Practical Medical Microbiology,
Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A.
(eds). 14th edition, Churchill Livingstone P.501-511.

- Bryce, E. A.; Spence, D. and Roberts, F. G. (2001):
An in-use evaluation of an alcohol-based pre-surgical
hand disinfection.
Infect. Control. Hosp. Epidemiol. 22(10): 613-617.

- Cheesbrough, M. (1994):
Medical Laboratory Manual for tropical countries, Vol.2:
Microbiology (published) by Tropical Health Technology,
Butterworth-Heinemann, Great Britain. Ch34&36 P.36&69.

- Cheung, A. H. and Wong, L. M. (2001):
Surgical infections in patients with chronic renal failure.
Infect. Dis. Clin. North. Am. 15(3):775-796.

- Circhton, P. B.; Logan, J. M. J. and Old, D. C. (1993):
A miniaturized biotyping system for strain discrimination
In *E. coli*.
Epidemiol. Infect. 111: 81-88.

- Circhton, P. B. (1996):
Enterobacteriaceae: Escherichia, Klebsiella, Proteus and
other genera. In: Mackie & McCartney, Practical Medical
Microbiology, Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and
Simmons, A. (eds). 14th edition, Churchill Livingstone
P.361 -384.

- Classen, D. C.; Evans, R. S.; Pestotnik, S. L.; Horn, S. D.; Menlove, R.
L. and Burke, J. P. (1992):
The timing of prophylactic administration of antibiotics and the
risk of surgical wound infection.
N. Engl. J. Med. 326: 281-286.

- Collee, J. G.; Duguid, J. P.; Fraser, A. G.; Marmion, B. P. and Simmons, A. (1996):
Laboratory strategy in the diagnosis of infective syndromes.
In: Mackie & McCartney, Practical Medical Microbiology,
Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A
(eds).14th edition, Churchill Livingstone P.53-94.

- Collee, J. G. and Marr, W. (1996):
Specimen collection, culture containers and media.
In: Mackie & McCartney, Practical Medical Microbiology,
Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A
(eds).14th edition, Churchill Livingstone P.95-112.

- Collee, J. G.; Miles, R. S. and Watt, B. (1996):
Tests for identification of bacteria.
In: Mackie & McCartney, Practical Medical Microbiology,
Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A
(eds).14th edition, Churchill Livingstone P.131-150.

- Coulster, W. A.; Chew-Graham, C. A.; Cheung, S. W. and Bruke, F.
J. (2001):
Autoclave performance and operator knowledge of autoclave
use in primary care: a survey
J. Hosp. Infect. 48(3): 180-185.

- Cruse, P. J. E. (1986):
Surgical infections: Incisional wound.
In Hospital Infection edited by Benett, J. F. and Brachman, P. S.
2nd edition, Little, Braun and Company, Boston. P.423.

- Cruse, P. J. E. and Ford, R. (1987):
The epidemiology of wound infection.
Surg. Clin. Dis., 60:72

- Davis, J. H. (1983):
Resistance to aminoglycosides: Mechanisms and frequency.
Rev. Infect. Dis. 5-261.

- Deglado, R. M.; Gomez, O. A.; Sillero, A. M.; Martinez, G.G.; Medina, M. and Liorca, J. (2001):
Efficacy of surveillance in nosocomial infection control in a surgical service.
Am. J. Infect. Control. 29(5): 289-294.

- Dellinger, P. E.; Peter, A. G.; Trisha, L.; Peter, J. K.; William, J. M.; John, E. M.; Richard, L. S. and Richard, P. W. (1994):
Quality standard for antimicrobial prophylaxis in surgical Procedures.
Clinical Infection Diseases (CID) 18: 422-427.

- Dellinger, P. E. (2001):
Preventing surgical site infection.
Infect. Control. Hosp. Epidemiol. 22: 604-606.

- Elaine, L. L.; Michele, L. P.; James, T. L. and Audrey, B. A. (1999):
CDC guideline for prevention of surgical site infection.
Hospital Infection Control Practices Advisory Commitee (HICPAC).
Infect. Control. Hosp. Epidem. 20 (4): 250-279.

- Emori, T. G. and Gaynes, R. P. (1993):
An overview of nosocomial infections including the role of the microbiology laboratory.
Clin. Microbiology. 6(4): 428-442.

- Gabriel, V. and Michael, J. (1997):
Gram negative rods.
In: Microbiology and Infectious Disease. Gabriel Virella (eds)
Third edition, William&Wilkins. P.140-159.

- Geffers, C.; Gastmeier, P.; Daschner, F. and Ruden, H. (2001):
Prevention of postoperative wound infections, Evidence-based recommendations.
Zentralbl-chir. 126 (2): 84-92.

- Gillespie, M. T.; Lyon, B. R. and Skurray, R. A. (1990):
 Typing of methicillin-resistant *Staphylococcus aureus* by
 antibiotic resistant phenotypes.
 Journal of Medical Microbiology. 31: 57-64.

- Goldman, D. A.; Weinstien, R. A.; Wenzel, R. P.; Toblan, O. C.;
 Duma, R. J. and Gaynes, R. P. (1996):
 Strategies to prevent and control the emergence and spread of
 antimicrobial-resistant microorganisms in hospitals.
 JAMA 275: 234-240.

- Govan, J. R. W. (1996):
Pseudomonas, Stenotrophomonas, Burkholderia.
 In: Mackie & McCartney, Practical Medical Microbiology,
 Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A
 (eds). 14th edition, Churchill Livingstone P.131-150.

- Greif, R.; Acsa, O.; Horn, E. P.; Kurz, A. and Sessler, D. I. (2000):
 Supplemental perioperative oxygen to reduce the incidence of
 surgical wound infection.
 N. Engl. J. Med. 342: 161-167.

- Hardy, J. (1981):
 Complications in surgery and their management 4th edition.
 Saunders Company, Philadelphia, London and Toronto: 18

- Hasper, C. M.; Mc Nicholas, T. and Gowrie-Mohan, S. (2003):
 Maintaining perioperative normothermia.
 BMJ 326: 721-722.

- Hentges, D. J. (1993):
 The anaerobic micro flora of the human body.
 Clin. Infect. Dis. 16(suppl-4): S175.

- Heritage, J. (2002):**
Tracing nosocomial infections
Infect. Control. Hosp. Epidemiol. 25: 426-439.

- Holloway, B. W. (1993):**
Genetics for all bacteria
Annu. Rev. Microbiol. 17: 215-221

- Holmes, B. and Gross, R. J. (1990):**
Coliform bacteria; various other members of the
Enterobacteriaceae.
In: Topley and Wilson's Principles of Bacteriology, Virology
and immunity, Parker, M. T. and Duren, B. I. (eds). 8th
edition. Vol 2. Edward Arnold, London, P 415-441.

- Hopkins, K. L. and Hilton, A. C. (2001):**
Restriction endonuclease analysis of RADP-PCR amplicons
derived from shiga -like toxin-producing *Escherichia coli*
O157 isolates
J. Med. Microbiol. 50(1): 90-95.

- Jensen, L. S.; Kissmeyer, N. P.; Wolf, B. and Qvist, N. (1996):**
Randomized comparison of leukocyte-depleted versus buffy-
coat poor blood transfusion and complication after colorectal
surgery.
Lancet 348: 841-845.

- Jett, B. D.; Huycke, M. M. and Gilmore, M. S. (1994):**
Virulence in *Enterococci*.
Clin. Microbiol. Rev. 7: 426.

- Johnson, J. R. (1991):**
Virulence factors of *E. coli* in urinary tract infection.
Clin. Microbiol. Rev. 4: 80.

- Jones, M. E. (1999):
Methods for typing organisms.
The European Network for Antimicrobial Resistant and
Epidemiology (ENARE) P 1-3.

- Jorgensen, L. N.; Kallehave, F.; Christensen, E.; Sana, J. E. and
Gottrup, F. (1998):
Less collagen production in smokers.
Surgery 132: 450-455.

- Kallen, J.; Driscoll, T. J.; Thornton, S.; Olson, P. E. and Wallace,
M. R. (2000):
Increase in community-acquired MRSA at Naval medical
centre.
Infect. Control. Hosp. Epidemiol. 21:223-226.

- Karim, H.; Chafik, K.; Karim, K.; Moez, H.; Makdi A. M.; Adnen, E.
H.; Morched, A. M. and Abdjellil, Z. (2000):
Risk factors for surgical wound infection in digestive surgery;
Retrospective study of 3000 surgical wounds.
Tunis. Med. 78(11): 634-640.

- Kasper, D. L. and Onderdonk, A. B. (1990):
International symposium on anaerobic bacteria and bacterial
Infection.
Rev. Infect. Dis. 16(suppl 4) S190.

- Kaul, A. F. and Jewett, J. F. (1981):
Agents and techniques for disinfection of the skin.
Surg. Gynecol. Obstet. , 152:677.

- Kenneth, T. (2002):
Pseudomonas aeruginosa.
Crit. Care. Med. 28(7):2573-2577.

- Khan, M. M. and Celik, Y. (2001):
Cost of nosocomial infection in Turkey: an estimate based on the University Hospital data.
Health. Serv. Manage. Res. 14(1):49-54.

- Killian, C. A.; Graffunder, E. M.; Vinciguersa, T. J. and Venezia, R. A. (2001):
Risk factors for surgical site infection following cesarean section.
Infect. Control. Hosp. Epidemiol. 22(10):613-617.

- Kluytmans, J. (1998):
Reduction of surgical site infections in major surgery by elimination of nasal carriage of staphylococcus aureus.
Journal of Hospital Infection suppl B: 25-29.

- Knight, R.; Charbonneau, P.; Ratzer, E.; Zeren, F.; Haun, W. and Clark, J. (2001):
Prophylactic antibiotic are not indicated in clean general surgery cases.
Am. J. Surgery 182(6):682-686.

- Kok, T. W.; Worswick, D. and Gowans, E. (1996):
Some serological techniques for microbial and viral infections.
In: Mackie & McCartney, Practical Medical Microbiology, Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A (eds). 14th edition, Churchill Livingstone P.179-204.

- Konowalchuck, J. Ispeirs, J. and Stavric, S. (1977):
Vero-response to the cytotoxin of Ecoli.
Inf. Immun. 775-779.

- Kurz, A.; Sessler, D. I. and Lenhardt, R. (1996):
Perioperative normothermia to reduce the incidence of Surgical wound infection and shorten hospitalization.
N. Eng. J. Med. 334:1209-1215.

- Kyle, J. (1934):**
Surgical handicraft.
21st edition; John Wright and Sons Ltd. Bristol pp.191.

- Latham, R.; Lancaster, A. D.; Covington, J. F.; Pirolo, J. S. and Thomas, S. (2001):**
The association of D.M. and glucose control with surgical -site infections among cardiothorathic surgery patients.
Infect Control. Hosp. Epidemiol. 22:607-612.

- Leclercq, R. and Courvalin, P. (1997):**
Resistance of *Enterococci* to glycopeptides.
Clin. Infect. Dis. 24:545.

- Leonard, R. B. (1995):**
Pulsed-field gel electrophoresis in characterizing strains of methicillin resistant *Staphylococcus aureus* from a recent hospital outbreak.
J. Clin. Microbiol. 33:2723-2726.

- Levy, D. A.; Bens, M. S.; Craun, G.F. (1998):**
Surveillance for water- borne disease outbreaks in United States,
Mor. Mortal. Wkly. Rep.CDC.11; 47(5):1-34.

- Lidwell, O. M. (1986):**
Clear air at operation and subsequent sepsis in the joints.
Clin. Orth. and related Res.211:91.

- Lipp, A. and Edwards, P. (2002):**
Disposable surgical face masks for preventing surgical wound infection in clean surgery.
Cochrane Database Syst. Rev. (1):CD002929.

- Lynch, W.; Malek, M.; Davey, P. G.; Byrne, D. J. and Napier, A. (1992):**
Costing wound infection in a Scottish hospital.
Pharmacoeconomics 2:163-170.

- Maclean, L. D.; Mulligan, W. G.; Maclean, A. P. and Duff, J. H. (1989):**
Pattern of septic shock in man: a detailed study of 56 patients.
Ann. Surg. 166:543.
- Mangram, A. J. Teresa, C. H.; and William, R. J. (1999):**
C.D.C. National Nosocomial Infections Surveillance (NNIS)
System report, data summery from January 1990-May 1999
Am. J. Infect. Control. 27:520-532.
- Marsou, R.; Idrissi, L.; Ben-Hammida, H.; Zouhdi, M.; Boudouma, M. and Goldner, M. (2001):**
Relationship of staphylococcal isolates in a Moroccan hospital
by comparing phenotypical and genotypical tests
Pathol-Biol (Paris) 49(2):109-114.
- Mashita, K.; Shinagawa, N.; Ish-Kawa, S.; Hirata, K.; Manabu, T.; Hiyama, E.; Yokoyama, T.; Ikeda, S. and Morimoto, K. (2001):**
Bacteria isolated from surgical infections and their
susceptibilities to antimicrobial agents.
Jpn. J. Antibiot. 54(10):497-530.
- Maslow, J. N.; Malligan, M. E. and Arbient, R. D. (1993):**
Molecular epidemiology for typing bacteria.
Clin. Inf. Dis. 17:153-162.
- Maslow, J. N. and Malligan, M. E. (1996):**
Epidemiologic typing systems.
Infect. Control. Hosp. Epidemiol. 17:595-604.
- Maurer, I. M. (1985):**
Hospital Hygiene (published) by Edward Arnold Ltd,
London, P.12.
-

- Mayer, L. W (1988):
Use of plasmid profiles in epidemiologic surveillance of disease outbreaks in tracing the transmission of antibiotic resistance.
Clin. Microbiol. 1:228-232.
- McCray, E.; Martone, W. J.; Wise, R. P. and Culver, D. H. (1986):
Risk factors after genitourinary reconstructive surgery.
Am. J. Epidemiol. 123:1026.
- McDougal, L. K.; Facklam, R.; Reeves, M.; Hunter, S.; Swenson, J. M.; Hill, B. and Tenover, F. C. (1992):
Analysis of multiply antimicrobial –resistant isolates of *Streptococcus pneumoniae* from the United States.
Antimicrob. Agents. Chemother. 36:2177-2184.
- Mead, P. B.; Pories, S. E. and Hall, P. (1986):
Decreasing the incidence of surgical wound infections: validation of a surveillance notification programme.
Arch. Surg. 121:458.
- Miles, R. S. and Amyes, S. G. B. (1996):
Laboratory control of antimicrobial therapy.
In: Mackie & McCartney, Practical Medical Microbiology, Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A (eds). 14th edition, Churchill Livingstone P.151-178.
- Nageeb, A. (1990):
Incidence of post-operative wound infection M. Sc. Thesis.
Bacteriology. Cairo University.
- Nicolle, L. E.; Huchcroft, S. A. and Cruse, P. J. (1992):
Risk factors for surgical wound infection among the elderly.
J. Clin. Epidemiol. 45(4):357-364.
- Nishith, B. and Kosloske, A. M. (1990):
Postoperative wound infection in pediatric surgical patients.
J. Ped. Surg. 25(1):125.

- Old, D. C. (1996):
Salmonella, In: Mackie & McCartney, Practical Medical Microbiology, Collee, J. G.; Fraser, A. G.; Marmion, B. P.; and Simmons, A. (eds). 14th edition, Churchill Livingstone P. 385-404.

- Page, C. P.; Bohnen, J. M.; Fletcher, J. R.; McManus, A. T.; Solomkin, J. S. and Hman, D. H. (1993):
Antimicrobial prophylaxis for surgical wounds: guidelines for clinical care. Arch. Surg. 128:79-88.

- Patacchiola, F.; Di-Paolantini, L.; Palermo, P.; Di-Stefano, L.; Mascaretti, G. and Moscarini, M. (2000):
Antibiotic prophylaxis of infective complications after cesarean section.
Minerva. Ginecol. 52(10):385-389.

- Platt, R.; Zucker, J. R. and Zalernik, D. F. (1992):
Prophylaxis against wound infection following herniorrhaphy or breast surgery. J. Infect. Dis. 166:556-560.

- Platt, C. and Hall, J. C. (2001):
The prevention of wound infection in patients undergoing colorectal surgery. J. Hosp. Infect. 49(4):233-238.

- Polk, H. C. and Malangoni, M. A. (1988):
Chemoprophylaxis of wound infection in: Howard, R. J. & Simmons, R. L. (eds). Surgical infectious disease, 2nd edition. Norwalk, C. T., Appleton & Lang P.351-361.

- Pollack, M. (2000):
Pseudomonas aeruginosa, in: Mandell, G. L.; Benett, J. E. and Dolin, R. Principles and practice of infectious disease. (eds). 5th edition. New York, NY: Churchill Livingstone. P.2310- 2327.

- Prevost, G.; Couppie, P. and Prevost, P. (1995):
Epidemiological data on *Staphylococcus aureus* strains
producing synergohymenotropic toxins.
J. Med. Microbiol. 237-242.

- Rains, A. J. H. and Richie, H. D. (1981):
Baily & Love's, Short practice of surgery, 18th edition. Lewis, H.
K. & Company Ltd. London.

- Reilly, J.; Twaddle, S.; McIntosh, J. and Kean, L. (2001):
An economic analysis of surgical wound infection.
J. Hosp. Infect. 49(4):245-249.

- Rios, A.; Rodriguez, J. M.; Munitiz, V.; Alcaraz, P.; Perez, F. and
Parrila, P. (2001):
Antibiotic prophylaxis in incisional hernia repair using prosthesis.
Hernia 5(3):148-152.

- Roherborn, A.; Thomas, A.; Potreck, O.; Ebener, C.; Ohmann, C.;
Goretzki, P. E. and Roher, H. D. (2001):
The microbiology of post operative peritonitis.
Clin. Infect. Dis. 33(9):1513-1519.

- Ross, P. W. (1996):
Streptococcus & Enterococcus. In: Mackie & McCartney, Practical
Medical Microbiology, Collee, J. G.; Fraser, A. G.; Marmion, B. P.;
and Simmons, A. (eds). 14th edition, Churchill Livingstone. P.263-274.

- Roth, A.; Fry, D. E. and Polk, H. C. (1986):
Infectious morbidity in extremity fractures.
J. Tr. 26:757.

- Rotstein, O. D. (1993):
Interaction between leukocyte and anaerobic bacteria in
polymicrobial surgical infections.
Clin. Infect. Dis. 16(suppl 4): S 190.

- Russel, G. V.; King, C.; May, C. G. and Pearsall, A. W. (2001):
Once daily high- dose gentamycin to prevent infection in
open fractures of the tibial shaft: a preliminary investigation.
South. Med. J. 94(12):1185-1191.

- Sartor, C.; Edward, J. R.; Gaynes, R. P. and Culver, D. (1995):
Evaluation of hospital participation in the National Nosocomial
Infectious Surveillance System.
Am. J. Infect. Control. 23:364-368.

- Schekler, W. E.; Brim hall, D.; Buck, A. S.; Farr, B. M.; Freidman, C.
and Garibald, R.A.(1998):
Requirements for infrastructure and essential activities of
infection control and epidemiology in hospitals: a consensus
panel report.
Infect. Control. Hosp. Epidem. 19:114-124.

- Scheel, O. and Stormark, M. (1999):
National prevalence survey on hospital infections in Norway.
J. Hosp. Infect. 41:331-335.

- Scott, J. D.; Forrest, A.; Feuerstein, S.; Fitzpatrick, P. and Schentag,
J. J. (2001):
Factors associated with postoperative infection.
Infect. Control. Hosp. Epidemiol. 22(6):347-351.

- Senior, B. W. (1990):
The Proteace: Proteus, Morganella and Providencia.
In: Parker, M. T. and Dureden, B. I. (eds). Topley and Wilson's
Principles of Bacteriology, Virology and Immunity, 8th edition
Vol 2. Edward Arnold, London. P.443-454.

- Schwarz, M.; Isenmann, R.; Thomsen, J.; Gaus, W. and Beger, H. G.
(2001):
Efficacy of oral ofloxacin for single dose perioperative
prophylaxis in general surgery: a controlled randomized clinical
study. Longenbecks.
Arch. Surg. 386(6):397-401.

- Simchen, E. (1984):**
 Determinations of wound infection after colon surgery.
 Ann. Surg., 199:260.

- Smith, H. R. and Scotland, S. M. (1993):**
 Isolation and identification methods for *Escherichia coli* O157 and
 other verocytotoxin producing strains.
 J. Clin. Path. 46:10-17.

- Tenover, F. C.; Arbeit, R.; Archer, G.; Biddle, J.; Bayne, S.;
 Georing, R.; Hancock, G. and Herbert, G. A. (1994):**
 Comparison of traditional and molecular methods of typing
 isolates of *Staphylococcus aureus*.
 J. Clin. Microbiol. 32:407-415.

- Tobin, G. R. (1984):**
 Closure of contaminated wounds, biologic and technical
 consideration.
 Surg. Clin. North Am. 61:639.

- Triulzi, D. J.; Vanek, K.; Ryan, D. H. and Blumberg, N. (1992):**
 A clinical and immunologic study of blood transfusion
 and postoperative bacterial infection in spinal surgery.
 Transfusion. 32:517-524.

- Troillet, N.; Petingat, C.; Matter, M.; Eisenring, M. C.; Mosimann, F.
 and Francioli, P. (2001):**
 Surgical site infection surveillance: an effective preventive
 measure.

- Truler, E.; Truler, A.; Ladurner, T.; Hirner, A. and Engels, H.
 (2000):**
 Surgical gown and patient drapes in routine practice: discrepancy
 between requirements and reality.
 Chirurg. 71(11):1409-1415.

- Twum, D. K.; Grant, C.; Al-Suleiman, S. A. (1992):
Microbiology of post-operative wound infection: a prospective study of 1770 wounds.
J. Hosp. Infect. 21(1):29-37.

- Vaudaux, P. E.; Lew, D. P. and Waldvogel, F. A. (1994):
Host factors predisposing to and influencing therapy of foreign body infection. In: Bisno, A. L. and Waldvogel, F. A. (eds):
Infections associated with indwelling medical devices, 2nd edition. American Society for Microbiology, Washinton, D.C. P.1.

- Vignali, A.; Braga, M.; Gianotti, L.; Radelli, G.; Orest, G.; Russo, A. and Dicalo, V. (1996)
A single unit of allogenic blood increases postoperative infections. Vox Sang. 71:170-175.

- Walter, J. B. and Israel, M. S. (1984):
Hospital infection, In: General pathology, 5th edition, Churchill Livingstone. P. 103.

- Walter, J. B. and Israel, M. S. (1984):
Hospital infection, In: General pathology, 5th edition, Churchill Livingstone. P. 233.

- Wong, J. T.; Chang, S. C.; Ko, W. J.; Chang, Y. Y.; Chen, M. L.; Pan, H. J. and Luh, K. T. (2001):
A hospital acquired outbreak of methicillin resistant *Staphylococcus aureus* infection initiated by a surgeon carrier.
J. Hosp. Infect. 47(2):104-109.

- Wood, C. A. (1992):
Significant infection caused by *Staphylococci*.
J. Clin. Microbiol. 30:2216-2217.

-Wurtz, R.; Wittrack, B.; Lavin, M. A. and Zawaki, A. (2001):
New surgeons have higher surgical site infection rates.
Infect. Control. Hosp. Epidemiol. 22(6):375-377.

-Yong, K. S.; Karcem, B. A.; Ruslan, G. N. and Harwant, S. (2001):
Risk factors for infection in total hip replacement surgery at
Kuala Lumpur hospital.
Med. J. Malaysia. 56 Suppl C: 57-60.

-Zoutman, D.; Mc Donald, S. and Vethanyagan, D. (1998):
Total and attributable costs of surgical wound infections at a
Canadian tertiary -care- center.
Infect. Control. Hosp. Epidemiol. 19:254-259.